



biblio.ugent.be

The UGent Institutional Repository is the electronic archiving and dissemination platform for all UGent research publications. Ghent University has implemented a mandate stipulating that all academic publications of UGent researchers should be deposited and archived in this repository. Except for items where current copyright restrictions apply, these papers are available in Open Access.

This item is the archived peer-reviewed author-version of:

Title: On a class of translation-invariant spaces of quasianalytic ultradistributions

Authors: Pavel Dimovski, Bojan Prangoski, and Jasson Vindas

In: Novi Sad Journal of Mathematics, Vol. 45 (1), 143-175, 2015.

Optional: http://www.dmi.uns.ac.rs/nsjom/Papers/45_1/NSJOM_45_1_143_175.pdf

To refer to or to cite this work, please use the citation to the published version:

P. Dimovski, B. Prangoski, and J. Vindas (2015). On a class of translation-invariant spaces of quasianalytic ultradistributions. *Novi Sad J. Math.* 45(1), 143-175.

ON A CLASS OF TRANSLATION-INVARIANT SPACES OF QUASIANALYTIC ULTRADISTRIBUTIONS

PAVEL DIMOVSKI, BOJAN PRANGOSKI, AND JASSON VINDAS

Dedicated to Professor B. Stanković on the occasion of his 90th birthday and to Professor J. Vickers on the occasion of his 60th birthday

ABSTRACT. A class of translation-invariant Banach spaces of quasianalytic ultradistributions is introduced and studied. They are Banach modules over a Beurling algebra. Based on this class of Banach spaces, we define corresponding test function spaces \mathcal{D}'_E and their strong duals \mathcal{D}'_{E^*} of quasianalytic type, and study convolution and multiplicative products on \mathcal{D}'_{E^*} . These new spaces generalize previous works about translation-invariant spaces of tempered (non-quasianalytic ultra-) distributions; in particular, our new considerations apply to the settings of Fourier hyperfunctions and ultrahyperfunctions. New weighted $\mathcal{D}'_{L^p_\eta}$ spaces of quasianalytic ultradistributions are analyzed.

1. INTRODUCTION

Recently, the authors and Pilipović have constructed and studied new classes of distribution and non-quasianalytic ultradistribution spaces in connection with translation-invariant Banach spaces [2, 4]. Those spaces generalize the concrete instances of weighted \mathcal{D}'_{L^p} and \mathcal{D}'_{L^p} spaces [1, 14] and have shown usefulness in the study of boundary values of holomorphic functions [3] and the convolution of generalized functions [4].

The aim of this article is to extend the theory of ultradistribution spaces associated to translation-invariant Banach spaces by considering mixed quasianalytic cases. We have been able here to transfer all results from [4] to this new setting with the aid of various new important results for quasianalytic ultradistribution spaces of type $\mathcal{S}'_{\dagger}(\mathbb{R}^d)$ (see Subsection 1.1 for the notation) from [10] concerning the construction of parametrices and the structure of these spaces. Such technical results will be stated in Section 2 without proofs, as details will be treated in [10]. Although our results in the present paper are

2010 *Mathematics Subject Classification*. Primary 46F05. Secondary 46H25; 46F10; 46F15; 46E10.

Key words and phrases. Quasianalytic ultradistributions; Convolution of ultradistributions; Translation-invariant Banach space of ultradistributions; Tempered ultradistributions; Beurling algebra; Hyperfunctions.

J. Vindas gratefully acknowledges support by Ghent University, through the BOF-grant 01N01014.

analogous to those from [4], new arguments and ideas have had to be developed here in order to deal with the quasianalytic case and achieve their proofs.

In Section 3 we study the class of translation-invariant Banach spaces of ultradistributions of class $*-\dagger$. These are translation-invariant Banach spaces satisfying $\mathcal{S}_\dagger^*(\mathbb{R}^d) \hookrightarrow E \hookrightarrow \mathcal{S}_\dagger'^*(\mathbb{R}^d)$ and having ultrapolynomially bounded weight function of class \dagger . Here $*$ and \dagger stand for the Beurling and Roumieu cases of sequences M_p and A_p , respectively. We would like emphasize that our considerations apply to hyperfunctions and ultra-hyperfunctions, which correspond to the symmetric choices $M_p = A_p = p!$; but more generally, our weight sequence M_p , measuring the ultradifferentiability, is allowed to satisfy the mild condition $p!^\lambda \subset M_p$ with $\lambda > 0$. The growth assumption on A_p is just $p! \subset A_p$, which also allows to deal with Banach spaces whose translation groups may have exponential growth.

Section 4 contains our main results. In analogy to [4], we introduce the test function spaces $\mathcal{D}_E^{(M_p)}$, $\mathcal{D}_E^{\{M_p\}}$, and $\tilde{\mathcal{D}}_E^{\{M_p\}}$. We prove that the following continuous and dense embeddings hold $\mathcal{S}_\dagger^*(\mathbb{R}^d) \hookrightarrow \mathcal{D}_E^* \hookrightarrow E \hookrightarrow \mathcal{S}_\dagger'^*(\mathbb{R}^d)$ and that \mathcal{D}_E^* are topological modules over the Beurling algebra L_ω^1 , where ω is the weight function of the translation group of E . We also prove the dense embedding $\mathcal{D}_E^* \hookrightarrow \mathcal{O}_{\dagger,C}^*(\mathbb{R}^d)$, where the spaces $\mathcal{O}_{\dagger,C}^*(\mathbb{R}^d)$ are defined in a similar way as in [4]. The space $\mathcal{D}_{E'_*}^*$ is defined as the strong dual of \mathcal{D}_E and various structural and topological properties of $\mathcal{D}_{E'_*}^*$ are obtained via the parametrix method (Lemma 2.2). We also prove that $\mathcal{D}_E^{\{M_p\}} = \tilde{\mathcal{D}}_E^{\{M_p\}}$, topologically.

As an application of our theory, we extend the theory of $\mathcal{D}_{L_\eta^p}^*$, \mathcal{B}_η^* , and $\dot{\mathcal{B}}_\eta^*$ spaces not only by considering quasianalytic cases of $*$ but also by allowing ultrapolynomially bounded weights η which may growth exponentially. We establish relations among them and make a detailed investigation of their topological properties. We would like to point out that applications of such results to the study of the general convolvability in the setting of quasianalytic ultradistributions will appear elsewhere [10]. We conclude this section with some results about convolution and multiplicative products on $\mathcal{D}_{E'_*}^*$.

1.1. Notation. Let $(M_p)_{p \in \mathbb{N}}$ and $(A_p)_{p \in \mathbb{N}}$ be two sequences of positive numbers such that $M_0 = M_1 = A_0 = A_1 = 1$. Throughout the article, we impose the following assumptions over these weight sequences. The sequence M_p satisfies the ensuing three conditions:

$$(M.1) \quad M_p^2 \leq M_{p-1}M_{p+1}, \quad p \in \mathbb{Z}_+;$$

$$(M.2) \quad M_p \leq c_0 H^p \min_{0 \leq q \leq p} \{M_{p-q}M_q\}, \quad p, q \in \mathbb{N}, \text{ for some } c_0, H \geq 1;$$

(M.5) there exists $s > 0$ such that M_p^s is strongly non-quasianalytic, i.e., there exists $c_0 \geq 1$ such that

$$\sum_{q=p+1}^{\infty} \frac{M_{q-1}^s}{M_q^s} \leq c_0 p \frac{M_p^s}{M_{p+1}^s}, \quad \forall p \in \mathbb{Z}_+.$$

It is clear that if M_p^s is strongly non-quasianalytic than for any $s' > s$, $M_p^{s'}$ is also strongly non-quasianalytic. One easily verifies that when M_p satisfies (M.5) there exists $\kappa > 0$ such that $p!^\kappa \subset M_p$, i.e. there exist $c_0, L_0 > 0$ such that $p!^\kappa \leq c_0 L_0^p M_p$, $p \in \mathbb{N}$ (cf. [7, Lemma 4.1]). Following Komatsu [7], for $p \in \mathbb{Z}_+$, we denote $m_p = M_p/M_{p-1}$ and for $\rho \geq 0$ let $m(\rho)$ be the number of $m_p \leq \rho$. As a consequence of [7, Proposition 4.4], by a change of variables, one verifies that M_p satisfies (M.5) if and only if

$$\int_\rho^\infty \frac{m(\lambda)}{\lambda^{s+1}} d\lambda \leq c \frac{m(\rho)}{\rho^s}, \quad \forall \rho \geq m_1.$$

A sufficient condition for M_p to satisfies (M.5) is if the sequence m_p/p^λ , $p \in \mathbb{Z}_+$, to be monotonically increasing for some $\lambda > 0$.

We assume that A_p satisfies (M.1) and (M.2). Of course, without losing generality, we can assume that the constants c_0 and H from the condition (M.2) are the same for M_p and A_p . Moreover, we also assume that A_p satisfies the following additional hypothesis:

(M.6) $p! \subset A_p$; i.e., there exist $c_0, L_0 > 0$ such that $p! \leq c_0 L_0^p A_p$, $p \in \mathbb{N}$.

Of course, the constants c_0 and L_0 in (M.6) can be chosen such that $c_0, L_0 \geq 1$. Although it is not part of our assumptions, we will be primarily interested in the quasianalytic case, i.e., $\sum_{p=1}^\infty \frac{M_{p-1}}{M_p} = \infty$.

We denote by $M(\cdot)$ and $A(\cdot)$ the associated functions of M_p and A_p , that is, $M(\rho) := \sup_{p \in \mathbb{N}} \ln_+ \frac{\rho^p}{M_p}$ and $A(\rho) := \sup_{p \in \mathbb{N}} \ln_+ \frac{\rho^p}{A_p}$ for $\rho > 0$, respectively. They are non-negative continuous increasing functions (cf. [7]). We denote by \mathfrak{R} the set of all positive monotonically increasing sequences which tend to infinity. For $(l_p) \in \mathfrak{R}$, denote by N_{l_p} and $B_{l_p}(\cdot)$ the associated functions of the sequences $M_p \prod_{j=1}^p l_j$ and $A_p \prod_{j=1}^p l_j$, respectively.

For $h > 0$ we denote by $\mathcal{S}_{A_p, h}^{M_p, h}$ the Banach spaces (in short (B)-space from now on) of all $\varphi \in C^\infty(\mathbb{R}^d)$ for which the norm

$$\sigma_h(\varphi) = \sup_\alpha \frac{h^{|\alpha|} \|e^{A(h|\cdot|)} D^\alpha \varphi\|_{L^\infty(\mathbb{R}^d)}}{M_\alpha}$$

is finite. One easily verifies that for $h_1 < h_2$ the canonical inclusion $\mathcal{S}_{A_p, h_2}^{M_p, h_2} \rightarrow \mathcal{S}_{A_p, h_1}^{M_p, h_1}$ is compact. As l.c.s., we define $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d) = \lim_{h \rightarrow \infty} \mathcal{S}_{A_p, h}^{M_p, h}$ and $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d) = \varprojlim_{h \rightarrow 0} \mathcal{S}_{A_p, h}^{M_p, h}$. Since for $h_1 < h_2$ the inclusion $\mathcal{S}_{A_p, h_2}^{M_p, h_2} \rightarrow \mathcal{S}_{A_p, h_1}^{M_p, h_1}$ is compact, $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d)$ is an (FS)-space and $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ is a (DFS)-space. In particular they are both Montel spaces.

For each $(r_p) \in \mathfrak{R}$, by $\mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)}$ we denote the space of all $\varphi \in C^\infty(\mathbb{R}^d)$ such that

$$\sigma_{(r_p)}(\varphi) = \sup_{\alpha} \frac{\|e^{B_{r_p}(\cdot)} D^\alpha \varphi\|_{L^\infty(\mathbb{R}^d)}}{M_\alpha \prod_{j=1}^{|\alpha|} r_j} < \infty.$$

Provided with the norm $\sigma_{(r_p)}$, the space $\mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)}$ becomes a (B) -space. Similarly as in [1, 9], one can prove that $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ is topologically isomorphic to $\varprojlim_{(r_p) \in \mathfrak{R}} \mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)}$.

In the future we shall employ $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ as a common notation for $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d)$ (Beurling case) and $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ (Roumieu case). It is clear that for each $h > 0$ and $(r_p) \in \mathfrak{R}$, the spaces $\mathcal{S}_{A_p, h}^{M_p, h}$ and $\mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)}$ are continuously injected into $\mathcal{S}(\mathbb{R}^d)$ (the Schwartz space).

We will often make use of the following technical result from [11].

Lemma 1.1 ([11]). *Let $(k_p) \in \mathfrak{R}$. There exists $(k'_p) \in \mathfrak{R}$ such that $k'_p \leq k_p$ and $\prod_{j=1}^{p+q} k'_j \leq 2^{p+q} \prod_{j=1}^p k'_j \cdot \prod_{j=1}^q k'_j$, for all $p, q \in \mathbb{Z}_+$.*

We adopt the following notations. The symbol “ \hookrightarrow ” stands for a continuous and dense inclusion between topological vector spaces. For $h \in \mathbb{R}^d$ and $f \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$ we denote as $T_h f$ translation by h , i.e., $T_h f = f(\cdot + h)$. We write $\langle x \rangle = (1 + |x|^2)^{1/2}$, $x \in \mathbb{R}^d$.

2. SOME IMPORTANT AUXILIARY RESULTS ON THE SPACE $\mathcal{S}_\dagger^*(\mathbb{R}^d)$

We collect in this section some important results on the nuclearity of $\mathcal{S}_\dagger^*(\mathbb{R}^d)$, the existence of parametrices as well as a characterisation of bounded sets in $\mathcal{S}_\dagger^*(\mathbb{R}^d)$. These are essential tools in the rest of the article. We refer to [10] for the proofs. Unless explicitly stated, we deal with the Beurling and Roumieu cases simultaneously. We follow the ensuing convention. We shall first state assertions for the $(M_p) - (A_p)$ case followed in parenthesis by the corresponding statements for the $\{M_p\} - \{A_p\}$ case.

Proposition 2.1. *The space $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is nuclear.*

Proposition 2.2. *For every $t > 0$ there exist $G \in \mathcal{S}_{A_p, t}^{M_p, t}$ and an ultradifferential operator $P(D)$ of class (M_p) (for every $(t_p) \in \mathfrak{R}$ there exist $G \in \mathcal{S}_{A_p, (t_p)}^{M_p, (t_p)}$ and an ultradifferential operator $P(D)$ of class $\{M_p\}$) such that $P(D)G = \delta$.*

Lemma 2.3. *Let $r > 0$ ($(r_p) \in \mathfrak{R}$).*

- i) For each $\chi, \varphi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$ and $\psi \in \mathcal{S}_{A_p, r}^{M_p, r}$ ($\psi \in \mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)}$), one has $\chi * (\varphi\psi) \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$.*

ii) Let $\varphi, \chi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$ with $\varphi(0) = 1$ and $\int_{\mathbb{R}^d} \chi(x) dx = 1$. For each $n \in \mathbb{Z}_+$ define $\chi_n(x) = n^d \chi(nx)$ and $\varphi_n(x) = \varphi(x/n)$. Then there exists $k \geq 2r$ ($(k_p) \in \mathfrak{R}$ with $(k_p) \leq (r_p/2)$) such that the operators $\tilde{Q}_n : \psi \mapsto \chi_n * (\varphi_n \psi)$, are continuous as mappings from $\mathcal{S}_{A_p, k}^{M_p, k}$ into $\mathcal{S}_{A_p, r}^{M_p, r}$ (from $\mathcal{S}_{A_p, (k_p)}^{M_p, (k_p)}$ into $\mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)}$), for all $n \in \mathbb{Z}_+$. Moreover $\tilde{Q}_n \rightarrow \text{Id}$ in $\mathcal{L}_b(\mathcal{S}_{A_p, k}^{M_p, k}, \mathcal{S}_{A_p, r}^{M_p, r})$ ($\mathcal{L}_b(\mathcal{S}_{A_p, (k_p)}^{M_p, (k_p)}, \mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)})$).

In the next proposition, given $t > 0$ ($(t_p) \in \mathfrak{R}$), we denote as $\overline{\mathcal{S}}_{A_p, t}^{M_p, t}$ (as $\overline{\mathcal{S}}_{A_p, (t_p)}^{M_p, (t_p)}$) the closure of $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d)$ in $\mathcal{S}_{A_p, t}^{M_p, t}$ (the closure of $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ in $\mathcal{S}_{A_p, (t_p)}^{M_p, (t_p)}$).

Proposition 2.4. Let B be a bounded subset of $\mathcal{S}_\dagger^*(\mathbb{R}^d)$. There exists $k > 0$ ($(k_p) \in \mathfrak{R}$) such that each $f \in B$ can be extended to a continuous functional \tilde{f} on $\overline{\mathcal{S}}_{A_p, k}^{M_p, k}$ (on $\overline{\mathcal{S}}_{A_p, (k_p)}^{M_p, (k_p)}$). Moreover, there exists $l \geq k$ ($(l_p) \in \mathfrak{R}$ with $(l_p) \leq (k_p)$) such that $\mathcal{S}_{A_p, l}^{M_p, l} \subseteq \overline{\mathcal{S}}_{A_p, k}^{M_p, k}$ ($\mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)} \subseteq \overline{\mathcal{S}}_{A_p, (k_p)}^{M_p, (k_p)}$) and $*$: $\mathcal{S}_{A_p, l}^{M_p, l} \times \mathcal{S}_{A_p, l}^{M_p, l} \rightarrow \overline{\mathcal{S}}_{A_p, k}^{M_p, k}$ ($*$: $\mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)} \times \mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)} \rightarrow \overline{\mathcal{S}}_{A_p, (k_p)}^{M_p, (k_p)}$) is a continuous bilinear mapping. Furthermore, there exist an ultradifferential operator $P(D)$ of class $*$ and $u \in \overline{\mathcal{S}}_{A_p, l}^{M_p, l}$ ($u \in \overline{\mathcal{S}}_{A_p, (l_p)}^{M_p, (l_p)}$) such that $P(D)u = \delta$ and $f = (P(D)u) * f = P(D)(u * \tilde{f})$ for each $f \in B$, where $u * \tilde{f}$ is the image of \tilde{f} under the transpose of the continuous mapping $\varphi \mapsto \check{u} * \varphi$, $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d) \rightarrow \overline{\mathcal{S}}_{A_p, k}^{M_p, k}$ ($\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d) \rightarrow \overline{\mathcal{S}}_{A_p, (k_p)}^{M_p, (k_p)}$). For $f \in B$, $u * \tilde{f} \in L_{e^{A(|\cdot|)}}^\infty \cap C(\mathbb{R}^d)$ ($u * \tilde{f} \in L_{e^{B_{l_p}(|\cdot|)}}^\infty \cap C(\mathbb{R}^d)$) and in fact $u * \tilde{f}(x) = \langle \tilde{f}, u(x - \cdot) \rangle$. The set $\{u * \tilde{f} | f \in B\}$ is bounded in $L_{e^{A(|\cdot|)}}^\infty$ (in $L_{e^{B_{l_p}(|\cdot|)}}^\infty$).

Lemma 2.5. Let $B \subseteq \mathcal{S}_\dagger^*(\mathbb{R}^d)$. The following statements are equivalent:

- i) B is bounded in $\mathcal{S}_\dagger^*(\mathbb{R}^d)$;
- ii) for each $\varphi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$, $\{f * \varphi | f \in B\}$ is bounded in $\mathcal{S}_\dagger^*(\mathbb{R}^d)$;
- iii) for each $\varphi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$ there exist $t, C > 0$ (there exist $(t_p) \in \mathfrak{R}$ and $C > 0$) such that $|(f * \varphi)(x)| \leq Ce^{A(t|x|)}$ ($|(f * \varphi)(x)| \leq Ce^{B_{t_p}(|x|)}$) for all $x \in \mathbb{R}^d$, $f \in B$;
- iv) there exist $C, t > 0$ (there exist $(t_p) \in \mathfrak{R}$ and $C > 0$) such that

$$|(f * \varphi)(x)| \leq Ce^{A(t|x|)} \sigma_t(\varphi) \quad (\text{resp. } |f * \varphi(x)| \leq Ce^{B_{t_p}(|x|)} \sigma_{(t_p)}(\varphi))$$

for all $\varphi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$, $x \in \mathbb{R}^d$, $f \in B$.

Lemma 2.6. Let $f \in \mathcal{S}'_{(p!)}(M_p)(\mathbb{R}^d)$ ($f \in \mathcal{S}'_{\{p!\}}^{\{M_p\}}(\mathbb{R}^d)$). Then $f \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$ if and only if there exists $t > 0$ (there exists $(t_p) \in \mathfrak{R}$) such that for every $\varphi \in \mathcal{S}_{(p!)}^{(M_p)}(\mathbb{R}^d)$ (for every $\varphi \in \mathcal{S}_{\{p!\}}^{\{M_p\}}(\mathbb{R}^d)$)

$$\sup_{x \in \mathbb{R}^d} e^{-A(t|x|)} |(f * \varphi)(x)| < \infty \quad \left(\sup_{x \in \mathbb{R}^d} e^{-B_{t_p}(|x|)} |(f * \varphi)(x)| < \infty \right).$$

3. TRANSLATION-INVARIANT BANACH SPACES OF QUASIANALYTIC ULTRADISTRIBUTIONS

We extend here the theory of translation-invariant Banach spaces of ultradistributions to the quasianalytic case. We closely follow the approach from [2, 4], where the distribution and non-quasianalytic ultradistribution cases were treated. We mention that some of the arguments below are similar to those from [4], but for the reader's convenience we include all details about the adaptations in the corresponding proofs.

Let E be a (B) -space. We call E a *translation-invariant (B) -space of ultradistributions of class $* - \dagger$* if it satisfies the following three axioms:

- (I) $\mathcal{S}'_{\dagger}(\mathbb{R}^d) \hookrightarrow E \hookrightarrow \mathcal{S}'_{\dagger}(\mathbb{R}^d)$.
- (II) $T_h(E) \subseteq E$ for each $h \in \mathbb{R}^d$.
- (III) There exist $\tau, C > 0$ (for every $\tau > 0$ there exists $C > 0$), such that

$$\|T_h g\|_E \leq C \|g\|_E e^{A(\tau|h|)}, \quad \forall h \in \mathbb{R}^d, \quad \forall g \in E.$$

Notice that the condition (III) implicitly makes use of the continuity of T_h . The next lemma shows that such a continuity is always ensured by the conditions (I) and (II).

Lemma 3.1. *Let E be a (B) -space satisfying (I) and (II). The translation operators $T_h : E \rightarrow E$ are bounded for all $h \in \mathbb{R}^d$.*

Proof. Observe that T_h is continuous as a mapping from E to $\mathcal{S}'_{\dagger}(\mathbb{R}^d)$ since it can be decomposed as $E \xrightarrow{\text{Id}} \mathcal{S}'_{\dagger}(\mathbb{R}^d) \xrightarrow{T_h} \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ and $T_h : \mathcal{S}'_{\dagger}(\mathbb{R}^d) \rightarrow \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ is continuous. Thus the graph of T_h is closed in $E \times \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ and since its image is in E its graph is also closed in $E \times E$ ($E \times E$ is continuously injected into $E \times \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ via the mapping $\text{Id} \times \text{Id}$). As E is a (B) -space, the closed graph theorem implies that T_h is continuous. \square

Lemma 3.2. *Let E be a translation-invariant (B) -space of ultradistributions of class $* - \dagger$. For every $g \in E$, $\lim_{h \rightarrow 0} \|T_h g - g\|_E = 0$. In particular for each $g \in E$ the mapping $h \mapsto T_h g, \mathbb{R}^d \rightarrow E$, is continuous at 0 (hence everywhere continuous).*

Proof. The proof is straightforward and we omit it. \square

Summarizing, Lemma 3.1 and Lemma 3.2 prove that a translation-invariant (B) -space of ultradistributions E of class $* - \dagger$ satisfies the following stronger condition than (II):

- ($\widetilde{\text{II}}$) for each $h > 0$, $T_h : E \rightarrow E$ is continuous and for each $g \in E$ the mapping $h \mapsto T_h g, \mathbb{R}^d \rightarrow E$, is continuous.

Clearly $T_0 = \text{Id}_E$, $T_{h_1+h_2} = T_{h_1} \circ T_{h_2} = T_{h_2} \circ T_{h_1}$. Next, we define *the weight function $\omega(h)$ of E* as

$$(3.1) \quad \omega(h) = \|T_{-h}\|_{\mathcal{L}(E)}.$$

Obviously the weight function is positive and $\omega(0) = 1$. Furthermore, since $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is separable (it is an (FS) -space or a (DFS) -space, respectively), so is E . Thus $\omega(h) = \|T_{-h}\|_{\mathcal{L}(E)}$ is the supremum of $\|T_{-h}g\|_E$ where g belongs to a countable dense subset of the closed unit ball of E . Since $h \mapsto \|T_{-h}g\|_E$ is continuous, ω is measurable. Clearly, the logarithm of ω is subadditive and there exist $C, \tau > 0$ (for every $\tau > 0$ there exists $C > 0$) such that $\omega(h) \leq Ce^{A(\tau|h|)}$.

Remark 3.3. In the Beurling case when $A_p = p!$, the assumption (III) is superfluous. In fact, assuming only (I) and (II), Lemma 3.1 implies that for each $h \in \mathbb{R}^d$, $T_h : E \rightarrow E$ is continuous. Additionally, one easily verifies that for each fixed $\varphi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$, one has $T_h\varphi \rightarrow \varphi$ as $h \rightarrow 0$ in $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ and consequently in E . Hence, employing the same reasoning as above, we obtain that ω is a measurable positive function with subadditive logarithm. Therefore, there exist $C, h > 0$ such that $\omega(h) \leq Ce^{k|h|}$, $\forall h \in \mathbb{R}^d$ (cf. [5, Sect 7.4]), which is in fact condition (III) in this case.

We will also give an alternative version of (III) in the Roumieu case which sometimes is easier to work with than (III). For this purpose we need the following technical result from [11].

Lemma 3.4 ([11]). *Let $g : [0, \infty) \rightarrow [0, \infty)$ be an increasing function that satisfies the following estimate:*

For every $L > 0$ there exists $C > 0$ such that $g(\rho) \leq A(L\rho) + \ln C$. Then there exists a subordinate function $\epsilon(\rho)$ such that $g(\rho) \leq A(\epsilon(\rho)) + \ln C'$, for some constant $C' > 1$.

See [7] for the definition of subordinate function.

Lemma 3.5. *In the Roumieu case condition (III) is equivalent to the following one:*

(III) *there exist $(l_p) \in \mathfrak{R}$ and $C > 0$ such that $\|T_h g\|_E \leq C \|g\|_E e^{B_{l_p}(|h|)}$, for all $g \in E$, $h \in \mathbb{R}^d$.*

Proof. The proof is analogous to that of (c) \Leftrightarrow (\tilde{c}) in [4, Theorem 4.2]. \square

The next theorem gives a weak criterion to conclude that a (B) -space E is a translation-invariant space of ultradistributions of class $* - \dagger$.

Theorem 3.6. *Let E be a (B) -space satisfying:*

- (I)' $\mathcal{S}_{(p)}^{(M_p)}(\mathbb{R}^d) \hookrightarrow E \hookrightarrow \mathcal{S}'_{(p!)}^{(M_p)}(\mathbb{R}^d)$ ($\mathcal{S}'_{\{p!\}}^{\{M_p\}}(\mathbb{R}^d) \hookrightarrow E \hookrightarrow \mathcal{S}'_{\{p!\}}^{\{M_p\}}(\mathbb{R}^d)$);
- (II) $T_h(E) \subseteq E$, for all $h \in \mathbb{R}^n$;
- (III)' *for any $g \in E$ there exist $C = C_g > 0$ and $\tau = \tau_g > 0$ (for every $\tau > 0$ there exists $C = C_{g,\tau} > 0$) such that $\|T_h g\|_E \leq Ce^{A(\tau|h|)}$, $\forall h \in \mathbb{R}^d$.*

Then E is a translation-invariant (B) -space of ultradistributions of class $ - \dagger$.*

Proof. Employing the same technique as in the proof of Lemma 3.1, one easily verifies that conditions (I)' and (II) imply the continuity of $T_h : E \rightarrow E$. The proof of (III) can be obtain by adapting the proof of (c) in [4, Theorem 4.2].

We now address (I). To prove $\mathcal{S}_\dagger^*(\mathbb{R}^d) \hookrightarrow E$, by (I)', it is enough to prove that $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is continuously injected into E . Pick $\psi_1 \in \mathcal{D}(\mathbb{R}^d)$ such that

$$\sum_{m \in \mathbb{Z}^d} \psi_1(x - m) = 1, \quad \forall x \in \mathbb{R}^d, \quad \text{supp } \psi_1 \in [-1, 1]^d$$

and ψ_1 is non-negative and even. Next, pick $\psi_2 \in \mathcal{S}_{(p!)}^{(M_p)}(\mathbb{R}^d)$ ($\psi_2 \in \mathcal{S}_{\{p!\}}^{\{M_p\}}(\mathbb{R}^d)$), such that $\int_{\mathbb{R}^d} \psi_2(x) dx = 1$ and ψ_2 is even. Set $\psi = \psi_1 * \psi_2$. One readily verifies that $\sum_{m \in \mathbb{Z}^d} \psi(x - m) = 1$ for all $x \in \mathbb{R}^d$ and $\psi \in \mathcal{S}_{(p!)}^{(M_p)}(\mathbb{R}^d)$ in the Beurling case and $\psi \in \mathcal{S}_{\{p!\}}^{\{M_p\}}(\mathbb{R}^d)$ in the Roumieu case, respectively. By (III), there exist $C, \tau > 0$ (for every $\tau > 0$ there exists $C > 0$) such that

$$(3.2) \quad \|\varphi T_{-m} \psi\|_E \leq C e^{-A(\tau|m|)} \|e^{2A(\tau|m|)} \psi T_m \varphi\|_E, \quad \forall \varphi \in \mathcal{S}_\dagger^*, \quad \forall m \in \mathbb{Z}^d.$$

For $m \in \mathbb{Z}^d$, consider the linear mapping $\rho_{m,\tau}(\varphi) = e^{2A(\tau|m|)} \psi T_m \varphi$, $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d) \rightarrow \mathcal{S}_{(p!)}^{(M_p)}(\mathbb{R}^d)$ ($\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d) \rightarrow \mathcal{S}_{\{p!\}}^{\{M_p\}}(\mathbb{R}^d)$). Clearly, it is well defined. Let B be a bounded subset of $\mathcal{S}_\dagger^*(\mathbb{R}^d)$. Then for every $h > 0$ (there exists $h > 0$) such that

$$(3.3) \quad \sup_{\varphi \in B} \sup_{\alpha \in \mathbb{N}^d} \frac{h^{|\alpha|} \|e^{A(h|\cdot|)} D^\alpha \varphi\|_{L^\infty}}{M_\alpha} < \infty$$

Now, [7, Lemma 3.6] implies

$$(3.4) \quad e^{2A(\tau|m|)} \leq 2c_0 e^{A(2H\tau|x+m|)} e^{A(2H\tau|x|)}.$$

In the Beurling case, let $h_1 > 0$ be arbitrary but fixed. Choose $h > 0$ such that $h \geq \max\{2H\tau, 2h_1\}$ and $e^{A(2H\tau\lambda)} \leq C' e^{h\lambda}$ for all $\lambda \geq 0$ (such an h exists because $p! \subset A_p$). By (3.3) and (3.4) we have

$$(3.5) \quad \frac{h_1^{|\alpha|} |D^\alpha(\psi(x) T_m \varphi(x))| e^{h_1|x|}}{M_\alpha} \leq C_2 e^{-2A(\tau|m|)},$$

for all $x \in \mathbb{R}^d$, $m \in \mathbb{Z}^d$, $\varphi \in B$. Hence $\{\rho_{m,\tau} | m \in \mathbb{Z}^d\}$ is uniformly bounded on B . In the Roumieu case there exist $\tilde{h}, \tilde{C} > 0$ such that $\tilde{h}^{|\alpha|} |D^\alpha \psi(x)| e^{\tilde{h}|x|} \leq \tilde{C} M_\alpha$ for all $x \in \mathbb{R}^d$, $\alpha \in \mathbb{N}^d$. For the $h > 0$ for which (3.3) holds choose $0 < \tau \leq h/(2H)$ such that $e^{A(2H\tau\lambda)} \leq C' e^{\tilde{h}\lambda/2}$ for all $\lambda \geq 0$ (such a τ exists because $p! \subset A_p$). Choose $h_1 \leq \min\{h/2, \tilde{h}/2\}$. Then, by using (3.3) and (3.4), similarly as in the Beurling case, we obtain (3.5), i.e., $\{\rho_{m,\tau} | m \in \mathbb{Z}^d\}$ is uniformly bounded on B . Now, (I)' implies that $\|\rho_{m,\tau}(\varphi)\|_E \leq C'_2$ for all $\varphi \in B$, $m \in \mathbb{Z}^d$. By using (3.2), we obtain that the sequence $\left\{ \sum_{|m| \leq N} \varphi T_{-m} \psi \right\}_{N=0}^\infty$ is a Cauchy sequence in E for each $\varphi \in B$. Since its limit is φ in $\mathcal{S}'_{(p!)}^{(M_p)}(\mathbb{R}^d)$ (in $\mathcal{S}'_{\{p!\}}^{\{M_p\}}(\mathbb{R}^d)$) it converges to $\varphi \in E$. Also $\|\varphi\|_E \leq C$ for all $\varphi \in B$. This implies that $\mathcal{S}_\dagger^*(\mathbb{R}^d) \subseteq E$ and the inclusion maps bounded sets into bounded sets. As $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is bornological, the inclusion is continuous. It remains to prove $E \subseteq \mathcal{S}_\dagger^*(\mathbb{R}^d)$. By (I)' for a bounded set B in $\mathcal{S}'_{(p!)}^{(M_p)}(\mathbb{R}^d)$ (in $\mathcal{S}'_{\{p!\}}^{\{M_p\}}(\mathbb{R}^d)$)

there exists $D > 0$ such that $|\langle g, \check{\varphi} \rangle| \leq D \|g\|_E$ for all $g \in E$ and $\varphi \in B$. Then (III) implies that there exist $C, \tau > 0$ (for every $\tau > 0$ there exists $C > 0$) such that

$$|(g * \varphi)(y)| \leq D \|T_y g\|_E \leq CD e^{A(\tau|y|)}, \text{ for all } y \in \mathbb{R}^d, \varphi \in B, g \in E.$$

In the Beurling case Lemma 2.6 implies $E \subseteq \mathcal{S}'_{(A_p)}^{(M_p)}(\mathbb{R}^d)$. In the Roumieu case Lemma 2.6 together with Lemma 3.5 implies $E \subseteq \mathcal{S}'_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$. Since $E \rightarrow \mathcal{S}'_{(p!)}^{(M_p)}(\mathbb{R}^d)$ is continuous ($E \rightarrow \mathcal{S}'_{\{p!\}}^{\{M_p\}}(\mathbb{R}^d)$ is continuous) it has a closed graph. Thus the inclusion $E \rightarrow \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ has a closed graph. As $\mathcal{S}'_{(A_p)}^{(M_p)}(\mathbb{R}^d)$ is a (DFS) -space ($\mathcal{S}'_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ is an (FS) -space), it is a Pták space (cf. [12, Sect. IV. 8, p. 162]). Thus the continuity of $E \rightarrow \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ follows from the Pták closed graph theorem (cf. [12, Thm. 8.5, p. 166]). \square

Throughout the rest of the article we shall *always assume* that E is a translation-invariant (B) -spaces of ultradistributions of class $* - \dagger$. Our next concern is the study of convolution structures on E . We need three technical lemmas.

Lemma 3.7. *Let $\varphi \in \mathcal{S}'_{\dagger}(\mathbb{R}^{2d})$. Then for each $y \in \mathbb{R}^d$, $\varphi(\cdot, y) \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ and the function $\psi(x) = \int_{\mathbb{R}^d} \varphi(x, y) dy$ is an element of $\mathcal{S}'_{\dagger}(\mathbb{R}^d)$. Moreover, the function $\mathbf{f} : \mathbb{R}^d \rightarrow E$, $y \mapsto \varphi(\cdot, y)$, is Bochner integrable and $\psi = \int_{\mathbb{R}^d} \mathbf{f}(y) dy$.*

Proof. The fact that $\varphi(\cdot, y) \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ for each $y \in \mathbb{R}^d$ and that $\psi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ is trivial. Thus \mathbf{f} is well defined on \mathbb{R}^d with values in E (in fact its values are in $\mathcal{S}'_{\dagger}(\mathbb{R}^d)$). One easily verifies that \mathbf{f} is continuous, hence strongly measurable. To prove that it is Bochner integrable it remains to prove that $y \mapsto \|\mathbf{f}(y)\|_E$ is in $L^1(\mathbb{R}^d)$. The condition (I) implies

$$\|\mathbf{f}(y)\|_E \leq C_1 \sup_{\alpha} \frac{m^{|\alpha|} \|e^{A(m|\cdot|)} D_x^{\alpha} \varphi(\cdot, y)\|_{L^{\infty}(\mathbb{R}^d)}}{M_{\alpha}} \leq C_2 \sigma_{mH}(\varphi) e^{-A(m|y|)}.$$

Thus \mathbf{f} is Bochner integrable. Now, for $n \in \mathbb{Z}_+$, denote $K_n = [-n, n]^d$. Since K_n is compact and \mathbf{f} is continuous there exists $l(n) \in \mathbb{Z}_+$ such that $l(n) \geq n$ and $\|\mathbf{f}(y) - \mathbf{f}(y')\|_E \leq 2^{-n}$ when $y, y' \in K_n$ and $|y_j - y'_j| \leq 1/l(n)$, $j = 1, \dots, d$. Of course we can take $l(n+1) > l(n)$, for all $n \in \mathbb{Z}_+$. Set $D_n = \{y \in K_n \mid y = (k_1/l(n), \dots, k_d/l(n)), k_j \in \mathbb{Z}, -nl(n) \leq k_j \leq nl(n) - 1, j = 1, \dots, d\}$ and let

$$L_n(x) = \sum_{t \in D_n} \varphi(x, t) l(n)^{-d}.$$

Clearly $L_n \in \mathcal{S}'_{\dagger}(\mathbb{R}^d) \subseteq E$. We prove that $L_n \rightarrow \psi$ when $n \rightarrow \infty$, in $\mathcal{S}'_{\dagger}(\mathbb{R}^d)$. We give the proof for the Roumieu case, the Beurling case being similar. There exists $m > 0$ such that $\varphi \in \mathcal{S}'_{A_p, m}^{M_p, m}(\mathbb{R}^d)$. Pick $m' > 0$ such that $m' \leq m/(2H^2)$. For each $t = (t_1, \dots, t_d) \in D_n$ denote $K_{n,t} = [t_1, t_1 + 1/l(n)] \times \dots \times [t_d, t_d + 1/l(n)]$.

Observe that

$$\begin{aligned} & |D^\alpha \psi(x) - D^\alpha L_n(x)| \\ & \leq \int_{\mathbb{R}^d \setminus K_n} |D_x^\alpha \varphi(x, y)| dy + \sum_{t \in D_n} \int_{K_{n,t}} |D_x^\alpha \varphi(x, y) - D_x^\alpha \varphi(x, t)| dy = S_1(x) + S_2(x). \end{aligned}$$

For $y \in K_{n,t}$, by Taylor expanding $D_x^\alpha \varphi(x, y)$ at (x, t) , we have

$$\begin{aligned} & |D_x^\alpha \varphi(x, y) - D_x^\alpha \varphi(x, t)| \\ & \leq \sum_{|\beta|=1} |(y-t)^\beta| \int_0^1 |D_x^\alpha D_y^\beta \varphi(x, t + s(y-t))| ds \\ & \leq \frac{d\sigma_m(\varphi)}{n} \cdot m^{-|\alpha|-1} M_{|\alpha|+1} \int_0^1 e^{-A(m|(x,t+s(y-t))|)} ds. \end{aligned}$$

By [7, Proposition 3.6] and the fact $e^{A(\rho+\mu)} \leq 2e^{A(2\rho)}e^{A(2\mu)}$, for $\rho, \mu > 0$ (which can be easily verified), we have

$$\begin{aligned} e^{A(m'|x|)} e^{A(m'|y|)} & \leq 2e^{A(m'|x|)} e^{A(2m'|t+s(y-t)|)} e^{A(2m'|(1-s)(y-t)|)} \\ & \leq c_1 e^{A(2m'H|(x,t+s(y-t))|)}. \end{aligned}$$

Hence

$$|D_x^\alpha \varphi(x, y) - D_x^\alpha \varphi(x, t)| \leq \frac{C_1}{n} \cdot \frac{H^{|\alpha|} M_\alpha}{m^{|\alpha|} e^{A(m'|x|)} e^{A(m'|y|)}} \leq \frac{C_1 M_\alpha}{nm^{|\alpha|} e^{A(m'|x|)} e^{A(m'|y|)}}.$$

Thus, for $S_2(x)$ we have the following estimate

$$(3.6) \quad S_2(x) \leq \frac{C_1 M_\alpha}{nm^{|\alpha|} e^{A(m'|x|)}} \int_{\mathbb{R}^d} e^{-A(m'|y|)} dy \leq \frac{C_2 M_\alpha}{nm^{|\alpha|} e^{A(m'|x|)}}.$$

To estimate S_1 , we proceed as follows

$$S_1(x) \leq \frac{\sigma_m(\varphi) M_\alpha}{m^{|\alpha|}} \int_{\mathbb{R}^d \setminus K_n} e^{-A(m|(x,y)|)} dy.$$

For $y \in \mathbb{R}^d \setminus K_n$, by [7, Proposition 3.6], we have

$$e^{A(m'n)} e^{A(m'|x|)} e^{A(m'|y|)} \leq c_0 e^{A(m'|x|)} e^{A(m'H|y|)} \leq c_0^2 e^{A(m'H^2|(x,y)|)}.$$

Hence

$$(3.7) \quad S_1(x) \leq \frac{C_3 M_\alpha}{m^{|\alpha|} e^{A(m'|x|)} e^{A(m'n)}} \int_{\mathbb{R}^d} e^{-A(m'|y|)} dy$$

$$(3.8) \quad \leq \frac{C_4 M_\alpha}{m^{|\alpha|} e^{A(m'|x|)} e^{A(m'n)}}.$$

Now, (3.6) and (3.8) imply that $L_n \rightarrow \psi$ in $\mathcal{S}_{A_p, m'}^{M_p, m'}(\mathbb{R}^d)$ and hence also in $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$. As we noted, the Beurling case is completely analogous. By (I)

this also implies $L_n \rightarrow \psi$ in E . Denote by $\chi_{n,t}$ the characteristic function of $K_{n,t}$ and define

$$\mathbf{L}_n(y) = \sum_{t \in D_n} \mathbf{f}(t) \chi_{n,t}(y), \quad y \in \mathbb{R}^d.$$

Then \mathbf{L}_n is a simple function on \mathbb{R}^d with values in E and $\int_{\mathbb{R}^d} \mathbf{L}_n(y) dy = L_n$.

By using the continuity of \mathbf{f} one easily verifies that \mathbf{L}_n converges pointwisely to \mathbf{f} . Moreover, by the definition of $K_{n,t}$ we have $\|\mathbf{L}_n(y)\|_E \leq \|\mathbf{f}(y)\|_E + 2^{-n}$, for $y \in K_n$ and for $y \notin K_n$, $\mathbf{L}_n(y) = 0$. Thus, by defining $g(y) = 1/2$ for $y \in K_1$ and $g(y) = 2^{-n}$ when $y \in K_n \setminus K_{n-1}$ for $n \in \mathbb{Z}_+$, $n \geq 2$, we obtain $\|\mathbf{L}_n(y)\|_E \leq \|\mathbf{f}(y)\|_E + g(y)$ for all $y \in \mathbb{R}^d$. Since $g \in L^1(\mathbb{R}^d)$ and \mathbf{f} is Bochner integrable, dominated convergence implies

$$\lim_{n \rightarrow \infty} L_n = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} \mathbf{L}_n(y) dy = \int_{\mathbb{R}^d} \mathbf{f}(y) dy,$$

which completes the proof. \square

Lemma 3.8. *The convolution mapping $(\varphi, \psi) \in \mathcal{S}_\dagger^*(\mathbb{R}^d) \times \mathcal{S}_\dagger^*(\mathbb{R}^d) \rightarrow \varphi * \psi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$ extends to a continuous bilinear mapping $\mathcal{S}_\dagger^*(\mathbb{R}^d) \times E \rightarrow E$. Furthermore, the following estimate holds*

$$(3.9) \quad \|\varphi * g\|_E \leq \|g\|_E \int_{\mathbb{R}^d} |\varphi(x)| \omega(x) dx.$$

Proof. Let $\varphi, \psi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$. One easily verifies that the function $f(x, y) = \varphi(y)\psi(x - y)$ is an element of $\mathcal{S}_\dagger^*(\mathbb{R}^{2d})$. Define $\mathbf{f} : \mathbb{R}^d \rightarrow E$, $\mathbf{f}(y) = f(\cdot, y) = \varphi(y)T_{-y}\psi$. Then, by Lemma 3.7, \mathbf{f} is Bochner integrable and

$$\varphi * \psi = \int_{\mathbb{R}^d} \mathbf{f}(y) dy.$$

Observe that $\|\mathbf{f}(y)\|_E \leq |\varphi(y)|\omega(y)\|\psi\|_E$. Thus, we have

$$\|\varphi * \psi\|_E \leq \int_{\mathbb{R}^d} \|\mathbf{f}(y)\|_E dy \leq \|\psi\|_E \int_{\mathbb{R}^d} |\varphi(y)|\omega(y) dy,$$

which proves (3.9) for $g \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$. For general $g \in E$, (3.9) follows from a standard density argument. The continuity of the convolution as a bilinear mapping $\mathcal{S}_\dagger^*(\mathbb{R}^d) \times E \rightarrow E$ in the Beurling case is an easy consequence of (3.9). In the Roumieu case, from (3.9) we can conclude separate continuity, but $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ and E are barreled (DF) -spaces, hence the separate continuity implies the continuity of the convolution. \square

Lemma 3.9. $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is dense in L_ω^1 .

Proof. Observe that $C_c(\mathbb{R}^d)$ (the space of continuous functions with compact support) is dense in L_ω^1 . Thus it is enough to prove that each $\psi \in C_c(\mathbb{R}^d)$ can be approximated by elements of $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ in L_ω^1 . Let $\psi \in L_\omega^1$. Select a nonzero

$\varphi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$ such that $\int_{\mathbb{R}^d} \varphi(x) dx = 1$. For $n \in \mathbb{Z}_+$, set $\varphi_n(x) = n^d \varphi(nx)$. One easily verifies that $\varphi_n * \psi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$. We prove $\varphi_n * \psi \rightarrow \psi$ in $L_\omega^1(\mathbb{R}^d)$. We consider the Roumieu case, as the Beurling case is analogous. By (III) there exist $(l_p) \in \mathfrak{R}$ and $C' > 0$ such that $\omega(x) \leq C' e^{B_{l_p}(|x|)}$. By Lemma 1.1 we can assume that (l_p) satisfies $\prod_{j=1}^{p+q} l_j \leq 2^{p+q} \prod_{j=1}^p l_j \cdot \prod_{j=1}^q l_j$, for all $p, q \in \mathbb{Z}_+$. Let $r_p = l_p/4H$, $p \in \mathbb{Z}_+$. Since $\varphi \in \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$, $|\varphi(x)| \leq C'' e^{-B_{r_p}(|x|)}$. Observe that

$$\begin{aligned} \omega(x) |(\varphi_n * \psi)(x) - \psi(x)| &\leq \omega(x) \int_{\mathbb{R}^d} |\varphi(y)| |\psi(x - y/n) - \psi(x)| dy \\ &\leq C e^{B_{l_p}(|x|)} \int_{\mathbb{R}^d} e^{-B_{r_p}(|y|)} |\psi(x - y/n) - \psi(x)| dy. \end{aligned}$$

Since ψ has compact support $e^{B_{l_p}(|x|)} e^{-B_{r_p}(|y|)} |\psi(x)| \in L^1(\mathbb{R}_{x,y}^{2d})$ and

$$e^{B_{l_p}(2|x|)} |\psi(x)| \leq C_1 \langle x \rangle^{-d-1}, \quad \forall x \in \mathbb{R}^d.$$

This inequality, together with $e^{B_{l_p}(\rho+\mu)} \leq 2e^{B_{l_p}(2\rho)} e^{B_{l_p}(2\mu)}$, $\rho, \mu > 0$, implies

$$\begin{aligned} e^{B_{l_p}(|x|)} |\psi(x - y/n)| &\leq 2e^{B_{l_p}(2|x-y/n|)} e^{B_{l_p}(2|y/n|)} |\psi(x - y/n)| \\ &\leq C_1 \langle x - y/n \rangle^{-d-1} e^{B_{l_p}(2|y|)} \leq C_2 \langle x \rangle^{-d-1} \langle y \rangle^{d+1} e^{B_{l_p}(2|y|)} \\ &\leq C_3 \langle x \rangle^{-d-1} \langle y \rangle^{-d-1} e^{2B_{l_p}(2|y|)}. \end{aligned}$$

Since the sequence $A_p \prod_{j=1}^p l_j$ satisfies (M.2) with the constant $2H$ instead of H , [7, Proposition 3.6] implies $e^{2B_{l_p}(2|y|)} \leq c' e^{B_{r_p}(|y|)}$ (by definition of (r_p)). Thus

$$e^{B_{l_p}(|x|)} e^{-B_{r_p}(|y|)} |\psi(x - y/n)| \leq C_4 \langle x \rangle^{-d-1} \langle y \rangle^{-d-1} \in L^1(\mathbb{R}_{x,y}^{2d}).$$

Since ψ is continuous, $e^{B_{l_p}(|x|)} e^{-B_{r_p}(|y|)} |\psi(x - y/n) - \psi(x)| \rightarrow 0$ as $n \rightarrow \infty$ pointwise. Hence, dominated convergence implies $\varphi_n * \psi - \psi \rightarrow 0$ as $n \rightarrow \infty$ in L_ω^1 . \square

Combining Lemmas 3.8 and 3.9, we immediately obtain the ensuing important proposition.

Proposition 3.10. *The convolution extends as a mapping $L_\omega^1 \times E \rightarrow E$ and E becomes a Banach module over the Beurling algebra L_ω^1 , i.e., $\|u * g\|_E \leq \|u\|_{1,\omega} \|g\|_E$.*

Corollary 3.11. *Let $g \in E$ and $\varphi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$. Set $\varphi_\varepsilon(x) = \varepsilon^{-d} \varphi(x/\varepsilon)$. Then,*

$$\lim_{\varepsilon \rightarrow 0^+} \|c g - \varphi_\varepsilon * g\|_E = 0,$$

where $c = \int_{\mathbb{R}^d} \varphi(x) dx$.

Proof. Let $0 < \varepsilon < 1$. We first consider the case when $\varphi, g \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$. Observe that

$$cg(x) - (\varphi_\varepsilon * g)(x) = \int_{\mathbb{R}^d} (g(x) - g(x - \varepsilon y)) \varphi(y) dy.$$

One easily verifies that the function $f_\varepsilon(x, y) = (g(x) - g(x - \varepsilon y)) \varphi(y)$ is in $\mathcal{S}_\dagger^*(\mathbb{R}^{2d})$. Define $\mathbf{f}_\varepsilon(y) = f_\varepsilon(\cdot, y) = (g - T_{-\varepsilon y}g) \varphi(y)$, $\mathbb{R}^d \rightarrow E$. Lemma 3.7 implies that \mathbf{f}_ε is Bochner integrable and

$$(3.10) \quad \|cg - \varphi_\varepsilon * g\|_E = \left\| \int_{\mathbb{R}^d} \mathbf{f}_\varepsilon(y) dy \right\|_E \leq \int_{\mathbb{R}^d} \|g - T_{-\varepsilon y}g\|_E |\varphi(y)| dy.$$

Clearly $\|g - T_{-\varepsilon y}g\|_E |\varphi(y)| \leq \|g\|_E (1 + Ce^{A(m|y|)}) |\varphi(y)|$ for some $C, m > 0$ (for each $m > 0$ and a corresponding $C = C_m$). Since the left hand side is in $L^1(\mathbb{R}^d)$ and, by Lemma 3.2, $\|g - T_{-\varepsilon y}g\|_E |\varphi(y)| \rightarrow 0$ for each fixed $y \in \mathbb{R}^d$ as $\varepsilon \rightarrow 0^+$, dominated convergence together with (3.10) proves the corollary. Due to the density of $\mathcal{S}_\dagger^*(\mathbb{R}^d) \hookrightarrow E$, the conclusion in the lemma for $g \in E$ and $\varphi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$ follows by using the estimate (3.9). \square

Proposition 3.12. *The space E' satisfies*

- a') $\mathcal{S}_\dagger^*(\mathbb{R}^d) \rightarrow E' \hookrightarrow \mathcal{S}_\dagger'^*(\mathbb{R}^d)$, with continuous embeddings.
- b') For each h , $T_h : E' \rightarrow E'$ is a bounded operator. The mappings $\mathbb{R}^d \rightarrow E'$, given by $h \mapsto T_h f$, are continuous for the weak* topology.

Moreover, the property (III) holds true when E is replaced by E' .

Proof. The proof is similar to that of [2, Proposition 2]. \square

We can now associate a Beurling algebra to E' . Set

$$\check{\omega}(h) := \|T_{-h}\|_{\mathcal{L}(E')} = \|T_h^\top\|_{\mathcal{L}(E')} = \omega(-h).$$

The very last equality follows from the well-known property $\|T_h^\top\|_{\mathcal{L}(E')} = \|T_h\|_{\mathcal{L}(E)}$, which is of course a consequence of the bipolar theorem (cf. [12, p. 160]). The associated Beurling algebra to the dual space E' is $L_{\check{\omega}}^1$. We define the convolution $u * f = f * u$ of $f \in E'$ and $u \in L_{\check{\omega}}^1$ via transposition:

$$(3.11) \quad \langle u * f, g \rangle := \langle f, \check{u} * g \rangle, \quad g \in E.$$

In view of Proposition 3.10, this convolution is well-defined because $\check{u} \in L_{\check{\omega}}^1$.

Corollary 3.13. *We have $\|u * f\|_{E'} \leq \|u\|_{1, \check{\omega}} \|f\|_{E'}$ and thus E' is a Banach module over the Beurling algebra $L_{\check{\omega}}^1$. In addition, if φ_ε and c are as in Corollary 3.11, then $\varphi_\varepsilon * f \rightarrow cf$ as $\varepsilon \rightarrow 0^+$ weakly* in E' for each fixed $f \in E'$.*

Proof. For $g \in E$ fixed we have $\langle \varphi_\varepsilon * f - cf, g \rangle = \langle f, \check{\varphi}_\varepsilon * g - cg \rangle$. \square

In general the embedding $\mathcal{S}_\dagger^*(\mathbb{R}^d) \rightarrow E'$ is not dense (consider for instance $E = L^1$). However, E' inherits the three properties (I), (II), and (III) whenever E is reflexive. The following result is a direct consequence of Proposition 3.12 and the Hahn-Banach theorem.

Proposition 3.14. *If E is reflexive, then its dual space E' is also a translation-invariant (B) -space of ultradistributions of class $* - \dagger$.*

The fact that the mappings $h \mapsto T_h f, \mathbb{R}^d \rightarrow E'$ do not have to be necessarily continuous in the non-reflexive case ($E = L^1(\mathbb{R}^d)$ is an example) causes various difficulties when dealing with this space. As in the non-quasianalytic case [2, 4], we will often work with the closed subspace E'_* of E' from the following definition rather than with E' itself.

Definition 3.15. The (B) -space E'_* stands for $L_\omega^1 * E'$.

Note that E'_* is a closed linear subspace of E' , due to the Cohen-Hewitt factorization theorem [6] and the fact that L_ω^1 possesses bounded approximation unities.

Remark 3.16. Observe that $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is a subset of the closure of $\text{span}(\mathcal{S}_\dagger^*(\mathbb{R}^d) * \mathcal{S}_\dagger^*(\mathbb{R}^d))$ in E' , where $\text{span}(A)$ denotes the linear span of a set. To see this, let $\varphi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$. Then, if $\chi_n, n \in \mathbb{Z}_+$, is a δ -sequence from $\mathcal{S}_\dagger^*(\mathbb{R}^d)$, $\chi_n * \varphi \rightarrow \varphi$ in $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ hence also in E' (by a') of Proposition 3.12). Whence we also obtain that $\mathcal{S}_\dagger^*(\mathbb{R}^d) \subseteq E'_*$.

The space E'_* will be of crucial importance throughout the rest of this work. It possesses richer properties than E' with respect to the translation group, as stated in the next theorem.

Theorem 3.17. *The space E'_* has the properties a'), (\widetilde{II}) and (III) . It is a Banach module over the Beurling algebra L_ω^1 . If φ_ε and c are as in Corollary 3.11, then, for each $f \in E'_*$,*

$$(3.12) \quad \lim_{\varepsilon \rightarrow 0^+} \|cf - \varphi_\varepsilon * f\|_{E'} = 0.$$

Furthermore, if E is reflexive, then $E'_* = E'$.

Proof. The proof goes in the same lines as that of [4, Theorem 4.4] □

We point out that (3.12) implies that $\mathcal{S}_\dagger^*(\mathbb{R}^d) * E' \subseteq L_\omega^1 * E'$ is dense in E'_* . In fact, E'_* is the biggest subspace of E' where the mappings $h \mapsto T_h f, \mathbb{R}^d \rightarrow E'$, are continuous. The proof of this result is essentially the same as that of [4, Theorem 4.4], so we omit it.

Proposition 3.18. *We have $E'_* = \left\{ f \in E' \mid \lim_{h \rightarrow 0} \|T_h f - f\|_{E'} = 0 \right\}$.*

In view of property b') from Proposition 3.12, we can naturally define a convolution mapping $E' \times \check{E} \rightarrow C(\mathbb{R}^d)$, where $\check{E} = \{g \in \mathcal{S}_\dagger^*(\mathbb{R}^d) \mid \check{g} \in E\}$ with norm $\|g\|_{\check{E}} := \|\check{g}\|_E$, via

$$(f * g)(x) = \langle f(t), g(x - t) \rangle = \langle f(t), T_{-x}\check{g}(t) \rangle.$$

Observe that if E is a translation invariant (B) -space of ultradistributions of class $* - \dagger$, then so is \check{E} . Clearly $\|T_h\|_{\mathcal{L}(E)} = \|T_{-h}\|_{\mathcal{L}(\check{E})}$. Hence the convolution

can be defined in the same way as a mapping from $\check{E}' \times E$ to $C(\mathbb{R}^d)$. We end this section with a simple proposition describing the mapping properties of this convolution. As usual, L_ω^∞ , the dual of the Beurling algebra L_ω^1 , is the (B) -space of all measurable functions satisfying

$$\|u\|_{\infty,\omega} = \operatorname{ess\,sup}_{x \in \mathbb{R}^d} \frac{|g(x)|}{\omega(x)} < \infty.$$

We need the following two closed subspaces of L_ω^∞ ,

$$(3.13) \quad UC_\omega := \left\{ u \in L_\omega^\infty \mid \lim_{h \rightarrow 0} \|T_h u - u\|_{\infty,\omega} = 0 \right\}$$

and

$$(3.14) \quad C_\omega := \left\{ u \in C(\mathbb{R}^d) \mid \lim_{|x| \rightarrow \infty} \frac{u(x)}{\omega(x)} = 0 \right\}.$$

The proof of the following proposition is simple and we thus omit it (the second part about the reflexive case follows from Proposition 3.14).

Proposition 3.19. *$E' * \check{E} \subseteq UC_\omega$ and $E' \times \check{E} \rightarrow UC_\omega$ is continuous. If E is reflexive, then $E' * \check{E} \subseteq C_\omega$. Similarly $\check{E}' * E \subseteq UC_{\check{\omega}}$ and $\check{E}' \times E \rightarrow UC_{\check{\omega}}$ is continuous. When E is reflexive, $E' * \check{E} \subseteq C_\omega$ and $\check{E}' * E \subseteq C_{\check{\omega}}$.*

We conclude this section with some examples of translation-invariant (B) -spaces of quasianalytic ultradistributions.

Example 3.20 (Weighted L_η^p spaces). Let η be an *ultrapolynomially bounded weight function of class \dagger* , that is, a (Borel) measurable function $\eta : \mathbb{R}^d \rightarrow (0, \infty)$ that fulfills the requirement $\eta(x+h) \leq C\eta(x)e^{A(\tau|h|)}$ for some $C, \tau > 0$ (for every $\tau > 0$ there exists $C > 0$). For $1 \leq p < \infty$ we denote as L_η^p the spaces of measurable functions g such that $\|g\|_{p,\eta} := \|\eta g\|_p < \infty$. Clearly L_η^p are translation-invariant (B) -spaces of ultradistributions of class $* - \dagger$ for $p \in [1, \infty)$ and for any sequence $(M_p)_{p \in \mathbb{N}}$. On the other hand, we make an exception and define L_η^∞ via the norm $\|g\|_{\infty,\eta} := \|g/\eta\|_\infty$. We also introduce the closed spaces UC_η and C_η of L_η^∞ as in (3.13) and (3.14) with ω replaced by η . Note that C_η is a translation-invariant (B) -spaces of ultradistributions of class $* - \dagger$ because $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is dense in it, while L_η^∞ and UC_η fail to have this property.

As usual, we write q for the conjugate index of p . As well known, $(L_\eta^p)' = L_{\eta^{-1}}^q$ if $1 < p < \infty$ and $(L_\eta^1)' = L_\eta^\infty$. In view of Proposition 3.17, the space E'_* corresponding to $E = L_{\eta^{-1}}^p$ is $E'_* = L_\eta^q$ whenever $1 < p < \infty$. On the other hand, Proposition 3.18 gives that $E'_* = UC_\eta$ for $E = L_\eta^1$. The Beurling algebra of L_η^p can be explicitly determined as in [2, Proposition 10], we state the result for the reader's convenience. Note that when $\log \eta$ is a subadditive function and $\eta(0) = 1$, the following proposition yields $\omega_\eta = \eta$ (a.e.).

Proposition 3.21. *Let $\omega_\eta(h) := \text{ess sup}_{x \in \mathbb{R}^d} \eta(x+h)/\eta(x)$. Then*

$$\|T_{-h}\|_{\mathcal{L}(L_\eta^p)} = \begin{cases} \omega_\eta(h) & \text{if } p \in [1, \infty), \\ \omega_\eta(-h) & \text{if } p = \infty. \end{cases}$$

Consequently, the Beurling algebra of L_η^p is $L_{\omega_\eta}^1$ if $p \in [1, \infty)$ and $L_{\tilde{\omega}_\eta}^1$ if $p = \infty$.

Clearly, the Beurling algebra of C_η is $L_{\omega_\eta}^1$. We now compute the space E'_* corresponding to $E = C_\eta$. Note that η can be assumed to be continuous (the continuous weight $\eta_1 = \eta * \varphi$ defines an equivalent norm if we choose $\varphi \in \mathcal{D}(\mathbb{R}^d)$ being non-negative with $\int_{\mathbb{R}^d} \varphi(x) dx = 1$). Thus $E = C_\eta$ is isometrically isomorphic to $C_0(\mathbb{R}^d)$, the isometry being $J_\eta : C_\eta \rightarrow C_0(\mathbb{R}^d)$, $J_\eta(\psi) = \psi/\eta$. Hence ${}^t J_\eta : \mathcal{M}^1 \rightarrow (C_\eta)'$ is isometric isomorphism and thus for each $f \in (C_\eta)'$ there exists a unique finite measure $\nu \in \mathcal{M}^1$ such that $\langle f, \psi \rangle = \int_{\mathbb{R}^d} \psi(x)/\eta(x) d\nu(x)$ for all $\psi \in C_\eta$. We will denote the dual of C_η by \mathcal{M}_η^1 . Now, one easily verifies that $L_{\omega_\eta}^1 * \mathcal{M}_\eta^1 \subseteq L_\eta^1$ and since $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is dense in L_η^1 (the proof is analogous to that of Lemma 3.9) and $\mathcal{S}_\dagger^*(\mathbb{R}^d) * \mathcal{S}_\dagger^*(\mathbb{R}^d)$ is dense in $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ (cf. Remark 3.16), we obtain that $E'_* = L_\eta^1$.

4. ULTRADISTRIBUTION SPACES OF CLASS $* - \dagger$ ASSOCIATED TO TRANSLATION-INVARIANT (B) -SPACES

In this section we construct and study test function and ultradistribution spaces associated to translation-invariant (B) -spaces of ultradistributions of class $* - \dagger$. The construction of such spaces is similar to the one given in [4] in the non-quasianalytic case; however, the study of their properties requires new non-trivial arguments. We recall that throughout the rest of the paper E stands for a tempered translation-invariant (B) -space of ultradistributions whose growth function of its translation group is ω (cf. (3.1)). The (B) -space $E'_* \subseteq E'$ was introduced in Definition 3.15.

4.1. The test function space \mathcal{D}_E^* . We begin by constructing our test space. Let

$$\mathcal{D}_E^{M_p, m} = \left\{ \varphi \in E \mid D^\alpha \varphi \in E, \forall \alpha \in \mathbb{N}^d, \|\varphi\|_{E, m} = \sup_{\alpha \in \mathbb{N}^d} \frac{m^\alpha \|D^\alpha \varphi\|_E}{M_\alpha} < \infty \right\}.$$

It is easy to verify that $\mathcal{D}_E^{M_p, m}$ is (B) -space with the norm $\|\cdot\|_{E, m}$. None of these spaces is trivial. To see this in the Beurling case one only needs to use the continuity of the inclusion $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d) \rightarrow E$ to obtain that $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d) \subseteq \mathcal{D}_E^{M_p, m}$ for each $m > 0$. In the Roumieu case observe that $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d)$ is continuously injected into $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$, hence we have the continuous inclusions $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d) \rightarrow E$. Now, similarly one proves that $\mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d) \subseteq \mathcal{D}_E^{M_p, m}$ for each $m > 0$. Obviously, $\mathcal{D}_E^{M_p, m_1} \subseteq \mathcal{D}_E^{M_p, m_2}$ for $m_2 < m_1$ and the inclusion mapping is continuous. As

l.c.s., we define

$$\mathcal{D}_E^{(M_p)} = \varprojlim_{m \rightarrow \infty} \mathcal{D}_E^{M_p, m}, \quad \mathcal{D}_E^{\{M_p\}} = \varinjlim_{m \rightarrow 0} \mathcal{D}_E^{M_p, m}.$$

Since $\mathcal{D}_E^{\{M_p\}, m}$ is continuously injected into E for each $m > 0$, $\mathcal{D}_E^{\{M_p\}}$ is indeed a (Hausdorff) l.c.s.. Moreover $\mathcal{D}_E^{\{M_p\}}$ is a barreled bornological (DF) -space, since it is an inductive limit of (B) -spaces. Obviously $\mathcal{D}_E^{(M_p)}$ is an (F) -space. Of course $\mathcal{D}_E^{(M_p)}$ and $\mathcal{D}_E^{\{M_p\}}$ are continuously injected into E .

Additionally, in the Roumieu case, for each fixed $(r_p) \in \mathfrak{R}$ we define the (B) -space

$$\mathcal{D}_E^{\{M_p\}, (r_p)} = \left\{ \varphi \in E \mid D^\alpha \varphi \in E, \forall \alpha \in \mathbb{N}^d, \|\varphi\|_{E, (r_p)} = \sup_{\alpha} \frac{\|D^\alpha \varphi\|_E}{M_\alpha \prod_{j=1}^{|\alpha|} r_j} < \infty \right\},$$

with norm $\|\cdot\|_{E, (r_p)}$. Since for $k > 0$ and $(r_p) \in \mathfrak{R}$, there exists $C > 0$ such that $k^{|\alpha|} \geq C / \left(\prod_{j=1}^{|\alpha|} r_j \right)$, $\mathcal{D}_E^{\{M_p\}, k}$ is continuously injected into $\mathcal{D}_E^{\{M_p\}, (r_p)}$. Define as l.c.s. $\tilde{\mathcal{D}}_E^{\{M_p\}} = \varprojlim_{(r_p) \in \mathfrak{R}} \mathcal{D}_E^{\{M_p\}, (r_p)}$. Then $\tilde{\mathcal{D}}_E^{\{M_p\}}$ is a complete l.c.s. and $\mathcal{D}_E^{\{M_p\}}$

is continuously injected into $\tilde{\mathcal{D}}_E^{\{M_p\}}$.

Lemma 4.1. *The space $\mathcal{D}_E^{\{M_p\}}$ is regular, i.e., every bounded set B in $\mathcal{D}_E^{\{M_p\}}$ is bounded in some $\mathcal{D}_E^{\{M_p\}, m}$. In addition $\mathcal{D}_E^{\{M_p\}}$ is complete.*

Proof. An adaptation of the proof of [4, Proposition 5.1] proves the lemma. \square

Similarly as in the first part of the proof of [4, Proposition 5.1] one can prove, by using [8, Lemma 3.4], that $\mathcal{D}_E^{\{M_p\}}$ and $\tilde{\mathcal{D}}_E^{\{M_p\}}$ are equal as sets, i.e., the canonical inclusion $\mathcal{D}_E^{\{M_p\}} \rightarrow \tilde{\mathcal{D}}_E^{\{M_p\}}$ is surjective.

The next proposition gives the relationship between $\mathcal{S}_\dagger^*(\mathbb{R}^d)$, \mathcal{D}_E^* and E . The proof is essentially the same as that of [4, Proposition 5.2].

Proposition 4.2. *The following dense inclusions hold $\mathcal{S}_\dagger^*(\mathbb{R}^d) \hookrightarrow \mathcal{D}_E^* \hookrightarrow E \hookrightarrow \mathcal{S}_\dagger^{*}(\mathbb{R}^d)$ and \mathcal{D}_E^* is a topological module over the Beurling algebra L_ω^1 , i.e., the convolution $*$: $L_\omega^1 \times \mathcal{D}_E^* \rightarrow \mathcal{D}_E^*$ is continuous. Moreover, in the Beurling case the following estimate*

$$(4.1) \quad \|u * \varphi\|_{E, m} \leq \|u\|_{1, \omega} \|\varphi\|_{E, m}, \quad m > 0$$

holds. In the Roumieu case, for each $m > 0$ the convolution is also continuous bilinear mapping $L_\omega^1 \times \mathcal{D}_E^{M_p, m} \rightarrow \mathcal{D}_E^{M_p, m}$ and the inequality (4.1) holds.

We will often use the following results on the action of ultradifferential operators on the test space \mathcal{D}_E^* (see [4] for their proofs).

Lemma 4.3. *If $P(D)$ is ultradifferential operator of $*$ type, then $P(D) : \mathcal{D}_E^* \rightarrow \mathcal{D}_E^*$ is continuous.*

Lemma 4.4. *Every ultradifferential operator $P(D)$ of $\{M_p\}$ class acts continuously on $\tilde{\mathcal{D}}_E^{\{M_p\}}$.*

It turns out that all elements of our test function space \mathcal{D}_E^* are ultradifferentiable functions of class $*$. We need the following lemmas in order to establish this fact.

Lemma 4.5. *There exists $l > 0$ (there exists $(l_p) \in \mathfrak{R}$) such that $\mathcal{S}_{A_p, l}^{M_p, l} \subseteq E \cap E'_*$ ($\mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)} \subseteq E \cap E'_*$). Moreover, the inclusion mappings $\mathcal{S}_{A_p, l}^{M_p, l} \rightarrow E$ and $\mathcal{S}_{A_p, l}^{M_p, l} \rightarrow E'_*$ ($\mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)} \rightarrow E$ and $\mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)} \rightarrow E'_*$) are continuous.*

Proof. We give the proof in the Roumieu case, the Beurling case is similar. Since the inclusion $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d) \rightarrow E$ is continuous and $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d) = \varprojlim_{(r_p) \in \mathfrak{R}} \mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)}$ there exist $C > 0$ and $(r_p) \in \mathfrak{R}$ such that $\|\varphi\|_E \leq C\sigma_{(r_p)}(\varphi)$, $\forall \varphi \in \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$. For this (r_p) , by Lemma 2.3, there exist $(k_p) \in \mathfrak{R}$ and $\chi_n, \varphi_n \in \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$, $n \in \mathbb{Z}_+$, such that $\chi_n * (\varphi_n \psi) \in \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ for each $n \in \mathbb{Z}_+$ and $\chi_n * (\varphi_n \psi) \rightarrow \psi$ when $n \rightarrow \infty$ in $\mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)}$ for all $\psi \in \mathcal{S}_{A_p, (k_p)}^{M_p, (k_p)}$. We have

$$(4.2) \quad \|\chi_n * (\varphi_n \psi)\|_E \leq C\sigma_{(r_p)}(\chi_n * (\varphi_n \psi)).$$

We obtain that $\chi_n * (\varphi_n \psi)$ is a Cauchy sequence in E , hence it converges. Since $\chi_n * (\varphi_n \psi) \rightarrow \psi$ in $\mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)}$ the convergence also holds in $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$. But E is continuously injected into $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ thus the limit of $\chi_n * (\varphi_n \psi)$ in E must be ψ . If we let $n \rightarrow \infty$ in (4.2) we have $\|\psi\|_E \leq C\sigma_{(r_p)}(\psi) \leq C\sigma_{(k_p)}(\psi)$, which gives the desired continuity of the inclusion $\mathcal{S}_{A_p, (k_p)}^{M_p, (k_p)} \rightarrow E$. Similarly, one obtains the continuous inclusion $\mathcal{S}_{A_p, (k'_p)}^{M_p, (k'_p)} \rightarrow E'_*$ possibly with another $(k'_p) \in \mathfrak{R}$. The conclusion of the lemma now follows by taking $(l_p) \in \mathfrak{R}$ defined as $l_p = \min\{k_p, k'_p\}$, $p \in \mathbb{Z}_+$. \square

Lemma 4.6. *Let $f \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$ be a continuous function such that for each $\beta \in \mathbb{N}^d$ the ultradistributional derivative $D^\beta f$ is a continuous function with ultrapolynomial growth of type \dagger . Then $f \in C^\infty(\mathbb{R}^d)$.*

Proof. Since f is continuous $f \in \mathcal{D}'(\mathbb{R}^d)$ (the Schwartz space of distributions). First we prove that the ultradistributional derivatives of f coincide with its distributional derivatives. We give the proof in the Roumieu case. The Beurling case is similar. Let $\beta \in \mathbb{N}^d$. Denote by f_β the distributional derivative $D^\beta f$ of f and by \tilde{f}_β the ultradistributional derivative $D^\beta f$ of f . Since f and \tilde{f}_β are continuous functions of ultrapolynomial growth $\{A_p\}$, similarly as in the proof of (c) \Leftrightarrow (\tilde{c}) in [4, Theorem 4.2], one can prove that there exist $(r_p) \in \mathfrak{R}$ and $C > 0$ such that $|f(x)| \leq Ce^{B_{r_p}(|x|)}$ and $|\tilde{f}_\beta(x)| \leq Ce^{B_{r_p}(|x|)}$. Pick $(k_p) \in \mathfrak{R}$ such that $(k_p) \leq (r_p)$ and $e^{B_{r_p}(|\cdot|)}e^{-B_{k_p}(|\cdot|)} \in L^1(\mathbb{R}^d)$. Fix $\psi \in \mathcal{D}(\mathbb{R}^d)$. Let

$\chi_n \in \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$, $n \in \mathbb{Z}_+$, be defined as in *ii*) of Lemma 2.3. One easily verifies that $\psi_n = \chi_n * \psi \in \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$. Let $\alpha \leq \beta$. Observe that

$$(4.3) \quad e^{B_{k_p}(|x|)} |D^\alpha \psi_n(x) - D^\alpha \psi(x)| \\ \leq 2 \int_{\mathbb{R}^d} |\chi(y)| e^{B_{k_p}(2|y|)} |D^\alpha \psi(x - y/n) - D^\alpha \psi(x)| e^{B_{k_p}(2|x-y/n|)} dy.$$

Let $\varepsilon > 0$. Since ψ is compactly supported,

$$|D^\alpha \psi(x - y/n) - D^\alpha \psi(x)| e^{B_{k_p}(2|x-y/n|)} \leq C_1 + |D^\alpha \psi(x)| e^{B_{k_p}(2|x-y/n|)} \\ \leq C_1 + C_2 e^{B_{k_p}(4|y|)}.$$

As $|\chi(y)| e^{B_{k_p}(2|y|)} (C_1 + C_2 e^{B_{k_p}(4|y|)}) \in L^1(\mathbb{R}^d)$, there exists $c_1 \geq 1$ such that

$$\int_{|y| \geq c_1} |\chi(y)| e^{B_{k_p}(2|y|)} (C_1 + C_2 e^{B_{k_p}(4|y|)}) dy \leq \varepsilon/4.$$

Of course, we can assume that c_1 is large enough such that $\text{supp } \psi \subseteq \{x \in \mathbb{R}^d \mid |x| \leq c_1\}$. Clearly $D^\alpha \psi(x) = 0$ and $D^\alpha \psi(x - y/n) = 0$ for all $n \in \mathbb{Z}_+$ when $|x| > 2c_1$ and $|y| \leq c_1$. Hence, for $|x| \leq 2c_1$, $|y| \leq c_1$ and $n \in \mathbb{Z}_+$ there exists C_2 such that $e^{B_{k_p}(2|x-y/n|)} \leq C_2$. Since $D^\alpha \psi$ is continuous, there exists $n_0 \in \mathbb{Z}_+$ such that for all $n \geq n_0$, $|x| \leq 2c_1$ and $|y| \leq c_1$

$$|D^\alpha \psi(x - y/n) - D^\alpha \psi(x)| \leq \varepsilon / \left(4C_2 \|\chi e^{B_{k_p}(2|\cdot|)}\|_{L^1(\mathbb{R}^d)} \right).$$

These estimates, together with (4.3), imply $\|e^{B_{k_p}(\cdot|\cdot)} (D^\alpha \psi_n - D^\alpha \psi)\|_{L^\infty(\mathbb{R}^d)} \leq \varepsilon$ for all $n \geq n_0$. We obtain that for each $\alpha \leq \beta$, $e^{B_{k_p}(\cdot|\cdot)} D^\alpha \psi_n \rightarrow e^{B_{k_p}(\cdot|\cdot)} D^\alpha \psi$ in $L^\infty(\mathbb{R}^d)$. Now, dominated convergence implies

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} \tilde{f}_\beta(x) \psi_n(x) dx = \int_{\mathbb{R}^d} \tilde{f}_\beta(x) \psi(x) dx, \\ \lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} f(x) (-D)^\beta \psi_n(x) dx = \int_{\mathbb{R}^d} f(x) (-D)^\beta \psi(x) dx.$$

Hence

$$\langle \tilde{f}_\beta, \psi \rangle = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} \tilde{f}_\beta(x) \psi_n(x) dx = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} f(x) (-D)^\beta \psi_n(x) dx = \langle f_\beta, \psi \rangle.$$

Since $\psi \in \mathcal{D}(\mathbb{R}^d)$ is arbitrary $\tilde{f}_\beta = f_\beta$. In other words f is a continuous function whose all distributional derivatives are continuous functions. Now the Sobolev imbedding theorem applied on a ball with center at a fixed point $x \in \mathbb{R}^d$ implies that f is C^∞ in that ball. As x is arbitrary, the assertion follows. \square

Define for every $m, h > 0$ the (B) -spaces

$$\mathcal{O}_{A_p, m, h}^{M_p} = \left\{ \varphi \in C^\infty(\mathbb{R}^d) \mid \|\varphi\|_{m, h} = \left(\sum_{\alpha \in \mathbb{N}^d} \frac{m^{2|\alpha|}}{M_\alpha^2} \|D^\alpha \varphi e^{-A(h|\cdot|)}\|_{L^2}^2 \right)^{1/2} < \infty \right\}.$$

Observe that for $m_1 \leq m_2$ we have the continuous inclusion $\mathcal{O}_{A_p, m_2, h}^{M_p} \rightarrow \mathcal{O}_{A_p, m_1, h}^{M_p}$ and for $h_1 \leq h_2$ the inclusion $\mathcal{O}_{A_p, m, h_1}^{M_p} \rightarrow \mathcal{O}_{A_p, m, h_2}^{M_p}$ is also continuous. As l.c.s. we define

$$\begin{aligned} \mathcal{O}_{(A_p), h}^{(M_p)} &= \varprojlim_{m \rightarrow \infty} \mathcal{O}_{A_p, m, h}^{M_p} \quad , \quad \mathcal{O}_{(A_p), C}^{(M_p)} = \varinjlim_{h \rightarrow \infty} \mathcal{O}_{(A_p), h}^{(M_p)} ; \\ \mathcal{O}_{\{A_p\}, h}^{\{M_p\}} &= \varinjlim_{m \rightarrow 0} \mathcal{O}_{A_p, m, h}^{M_p} \quad , \quad \mathcal{O}_{\{A_p\}, C}^{\{M_p\}} = \varprojlim_{h \rightarrow 0} \mathcal{O}_{\{A_p\}, h}^{\{M_p\}} . \end{aligned}$$

Observe that $\mathcal{O}_{(A_p), h}^{(M_p)}$ is an (F) -space and since all inclusions $\mathcal{O}_{(A_p), h}^{(M_p)} \rightarrow C^\infty(\mathbb{R}^d)$ are continuous (by the Sobolev imbedding theorem), $\mathcal{O}_{(A_p), C}^{(M_p)}$ is indeed a (Hausdorff) l.c.s.. Moreover, as an inductive limit of barreled and bornological spaces $\mathcal{O}_{(A_p), C}^{(M_p)}$ is barreled and bornological. Also $\mathcal{O}_{\{A_p\}, h}^{\{M_p\}}$ is (Hausdorff) l.c.s., because all inclusions $\mathcal{O}_{A_p, m, h}^{M_p} \rightarrow C^\infty(\mathbb{R}^d)$ are continuous (by the Sobolev imbedding theorem). Hence $\mathcal{O}_{\{A_p\}, C}^{\{M_p\}}$ is indeed a (Hausdorff) l.c.s.. Furthermore, $\mathcal{O}_{\{A_p\}, h}^{\{M_p\}}$ is a barreled and bornological (DF) -space, as inductive limit of (B) -spaces. By this considerations it also follows that $\mathcal{O}_{\dagger, C}^*$ is continuously injected into $C^\infty(\mathbb{R}^d)$. One easily verifies that $\mathcal{S}_{\dagger}^*(\mathbb{R}^d)$ is continuously and densely injected into $\mathcal{O}_{\dagger, C}^*$. We mention that $\mathcal{O}_{\dagger, C}^*$ was introduced and studied in [4] in the non-quasianalytic case.

Proposition 4.7. *The embedding $\mathcal{D}_E^* \hookrightarrow \mathcal{O}_{\dagger, C}^*(\mathbb{R}^d)$ holds. Furthermore, for $\varphi \in \mathcal{D}_E^*$, $D^\alpha \varphi \in C_{\tilde{\omega}}$ for all $\alpha \in \mathbb{N}^d$ and they satisfy the following growth condition: For every $m > 0$ (for some $m > 0$)*

$$(4.4) \quad \sup_{\alpha \in \mathbb{N}^d} \frac{m^{|\alpha|}}{M_\alpha} \|D^\alpha \varphi\|_{L_\omega^\infty(\mathbb{R}^d)} < \infty .$$

Proof. Let $r > 0$ ($(r_p) \in \mathfrak{R}$) be as in Lemma 4.5, that is, $\mathcal{S}_{A_p, r}^{M_p, r} \subseteq E \cap E'_*$ ($\mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)} \subseteq E \cap E'_*$) and the inclusion mappings $\mathcal{S}_{A_p, r}^{M_p, r} \rightarrow E$ and $\mathcal{S}_{A_p, r}^{M_p, r} \rightarrow E'_*$ ($\mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)} \rightarrow E$ and $\mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)} \rightarrow E'_*$) are continuous. By Proposition 2.2, there exist $u \in \mathcal{S}_{A_p, r}^{M_p, r}$ and $P(D)$ of type (M_p) ($u \in \mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)}$ and $P(D)$ of type $\{M_p\}$) such that $P(D)u = \delta$. Let $f \in \mathcal{D}_E^*$. Then $f = (P(D)u) * f$. We first prove that

$$(4.5) \quad f = (P(D)u) * f = P(D)(u * f) = u * (P(D)f) .$$

Since $\tilde{u} \in \mathcal{S}_{A_p, r}^{M_p, r} \subseteq E'$ ($\tilde{u} \in \mathcal{S}_{A_p, (r_p)}^{M_p, (r_p)} \subseteq E'$) and $f \in \mathcal{D}_E^* \subseteq E$, Proposition 3.19 implies that $u * f \in UC_{\tilde{\omega}} \subseteq \mathcal{S}_{\dagger}^*(\mathbb{R}^d)$, hence $P(D)(u * f)$ is well a defined element of $\mathcal{S}_{\dagger}^*(\mathbb{R}^d)$. Similarly, by Lemma 4.3, $P(D)f \in \mathcal{D}_E^* \subseteq E$, hence Proposition 3.19 implies $u * (P(D)f)$ is well defined element of $\mathcal{S}_{\dagger}^*(\mathbb{R}^d)$. By Proposition 4.2 there exists a net $f_\nu \in \mathcal{S}_{\dagger}^*(\mathbb{R}^d)$ which converges to f in \mathcal{D}_E^* . Then

$$(4.6) \quad f_\nu = \delta * f_\nu = (P(D)u) * f_\nu = P(D)(u * f_\nu) = u * (P(D)f_\nu) .$$

Now, since $f_\nu \rightarrow f$ in \mathcal{D}_E^* the convergence also holds in E and thus Proposition 3.19 implies $u * f_\nu \rightarrow u * f$ in $UC_{\tilde{\omega}}$ and therefore also in $\mathcal{S}_\dagger^*(\mathbb{R}^d)$. Hence $P(D)(u * f_\nu) \rightarrow P(D)(u * f)$ in $\mathcal{S}_\dagger^*(\mathbb{R}^d)$. Next $P(D)f_\nu \rightarrow P(D)f$ in \mathcal{D}_E^* (cf. Lemma 4.3) consequently also in E . Again, Proposition 3.19 implies $u * (P(D)f_\nu) \rightarrow u * (P(D)f)$ in $UC_{\tilde{\omega}}$, hence also in $\mathcal{S}_\dagger^*(\mathbb{R}^d)$. Now after taking limit in (4.6), we obtain (4.5). For $\beta \in \mathbb{N}^d$, since $D^\beta f \in \mathcal{D}_E^*$, (4.5) implies $D^\beta f = u * D^\beta P(D)f$. Since $D^\beta P(D)f \in \mathcal{D}_E^* \subseteq E$, Proposition 3.19 and the discussion preceding it imply that $D^\beta f$ is continuous function and $D^\beta f \in UC_{\tilde{\omega}}$ for each $\beta \in \mathbb{N}^d$. Thus, Lemma 4.6 implies that $f \in C^\infty(\mathbb{R}^d)$. To prove the inclusion $\mathcal{D}_E^* \rightarrow \mathcal{O}_{\dagger,C}^*(\mathbb{R}^d)$, we consider first the (M_p) case. Let $m > 0$ be arbitrary but fixed. Since $P(D) = \sum_\alpha c_\alpha D^\alpha$ is of (M_p) type, there exist $m_1, C' > 0$ such that $|c_\alpha| \leq C' m_1^{|\alpha|} / M_\alpha$. Let $m_2 = 4 \max\{m, m_1\}$. By Lemma 4.3 (and its proof), we have

$$|D^\beta f(x)| \leq \|u\|_{\tilde{E}'} \|D^\beta P(D)f(x)\|_E \omega(-x) \leq C_2 \omega(-x) \|\check{u}\|_{E'} \|f\|_{E, m_2 H} \frac{M_\beta}{(2m)^{|\beta|}}.$$

Hence

$$(4.7) \quad \frac{(2m)^{|\beta|} |D^\beta f(x)|}{M_\beta \omega(-x)} \leq C'' \|\check{u}\|_{E'} \|f\|_{E, m_2 H}.$$

Since there exist $\tau, C''' > 0$ such that $\omega(x) \leq C''' e^{A(\tau|x|)}$, by using [7, Proposition 3.6], we obtain $\omega(-x) e^{A(\tau|x|)} \leq C_4 e^{A(\tau H|x|)}$. Hence

$$\begin{aligned} \left(\sum_\alpha \frac{m^{2|\alpha|}}{M_\alpha^2} \|D^\alpha f e^{-A(\tau H|\cdot|)}\|_{L^2}^2 \right)^{1/2} &\leq C_5 \left(\sum_\alpha \frac{m^{2|\alpha|}}{M_\alpha^2} \left\| \frac{D^\alpha f}{\omega(-\cdot)} \right\|_{L^\infty}^2 \right)^{1/2} \\ &\leq C \|\check{u}\|_{E'} \|f\|_{E, m_2 H}, \end{aligned}$$

which proves the continuity of the inclusion $\mathcal{D}_E^{(M_p)} \rightarrow \mathcal{O}_{(A_p), \tau H}^{(M_p)}$ and hence also the continuity of the inclusion $\mathcal{D}_E^{(M_p)} \rightarrow \mathcal{O}_{(A_p), C}^{(M_p)}$.

In order to prove that the inclusion $\mathcal{D}_E^{\{M_p\}} \rightarrow \mathcal{O}_{\{A_p\}, C}^{\{M_p\}}$ is continuous, it is enough to prove that for each $h > 0$, $\mathcal{D}_E^{\{M_p\}} \rightarrow \mathcal{O}_{\{A_p\}, h}^{\{M_p\}}$ is continuous. And in order to prove this it is enough to prove that for every $m > 0$ there exists $m' > 0$ such that we have the continuous inclusion $\mathcal{D}_E^{M_p, m} \rightarrow \mathcal{O}_{A_p, m', h}^{M_p}$. So, let $h, m > 0$ be arbitrary but fixed. Take $m' \leq m/(4H)$. For $f \in \mathcal{D}_E^{M_p, m}$, keeping notations as above, by Lemma 4.3 (and its proof), we have

$$|D^\beta f(x)| \leq \|\check{u}\|_{E'} \|D^\beta P(D)f(x)\|_E \omega(-x) \leq C_2 \omega(-x) \|\check{u}\|_{E'} \|f\|_{E, m} \frac{M_\beta}{(2m')^{|\beta|}},$$

namely,

$$(4.8) \quad \frac{(2m')^{|\beta|} |D^\beta f(x)|}{M_\beta \omega(-x)} \leq C'' \|\check{u}\|_{E'} \|f\|_{E, m}.$$

For the fixed h take $\tau > 0$ such that $\tau H \leq h$. Then there exists $C''' > 0$ such that $\omega(x) \leq C''' e^{A(\tau|x|)}$ and by using [7, Proposition 3.6] we obtain $\omega(x)e^{A(\tau|x|)} \leq C_4 e^{A(\tau H|x|)}$. Similarly as above, we have

$$\left(\sum_{\alpha} \frac{m'^{2|\alpha|}}{M_{\alpha}^2} \|D^{\alpha} f e^{-A(h|\cdot|)}\|_{L^2}^2 \right)^{1/2} \leq C \|\check{u}\|_{E'} \|f\|_{E,m},$$

which proves the continuity of the inclusion $\mathcal{D}_E^{\{M_p\},m} \rightarrow \mathcal{O}_{A_p,m',h}^{M_p}$.

Observe that (4.4) follows from (4.7) and (4.8), respectively. It remains to prove that $D^{\alpha} f \in C_{\check{\omega}}$. We will prove this in the Roumieu case as the Beurling case is similar. By using Lemma 4.4, with a similar technique as above, one can prove that for every $(k_p) \in \mathfrak{R}$ there exists $(l_p) \in \mathfrak{R}$ such that for $f \in \mathcal{D}_E^{\{M_p\}}$ we have

$$(4.9) \quad \frac{|D^{\beta} f(x)|}{w(-x)M_{\beta} \prod_{j=1}^{|\beta|} k_j} \leq C'' \|\check{u}\|_{E'} \|f\|_{E,(l_p)}.$$

Let $\varepsilon > 0$. Since $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ is dense in $\mathcal{D}_E^{\{M_p\}}$ (cf. Proposition 4.2), it is dense in $\tilde{\mathcal{D}}_E^{\{M_p\}}$. Pick $\chi \in \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ such that $\|f - \chi\|_{E,(l_p)} \leq \varepsilon / (2C'' \|\check{u}\|_{E'})$. Since $1 = \omega(0) \leq \omega(-x)\omega(x)$, by (III) there exist $(l'_p) \in \mathfrak{R}$ and $C_0 > 0$ such that $1/\omega(-x) \leq C_0 e^{B_{l'_p}(|x|)}$. Thus, as $\chi \in \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$, there exists $K \subset\subset \mathbb{R}^d$ such

that $\frac{|D^{\beta} \chi(x)|}{w(-x)M_{\beta} \prod_{j=1}^{|\beta|} k_j} \leq \varepsilon/2$ for all $x \in \mathbb{R}^d \setminus K$ and $\beta \in \mathbb{N}^d$. Then, by (4.9),

for $x \in \mathbb{R}^d \setminus K$ and $\beta \in \mathbb{N}^d$, we have

$$\frac{|D^{\beta} f(x)|}{w(-x)M_{\beta} \prod_{j=1}^{|\beta|} k_j} \leq \frac{|D^{\beta} (f(x) - \chi(x))|}{w(-x)M_{\beta} \prod_{j=1}^{|\beta|} k_j} + \frac{|D^{\beta} \chi(x)|}{w(-x)M_{\beta} \prod_{j=1}^{|\beta|} k_j} \leq \varepsilon,$$

which proves that $D^{\beta} f \in C_{\check{\omega}}$. \square

Remark 4.8. If $f \in \mathcal{S}_{\dagger}^*(\mathbb{R}^d)$, the proof of the previous proposition (combined with Proposition 3.10) yields $\|D^{\beta} f\|_E \leq \|u\|_E \|D^{\beta} P(D)f\|_{1,\omega}$, since $u \in E$. Employing a similar technique as in the proof of Lemma 4.3 (Lemma 4.4), we obtain that for every $m > 0$ there exist $\tilde{m} > 0$ and $C_1 > 0$ (for every $(k_p) \in \mathfrak{R}$ there exist $(l_p) \in \mathfrak{R}$ and $C_1 > 0$) such that

$$(4.10) \quad \|f\|_{E,m} \leq C_1 \sup_{\alpha} \frac{\tilde{m}^{|\alpha|} \|D^{\alpha} f\|_{1,\omega}}{M_{\alpha}} \left(\|f\|_{E,(k_p)} \leq C_1 \sup_{\alpha} \frac{\|D^{\alpha} f\|_{1,\omega}}{M_{\alpha} \prod_{j=1}^{|\alpha|} l_j} \right).$$

4.2. The ultradistribution space $\mathcal{D}'_{E'_*}$. We can now define our new distribution space. We denote by $\mathcal{D}'_{E'_*}$ the strong dual of $\mathcal{D}^*_{E'}$. Then, $\mathcal{D}'_{E'_*}{}^{(M_p)}$ is a complete (DF) -space because $\mathcal{D}_E^{(M_p)}$ is an (F) -space. Also, $\mathcal{D}'_{E'_*}{}^{(M_p)}$ is an (F) -space as the strong dual of a (DF) -space. When E is reflexive, we write $\mathcal{D}'_{E'_*} = \mathcal{D}'_{E'_*}$ in accordance with the last assertion of Theorem 3.17. The notation $\mathcal{D}'_{E'_*} = (\mathcal{D}^*_{E'})'$ is motivated by the next structural theorem which characterizes

the elements of this dual space and bounded sets in two ways, in terms of convolution averages and as the product of ultradifferential operators acting on elements of E'_* .

Theorem 4.9. *Let $B \subseteq \mathcal{S}'_{\dagger}(\mathbb{R}^d)$. The following statements are equivalent:*

- (i) B is a bounded subset of $\mathcal{D}'_{E'_*}$.
- (ii) for each $\psi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$, $\{f * \psi \mid f \in B\}$ is a bounded subset of E' .
- (iii) for each $\psi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$, $\{f * \psi \mid f \in B\}$ is a bounded subset of E'_* .
- (iv) there exist a bounded subset B_1 of E' and an ultradifferential operator $P(D)$ of class $*$ such that each $f \in B$ can be expressed as $f = P(D)g$ with $g \in B_1$.
- (v) there exist $B_2 \subseteq E'_* \cap UC_{\omega}$ which is bounded in E'_* and in UC_{ω} and an ultradifferential operator $P(D)$ of class $*$ such that each $f \in B$ can be expressed as $f = P(D)g$ with $g \in B_2$. Moreover, if E is reflexive, we may choose $B_2 \subseteq E' \cap C_{\omega}$.

Proof. We denote $B_E = \{\varphi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d) \mid \|\varphi\|_E \leq 1\}$.

(i) \Rightarrow (ii). Fix first $\psi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$. By Proposition 3.10 the set $\check{\psi} * B_E = \{\check{\psi} * \varphi \mid \varphi \in B_E\}$ is bounded in \mathcal{D}'_E . As \mathcal{D}'_E is barreled, B is equicontinuous. Hence, $|\langle f * \psi, \varphi \rangle| = |\langle f, \check{\psi} * \varphi \rangle| \leq C_{\psi}, \forall \varphi \in B_E, \forall f \in B$. So, $|\langle f * \psi, \varphi \rangle| \leq C_{\psi} \|\varphi\|_E, \forall \varphi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d), \forall f \in B$. Since $\mathcal{S}'_{\dagger}(\mathbb{R}^d)$ is dense in E , we obtain $\{f * \psi \mid f \in B\}$ is a bounded subset of E' , for each $\psi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$.

We prove (ii) \Rightarrow (iv) and (ii) \Rightarrow (v) simultaneously. Let (ii) hold. For arbitrary but fixed $\psi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ we have $\langle f * \check{\varphi}, \check{\psi} \rangle = \langle f * \psi, \varphi \rangle$. We obtain that the set $\{|\langle f * \check{\varphi}, \check{\psi} \rangle| \mid \varphi \in B_E, f \in B\}$ is bounded in \mathbb{C} , i.e., $\{f * \check{\varphi} \mid \varphi \in B_E, f \in B\}$ is weakly bounded in $\mathcal{S}'_{\dagger}(\mathbb{R}^d)$, hence it is equicontinuous. Moreover, Lemma 2.5 implies that B is bounded in $\mathcal{S}'_{\dagger}(\mathbb{R}^d)$. We continue the proof in the Roumieu case. The Beurling case is similar. For $(t_p) \in \mathfrak{R}$, denote by $X_{(t_p)}$ the closure of $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ in $\mathcal{S}_{A_p, (t_p)}^{M_p, (t_p)}$. The equicontinuity of the set $\{f * \check{\varphi} \mid \varphi \in B_E, f \in B\}$ implies that there exist $(r_p) \in \mathfrak{R}$ and $C > 0$ such that

$$(4.11) \quad |\langle f * \psi, \varphi \rangle| \leq C \sigma_{(r_p)}(\psi), \quad \forall \psi \in \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d), \quad \forall \varphi \in B_E, \quad \forall f \in B.$$

By Lemma 4.5, there exists $(r'_p) \in \mathfrak{R}$ such that $\mathcal{S}_{A_p, (r'_p)}^{M_p, (r'_p)} \subseteq E \cap E'_*$ and the inclusion mappings $\mathcal{S}_{A_p, (r'_p)}^{M_p, (r'_p)} \rightarrow E$ and $\mathcal{S}_{A_p, (r'_p)}^{M_p, (r'_p)} \rightarrow E'_*$ are continuous. Of course, we can take $(r'_p) \leq (r_p)$. Since B is bounded in $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$, Proposition 2.4 implies that there exist $(l_p), (k_p) \in \mathfrak{R}$ with $(l_p) \leq (k_p)$ such that f can be extended to $X_{(k_p)}$, $\mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)} \subseteq X_{(k_p)}$, the convolution is a continuous bilinear mapping from $\mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)} \times \mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)}$ into $X_{(k_p)}$ and there exists $u \in X_{(l_p)}$ and $P(D)$ of class $\{M_p\}$ such that $P(D)u = \delta$ and $f = P(D)(u * \tilde{f})$, where \tilde{f} is the extension of $f \in B$ to $X_{(k_p)}$ and $u * \tilde{f}$ is the transpose of the continuous mapping $\psi \mapsto \check{u} * \psi, \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d) \rightarrow X_{(k_p)}$. We may assume that $(k_p) \leq (r'_p)$.

Let $u_n \in \mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$, $n \in \mathbb{Z}_+$, be such that $u_n \rightarrow u$ in $X_{(l_p)}$. The continuity of the convolution $*$: $\mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)} \times \mathcal{S}_{A_p, (l_p)}^{M_p, (l_p)} \rightarrow X_{(k_p)}$, together with (4.11), implies

$$\left| \langle u * \tilde{f}, \varphi \rangle \right| \leq C', \quad \forall \varphi \in B_E, \forall f \in B,$$

i.e., $\{u * \tilde{f} | f \in B\}$ is a bounded subset of E' . Now, $f = P(D)(u * \tilde{f})$ hence (iv) is proved. Since $\{u * \tilde{f} | f \in B\}$ is bounded in E' , therefore so is it in $\mathcal{S}'_{\{A_p\}}(\mathbb{R}^d)$, Proposition 2.4 again implies that there exist $(l'_p), (k'_p) \in \mathfrak{R}$ with $(l'_p) \leq (k'_p)$ such that $u * \tilde{f}$ can be extended to $X_{(k'_p)}$, $\mathcal{S}_{A_p, (l'_p)}^{M_p, (l'_p)} \subseteq X_{(k'_p)}$, the convolution is continuous bilinear mapping from $\mathcal{S}_{A_p, (l'_p)}^{M_p, (l'_p)} \times \mathcal{S}_{A_p, (l'_p)}^{M_p, (l'_p)}$ into $X_{(k'_p)}$ and there exists $v \in X_{(l'_p)}$ and $P_1(D)$ of class $\{M_p\}$ such that $P_1(D)v = \delta$ and $u * \tilde{f} = P_1(D)(v * (u * \tilde{f}))$, where $v * (u * \tilde{f})$ is the transpose of the continuous mapping $\psi \mapsto \check{v} * \psi$, $\mathcal{S}'_{\{A_p\}}(\mathbb{R}^d) \rightarrow X_{(k'_p)}$. We can suppose that $(k'_p) \leq (l_p)$. Moreover, by Lemma 3.5, there exist $(t_p) \in \mathfrak{R}$ and $C > 0$ such that $\omega(x) \leq Ce^{Bt_p(|x|)}$ and by Lemma 1.1 we can assume that $\prod_{j=1}^{p+q} t_j \leq 2^{p+q} \prod_{j=1}^p t_j \cdot \prod_{j=1}^q t_j$, $\forall p, q \in \mathbb{Z}_+$. Hence by choosing $(k'_p) \leq (t_p/2H)$, it follows $v \in L_\omega^1 \cap L_\omega^1$. Now $f = P(D)(u * \tilde{f}) = P(D)(P_1(D)(v * (u * \tilde{f})))$. But the composition of two ultradifferential operators is again an ultradifferential operator, hence $f = P_2(D)(v * (u * \tilde{f}))$, where $P_2(D) = P(D) \circ P_1(D)$. Since $v \in L_\omega^1 \cap L_\omega^1$ and $\{u * \tilde{f} | f \in B\}$ is a bounded subset of E' , $v * (u * \tilde{f}) \in E'_*$, and Corollary 3.13 implies that $\{v * (u * \tilde{f}) | f \in B\}$ is bounded in E'_* . Furthermore, since $\check{v} \in X_{(l_p)} \subseteq X_{(l_p)} \subseteq \mathcal{S}_{A_p, (r'_p)}^{M_p, (r'_p)} \subseteq E$, Proposition 3.19 implies that $\{v * (u * \tilde{f}) | f \in B\}$ is a bounded subset of UC_ω and if E is reflexive, also in C_ω . Thus (v) also holds.

The implications (iv) \Rightarrow (i), (v) \Rightarrow (i), (iii) \Rightarrow (ii) and (v) \Rightarrow (iii) are obvious. \square

Proposition 4.10. *Let $\mathbf{f} : \mathcal{S}'_{\dagger}(\mathbb{R}^d) \rightarrow \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ be continuous. The following statements are equivalent:*

- i) \mathbf{f} commutes with every translation, i.e., $\langle \mathbf{f}, T_{-h}\varphi \rangle = T_h \langle \mathbf{f}, \varphi \rangle$, for all $h \in \mathbb{R}^d$ and $\varphi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$.
- ii) \mathbf{f} commutes with every convolution, i.e., $\langle \mathbf{f}, \psi * \varphi \rangle = \check{\psi} * \langle \mathbf{f}, \varphi \rangle$, for all $\psi, \varphi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$.
- iii) There exists $f \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ such that $\langle \mathbf{f}, \varphi \rangle = f * \check{\varphi}$ for every $\varphi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$.

Proof. i) \Rightarrow ii). Let $\varphi, \psi \in \mathcal{S}'_{\dagger}(\mathbb{R}^d)$. Then $\tilde{\varphi}(x, y) = \varphi(x - y)\psi(y) \in \mathcal{S}'_{\dagger}(\mathbb{R}^{2d})$. By carefully examining the first part of the proof of Lemma 3.7, one can verify that

$$\mathcal{S}'_{\dagger}(\mathbb{R}^d) \ni L_{\psi, n}(x) = \sum_{t \in D_n} \tilde{\varphi}(x, t) l(n)^{-d} = \sum_{t \in D_n} \varphi(x - t) \psi(t) l(n)^{-d} \rightarrow \psi * \varphi,$$

in $\mathcal{S}_\dagger^*(\mathbb{R}^d)$, where $l(n)$ can be taken to be equal to n (there, the specific definition of $l(n)$ was only needed for the second part of the proof). The continuity of \mathbf{f} implies

$$\begin{aligned} \langle \mathbf{f}, \psi * \varphi \rangle &= \lim_{n \rightarrow \infty} \left\langle \mathbf{f}, \sum_{t \in D_n} \varphi(x-t) \psi(t) n^{-d} \right\rangle \\ &= \lim_{n \rightarrow \infty} \sum_{t \in D_n} \psi(t) \langle \mathbf{f}, T_{-t} \varphi \rangle n^{-d} = \lim_{n \rightarrow \infty} \sum_{t \in D_n} \psi(t) T_t \langle \mathbf{f}, \varphi \rangle n^{-d}, \end{aligned}$$

in $\mathcal{S}'_*(\mathbb{R}^d)$. Let $\chi \in \mathcal{S}'_*(\mathbb{R}^d)$. Then

$$\begin{aligned} \left\langle \lim_{n \rightarrow \infty} \sum_{t \in D_n} \psi(t) T_t \langle \mathbf{f}, \varphi \rangle n^{-d}, \chi \right\rangle &= \left\langle \langle \mathbf{f}, \varphi \rangle, \lim_{n \rightarrow \infty} \sum_{t \in D_n} \psi(t) T_{-t} \chi n^{-d} \right\rangle \\ &= \langle \langle \mathbf{f}, \varphi \rangle, \check{\psi} * \chi \rangle = \langle \check{\psi} * \langle \mathbf{f}, \varphi \rangle, \chi \rangle. \end{aligned}$$

ii) \Rightarrow iii). Take $\chi_n \in \mathcal{S}'_*(\mathbb{R}^d)$, $n \in \mathbb{Z}_+$, to be as in *ii)* of Lemma 2.3. Then, for every $\psi \in \mathcal{S}'_*(\mathbb{R}^d)$ we have that $\psi * \chi_n \rightarrow \psi$ in $\mathcal{S}'_*(\mathbb{R}^d)$ as $n \rightarrow \infty$ and hence,

$$(4.12) \quad \check{\psi} * \langle \mathbf{f}, \chi_n \rangle = \langle \mathbf{f}, \psi * \chi_n \rangle \rightarrow \langle \mathbf{f}, \psi \rangle \text{ as } n \rightarrow \infty.$$

Thus $\{\check{\psi} * \langle \mathbf{f}, \chi_n \rangle \mid n \in \mathbb{Z}_+\}$ is bounded in $\mathcal{S}'_*(\mathbb{R}^d)$. Lemma 2.5 implies that $B = \{\langle \mathbf{f}, \chi_n \rangle \mid n \in \mathbb{Z}_+\}$ is bounded in $\mathcal{S}'_*(\mathbb{R}^d)$. As $\mathcal{S}'_*(\mathbb{R}^d)$ is Montel, its closure \overline{B} is compact and the weak and the strong topologies on \overline{B} coincide. As \overline{B} is equicontinuous and $\mathcal{S}'_*(\mathbb{R}^d)$ is separable, the weak topology on \overline{B} is metrizable (cf. [12, Theorem 4.7, p. 87]) hence also the strong topology. Thus, there exists a subsequence $\langle \mathbf{f}, \chi_{n_k} \rangle \in B$, $k \in \mathbb{Z}_+$, which converges to $f \in \mathcal{S}'_*(\mathbb{R}^d)$. Now, (4.12) implies that $\langle \mathbf{f}, \psi \rangle = \check{\psi} * f$.

The implication *iii) \Rightarrow i)* is clear. \square

If F is a l.c.s., as in [8], we define

$$\mathcal{S}'_*(\mathbb{R}^d, F) = \mathcal{S}'_*(\mathbb{R}^d) \varepsilon F = \mathcal{L}_\epsilon \left((\mathcal{S}'_*(\mathbb{R}^d))'_c, F \right) = \mathcal{L}_b(\mathcal{S}'_*(\mathbb{R}^d), F),$$

where the indices ϵ and c stand for the topology of equicontinuous convergence and the topology of compact convex circled convergence, respectively; the last equality follows from the fact that $\mathcal{S}'_*(\mathbb{R}^d)$ and $\mathcal{S}'_*(\mathbb{R}^d)$ are complete Montel spaces. If F is complete, since $\mathcal{S}'_*(\mathbb{R}^d)$ is nuclear, it satisfies the weak approximation property, hence $\mathcal{S}'_*(\mathbb{R}^d) \varepsilon F \cong \mathcal{S}'_*(\mathbb{R}^d) \hat{\otimes} F$, cf. [8, Proposition 1.4] (for the definition of the ε tensor product; for the definition of the weak approximation property and their connection we refer to [13] and [8]).

Corollary 4.11. *Let $\mathbf{f} \in \mathcal{S}'_*(\mathbb{R}^d, E'_{\sigma(E', E)})$. If \mathbf{f} commutes with every translation in sense of Proposition 4.10, then there exists $f \in \mathcal{D}'_{E'_*}$ such that \mathbf{f} is of the form*

$$(4.13) \quad \langle \mathbf{f}, \varphi \rangle = f * \check{\varphi}, \quad \varphi \in \mathcal{S}'_*(\mathbb{R}^d).$$

Proof. The proof is analogous to the proof of [4, Corollary 6.4]. \square

Our results from above implicitly suggest to embed the ultradistribution space $\mathcal{D}'_{E'_*}$ into the space of E' -valued ultradistributions as follows. Define first the continuous injection $\iota : \mathcal{S}'_{\dagger}(\mathbb{R}^d) \rightarrow \mathcal{S}'_{\dagger}(\mathbb{R}^d, \mathcal{S}'_{\dagger}(\mathbb{R}^d))$, where $\iota(f) = \mathbf{f}$ is given by (4.13). Consider the restriction of ι to $\mathcal{D}'_{E'_*}$,

$$(4.14) \quad \iota : \mathcal{D}'_{E'_*} \rightarrow \mathcal{S}'_{\dagger}(\mathbb{R}^d, E'_*),$$

(for $f \in \mathcal{D}'_{E'_*}$, the range of $\iota(f)$ is subset of E'_* by Theorem 4.9). Let B_1 be an arbitrary bounded subset of $\mathcal{S}'_{\dagger}(\mathbb{R}^d)$. The set $B = \{\psi * \varphi \mid \varphi \in B_1, \|\psi\|_E \leq 1\}$ is bounded in $\mathcal{D}'_{E'_*}$ (by Lemma 3.8). For $f \in \mathcal{D}'_{E'_*}$,

$$\sup_{\varphi \in B_1} \|\langle \mathbf{f}, \varphi \rangle\|_{E'} = \sup_{\varphi \in B_1} \|f * \check{\varphi}\|_{E'} = \sup_{\varphi \in B_1} \sup_{\|\psi\|_E \leq 1} |\langle f, \psi * \varphi \rangle| = \sup_{\chi \in B} |\langle f, \chi \rangle|.$$

Hence, $\iota(f) \in \mathcal{S}'_{\dagger}(\mathbb{R}^d, E'_*)$ ($\mathcal{S}'_{\dagger}(\mathbb{R}^d)$ is bornological) and ι is continuous. Furthermore, Proposition 4.10 tells us that $\iota(\mathcal{D}'_{E'_*})$ is precisely the subspace of $\mathcal{S}'_{\dagger}(\mathbb{R}^d, E'_*)$ consisting of those \mathbf{f} which commute with all translations in the sense of Proposition 4.10. Since the translations T_h are continuous operators on E'_* , we actually obtain that the range $\iota(\mathcal{D}'_{E'_*})$ is a closed subspace of $\mathcal{S}'_{\dagger}(\mathbb{R}^d, E'_*)$.

Corollary 4.12. *For $B \subseteq \mathcal{S}'_{\dagger}(\mathbb{R}^d)$ the equivalent conditions from Theorem 4.9 are equivalent to the following:*

(vi) $\iota(B)$ is a bounded subset of $\mathcal{S}'_{\dagger}(\mathbb{R}^d, E'_*)$ (or equivalently of $\mathcal{S}'_{\dagger}(\mathbb{R}^d, E'_*)$)

Proof. (i) \Rightarrow (vi) and (vi) \Rightarrow (ii) are trivial. \square

Corollary 4.13. *Let $\{f_{\lambda}\}_{\lambda \in \Lambda} \subseteq \mathcal{D}'_{E'_*}$ be a bounded net (or similarly, a sequence). The following statements are equivalent:*

- (i) $\{f_{\lambda}\}_{\lambda \in \Lambda}$ is convergent in $\mathcal{D}'_{E'_*}$.
- (ii) $\{\iota(f_{\lambda})\}_{\lambda \in \Lambda}$ is convergent in $\mathcal{S}'_{\dagger}(\mathbb{R}^d, E'_*)$ (or equivalently in $\mathcal{S}'_{\dagger}(\mathbb{R}^d, E'_*)$).
- (iii) There exists a convergent bounded net $\{g_{\lambda}\}_{\lambda \in \Lambda}$ in E' and an ultradifferential operator $P(D)$ of class $*$ such that each $f_{\lambda} = P(D)g_{\lambda}$.
- (iv) There exists a net $\{g_{\lambda}\}_{\lambda \in \Lambda} \subseteq E'_* \cap UC_{\omega}$ which is convergent and bounded in E'_* and in UC_{ω} and an ultradifferential operator $P(D)$ of class $*$ such that $f_{\lambda} = P(D)g_{\lambda}$; if E is reflexive one may choose $\{g_{\lambda}\}_{\lambda \in \Lambda} \subseteq E' \cap C_{\omega}$.

Proof. We consider the Roumieu case as the Beurling case is similar. Let (ii) hold. Since the image of $\mathcal{D}'_{E'_*}^{\{M_p\}}$ under ι is closed subspace of $\mathcal{S}'_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d, E'_*)$, $\iota(f_{\lambda}) \rightarrow \iota(f)$, for $f \in \mathcal{D}'_{E'_*}^{\{M_p\}}$. As $B = \{\iota(f)\} \cup \{\iota(f_{\lambda}) \mid \lambda \in \Lambda\}$ is bounded in $\mathcal{L}_b(\mathcal{S}'_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d), E'_*)$, it is equicontinuous ($\mathcal{S}'_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ is barreled) and thus, there exists $(r_p) \in \mathfrak{R}$ such that the elements of B can be extended to a bounded subset $\tilde{B} = \{\iota(\widetilde{f})\} \cup \{\iota(\widetilde{f_{\lambda}}) \mid \lambda \in \Lambda\}$ of $\mathcal{L}_b(X_{(r_p)}, E'_*)$. Moreover, $\iota(\widetilde{f_{\lambda}})(\varphi) \rightarrow \iota(\widetilde{f})(\varphi)$ for each φ in the dense subset $\mathcal{S}'_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ of $X_{(r_p)}$. Since

\widetilde{B} is bounded in $\mathcal{L}_b(X_{(r_p)}, E'_*)$, $\widetilde{\iota}(f_\lambda) \rightarrow \widetilde{\iota}(f)$ in $\mathcal{L}_\sigma(X_{(r_p)}, E'_*)$, the Banach-Steinhaus theorem implies that it is also bounded in $\mathcal{L}_p(X_{(r_p)}, E'_*)$. Pick now, $(r'_p) \in \mathfrak{R}$ with $(r'_p) \leq (r_p)$ such that the inclusion $X_{(r'_p)} \rightarrow X_{(r_p)}$ is compact. Then the inclusion $\mathcal{L}_p(X_{(r_p)}, E'_*) \rightarrow \mathcal{L}_b(X_{(r'_p)}, E'_*)$ is continuous. Thus $\widetilde{\iota}(f_\lambda) \rightarrow \widetilde{\iota}(f)$ in $\mathcal{L}_b(X_{(r'_p)}, E'_*)$. Now one can use a similar technique as in the proof of (ii) \Rightarrow (iv) of Theorem 4.9 to conclude (iii) and similar technique as in the proof of (ii) \Rightarrow (v) of Theorem 4.9 to conclude (iv). The implications (i) \Rightarrow (ii), (iii) \Rightarrow (i) and (iv) \Rightarrow (i) are obvious. \square

This corollary implies that the restriction of ι on each bounded subset B of $\mathcal{D}_{E'_*}^*$ is topological homeomorphism between B and $\iota(B)$.

For the proof of the following two results we refer to [4, Proposition 6.7, Proposition 6.8]

Theorem 4.14. *The spaces $\mathcal{D}_E^{\{M_p\}}$ and $\widetilde{\mathcal{D}}_E^{\{M_p\}}$ are isomorphic as l.c.s..*

Proposition 4.15. *If E is reflexive, then $\mathcal{D}_E^{(M_p)}$ and $\mathcal{D}_{E'}^{\{M_p\}}$ are (FS^*) -spaces, $\mathcal{D}_E^{\{M_p\}}$ and $\mathcal{D}_E^{(M_p)}$ are (DFS^*) -spaces. Consequently, they are reflexive. In addition, $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is dense in $\mathcal{D}_{E'}^*$.*

4.3. Weighted $\mathcal{D}_{L_\eta^p}^*$ spaces. In this subsection we discuss some important examples of the spaces \mathcal{D}_E^* and $\mathcal{D}_{E'}^*$, where E is taken as a weighted L_η^p spaces. In the next considerations we retain the notation exactly as in Example 3.20. In particular, η is ultrapolynomially bounded weight of class \dagger and the number q always stands for $p^{-1} + q^{-1} = 1$ ($p \in [1, \infty]$). It should be mentioned that in the case $\eta = 1$ and $M_p = A_p$ satisfying (M.3) the spaces we study below were considered in [9] (see also [1]). The non-quasianalytic case with $A_p = M_p$ for general weights η was studied in detail in [4].

Consider now the spaces $\mathcal{D}_{L_\eta^p}^*$ for $p \in [1, \infty]$ and $\widetilde{\mathcal{D}}_{L_\eta^\infty}^{\{M_p\}}$ defined as in Section 4 by taking $E = L_\eta^p$. We also treat \mathcal{D}_{C_η} defined via $E = C_\eta$. Once again, the case $p = \infty$ is an exception since $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is not dense in $\mathcal{D}_{L_\eta^\infty}^*$ nor in $\widetilde{\mathcal{D}}_{L_\eta^\infty}^{\{M_p\}}$. Nonetheless, we can repeat the proof of Lemma 4.1 to prove that $\mathcal{D}_{L_\eta^\infty}^{\{M_p\}}$ is regular and complete. Also, similarly as in Lemmas 4.3 and 4.4, one can obtain that each ultradifferential operator of $*$ class acts continuously on $\mathcal{D}_{L_\eta^\infty}^*$ and each ultradifferential operator of $\{M_p\}$ class acts continuously on $\widetilde{\mathcal{D}}_{L_\eta^\infty}^{\{M_p\}}$. Obviously $\mathcal{D}_{L_\eta^\infty}^{\{M_p\}}$ is continuously injected into $\widetilde{\mathcal{D}}_{L_\eta^\infty}^{\{M_p\}}$ and by using [8, Lemma 3.4] and employing a similar technique as in the proof of Lemma 4.1, one can prove that this inclusion is in fact surjective. We denote by \mathcal{B}_η^* the space $\mathcal{D}_{L_\eta^\infty}^*$ and by $\dot{\mathcal{B}}_\eta^*$ the closure of $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ in \mathcal{B}_η^* . We denote by $\dot{\mathcal{B}}_\eta^{\{M_p\}}$ the closure of $\mathcal{S}_{\{A_p\}}^{\{M_p\}}(\mathbb{R}^d)$ in $\widetilde{\mathcal{D}}_{L_\eta^\infty}^{\{M_p\}}$. We immediately see that $\dot{\mathcal{B}}_\eta^{\{M_p\}} = \mathcal{D}_{C_\eta}^{(M_p)}$. In the Roumieu case this is result is given by the following theorem. Its proof is analogous to that of [4, Theorem 7.2] and we omit it.

Theorem 4.16. *The spaces $\mathcal{D}_{C_\eta}^{\{M_p\}}$, $\dot{\mathcal{B}}_\eta^{\{M_p\}}$ and $\check{\mathcal{B}}_\eta^{\{M_p\}}$ are isomorphic one to another as l.c.s..*

Proposition 4.7 together with the estimate (4.7) (resp. Proposition 4.7 together with (4.8)) imply $\mathcal{D}_{L_\eta^p}^* \hookrightarrow \dot{\mathcal{B}}_{\omega_\eta}^*$ for every $p \in [1, \infty)$. It follows from Proposition 4.15 that $\mathcal{D}_{L_\eta^p}^*$ is reflexive when $p \in (1, \infty)$.

In accordance to Subsection 4.2, the weighted spaces $\mathcal{D}_{L_\eta^p}^{\prime*}$ are defined as $\mathcal{D}_{L_\eta^p}^{\prime*} = (\mathcal{D}_{L_{\eta^{-1}}^q}^*)'$ where $p^{-1} + q^{-1} = 1$ if $p \in (1, \infty)$; if $p = 1$, $\mathcal{D}_{L_\eta^1}^{\prime*} = (\mathcal{D}_{C_\eta}^*)' = (\dot{\mathcal{B}}_\eta^*)'$ and for $p = \infty$ we define $\mathcal{D}_{L_\eta^\infty}^{\prime*} := \mathcal{D}_{UC_\eta}^{\prime*} = (\mathcal{D}_{L_\eta^1}^*)'$. We write $\mathcal{B}_\eta^{\prime*} = \mathcal{D}_{L_\eta^\infty}^{\prime*}$ and $\dot{\mathcal{B}}_\eta^{\prime*}$ for the closure of $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ in $\mathcal{B}_\eta^{\prime*}$.

For $f \in \mathcal{D}_{L_\eta^1}^{\prime*}$, by Theorem 4.9, there exist an ultradifferential operator $P(D)$ of class $*$ and $g \in L_\eta^1$ such that $f = P(D)g$. But, since $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is dense in L_η^1 , there exists a sequence $g_n \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$, $n \in \mathbb{Z}_+$, such that $g_n \rightarrow g$ in L_η^1 . Hence $\mathcal{S}_\dagger^*(\mathbb{R}^d) \ni f_n = P(D)g_n \rightarrow f$ in $\mathcal{D}_{L_\eta^1}^{\prime*}$, i.e., $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is sequentially dense in $\mathcal{D}_{L_\eta^1}^{\prime*}$. Moreover, as an easy consequence of the Sobolev imbedding theorem, we obtain that $\mathcal{D}_{L_\eta^p}^*$ is continuously injected into $C^\infty(\mathbb{R}^d)$ for each $p \in [1, \infty)$. Since $\mathcal{S}_\dagger^*(\mathbb{R}^d) \hookrightarrow \mathcal{S}(\mathbb{R}^d) \hookrightarrow C^\infty(\mathbb{R}^d)$, $\mathcal{D}_{L_\eta^p}^*$ is dense in $C^\infty(\mathbb{R}^d)$, hence $\mathcal{E}'(\mathbb{R}^d)$ is continuously injected into $\mathcal{D}_{L_{1/\eta}^q}^{\prime*}$. In particular the delta (ultra)distribution belongs to $\mathcal{D}_{L_{1/\eta}^q}^{\prime*}$.

Theorem 4.17. *The strong bidual of $\dot{\mathcal{B}}_\eta^*$ is isomorphic to $\mathcal{D}_{L_\eta^\infty}^*$ as l.c.s.. In the Roumieu case $\mathcal{D}_{L_\eta^\infty}^{\{M_p\}}$ and $\tilde{\mathcal{D}}_{L_\eta^\infty}^{\{M_p\}}$ are isomorphic l.c.s.. Moreover $\dot{\mathcal{B}}_\eta^{\{M_p\}}$ is a distinguished (F)-space and consequently $\mathcal{D}_{L_\eta^1}^{\prime(M_p)}$ is barreled and bornological.*

Proof. We may assume that η is continuous (cf. Example 3.20). Let $\tilde{\eta}(x) = 1/(\eta(x)\langle x \rangle^{d+1})$. Then, clearly $\tilde{\eta}(x)$ is a continuous ultrapolynomially bounded weight of class \dagger and $\dot{\mathcal{B}}_\eta^* \hookrightarrow \mathcal{D}_{L_\eta^2}^*$. Since $\mathcal{S}_\dagger^*(\mathbb{R}^d)$ is dense in $\mathcal{D}_{L_\eta^1}^*$, we have $\mathcal{D}_{L_{1/\tilde{\eta}}^2}^{\prime*} \hookrightarrow \mathcal{D}_{L_\eta^1}^*$. This, together with Proposition 4.15, implies that $(\mathcal{D}_{L_\eta^1}^*)'_b$ (where b stands for the strong topology) is continuously injected into $\mathcal{D}_{L_\eta^2}^*$, i.e., the elements of $(\mathcal{D}_{L_\eta^1}^*)'_b$ are smooth functions. In the Roumieu case, we already saw that $\mathcal{D}_{L_\eta^\infty}^{\{M_p\}}$ and $\tilde{\mathcal{D}}_{L_\eta^\infty}^{\{M_p\}}$ are equal as sets. First we prove that the bidual of $\dot{\mathcal{B}}_\eta^*$ is isomorphic to $\mathcal{D}_{L_\eta^\infty}^{\{M_p\}}$, and to $\tilde{\mathcal{D}}_{L_\eta^\infty}^{\{M_p\}}$ respectively. Let $r > 0$ (let $(r_p) \in \mathfrak{R}$ and set $R_\alpha = \prod_{j=1}^{|\alpha|} r_j$). Consider the set

$$B_r = \left\{ \frac{(\eta(a))^{-1} r^{|\alpha|} D^\alpha \delta_a}{M_\alpha} \mid a \in \mathbb{R}^d, \alpha \in \mathbb{N}^d \right\}$$

$$\left(\text{resp. } B_{(r_p)} = \left\{ \frac{(\eta(a))^{-1} D^\alpha \delta_a}{M_\alpha R_\alpha} \mid a \in \mathbb{R}^d, \alpha \in \mathbb{N}^d \right\} \right).$$

One easily proves that B_r is a bounded subset of $\mathcal{D}'_{L_\eta^1(M_p)}$ ($B_{(r_p)}$ is a bounded subset of $\mathcal{D}'_{L_\eta^1\{M_p\}}$). Hence if $\psi \in \left(\mathcal{D}'_{L_\eta^1}\right)'_b$, $\psi(B_r)$ (resp. $\psi(B_{(r_p)})$) is bounded in \mathbb{C} and thus

$$\sup_{a,\alpha} \frac{|(\eta(a))^{-1}r^{|\alpha|}D^\alpha\psi(a)|}{M_\alpha} = \sup_{f \in B_r} |\langle \psi, f \rangle| < \infty$$

$$\left(\text{resp. } \sup_{a,\alpha} \frac{|(\eta(a))^{-1}D^\alpha\psi(a)|}{M_\alpha R_\alpha} = \sup_{f \in B_{(r_p)}} |\langle \psi, f \rangle| < \infty \right).$$

We obtain that $\left(\mathcal{D}'_{L_\eta^1}\right)' \subseteq \mathcal{D}'_{L_\eta^\infty}$ and the inclusion $\left(\mathcal{D}'_{L_\eta^1(M_p)}\right)' \rightarrow \mathcal{D}'_{L_\eta^\infty(M_p)}$, $\left(\left(\mathcal{D}'_{L_\eta^1\{M_p\}}\right)'\right)' \rightarrow \tilde{\mathcal{D}}'_{L_\eta^\infty\{M_p\}}$ is continuous.

Let $\psi \in \mathcal{D}'_{L_\eta^\infty}$. If $f \in \mathcal{D}'_{L_\eta^1}$, by Theorem 4.9 there exist an ultradifferential operator $P(D)$ of class $*$ and $g \in L_\eta^1$ such that $f = P(D)g$. Define S_ψ by

$$S_\psi(f) = \int_{\mathbb{R}^d} g(x)P(-D)\psi(x)dx.$$

Obviously, the integral on the right hand side is absolutely convergent. We will prove that S_ψ is well defined element of $\left(\mathcal{D}'_{L_\eta^1}\right)'$. Let $\tilde{P}(D), \tilde{g} \in L_\eta^1$ be such that $f = \tilde{P}(D)\tilde{g}$. Let $\varphi_n \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$, $n \in \mathbb{Z}_+$, be as in *ii*) of Lemma 2.3. Then it is easy to verify that

$$\int_{\mathbb{R}^d} P(-D)(\varphi_n(x)\psi(x))g(x)dx \rightarrow \int_{\mathbb{R}^d} P(-D)\psi(x)g(x)dx,$$

$$\int_{\mathbb{R}^d} \tilde{P}(-D)(\varphi_n(x)\psi(x))\tilde{g}(x)dx \rightarrow \int_{\mathbb{R}^d} \tilde{P}(-D)\psi(x)\tilde{g}(x)dx,$$

as $n \rightarrow \infty$. Also, observe that for each $n \in \mathbb{Z}_+$, $\varphi_n\psi \in \mathcal{S}_\dagger^*(\mathbb{R}^d)$ and thus

$$\int_{\mathbb{R}^d} P(-D)(\varphi_n(x)\psi(x))g(x)dx = \mathcal{S}_\dagger^*\langle f, \varphi_n\psi \rangle \mathcal{S}_\dagger^*$$

$$= \int_{\mathbb{R}^d} \tilde{P}(-D)(\varphi_n(x)\psi(x))\tilde{g}(x)dx.$$

Hence, S_ψ is a well defined mapping $\mathcal{D}'_{L_\eta^1\{M_p\}} \rightarrow \mathbb{C}$, since it does not depend on the representation of f . To prove that it is continuous we consider first the Beurling case. The space $\mathcal{D}'_{L_\eta^1(M_p)}$ is a complete (DF) -space. Thus it is enough to prove that the restriction of S_ψ on each bounded subset of $\mathcal{D}'_{L_\eta^1(M_p)}$ is continuous (see the corollary to [12, Theorem 6.7, p. 154]), i.e., we have to prove that if $\{f_\lambda\}_{\lambda \in \Lambda}$ is bounded net which converges to f in $\mathcal{D}'_{L_\eta^1(M_p)}$, then $S_\psi(f_\lambda) \rightarrow S_\psi(f)$. If $\{f_\lambda\}_{\lambda \in \Lambda}$ is such net, Corollary 4.13 implies that there exists a net $\{g_\lambda\}_{\lambda \in \Lambda} \subseteq L_\eta^1$ which is bounded and convergent in L_η^1 and an ultradifferential operator $P(D)$ of class (M_p) such that $f_\lambda = P(D)g_\lambda$ and

$f = P(D)g$ where $g \in L_\eta^1$ is the limit of $\{g_\lambda\}_{\lambda \in \Lambda}$. But then one easily verifies that $g_\lambda P(-D)\psi \rightarrow gP(-D)\psi$ in L^1 , hence $S_\psi(f_\lambda) \rightarrow S_\psi(f)$. Thus $S_\psi \in \left(\mathcal{D}'_{L_\eta^1}\right)'$. In the Roumieu case, as $\mathcal{D}'_{L_\eta^1}\{M_p\}$ is an (F) -space one can similarly prove that $S_\psi \in \left(\mathcal{D}'_{L_\eta^1}\{M_p\}\right)'$. We obtain that $\left(\mathcal{D}'_{L_\eta^1}\{M_p\}\right)' = \mathcal{D}_{L_\eta^\infty}\{M_p\}$ (resp. $\left(\mathcal{D}'_{L_\eta^1}\{M_p\}\right)' = \tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$) as sets and $\left(\mathcal{D}'_{L_\eta^1}\{M_p\}\right)'$ has stronger topology than the latter. In the Beurling case, $\left(\mathcal{D}'_{L_\eta^1}\{M_p\}\right)'_b$ is an (F) -space as the strong dual of the (DF) -space $\mathcal{D}'_{L_\eta^1}\{M_p\}$. Hence the open mapping theorem proves that $\left(\mathcal{D}'_{L_\eta^1}\{M_p\}\right)'_b = \mathcal{D}_{L_\eta^\infty}\{M_p\}$ as l.c.s.. In the Roumieu case, let $V = B^\circ$ be a neighborhood of zero $\left(\mathcal{D}'_{L_\eta^1}\{M_p\}\right)'_b$ for B a bounded subset of $\mathcal{D}'_{L_\eta^1}\{M_p\}$. By Theorem 4.9, there exist an ultradifferential operator $P(D)$ of class $\{M_p\}$ and bounded subset B_1 of L_η^1 such that each $f \in B$ can be represented by $f = P(D)g$ for some $g \in B_1$. There exists $C_1 \geq 1$ such that $\|g\|_{L_\eta^1} \leq C_1$ for all $g \in B_1$. Also, since $P(D) = \sum_\alpha c_\alpha D^\alpha$ is of $\{M_p\}$ class, there exist $(r_p) \in \mathfrak{R}$ and $C_2 \geq 1$ such that $|c_\alpha| \leq C_2/(M_\alpha R_\alpha)$ (see the proof of Lemma 4.4). Observe the neighborhood of zero $W = \left\{ \psi \in \tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\} \mid \sup_{x, \alpha} \frac{(\eta(x))^{-1} |D^\alpha \psi(x)|}{M_\alpha \prod_{j=1}^{|\alpha|} (r_j/2)} \leq \frac{1}{2C_1 C_2 C_3} \right\}$ in $\tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$, where we put $C_3 = \sum_\alpha 2^{-|\alpha|}$. One easily verifies that $W \subseteq V$. We obtain that $\left(\mathcal{D}'_{L_\eta^1}\{M_p\}\right)'_b$ and $\tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$ are isomorphic l.c.s.. Hence $\tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$ is a complete (DF) -space (since $\mathcal{D}'_{L_\eta^1}\{M_p\}$ is an (F) -space). As the identity mapping $\mathcal{D}'_{L_\eta^1}\{M_p\} \rightarrow \tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$ is continuous and bijective, it remains to prove that the inverse is continuous. Since $\tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$ is a (DF) -space, to prove the continuity of the inverse mapping it is enough to prove that its restriction to every bounded subset of $\tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$ is continuous (see the corollary to [12, Theorem 6.7, p. 154]). If B is a bounded subset of $\tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$ then for every $(r_p) \in \mathfrak{R}$, $\sup_{\psi \in B} \sup_\alpha \frac{\|D^\alpha \psi\|_{L_\eta^\infty(\mathbb{R}^d)}}{M_\alpha R_\alpha} < \infty$. Hence, by [8, Lemma 3.4], there exists $h > 0$ such that $\sup_{\psi \in B} \sup_\alpha \frac{h^{|\alpha|} \|D^\alpha \psi\|_{L_\eta^\infty(\mathbb{R}^d)}}{M_\alpha} < \infty$, i.e., B is bounded in $\mathcal{D}_{L_\eta^\infty}\{M_p\}$. Since every bounded subset of $\mathcal{D}_{L_\eta^\infty}\{M_p\}$ is obviously bounded in $\tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$, $\mathcal{D}_{L_\eta^\infty}\{M_p\}$ and $\tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$ have the same bounded sets. Let ψ_λ be a bounded net in $\tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$ which converges to ψ in $\tilde{\mathcal{D}}_{L_\eta^\infty}\{M_p\}$. Then there exist $0 < h \leq 1$ and $C > 0$ such that

$$\sup_\lambda \sup_\alpha \frac{h^{|\alpha|} \|D^\alpha \psi_\lambda\|_{L_\eta^\infty}}{M_\alpha} \leq C \text{ and } \sup_\alpha \frac{h^{|\alpha|} \|D^\alpha \psi\|_{L_\eta^\infty}}{M_\alpha} \leq C.$$

Choose $0 < h_1 < h$. Let $\varepsilon > 0$ be arbitrary but fixed. Take $p_0 \in \mathbb{Z}_+$ such that $(h_1/h)^{|\alpha|} \leq \varepsilon/(2C)$ for all $|\alpha| \geq p_0$. Since $\psi_\lambda \rightarrow \psi$ in $\tilde{\mathcal{D}}_{L_\infty^\eta}^{\{M_p\}}$, for the sequence $r_p = p$, $p \in \mathbb{Z}_+$, there exists λ_0 such that for all $\lambda \geq \lambda_0$ we have

$$\sup_\alpha \frac{\|D^\alpha(\psi_\lambda - \psi)\|_{L_\infty^\eta}}{M_\alpha R_\alpha} \leq \frac{\varepsilon}{p_0!}. \text{ Then for } |\alpha| < p_0, \text{ we have } \frac{h_1^{|\alpha|} \|D^\alpha(\psi_\lambda - \psi)\|_{L_\infty^\eta}}{M_\alpha} \leq \varepsilon.$$

For $|\alpha| \geq p_0$, we have

$$\frac{h_1^{|\alpha|} \|D^\alpha(\psi_\lambda - \psi)\|_{L_\infty^\eta}}{M_\alpha} \leq 2C \left(\frac{h_1}{h}\right)^{|\alpha|} \leq \varepsilon.$$

It follows that $\psi_\lambda \rightarrow \psi$ in $\mathcal{D}_{L_\infty^\eta}^{\{M_p\}, h_1}$ and hence in $\mathcal{D}_{L_\infty^\eta}^{\{M_p\}}$. We obtain that the induced topology by $\tilde{\mathcal{D}}_{L_\infty^\eta}^{\{M_p\}}$ on every bounded subset of $\tilde{\mathcal{D}}_{L_\infty^\eta}^{\{M_p\}}$ is stronger than the induced topology by $\mathcal{D}_{L_\infty^\eta}^{\{M_p\}}$. Hence the identity mapping $\tilde{\mathcal{D}}_{L_\infty^\eta}^{\{M_p\}} \rightarrow \mathcal{D}_{L_\infty^\eta}^{\{M_p\}}$ is continuous.

It remains to prove that $\dot{\mathcal{B}}_\eta^{(M_p)}$ is distinguished. Denote by $\mathcal{D}_{L_\infty^\eta, \sigma}^{(M_p)}$ the space $\mathcal{D}_{L_\infty^\eta}^{(M_p)}$ equipped with the weak topology from the duality $\langle \mathcal{D}'_{L_\eta^1}, \mathcal{D}_{L_\infty^\eta}^{(M_p)} \rangle$. We have to prove that each bounded subset of $\mathcal{D}_{L_\infty^\eta}^{(M_p)}$ (the strong bidual of $\dot{\mathcal{B}}_\eta^{(M_p)}$) is contained in the closure of a bounded subset of $\dot{\mathcal{B}}_\eta^{(M_p)}$ in $\mathcal{D}_{L_\infty^\eta, \sigma}^{(M_p)}$. Let B be a bounded subset of $\mathcal{D}_{L_\infty^\eta}^{(M_p)}$. Let $\varphi_n \in \mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d)$, $n \in \mathbb{Z}_+$, be the sequence from *ii*) of Lemma 2.3. Then, $\varphi_n \psi \in \mathcal{S}_{(A_p)}^{(M_p)}(\mathbb{R}^d)$ for each $n \in \mathbb{Z}_+$, $\psi \in B$. For $r > 0$ one easily verifies that $\|\varphi_n \psi\|_{L_\infty^\eta, r} \leq \|\varphi\|_{L_\infty^\eta, 2r} \|\psi\|_{L_\infty^\eta, 2r}$. Hence the set $\tilde{B} = \{\varphi_n \psi \mid n \in \mathbb{Z}_+, \psi \in B\}$ is bounded subset of $\dot{\mathcal{B}}_\eta^{(M_p)}$. Let $\psi \in B$ and $f \in \mathcal{D}'_{L_\eta^1}$. By Theorem 4.9, there exist an ultradifferential operator $P(D)$ of class (M_p) and $g \in L_\eta^1$ such that $f = P(D)g$. Then one easily verifies that $gP(-D)(\varphi_n \psi) \rightarrow gP(-D)\psi$ in L^1 , thus $\langle \varphi_n \psi, f \rangle \rightarrow \langle \psi, f \rangle$, i.e., $\varphi_n \psi \rightarrow \psi$ in $\mathcal{D}_{L_\infty^\eta, \sigma}^{(M_p)}$, which proves that B belongs in the closure of \tilde{B} in $\mathcal{D}_{L_\infty^\eta, \sigma}^{(M_p)}$. \square

4.4. Convolution and multiplication. Our previous work allows us to extend all results on convolution and multiplicative products on \mathcal{D}'_{E^*} from [2, 4] to our spaces. We omit the proofs of the following propositions because they go in the same lines as those of [2, Theorem 4 and Proposition 11] (adapting them with the aid of our results from the previous subsections).

Proposition 4.18. *We have the (continuous) inclusions $\mathcal{D}_{L_\omega^1}^* \hookrightarrow \mathcal{D}_E^* \hookrightarrow \dot{\mathcal{B}}_\omega^*$ and $\mathcal{D}'_{L_\omega^1} \rightarrow \mathcal{D}'_{E^*} \rightarrow \mathcal{B}_\omega^*$. If E is reflexive, one has $\mathcal{D}'_{L_\omega^1} \hookrightarrow \mathcal{D}'_{E^*} \hookrightarrow \dot{\mathcal{B}}_\omega^*$.*

In particular, we have $\mathcal{D}_{L_{\omega_\eta}^1}^* \hookrightarrow \mathcal{D}_{L_\eta^p}^* \hookrightarrow \dot{\mathcal{B}}_{\omega_\eta}^*$ and $\mathcal{D}'_{L_{\omega_\eta}^1} \hookrightarrow \mathcal{D}'_{L_\eta^p} \hookrightarrow \dot{\mathcal{B}}_{\omega_\eta}^*$ for $1 \leq p < \infty$ (for $p = 1$ in the latter dense inclusion we have used the fact $\mathcal{S}_\dagger^*(\mathbb{R}^d) \hookrightarrow \mathcal{D}'_{L_\eta^1}$). In addition, $\dot{\mathcal{B}}_\eta^* \hookrightarrow \dot{\mathcal{B}}_{\omega_\eta}^*$ and $\dot{\mathcal{B}}_\eta^* \hookrightarrow \dot{\mathcal{B}}_{\omega_\eta}^*$. We can now define multiplicative and convolution operations on \mathcal{D}'_{E^*} . In the next proposition we

denote by $\mathcal{O}'_{\dagger,C,b}$ the space $\mathcal{O}'_{\dagger,C}$ equipped with the strong topology from the duality $\langle \mathcal{O}'_{\dagger,C}, \mathcal{O}'_{\dagger,C} \rangle$.

Proposition 4.19. *The convolution mappings $*$: $\mathcal{D}'_{E'_*} \times \mathcal{D}'_{L^1_\omega} \rightarrow \mathcal{D}'_{E'_*}$ and $*$: $\mathcal{D}'_{E'_*} \times \mathcal{O}'_{\dagger,C,b}(\mathbb{R}^d) \rightarrow \mathcal{D}'_{E'_*}$ are continuous. The convolution and multiplicative products are hypocontinuous in the following cases: \cdot : $\mathcal{D}'_{E'_*} \times \mathcal{D}'_{L^1_\omega} \rightarrow \mathcal{D}'_{L^1}$, \cdot : $\mathcal{D}'_{L^1_\omega} \times \mathcal{D}^*_E \rightarrow \mathcal{D}'_{L^1}$. and $*$: $\mathcal{D}'_{E'_*} \times \mathcal{D}^*_E \rightarrow \mathcal{B}^*_\omega$. When E is reflexive, we have $*$: $\mathcal{D}'_{E'_*} \times \mathcal{D}^*_E \rightarrow \dot{\mathcal{B}}^*_\omega$.*

REFERENCES

- [1] R. Carmichael, A. Kamiński, S. Pilipović, *Boundary values and convolution in ultradistribution spaces*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007.
- [2] P. Dimovski, S. Pilipović, J. Vindas, *New distribution spaces associated to translation-invariant Banach spaces*, *Monatsh. Math.*, in press (doi:10.1007/s00605-014-0706-3).
- [3] P. Dimovski, S. Pilipović, J. Vindas, *Boundary values of holomorphic functions and heat kernel method in translation-invariant distribution spaces*, *Complex Var. Elliptic Equ.*, in press (doi:10.1080/17476933.2014.1002399).
- [4] P. Dimovski, S. Pilipović, B. Prangoski, J. Vindas, *Convolution of ultradistributions and ultradistribution spaces associated to translation-invariant Banach spaces*, *Kyoto J. Math.*, to appear.
- [5] E. Hille, R. S. Phillips, *Functional analysis and semi-groups*, American Mathematical Society, Providence, R.I, 1974.
- [6] J. Kisyński, *On Cohen's proof of the factorization theorem*, *Ann. Polon. Math.* **75** (2000), 177–192.
- [7] H. Komatsu, *Ultradistributions, I: Structure theorems and a characterization*, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **20** (1973), 25–105.
- [8] H. Komatsu, *Ultradistributions, III: Vector valued ultradistributions and the theory of kernels*, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **29** (1982), 653–717.
- [9] S. Pilipović, *Characterizations of bounded sets in spaces of ultradistributions*, *Proc. Amer. Math. Soc.* **120** (1994), 1191–1206.
- [10] S. Pilipović, B. Prangoski, J. Vindas, *On quasianalytic classes of Gelfand-Shilov type. Parametrix and convolution*, preprint.
- [11] B. Prangoski, *Laplace transform in spaces of ultradistributions*, *Filomat* **27** (2013), 747–760.
- [12] H. H. Schaefer, *Topological vector spaces*, Springer-Verlag, New York-Heidelberg-Berlin, 1970.
- [13] L. Schwartz, *Théorie des distributions à valeurs vectorielles. I*, *Ann. Inst. Fourier* **7** (1957), 1–141.
- [14] L. Schwartz, *Théorie des distributions*, Hermann, Paris, 1966.

P. DIMOVSKI, FACULTY OF TECHNOLOGY AND METALLURGY, UNIVERSITY SS. CYRIL AND METHODIUS, RUGER BOSKOVIC 16, 1000 SKOPJE, MACEDONIA
E-mail address: dimovski.pavel@gmail.com

B. PRANGOSKI, FACULTY OF MECHANICAL ENGINEERING, UNIVERSITY SS. CYRIL AND METHODIUS, KARPOS II BB, 1000 SKOPJE, MACEDONIA
E-mail address: bprangoski@yahoo.com

J. VINDAS, DEPARTMENT OF MATHEMATICS, GHENT UNIVERSITY, KRIJGSLAAN 281 GEBOUW S22, 9000 GENT, BELGIUM
E-mail address: jvindas@cage.UGent.be