

Miserez, A., Colot, C., Nyssen, J., Enyew Adgo, Verhoest, N., Poesen, J., Deckers, J., 2015. Spatial patterns of soil erodibility through soils in land systems of Lake Tana catchment (Ethiopia). In: Lanckriet, S., Enyew Adgo, Nyssen, J., 2015. TropiLakes2015 international conference on tropical lakes in a changing environment: water, land, biology, climate and humans, Bahir Dar (Ethiopia), 23-29 September 2015. Bahir Dar, Ethiopia: Bahir Dar University, Book of Abstracts: 126.

### **Spatial patterns of soil erodibility through soils in land systems of Lake Tana catchment (Ethiopia)**

Audrey Miserez (1), Charlotte Colot (2), Jan Nyssen (3), Enyew Adgo(4), Niko Verhoest (5), Jean Poesen (1), Jozef Deckers (1)

(1) Department of Earth and environmental Science, KU Leuven

(2) ARCADIS, Belgium sa, Rue Royale 80, 1000 Bruxelles

(3) Department of Geography, Ghent University, Krijgslaan 281 (S8), B-9000 Gent, Belgium

(4) College of Agriculture and Environmental Sciences, Bahir Dar University, P.O. Box 79, Ethiopia

(5) Laboratory of Hydrology and Water Management, Ghent University, Coupure links 653, B-9000 Gent

#### **Abstract**

This presentation draws up the logic of the soils in different land systems in Lake Tana basin. It aims at answering three questions: “Which major soil types are found at which positions in the landscape”, “How can this distribution be explained”, and “How is the spatial pattern of soil erodibility in these soils?” .

The soils and geomorphology of the basin are characterized by a volcanic past. Basaltic and lacustrine parent materials are dominating the basin and influencing the soil-landscape. Most Vertisols and Fluvisols developed on the lacustrine deposits or in the alluvial plains. The alkali basalts from the Oligo-pliocene (Tertiary trap series) are characterized by different soil types with Nitisols and Leptosols as dominating groups. Nitisols are also the dominant soil type on the alkali basalts from the Quaternary (Quaternary Aden series). Those Nitisols are typical for slopes and higher well-drained positions of the landscapes.

We use the knowledge on soil distribution to study the K-factor in the Revised Universal Soil Loss Equation (RUSLE). The soil erodibility was estimated using (1) the approach of Renard et al. (1997) based on soil texture, organic carbon content, soil structure and permeability and (2), by Römken et al. (1997) using only texture as input. A logarithmic relationship was found between rock fragment cover (%) and slope gradient (%) for cultivated land ( $R^2=0.58$ ). Rock fragment cover protects the soil against aggregate breakdown, raindrop impact and supports the soil structure as well, lowering the effect of increased soil erosion on slopes.

Structure and rock fragment cover seemed to be the most important soil properties influencing their erodibility. There is no evidence that soil texture and organic matter are contributing to a statistical difference of K-factors between soil group or geomorphological units in this basin. This fact is not related to the smaller importance of soil texture but rather to the relatively homogeneous soil texture composition of the basin.

According to the formula of Renard et al., accounting for the Nitisols revealed to be significantly more erodible than the other soil types of the basin ( $p=2.10^{-6}$ ), for two main reasons being (1) the nutty, blocky structure and (2) the low rock fragment cover. No statistical differences were measured when using the formula based on textural data only.