

Psychiatr Q (2010) 81:227–238  
DOI 10.1007/s11126-010-9132-4

ORIGINAL PAPER

# ‘The Human Prerogative’: A Critical Analysis of Evidence-Based and Other Paradigms of Care in Substance Abuse Treatment

Eric Broekaert · Mieke Autrique · Wouter Vanderplasschen · Kathy Colpaert

Published online: 2 April 2010  
© Springer Science+Business Media, LLC 2010

**Abstract** Present-day substance abuse treatment is characterized by a compelling demand for applying evidence-based interventions. Vehement discussions between policymakers, practitioners and researchers illustrate this clash of differing paradigms. The aim of this article is to situate evidence-based practice among the leading paradigms of care and to elucidate its implicit assumptions and potential implications. Evidence-based practice is inherent in the empirical-analytical paradigm of care and science, founded upon randomized and controlled studies. This paradigm is compared with the phenomenological-existential and the critical post-structural paradigm, which focus on elaborating the human potential and exploring individuals’ subjective interpretations, and on criticizing social inequalities and striving for compliance with human rights, respectively. Evidence-based practice and the methodological rigidity in each paradigm are analyzed critically. We conclude that through the dialectical integration of these diverse approaches, evidence, existence/humanism and social emancipation can be combined for the benefit of the human prerogative of care.

**Keywords** Addiction · Treatment · Research paradigm · Evidence-based medicine · Therapeutic communities · Harm reduction

## Introduction

Evidence-based practice is currently considered to be an important prerequisite to realizing quality of care and has also found its way into substance abuse treatment. In various countries, the need for the implementation of evidence-based interventions and guidelines is emphasized repeatedly and increasingly by researchers and policymakers

---

E. Broekaert (✉) · W. Vanderplasschen · K. Colpaert  
Department of Orthopedagogics, Ghent University, Henri Dunantlaan 2, 9000 Ghent, Belgium  
e-mail: Eric.Broekaert@UGent.be

M. Autrique  
Association for Alcohol and Other Drug Problems (VAD), Vanderlindenstraat 15, 1030 Brussels, Belgium

[1–6]. According to some authors [7, 8], the field of substance abuse treatment is even moving towards a binding demand for the use of evidence-based standards in which the reimbursement of costs will gradually be tied to the delivery of evidence-based interventions.

This tendency may have far-reaching consequences for daily practice since practitioners are afraid that substance abuse treatment will be limited to interventions which have been proven to be effective in randomized and controlled trials or will wipe out the subjective component in treatment [9]. Such concerns were also expressed by clinicians, treatment providers and researchers at various national and international conferences in Brussels (Belgian Federal Science Policy, March 27, 2007), Thessaloniki (Society & Mental Health Conference, October 20–21, 2007<sup>1</sup>), Oslo (European Working Group on Drug-Oriented Policy Research, October 1–3, 2007) and Oxford (Oxford Science Meeting for Therapeutic Communities, March 31–April 1, 2008). Similar reactions were observed during a study we made concerning the implementation and application of evidence-based interventions and guidelines in Belgian substance abuse treatment [10]. Strikingly, in the French-speaking part of Belgium (the Walloon provinces), which has a strong psychodynamic treatment tradition and is more closely oriented to the Latin tradition and the Southern European countries, the opposition towards the evidence-based paradigm is much stronger than in the Dutch-speaking part of the country (Flanders). The latter region is more connected to the Anglo-Saxon countries and their traditions [11].

The afore-mentioned discussions between practitioners, policymakers and researchers clearly illustrate a clash of treatment and research paradigms. Consequently, the aim of this article is to elucidate implicit assumptions of evidence-based medicine and what may be its implications for substance abuse treatment. We want to clarify which paradigm is behind the evidence-based approach and whether, by the dominance of this paradigm, some treatment approaches are over- or undervalued. First, we will examine the origins of evidence-based medicine and its underpinning paradigm of care. Second, we will analyze this and other paradigms and the criticism they inflict, and discuss whether an integration of different paradigms is necessary to benefit the ‘human prerogative’ of care. This ‘human prerogative’ implies that a modernist, humanistic approach is seen as a prerequisite for every action and intervention in substance abuse treatment [12] and may offer opportunities to demonstrate similarities rather than dogmatic antitheses in the basic assumptions about evidence-based practice.

## Evidence-Based Medicine

Evidence-based medicine is a movement that originated under the impetus of a large coalition of physicians, researchers, professors and policymakers to improve the application of results from experimental scientific research in clinical practice [13–15]. It has been defined as ‘... *the conscientious, explicit, and judicious use of current best evidence in making decisions about the care of individual patients. The practice of evidence-based medicine means integrating best research evidence with clinical expertise and patient*

<sup>1</sup> During this conference a first draft of this paper has been presented and was published in Greek in the Proceedings of the Society & Mental Health Conference, pp. 53–62.

values' [16, p. 1]. An important exponent of this movement is the international Cochrane Collaboration, which was founded by the Scottish professor Archie Cochrane [17]. His ideas were at the basis of the establishment of the Cochrane Library, a database of systematic reviews concerning a wide range of interventions for any kind of health problem [18].

In the evidence-based discourse, evidence is graded according to the methods used to collect it. This resulted in criteria to assign a level of evidence (e.g. 'strong', 'some' or 'no' evidence) for the effectiveness of interventions, based on the number, nature and quality of studies about this intervention (cf. [8, 19, 20]). According to these criteria, RCTs or 'randomized controlled trials' are considered to be the 'gold standard', followed by non- or partially randomized quasi-experimental studies, other quantitative research designs (e.g. correlational studies) and qualitative research (e.g. case studies). In various review studies, publications have been selected and classified on the basis of such criteria, leading to conclusions concerning the evidence for the effectiveness of interventions [20–22]. For example, the 'Mesa Grande' study [23], a meta-analysis of clinical trials, has compared various treatment interventions for alcohol abusers and classified these according to their degree of evidence.

Although such classifications are not always univocal, it can be concluded that there is mainly evidence for the effectiveness of various pharmacological interventions (e.g. benzodiazepines and carbamazepine; methadone and buprenorphine) for the treatment of alcohol and opiate abuse [21, 24–28]. Further, there is evidence for the effectiveness of some psychosocial treatments, mainly specific, brief and behavioral interventions (such as cognitive behavioral therapy, community reinforcement approach, motivational interviewing, contingency management, brief interventions and multidimensional family therapy) [11, 20]. On the other hand, there is not (yet) much evidence for the effectiveness of other widely implemented psychosocial interventions, such as therapeutic communities, psycho-education, relapse prevention, case management, social skills training, psychotherapy, counseling or psychodynamic therapy. No evidence is available for the effectiveness of acupuncture, educational lectures, confrontational approaches and compulsory participation in self-help groups [2, 29].

The assumption behind these classifications is that treatment approaches and aspects can be assessed—and even compared and classified—based on experimental evidence. This tendency is closely related to the empirical-analytical paradigm of care, which will be discussed and compared in the next section with other paradigms of care, namely the phenomenological-existential and the critical post-structural paradigm of care.

It is our basic assumption that research and science have to contribute to the further development of mankind. Each of these different paradigms offers a specific contribution to some specific problems and challenges. The human condition is the prerogative of free, but responsible decision-making from scientific paradigms that underpin treatment systems. This process does not exclude one or another option, but examines carefully which paradigm(s) contribute(s) in what situation to the best solution. From this point of view, the different paradigms do not exclude, but rather complement each other: in their search for the best answer in a given situation, they try to find the most suitable treatment and do the best for mankind. Whatever 'the best' may be, it has to be the exponent of a meaningful, significant flexible process of a methodical and systematic search or action for an expected valuable solution, free of dogmatic premises. This is what we call the 'human prerogative of care' and various research paradigms may be used to achieve it.

## Diverse Paradigms of Care

### The Empirical-Analytical Paradigm of Care

In the empirical-analytical paradigm of care, treatment is conceived as a coherent totality of causal conditional relations which can be analyzed in their composing parts and can be defined separately. Evidence-based medicine is underpinned by this paradigm. In its purest form, it raises experimental evidence to rule over other forms of knowledge. Certain study designs, such as meta-analyses and randomized trials, are thought to be less vulnerable to bias and, therefore, provide ‘superior’ evidence [30]. Interventions in substance abuse treatment are subdivided into (good) practices that have been proven to work (or not) and that are based on evidence (or not). This paradigm of care is most closely related to behavioral treatment approaches.

### The Phenomenological-Existential Paradigm of Care

From a phenomenological-existential perspective, treatment is considered to be a methodical approach that involves the whole person and his or her social network and that aims at facilitating (mental) health and well-being. Or, as treatment was traditionally defined, as ‘a range of interventions which are intended to remedy an identified drug-related problem or condition with a view to a person’s physical, psychological or social well-being’ [31]. Referring to interventions intended to improve someone’s well-being implies per definition meaningful action (‘to become well through the act of intervention’). In this context, treatment consists of taking adequate action in special situations for helping or supporting individuals.

In the phenomenological-existential paradigm, treatment success is measured based on the reported experiences (of well-being) concerning the intervention. It is argued that human motives, social interactions and beliefs—in short, human life and activities in general—are far too complex to be reduced to statistical analyses that aim at universally valid evidence. In order to reach a scientific and causal explanation of proceedings and effects, a rational and/or empathic interpretative understanding is needed [32]. This phenomenological-existential paradigm of care is clearly connected to the therapeutic community and self-help approaches in substance abuse treatment; it stresses self-actualization and growth and is inherent in humanist psychology and pedagogy.

### The Critical Post-Structural Paradigm of Care

From a third, post-modern, post-structural perspective, treatment should be discussed from a critical position towards society, in which concepts like inclusion, self-advocacy and emancipation of the poor and weakest prevail [33]. The grand theory of science and its associated methodologies no longer make sense, but should be replaced by narratives [34]. As long as we are captured by our genealogy and mental (language) structures, the subject cannot become an actualized self [35]. Humans have to deconstruct their world to get some grip on reality (cf. [36]), which is characterized by uncertainty and relativity. Human rights for every citizen and the rights to quality of life and quality of care are guiding principles in disability studies [37], including the support for and treatment of substance abusers. This critical post-structural paradigm of care is closely linked with the harm-reduction

movement, in which social inclusion, (self-) advocacy and emancipation are central concepts, and that takes a critical position towards the conservative values of society.

### Critical Reactions to Evidence-Based Medicine

#### From an Empirical-Analytical Point of View

From an empirical-analytical point of view, mainly methodological questions and concerns are expressed concerning the actual tendency towards more evidence-based practice, although it starts from the same fundamental ideas. Some authors emphasize statistical problems when searching for convincing evidence in comprehensive ‘box-score reviews’ [38–40]. They refer to issues such as low/variable statistical power to demonstrate treatment effects, varying statistical tests, diverse comparison groups across studies, and lack of consistent data on client characteristics.

Because some insurmountable problems are intrinsic to the experimental, randomized and controlled setup, several researchers choose ‘minimal bias designs’ (e.g. [41–43]). Still, these criticisms are of secondary nature and do not touch ‘the heart’ of the approach itself.

#### From a Phenomenological Point of View

Phenomenologists criticize the lack of distinction between professional treatment as a compilation of good practices and evidence-based treatment as one of its constituting elements [44]. They call attention to the constriction and simplification of various treatment problems into elements that can be controlled by an empirical-analytical research design [38]. The evidence-based paradigm may underestimate the essence of treatment as meaningful action and deny its phenomenological nature that requires qualitative (empirical) research [31]. They state that the use of evidence-based practice and research findings requires an understanding of the processes by which implicit and explicit knowledge and beliefs are constructed and that guide the daily practice of treatment [45]. They further stress the importance of clinical freedom and the need to find a balance between the ‘art of medicine’ and evidence-based interventions, even if this might be a false dichotomy [46]. Evidence-based medicine does not provide a means to integrate other, non-statistical sources of medical information, such as professional experience and patient-specific factors [30], although their importance is recognized by proponents of EBM.

Phenomenologists argue that, if evidence is used as part of action, it inevitably becomes an integral part of meaningful action. The ‘act’ of using evidence-based approaches is a prerequisite for the method itself. According to De Leon [47, 48], action in the field of substance abuse treatment unarguably implies self-selection. It is not possible to enjoy the benefits of a remedy, such as medication, without the act of taking it (or not taking it, if one wants to falsify this induction or hypothesis). According to this rationale, the ‘act of taking’ is of greater importance than the evidence itself. From a more global perspective, one can also question the idea that addiction is a chronic relapsing disorder (cf. [49]). For, not all substance abusers relapse, but some stay sober. Moreover, addiction is not only a medical condition but is determined by a complex interaction of psychological, social, hereditary and biological factors [50, 51].

The ultimate consequence is that *neither* psychiatry *nor* therapeutic communities *nor* self-help groups *nor* psychological counseling can be evidence-based, but only the interventions that are part of it. However, these interventions do not determine the complexity of the treatment approaches they are part of.

#### From a Social Critical Point of View

Poststructuralists criticize the denial of post-modern approaches and the lack of collaborative, empowering and inclusive methodologies in evidence-based medicine [52, 53]. They state that the negation of post-structural approaches in science stands for conservatism and a lack of social activism and compassion for the underprivileged [54]. According to social critical practitioners, the underprivileged have to be empowered in order to achieve the full civil right of free choice and self-selection of treatment. The act of self-selection is seen as an integral part of the act of treatment but is in essential contradiction with the principle of randomization. Alternatively, they advocate for a replacement of the search for effectiveness and efficiency by assessing and monitoring quality of care and quality of life [55]. It should be clear that both quality concepts do not necessarily correlate—on the contrary [37, 56]. Last but not least, the critical social perspective questions evidence-based medicine as it is adopted by the current neo-liberal movement towards managed care and compartmentalization of care into units, modules and functions [57]. This may lead to dramatic cutbacks in funding of health care and preserve evidence-based interventions for the ‘higher’ social classes, although this was not the intention of the founders and proponents of evidence-based medicine. It could go so far that health resources for the rich are preserved—due to governmental priorities—at the cost of essential needs of the poor and underprivileged.

The social critical point of view can be combined with the evidence-based paradigm, as is illustrated in controlled heroin trials in the Netherlands [58] and Germany [59] in which evidence is applied as part of a critical social commitment against traditional and conservative societal forces.

### Consequences of Methodo- and Ethnocentrism for Substance Abuse Treatment

#### From a Methodocentric Point of View

The methodocentric scientific point of view rigorously uses one specific methodology for evaluating any type of intervention and promotes the exclusive use of one ‘scientific’ paradigm [60]. In the case of empirical-analytical (c.q. evidence-based) research, the guiding principles are observable behavior, objectivity and generalization, falsification of hypotheses, explanation, prediction and control based on facts. Phenomenological research concerns meaningful action, critical dialogue, inter-subjectivity and particularity, and a cycle of planning, action, evaluation, change and improvement, based on subjective interpretations [61]. Post-modern post-structuralism is about social criticism and emancipation, deconstruction, inclusion and diversity, uncertainty, narratives and relativity. Consequently, clear-cut interventions (such as contingency management or CBT), global approaches (such as therapeutic communities or Alcoholics Anonymous) and progressive social practices (such as needle exchange or controlled heroin trials) and alienating social institutions (such as prisons) are studied solely from an empirical-analytical or a phenomenological or a post-structural point of view.

This kind of methodocentrism is typical for evidence-based medicine, in which the effectiveness of all treatment approaches and interventions is assessed according to the same criteria and standards, thus elevating experimental research to a higher level than any other type of research.

#### From an Ethnocentric Point of View

The ethnocentric scientific point of view considers the method used as a social construction and promotes the application of a methodology that tight-fits the nature of the type of intervention [62]. Consequently, behavioral approaches (with their focus on observable behavior) can exclusively be investigated from an empirical-analytical point of view, while comprehensive approaches that link various interventions into ‘action’ require a phenomenological point of view. Progressive social practices and alienating social institutions should, from this ethnocentric perspective, be evaluated from a post-structural critical point of view.

#### From an Integrative Point of View

The integrative scientific point of view searches for the integration of diverse types of interventions as well as methodological approaches. It means that various treatment modalities, paradigms of care and research methodologies can alternatively go together [57]. Consequently, behavioral interventions, therapeutic communities, self-help groups, harm reduction and prison treatment for substance abusers can complement each other and all have their own specific value. Therefore, clear-cut interventions, global approaches and progressive social practices need to be evaluated from an analytical-empirical point of view, as well as from a phenomenological and post-structuralist critical point of view.

In the following section, we discuss why the dialectical integration of diverse approaches (empirical-analytical, phenomenological, and social critical)—and thus of evidence, existence and social emancipation—could benefit the human prerogative of care. By doing so, we want to offer practitioners and researchers an alternative framework for addressing the ever-increasing demand for evidence-based practice.

### **Towards the Integration of Paradigms and the Search for a Human Quality of Care and Life: Actual Paradoxes and Future Options**

Present-day substance abuse treatment is characterized by a compelling demand for evidence-based practice that applies and promotes the empirical-analytical paradigm of care and science. Still, practitioners appear to be rather hesitant or even reluctant to adopt this tendency [1]. The predominance of one paradigm implicitly—and maybe not intentionally—leads to a hierarchy among treatment modalities, since these are underpinned by different methodological paradigms. Such a hierarchy implies the loss of self-help and therapeutic community approaches, but also—and at first sight paradoxically—the promotion of harm reduction approaches. The marriage between two orthodoxies—experimentalism and harm reduction—can mainly be explained by the search for certainty (evidence) in an uncertain post-modern world (i.e. the search for reduction of harm). The former is denominated as ‘managed health care’, the latter as ‘public safety’, illustrating a neo-liberal approach which at the same time promotes individual freedom, hedonism and

even the right to controlled drug use, and on the other hand stresses the need for control, security and rationalization.

However, in their search for a better quality of care and life, evidence and emancipation miss their third partner: existence and humanism. If we want to enhance human quality of life and care for substance (ab)users, a further integration of the different paradigms of care is needed. In order to elevate the quality of care as well as the quality of life of substance (ab)users, evidence from experimental research, the self-actualizing existence of people and the search for social emancipation should be combined. In this new search for an integration between parts and totality, various complementing (partial) approaches alternatively go together in their never-ending pursuit of unity and transformation of actual conditions.

In such an integrated model, the going together of each element's excluding aspects of reality is referred to as *complementarity* [63]. The opposites are interchangeable and may replace each other in a series of transformations. This means that the "system as a whole shifts to a new, higher order, a more complex structural form of which the parts are governed by a new set of functional properties and are characterized by a new set of statistical parameters" [64, p. 380]. This dynamic dialectical process includes cycles of disorganization and the transformation of activities, meanings, self-development and relations. This understanding of reality is founded on two premises: 1) that everything is interconnected; and 2) that a living organism cannot be reduced to its physical parts. The opposing terms *sum* and *parts* are in fact misleading, because their relation is intrinsic as well as external.

The mechanism behind this search for holistic unity is called *integration* [65]. It transforms 'thesis' and 'anti-thesis' (part and counterpart) to a new unity or synthesis. In other words, once unity is reached, a new process starts, a transformation is realized and a higher order reached. Consequently, from this point of view, evidence-based medicine (the empirical-analytical paradigm), therapeutic communities (the phenomenological-existentialist paradigm) and harm reduction (the social critical paradigm) exclude each others' reality only at first sight. In fact, they are internally connected. This interconnection is the human interconnection—the human prerogative—the human dedication to achieving the best solution for persons with substance (ab)use problems.

Once we accept this internal link, we can start an open, methodical, meaningful and valuable search for the best answer to this or that problem, based on the well-being and human dignity of the substance (ab)using person. As a human being, the latter is also an interconnected and integral part of this 'best' choice. The underlying paradigms of care are no longer conflicting; instead they complete each other as a human endeavour. By doing so, we can transcend postulated and presupposed solutions.

In summary, various treatment approaches and methodologies simultaneously co-exist, go alternately together and strive for unity through integration, but once unity or synthesis is reached they dissipate and transform the closed system into a new open one. In other words, if open systems such as therapeutic communities or psychiatric hospitals use motivational interviewing, they do so in a process of transforming treatment towards a higher level of functioning. Psychiatric treatment and therapeutic communities interact as a whole with their composing parts. There is no reason to study therapeutic communities or AA group meetings as 'methods' and compare them with other 'partial interventions', as there is no reason to compare psychiatry with other 'global interventions' such as TC or AA. Within this context, 'medical treatment as an art' versus 'evidence-based medicine' is just a false dichotomy: they simply complete each other in the search for a better integrated medicine. Similarly, the ethical questions concerning 'at random' or 'minimal bias' methodologies can be considered as two sides of the same coin in the search for the best evidence for current practice.



Our position on the ‘integration of paradigms’ might have important consequences for substance abuse treatment and it provides a theoretical framework underpinning the application of ‘integrated treatment systems’ [57]. Such an integrated and comprehensive system of treatment modalities (including drug-free treatment, substitution therapy, harm reduction initiatives, and other health, psychiatric and social services) provides the opportunity to tailor services according to individuals’ needs and expectations. Case management is an important method in such an integrated network of services, as it provides the ability to link people to appropriate services [66] and to guarantee coordination and continuity of care [67]. One of the prerequisites of an integrated treatment system is that no treatment modality is in essence better than another [57]: various treatment approaches are not competing, but may all have their value for someone with substance abuse problems at a certain moment in a certain situation throughout their lifespan.

## Conclusion

Ultimately, our critical analysis of evidence-based and other paradigms of care in substance abuse treatment led to the conclusion that each of these approaches is underpinned by another scientific paradigm. We distinguished between the empirical-analytical (e.g., evidence-based medicine), phenomenological-existentialist (e.g., therapeutic communities) and social critical paradigm (e.g., harm reduction). We examined the consequences of methodo- and ethnocentrism for substance abuse treatment and found that each of these paradigms is to a certain extent reductionist. Therefore, we chose an integrated paradigm of care. We argued that care approaches are interconnected by a common human dedication to achieving the best answer for substance abusers’ treatment demands. We called this interconnection ‘the human prerogative of care’ and accepted as a logical consequence that no single treatment system can solve all substance abuse problems. Real human commitment consists of an open, methodical, meaningful search for the best solution for a certain problem. This process implies doubt and uncertainty, but is ultimately based on free and responsible decision-making, in interaction with the client. It assumes a treatment system that gives full opportunity and authority to a broad spectrum of interconnected treatment modalities and services, all with their own explicit identity, strengths and weaknesses (i.e., an integrated treatment system).

All of this requires an interaction between evidence, facts and methodological security and the uncertainty of scientific doubt. Scientific evaluation of substance abuse treatment from diverse points of view can contribute greatly to the improvement of the quality of care and life of substance users and to the effectiveness and efficiency of substance abuse treatment. It is in fact this alternately going together of evidence, existence and emancipation that constitutes the ‘human prerogative of care’.

## References

1. Autrique M, Vanderplasschen W, Broekaert E, et al.: Practitioners’ attitudes concerning evidence-based guidelines in substance abuse treatment. *European Addiction Research* 15:47–55, 2009
2. Miller WR, Sorensen JL, Selzer JA, et al.: Disseminating evidence-based practices in substance abuse treatment: A review with suggestions. *Journal of Substance Abuse Treatment* 31:25–39, 2006

3. Ravndal E, Vaglum P, Lauritzen G: Completion of long-term inpatient treatment of drug abusers: A prospective study from 13 different units. *European Addiction Research* 11:180–185, 2005
4. McGovern MP, Fox TS, Xie HY, et al.: A survey of clinical practices and readiness to adopt evidence-based practices: Dissemination research in an addiction treatment system. *Journal of Substance Abuse Treatment* 26:305–312, 2004
5. Berglund M, Thelander S, Jonsson E: *Treating Alcohol and Drug Abuse: An Evidence Based Review*. Weinheim, Wiley-VCH, 2003
6. Schippers GM, Schramade M, Walburg JA: Reforming Dutch substance abuse treatment services. *Addictive Behaviors* 27:995–1007, 2002
7. Brown BS: Commentary on 'Evidence-based treatment: Why, what, where, and how?' *Journal of Substance Abuse Treatment* 30:87–89, 2006
8. Miller WR, Zweben J, Johnson WR: Evidence-based treatment: Why, what, where, and how? *Journal of Substance Abuse Treatment* 29:267–276, 2005
9. Petrova M, Dale J, Fulford B: Value-based practice in primary care: Easing the tensions between individual values, ethical principles and best evidence. *British Journal of General Practice* 56:703–709, 2006
10. Autrique M, Vanderplasschen W, Pham TH, et al.: *Evidence-based werken in de verslavingszorg: een stand van zaken*. Reeks Wetenschap en Maatschappij, Antwerpen-Apeldoorn, Garant, 2007
11. Autrique M, Vanderplasschen W, Broekaert E, et al.: The drug-free therapeutic community: Findings and reflections in an evidence-based era. *Therapeutic Communities* 29:5–15, 2008
12. Broekaert E, Vanderplasschen W, Colpaert K, et al.: The Human Prerogative: Questions on Integration of Evidence-Based, Existential and Social Critical Paradigms of Care in Substance Abuse Treatment. In: *Proceedings of the Society & Mental Health Conference, Thessaloniki, 2008*
13. Haynes RB, Sackett DL, Gray JM, et al.: Transferring evidence from research into practice: 1. The role of clinical care research evidence in clinical decisions. *Evidence-based Medicine* 1:196–198, 1996
14. Bloch RM, Saeed SA, Rivard JC, et al.: Lessons learned in implementing evidence-based practices: Implications for psychiatric administrators. *Psychiatric Quarterly* 77:309–318, 2006
15. Gray GE, Pinson LA: Evidence-based medicine and psychiatric practice. *Psychiatric Quarterly* 74:387–399, 2003
16. Sackett DL: *Evidence-Based Medicine: How to Practice and Teach EBM*, 2nd ed. New York, Churchill Livingstone, 2000
17. Chalmers I, Haynes B: Systematic reviews: Reporting, updating and correcting systematic reviews of effects of health care. *British Medical Journal* 309:862–865, 1994
18. The Cochrane Library: The Cochrane Library. Evidence for healthcare decision- making [in <http://www.theCochraneLibrary.com>, December 2008]
19. Kwaliteitsinstituut voor de Gezondheidszorg (CBO): *Evidence-based richtlijn ontwikkeling: Handleiding voor werkgroepen*. Utrecht, Kwaliteitsinstituut voor de gezondheidszorg CBO, 2005
20. Rigter H, Van Gageldonk A, Ketelaars T, et al.: *Hulp bij probleemgebruik van drugs*. Utrecht, Trimbos-instituut, 2004
21. Lingford-Hughes AR, Welch S, Nutt DJ: Evidence-based guidelines for the pharmacological management of substance misuse, addiction and comorbidity: Recommendations from the British Association for Psychopharmacology. *Journal of Psychopharmacology* 18:293–335, 2004
22. Van Gageldonk A, De Zwart W, Van der Stel J, et al.: *De Nederlandse Verslavingszorg: overzicht van de kennis over aanbod, vraag en effect*. Utrecht, Trimbos-instituut, 1997
23. Miller WR, Wilbourne PL: Mesa Grande: A methodological analysis of clinical trials of treatment for alcohol use disorders. *Addiction* 97:265–277, 2002
24. Gowing L, Ali R, White J: Buprenorphine for the management of opioid withdrawal. *Cochrane Database Systematic Reviews* 2:1–49, 2006
25. Mattick RP, Breen C, Kimber J, et al.: Methadone maintenance therapy versus no opioid replacement therapy for opioid dependence. *Cochrane Database Systematic Reviews* 2:1–19, 2003
26. Malcolm R, Myrick H, Roberts J, et al.: The effects of carbamazepine and lorazepam on single versus multiple previous alcohol withdrawals in an outpatient randomized trial. *Journal of General Internal Medicine* 17:349–355, 2002
27. Holbrook AM, Crowther R, Lotter A, et al.: Meta-analysis of benzodiazepine use in the treatment of acute alcohol withdrawal. *Canadian Medical Association Journal* 160:649–655, 1999
28. Mayo-Smith M: Pharmacological management of alcohol withdrawal. *The Journal of the American Medical Association* 278:144–151, 1997
29. Miller WR, Wilbourne PL, Hettema JE: *What Works? A Summary of Alcohol Treatment Outcome Research*. In: Hester RK, Miller WR (Eds) *Handbook of Alcoholism Treatment Approaches*, 3rd edn. Boston, MA, Pearson Education, 2003

30. Cohen AM, Stavri PZ, Hersh WR: A categorization and analysis of the criticisms of evidence-based medicine. *International Journal of Medical Informatics* 73:35–43, 2004
31. Broekaert E, De Wilde J: The Construction of the VACT for Women: A Flexible Tool for the Treatment of Female Substance Abusers. In: Pedersen MU, Segraeus V, Helleman M (Eds) *Evidence-Based Practice? Challenges in Substance Abuse Treatment*. Helsinki, NAD Publication 47, 2005
32. Colins O, Broekaert E, Vandeveldde S, et al.: Max Weber and Alfred Schutz: the theoretical and methodological background of the case-oriented quantification approach behind winMAX. *Social Science Computer Review* 26:369–378, 2008
33. Broekaert E, D'Oosterlinck F, Van Hove G, et al.: The search for an integrated paradigm of care models for people with handicaps, disabilities and behavioural disorders at the department of Orthopedagogy of Ghent University. *Education and Training in Developmental Disabilities* 39:206–216, 2004
34. Lyotard JF: *The Post-Modern Condition: A Report on Knowledge*. Minneapolis, University of Minnesota, 1979
35. Foucault M: *The Archaeology of Knowledge*. London, Tavistock Publications, 1972
36. Derrida J: *De la Gramatologie*. Paris, Editions de minuit, 1967
37. De Waele I, Van Hove G: Modern times: An ethnographic study on the quality of life of people with a high support need in a Flemish residential facility. *Disability & Society* 20:625–639, 2005
38. Bergmark A: Evidence-Based Practice—More Control or More Uncertainty. In: Pedersen MU, Segraeus V, Helleman M (Eds) *Evidence-Based Practice? Challenges in Substance Abuse Treatment*. Helsinki, NAD Publication 47, 2005
39. Bergmark A: On treatment mechanisms—What can we learn from the COMBINE study? *Addiction* 103:703–705, 2008
40. Finney JW: Limitations in using existing alcohol treatment trials to develop practice guidelines. *Addiction* 95:1491–1500, 2000
41. Welsh WN: A multisite evaluation of prison-based therapeutic community drug treatment. *Criminal Justice and Behavior* 34:1481–1498, 2007
42. Soyeze V, De Leon G, Rosseel Y, et al.: The impact of a social network intervention on retention in Belgian therapeutic communities: A quasi experimental study. *Addiction* 101:1027–1034, 2006
43. McLellan AT, Hagan TA, Levine M, et al.: Does clinical case management improve outpatient addiction treatment. *Drug and Alcohol Dependence* 55:91–103, 1999
44. Pedersen MU: Drug-Free Treatment of Substance Misusers: Where Are We Now, Where Are We Heading. In: Pedersen MU, Segraeus V, Helleman M (Eds) *Evidence-Based Practice? Challenges in Substance Abuse Treatment*. Helsinki, NAD Publication 47, 2005
45. Gabbay J, May A: Evidence based guidelines or collectively constructed “mindlines?” Ethnographic study of knowledge management in primary care. *British Medical Journal* 329:1013–1017, 2004
46. Parker M: False Dichotomies: EBM, clinical freedom, and the art of medicine. *Medical Humanities* 31:23–30, 2005
47. De Leon G: Commentary: Reconsidering the self-selection factor in addiction treatment research. *Psychology of Addictive Behaviors* 12:71–77, 1998
48. De Leon G: Commentary on “Self-help organizations for alcohol and drug problems: Toward evidence-based practice and policy”. *Journal of Substance Abuse Treatment* 26:163–165, 2004
49. McLellan AT, Lewis DC, O'Brien CP, et al.: Drug Dependence, a Chronic Medical Illness: Implications for treatment, insurance and outcomes evaluation. *JAMA* 284:1689–1695, 2000
50. Kooyman M: The Different Aspects of Addiction and the Therapeutic Community Approach. Paper presented at the 12th EFTC Conference in Ljubljana, Slovenia, June 6, 2007
51. Marsh DC, Fair BR: Addiction treatment in Vancouver. *International Journal of Drug Policy* 17:137–141, 2006
52. Braidotti R: Feminist epistemology after postmodernism: Critiquing science, technology and globalisation. *Interdisciplinary Science Reviews* 32:65–74, 2007
53. Fisher P, Goodley D: The linear medical model of disability: Mothers of disabled babies resist with counter-narratives. *Sociology of Health & Illness* 29:66–81, 2007
54. Roets G, Van de Perre D, Van Hove G, et al.: One for All—All for One! An account of the joint fight for human rights by Flemish Musketeers and their Tinker Ladies. *British Journal of Learning Disabilities* 32:54–64, 2004
55. Broekaert E: What future for the Therapeutic Community in the field of addiction? A view from Europe. *Addiction* 101:1677–1678, 2006
56. De Maeyer J, Vanderplasschen W, Broekaert E: Exploratory study on drug users' perspectives on quality of life: More than health-related quality of life? *Social Indicators Research* 90:107–126, 2009

57. Broekaert E, Vanderplasschen W: Towards the integration of treatment systems for substance abusers: Report on the second international symposium on substance abuse treatment and special target groups. *Journal of Psychoactive Drugs* 35:237–345, 2003
58. van den Brink W, Hendriks VM, Blanken P, et al.: Medical prescription of heroin to treatment resistant heroin addicts: Two randomised controlled trials. *British Medical Journal* 327:310B–312B, 2003
59. Verthein U, Bonorden-Kleij K, Degkwitz P, et al.: Long-term effects of heroin-assisted treatment in Germany. *Addiction* 103:960–966, 2008
60. Green RT: Constitutional jurisprudence—Reviving praxis in public administration. *Administration & Society* 24:3–21, 1992
61. Gieles F: Conflict en Contact. Een onderzoek naar handelingsmogelijkheden voor groepsleiders bij botsingen en conflicten in de dagelijkse leefsituatie. Proefschrift Universiteit Groningen, 1992
62. Garfinkel F: *Studies in Ethnomethodology*. Englewood Cliffs, NJ, Prentice Hall, 1967
63. Capra F: *The Tao of Physics: An Exploration of the Parallels Between Modern Physics and Eastern Mysticism*. Berkeley, CA, Shambhala Publications, 1975
64. Brent SB: Prigogine's model for self-organization in no equilibrium systems: Its relevance for developmental psychology. *Human Development* 21:374–387, 1978
65. Rucker R: *Oneindigheid, filosofie en wetenschap van het oneindige*. Amsterdam, Contact, 1986
66. Hesse M, Vanderplasschen W, Rapp RC, et al.: Case management for persons with substance use disorders. *Cochrane Database of Systematic Reviews* 4:1–82, 2007
67. Vanderplasschen W, Wolf J, Rapp RC, et al.: Effectiveness of different models of case management for substance-abusing populations. *Journal of Psychoactive Drugs* 39:81–93, 2007

## Author Biographies

**Eric Broekaert, PhD**, is a professor of orthopedagogics at Ghent University (Belgium). He is honoree vice president of the European Federation of Therapeutic Communities (EFTC) and member of the editorial collective of *The International Journal of Therapeutic Communities*.

**Mieke Autrique, MEd**, is working as a staff member at the Association for Alcohol and Other Drug Problems (VAD) in Brussels (Belgium). Her areas of interest are evidence-based guidelines, outcome management and evaluation studies.

**Wouter Vanderplasschen, PhD**, is working as a Ph.D. research and teaching assistant at the Department of Orthopedagogics (Ghent University, Belgium). His main research subjects are: substance abuse, assessment and treatment planning, organisation of care, case management and treatment evaluation.

**Kathy Colpaert, MEd**, is working as a research and teaching assistant at the Department of Orthopedagogics (Ghent University, Belgium). Her areas of interest are addiction severity, psychiatric characteristics and quality of life of poly-substance abusers in substance abuse treatment.