

The kernel of the generalized Clifford-Fourier transform and its generating function

Pan Lian^{1,2*} Gejun Bao^{1 †} Hendrik De Bie^{2‡} Denis Constaes^{2 §}

1: Department of Mathematics - Harbin Institute of Technology
West Da-Zhi Street 92, 150001 Harbin, P.R.China

2: Department of Mathematical Analysis
Faculty of Engineering and Architecture – Ghent University
Galglaan 2, 9000 Gent, Belgium

Abstract

In this paper, we study the generalized Clifford-Fourier transform introduced in [7] using the Laplace transform technique. We give explicit expressions in the even dimensional case, we obtain polynomial bounds for the kernel functions and establish a generating function.

Keywords: Clifford-Fourier transform, Laplace transform, Bessel function, generalized Fourier transform

Mathematics Subject Classification: 42B10, 30G35, 15A66, 44A10

Contents

1	Introduction	1
2	Preliminaries	2
2.1	The Laplace transform	2
2.2	Clifford analysis and generalized Fourier transforms	3
3	Generalized kernel in the Laplace domain	5
3.1	Closed expression for $e^{i\frac{\pi}{2}\Gamma_y^2}e^{-i(x,y)}$	5
3.2	Closed expression for $e^{i\frac{\pi}{2}G(\Gamma_y)}e^{-i(x,y)}$	8
4	Conclusion	12

1 Introduction

In recent years, quite some attention has been devoted to the study of hypercomplex Fourier transforms. For the historical development of quaternion and Clifford-Fourier transforms we refer to [4]. In the present paper, we consider the Clifford-Fourier transform first established in [2, 3]. This is a genuinely non-scalar

*E-mail: pan.lian@ugent.be (corresponding author)

†E-mail: baogj@hit.edu.cn

‡E-mail: hendrik.debie@ugent.be

§E-mail: denis.constaales@ugent.be

32 generalization of the Fourier transform, developed within the framework of Clifford analysis [9]. Indeed,
 33 it can be written as

$$F_-(f)(y) = (2\pi)^{-m/2} \int_{\mathbb{R}^m} K_m(x, y) f(x) dx$$

34 with

$$K_m(x, y) = e^{i\frac{\pi}{2}\Gamma_y} e^{-i(x, y)}$$

35 with Γ_y the spherical Dirac operator (see equation (7)).

36 It turned out to be a difficult problem to determine the kernel $K_m(x, y)$ explicitly. This was first
 37 achieved in [8] using plane wave decompositions. Later, in [6] a different method using wave equations
 38 was established. In [5], a short proof was obtained by considering the Clifford-Fourier kernel in the
 39 Laplace domain, where it takes on a much simpler form.

40 Our aim in the present paper is to develop the Laplace transform method for a much wider class of
 41 generalized Fourier transforms. According to investigations in [7] using the representation theory for the
 42 Lie superalgebra $\mathfrak{osp}(1|2)$, the following expression

$$e^{i\frac{\pi}{2}G(\Gamma_y)} e^{-i(x, y)} \tag{1}$$

43 where G is an integer-valued polynomial can be used as the kernel for a generalized Fourier transform
 44 that still satisfies properties very close to that of the classical transform. The extension of the Laplace
 45 transform technique to kernels of type (1) will allow us to find explicit expressions for the kernel. We
 46 will moreover determine which polynomials G give rise to polynomially bounded kernels and we will
 47 determine the generating function corresponding to a fixed polynomial G .

48 The paper is organized as follows. In order to make the exposition self-contained, in Section 2, we
 49 recall basic facts of the Laplace transform, Clifford analysis and the generalized Clifford-Fourier transform.
 50 Section 3 is devoted to establishing the connection between the kernel of the fractional Clifford-Fourier
 51 transform [5] and the generalized Clifford-Fourier transform. We first compute a special case in Section
 52 3.1. Then the method is generalized to the case in which the polynomial has integer coefficients in Section
 53 3.2. The kernel and the generating function in the even dimensional case are given. We also discuss which
 54 kernels are polynomially bounded.

55 2 Preliminaries

56 2.1 The Laplace transform

57 The Laplace transform of a real or complex valued function f which has exponential order α , i.e. $|f(t)| \leq$
 58 $Ce^{\alpha t}, t \geq t_0$ is defined as

$$F(s) = \mathcal{L}(f(t)) = \int_0^\infty e^{-st} f(t) dt.$$

59 By Lerch's theorem [16], the inverse transform

$$\mathcal{L}^{-1}(F(s)) = f(t)$$

60 is uniquely defined when we restrict to functions which are continuous on $[0, \infty)$. Usually, we can use
 61 integral transform tables (see e.g. [11]) and the partial fraction expansion to compute the Laplace
 62 transform and its inverse. We list some which will be used in this paper:

$$\mathcal{L}(e^{-\alpha t}) = \frac{1}{s + \alpha}; \tag{2}$$

$$\mathcal{L}(t^{k-1} e^{-\alpha t}) = \frac{\Gamma(k)}{(s + \alpha)^k}, \quad k > 0. \tag{3}$$

63 We also need the convolution formula and the inverse Laplace transform. Denote by $r = (s^2 + a^2)^{1/2}$,
64 $R = s + r$, $G(s) = \mathcal{L}(g(t))$ and $F(s) = \mathcal{L}(f(t))$. We have

$$G(s)F(s) = \mathcal{L}\left(\int_0^t g(t-\tau)f(\tau)d\tau\right); \quad (4)$$

$$\mathcal{L}^{-1}(a^\nu r^{-2\nu-1}) = 2^\nu \pi^{-1/2} \Gamma(\nu + \frac{1}{2}) t^\nu J_\nu(at), \quad \text{Re}(\nu) > -1/2, \text{Re}(s) > |\text{Im}(a)|. \quad (5)$$

65 2.2 Clifford analysis and generalized Fourier transforms

66 In this section, we give a quick review of the basic concepts in Clifford analysis and generalized Fourier
67 transforms. Denoting by $\{e_1, e_2, \dots, e_m\}$ the orthonormal basis of \mathbb{R}^m , the Clifford algebra $\mathcal{C}\ell_{0,m}$ over
68 \mathbb{R}^m is spanned by the reduced products

$$\bigcup_{j=1}^m \{e_\alpha = e_{i_1} e_{i_2} \dots e_{i_j} : \alpha = \{i_1, i_2, \dots, i_j\}, \quad 1 \leq i_1 < i_2 < \dots < i_j \leq m\}$$

69 with the relations $e_i e_j + e_j e_i = -2\delta_{ij}$. We identify the point $x = (x_1, \dots, x_m)$ in \mathbb{R}^m with the vector
70 variable $x = \sum_{j=1}^m e_j x_j$. The inner product and the wedge product of two vectors $x, y \in \mathbb{R}^m$ can be
71 defined by the Clifford product:

$$(x, y) := \sum_{j=1}^m x_j y_j = -\frac{1}{2}(xy + yx);$$

72

$$x \wedge y := \sum_{j < k} e_j e_k (x_j y_k - x_k y_j) = \frac{1}{2}(xy - yx).$$

73 We can find the Clifford product $xy = -(x, y) + x \wedge y$, and $(x \wedge y)^2 = -|x|^2 |y|^2 + (x, y)^2$ (see [8]). The
74 complexified Clifford algebra $\mathcal{C}\ell_{0,m}^c$ is defined as $\mathbb{C} \otimes \mathcal{C}\ell_{0,m}$.

75 The conjugation is defined by $\overline{(e_{j_1} \dots e_{j_l})} = (-1)^l e_{j_l} \dots e_{j_1}$ as a linear mapping. For $x, y \in \mathcal{C}\ell_{0,m}^c$, we
76 have $\overline{(xy)} = \bar{y}\bar{x}$, $\bar{\bar{x}} = x$, and $\bar{\bar{i}} = i$ which is not the usual complex conjugation. We define the Clifford
77 norm of x by $|x|^2 = x\bar{x}$, $x \in \mathcal{C}\ell_{0,m}^c$.

78 The Dirac operator is given by $D = \sum_{j=1}^m e_j \partial_{x_j}$. Together with the vector variable x , they satisfy the
79 relations

$$D^2 = -\Delta, \quad x^2 = -|x|^2, \quad \{x, D\} = -2\mathbb{E} - m,$$

80 where $\{a, b\} = ab + ba$ and $\mathbb{E} = \sum_{j=1}^m x_j \partial_{x_j}$ is the Euler operator and hence they generate a realization
81 of the Lie superalgebra $\mathfrak{osp}(1|2)$, which contains the Lie algebra $\mathfrak{sl}_2 = \text{span}\{\Delta, |x|^2, [\Delta, |x|^2]\}$ as its even
82 part. A function $u(x)$ is called monogenic if $Du = 0$. An important example of monogenic functions is
83 the generalized Cauchy kernel

$$G(x) = \frac{1}{\omega_m} \frac{\bar{x}}{|x|^m}$$

84 where ω_m is the surface area of the unit ball in \mathbb{R}^m . It is the fundamental solution of the Dirac operator
85 [9]. Note that the norm here is $|x| = (\sum_{i=1}^m x_i^2)^{1/2}$ and coincides with Clifford norm.

86 Denote by \mathcal{P} the space of polynomials taking values in $\mathcal{C}\ell_{0,m}$, i.e. $\mathcal{P} := \mathbb{R}[x_1, \dots, x_m] \otimes \mathcal{C}\ell_{0,m}$. The
87 space of homogeneous polynomials of degree k is then denoted by \mathcal{P}_k . The space $\mathcal{M}_k := (\ker D) \cap \mathcal{P}_k$, is
88 called the space of homogeneous monogenic polynomials of degree k . An arbitrary element of it is called
89 a spherical monogenic of degree k [9].

90 The local behaviour of a monogenic function near a point can be investigated by the polynomials
91 introduced above. The following theorem is the analogue of the Taylor series in complex analysis.

92 **Theorem 1.** [9] *Suppose f is monogenic in an open set Ω containing the origin. Then there exists an*
93 *open neighbourhood Λ of the origin in which f can be developed into a normally convergent series of*
94 *spherical monogenics $M_k f(x)$, i.e.*

$$f(x) = \sum_{k=0}^{\infty} M_k f(x),$$

95 with $M_k f(x) \in \mathcal{M}_k$.

96 The classical Fourier transform

$$\mathcal{F}(f)(y) = (2\pi)^{-m/2} \int_{\mathbb{R}^m} e^{-i(x,y)} f(x) dx,$$

97 with (x, y) the usual inner product can be represented by the operator exponential [14], [15]

$$\mathcal{F} = e^{-i\frac{\pi}{4}(\Delta - |x|^2 - m)}.$$

98 The Clifford-Hermite functions

$$\psi_{2p,k,l}(x) := 2^p p! L_p^{\frac{m}{2} + k - 1}(|x|^2) M_k^l e^{-|x|^2/2},$$

99

$$\psi_{2p+1,k,l}(x) := 2^p p! L_p^{\frac{m}{2} + k}(|x|^2) x M_k^l e^{-|x|^2/2},$$

100 where $p, k \in \mathbb{Z}_{\geq 0}$ and $\{M_k^l | l = 1, \dots, \dim(\mathcal{M}_k)\}$ form a basis for \mathcal{M}_k , the space of spherical monogenics of
 101 degree k . They moreover realize the complete decomposition of the rapidly decreasing functions $\mathcal{S}(\mathbb{R}^m) \otimes$
 102 $\mathcal{C}\ell_m \subset L^2(\mathbb{R}^m) \otimes \mathcal{C}\ell_m$ in irreducible subspaces under the action of the dual pair $(Spin(m), \mathfrak{osp}(1|2))$. The
 103 action of the regular Fourier transform on this basis is given by

$$\mathcal{F}\psi_{j,k,l} = e^{-i\frac{\pi}{2}(j+k)} \psi_{j,k,l} = (-i)^{j+k} \psi_{j,k,l}. \quad (6)$$

104 We further introduce the Gamma operator or the angular Dirac operator (see [9])

$$\Gamma_x := - \sum_{j < k} e_j e_k (x_j \partial_{x_k} - x_k \partial_{x_j}) = -x D_x - \mathbb{E}_x = -x \wedge D_x, \quad (7)$$

105 here $\mathbb{E}_x = \sum_{i=1}^m x_i \partial_{x_i}$ is the Euler operator. Note that Γ_x commutes with scalar radial functions. The
 106 operator Γ_x has two important eigenspaces:

$$\Gamma_x \mathcal{M}_k = -k \mathcal{M}_k, \quad (8)$$

107

$$\Gamma_x (x \mathcal{M}_{k-1}) = (k + m - 2) x \mathcal{M}_{k-1} \quad (9)$$

108 which follows from the definition of Γ_x . The Scasimir S in our operator realization of $\mathfrak{osp}(1|2)$ is related
 109 to the angular Dirac operator by $S = -\Gamma_x + \frac{m-1}{2}$, see [12]. The Casimir element $C = S^2$ acts on the
 110 Clifford-Hermite function by

$$C\psi_{j,k,l} = \left(k + \frac{m-1}{2}\right)^2 \psi_{j,k,l}.$$

111 In [7], the authors studied the full class of integral transforms which satisfy the conditions stated in the
 112 following theorem.

113 **Theorem 2.** *The properties*

114 (1) *the Clifford-Helmholtz relations*

$$T \circ D_x = -iy \circ T,$$

115

$$T \circ x = -iD_y \circ T,$$

116 (2) $T\psi_{j,k,l} = \mu_{j,k} \psi_{j,k,l}$ with $\mu_{j,k} \in \mathbb{C}$,

117 (3) $T^4 = id$

118 are satisfied by the operators T of the form

$$T = e^{i\frac{\pi}{2}F(C)} e^{i\frac{\pi}{4}(\Delta - |x|^2 - m)} \in e^{i\frac{\pi}{2}\bar{U}(\mathfrak{osp}(1|2))}$$

119 where $F(C)$ is a power series in C that takes integer values when evaluated in the eigenvalues of C and
 120 $\bar{U}(\mathfrak{osp}(1|2))$ is the extension of the universal enveloping algebra that allows infinite power series in the
 121 elements of \mathfrak{sl}_2 .

122 The integral kernel of the generalized Fourier transform T can be expressed as $e^{i\frac{\pi}{2}F(C)}e^{-i(x,y)}$. We are
 123 in particular interested in the case where $F(C)$ reduces to a polynomial $G(\Gamma_y)$ with integer coefficients.

124 **Remark 1.** *In general, when $G(x) \neq 0$, the generalized Fourier transform T and the Clifford fractional
 125 Fourier transform in [1] are two different classes of transforms because their eigenvalues on the Clifford-
 126 Hermite functions are different.*

127 **Remark 2.** *The Clifford Fourier transform in $Cl_{(3,0)}$ can also be expressed by operator exponential,
 128 see e.g. [10].*

129 3 Generalized kernel in the Laplace domain

130 3.1 Closed expression for $e^{i\frac{\pi}{2}\Gamma_y^2}e^{-i(x,y)}$

131 In this subsection, we use the Laplace transform method to compute $e^{i\frac{\pi}{2}\Gamma_y^2}e^{-i(x,y)}$. The trick here will be
 132 used to compute the more general case in next subsection. We use the notation $\sqrt{\mp} := \sqrt{s^2 + |x|^2|y|^2}$.
 133 The following lemma was obtained in [5].

134 **Lemma 1.** *The Laplace transform of $t^{m/2-1}e^{-it(x,y)}$ can be expressed as*

$$\mathcal{L}(t^{m/2-1}e^{-it(x,y)}) = \frac{2^{m/2-1}\Gamma(m/2)}{\sqrt{\mp}(s + \sqrt{\mp})^{m/2-1}} \frac{1 - \frac{iyx}{s + \sqrt{\mp}} + \frac{iy(1 - \frac{iyx}{s + \sqrt{\mp}})x}{s + \sqrt{\mp}}}{\left|1 - \frac{iyx}{s + \sqrt{\mp}}\right|^m}. \quad (10)$$

135 In the following, we will act with $e^{i\frac{\pi}{2}\Gamma_y^2}$ on both sides of (10) to obtain the integral kernel in the
 136 Laplace domain. Denote by

$$f(y) = \frac{2^{\frac{m}{2}}}{\sqrt{\mp}(s + \sqrt{\mp})^{m/2-1}} \frac{1 - \frac{iyx}{s + \sqrt{\mp}}}{\left|1 - \frac{iyx}{s + \sqrt{\mp}}\right|^m} = \frac{s + \sqrt{\mp} - iyx}{\sqrt{\mp}(s + i(x,y))^{m/2}},$$

137 and

$$g(y) = \frac{2^{\frac{m}{2}}}{\sqrt{\mp}(s + \sqrt{\mp})^{m/2-1}} \frac{\frac{iy(1 - \frac{iyx}{s + \sqrt{\mp}})x}{s + \sqrt{\mp}}}{\left|1 - \frac{iyx}{s + \sqrt{\mp}}\right|^m} = \frac{iy}{s + \sqrt{\mp}} f(y)x = \frac{\sqrt{\mp} - s + iyx}{\sqrt{\mp}(s + i(x,y))^{m/2}}.$$

138 In [5], it has been proved that $f(y)$ has a series expansion as

$$f(y) = \frac{2^{\frac{m}{2}}}{\sqrt{\mp}(s + \sqrt{\mp})^{m/2-1}} \sum_{k=0}^{\infty} \frac{M_k(y)}{(s + \sqrt{\mp})^k}.$$

139 Here we rewrite

$$f(y) = f_0(y) + f_1(y) + f_2(y) + f_3(y),$$

140 with

$$f_k(y) = \frac{2^{\frac{m}{2}}}{\sqrt{\mp}(s + \sqrt{\mp})^{m/2-1}} \sum_{n=0}^{\infty} \frac{M_{4n+k}(y)}{(s + \sqrt{\mp})^{4n+k}}, \quad k = 0, 1, 2, 3. \quad (11)$$

141 Each f_k is an eigenfunction of the operator $e^{i\frac{\pi}{2}\Gamma^2}$. In fact, by (8), we have

$$e^{i\frac{\pi}{2}\Gamma_y^2}M_k(y) = e^{i\frac{\pi}{2}(-k)^2}M_k(y),$$

142 so

$$\begin{aligned} e^{i\frac{\pi}{2}\Gamma_y^2}M_{4n}(y) &= M_{4n}(y); \\ e^{i\frac{\pi}{2}\Gamma_y^2}M_{4n+1}(y) &= iM_{4n+1}(y); \\ e^{i\frac{\pi}{2}\Gamma_y^2}M_{4n+2}(y) &= M_{4n+2}(y); \\ e^{i\frac{\pi}{2}\Gamma_y^2}M_{4n+3}(y) &= iM_{4n+3}(y), \end{aligned} \tag{12}$$

143 here $n = 0, 1, 2, \dots$. Since the operator Γ commutes with radial functions, we know that each f_k is an
144 eigenfunction of $e^{i\frac{\pi}{2}\Gamma^2}$ and the eigenvalues are given in (12). In the following, we denote

$$f_\alpha(y) = \frac{2^{\frac{m}{2}}}{\sqrt{+}(s + \sqrt{+})^{m/2-1}} \sum_{k=0}^{\infty} \frac{M_k(iy)}{(s + \sqrt{+})^k} = \frac{s + \sqrt{+} + yx}{\sqrt{+}(\sqrt{+} - (x, y))^{m/2}},$$

145

$$f_\beta(y) = \frac{2^{\frac{m}{2}}}{\sqrt{+}(s + \sqrt{+})^{m/2-1}} \sum_{k=0}^{\infty} \frac{M_k(-y)}{(s + \sqrt{+})^k} = \frac{s + \sqrt{+} + iyx}{\sqrt{+}(s - i(x, y))^{m/2}},$$

146

$$f_\gamma(y) = \frac{2^{\frac{m}{2}}}{\sqrt{+}(s + \sqrt{+})^{m/2-1}} \sum_{k=0}^{\infty} \frac{M_k(-iy)}{(s + \sqrt{+})^k} = \frac{s + \sqrt{+} - yx}{\sqrt{+}(\sqrt{+} + (x, y))^{m/2}}$$

147 as well as

$$\begin{aligned} g_\alpha(y) &= \frac{iy}{s + \sqrt{+}} f_\alpha(y)x = \frac{i(\sqrt{+} - s) + iyx}{\sqrt{+}(\sqrt{+} - (x, y))^{m/2}}, \\ g_\beta(y) &= \frac{iy}{s + \sqrt{+}} f_\beta(y)x = \frac{s - \sqrt{+} + iyx}{\sqrt{+}(s - i(x, y))^{m/2}}, \\ g_\gamma(y) &= \frac{iy}{s + \sqrt{+}} f_\gamma(y)x = \frac{i(s - \sqrt{+}) + iyx}{\sqrt{+}(\sqrt{+} + (x, y))^{m/2}}. \end{aligned}$$

148 **Remark 3.** Comparing with Theorem 3 in [5], $\frac{\Gamma(m/2)}{2}(f_\gamma + g_\alpha)$ is the Clifford-Fourier kernel of dimension
149 $m = 4n + 1, n \in \mathbb{N}$ in the Laplace domain. Denote the first part of the fractional Clifford-Fourier kernel
150 as

$$F_p(x, y) = \frac{s + \sqrt{+} - ie^{-ip}yx}{\sqrt{+}(e^{-ip}(s \cos p + i\sqrt{+} \sin p + i(x, y)))^{m/2}}$$

151 and the second part of the kernel as

$$G_p(x, y) = -e^{ip} \frac{s - \sqrt{+} - ie^{-ip}yx}{\sqrt{+}(e^{ip}(s \cos p - i\sqrt{+} \sin p + i(x, y)))^{m/2}}.$$

152 We find that $f(y) = F_0(x, y)$, $f_\alpha(y) = F_{-\frac{\pi}{2}}(x, y)$, $f_\beta(y) = F_\pi(x, y)$, $f_\gamma(y) = F_{\frac{\pi}{2}}(x, y)$, $g(y) = G_0(x, y)$,
153 $g_\alpha(y) = G_{\frac{\pi}{2}}(x, y)$, $g_\beta(y) = G_\pi(x, y)$ and $g_\gamma(y) = G_{-\frac{\pi}{2}}(x, y)$. We could get the plane wave expansion and
154 integral expression of $f, f_\alpha, f_\beta, f_\gamma$ and $g, g_\alpha, g_\beta, g_\gamma$ from [5].

155 As M_k is a polynomial of degree k , we have the following relations,

$$\begin{cases} f(y) = f_0(y) + f_1(y) + f_2(y) + f_3(y); \\ f_\alpha(y) = f_0(y) + if_1(y) - f_2(y) - if_3(y); \\ f_\beta(y) = f_0(y) - f_1(y) + f_2(y) - f_3(y); \\ f_\gamma(y) = f_0(y) - if_1(y) - f_2(y) + if_3(y). \end{cases}$$

156 Each $f_k(y)$ can be obtained as follows:

$$\begin{cases} 4f_0(y) = f(y) + f_\alpha(y) + f_\beta(y) + f_\gamma(y); \\ 4f_1(y) = f(y) - if_\alpha(y) - f_\beta(y) + if_\gamma(y); \\ 4f_2(y) = f(y) - f_\alpha(y) + f_\beta(y) - f_\gamma(y); \\ 4f_3(y) = f(y) + if_\alpha(y) - f_\beta(y) - if_\gamma(y). \end{cases} \quad (13)$$

157 Now the action of $e^{i\frac{\pi}{2}\Gamma_y^2}$ on $f(y)$ is known through its eigenfunctions,

$$\begin{aligned} e^{i\frac{\pi}{2}\Gamma_y^2}f(y) &= e^{i\frac{\pi}{2}\Gamma_y^2}\left(f_0(y) + f_1(y) + f_2(y) + f_3(y)\right) \\ &= f_0(y) + if_1(y) + f_2(y) + if_3(y) \\ &= \frac{1}{2}\left(f(y) + f_\beta(y) + if(y) - if_\beta(y)\right). \end{aligned}$$

158 The case $e^{i\frac{\pi}{2}\Gamma_y^2}g(y)$ can be treated similarly, using (12) and

$$\begin{aligned} e^{i\frac{\pi}{2}\Gamma_y^2}(yM_k(y)) &= e^{i\frac{\pi}{2}(m-1+k)^2}(yM_k(y)) \\ &= e^{i\frac{\pi}{2}(m-1)^2}e^{i\frac{\pi}{2}k^2}(yM_k(e^{i\pi(m-1)}y)) \\ &= e^{i\frac{\pi}{2}(m-1)^2}ye^{i\frac{\pi}{2}k^2}(M_k(e^{i\pi(m-1)}y)). \end{aligned}$$

159 Collecting everything, we have

160 **Theorem 3.** *The kernel $t^{m/2-1}e^{i\frac{\pi}{2}\Gamma_y^2}e^{-i(x,y)}$ in the Laplace domain is*

$$\begin{aligned} &\mathcal{L}(t^{m/2-1}e^{i\frac{\pi}{2}\Gamma_y^2}e^{-it(x,y)}) \\ &= \frac{\Gamma(m/2)}{4\sqrt{+}}\left((1+i)U_m^1 + (1-i)U_m^2 + e^{i\frac{\pi}{2}(m-1)^2}((1+i)U_m^3 + (1-i)U_m^4)\right), \end{aligned}$$

161 *with*

$$\begin{aligned} 162 \quad U_m^1 &= \frac{s + \sqrt{+} - iyx}{(s + i(x,y))^{m/2}}; & U_m^2 &= \frac{s + \sqrt{+} + iyx}{(s - i(x,y))^{m/2}}; \\ 163 \quad U_m^3 &= \frac{(-1)^{m-1}(\sqrt{+} - s) + iyx}{(s + (-1)^{m-1}i(x,y))^{m/2}}; & U_m^4 &= \frac{(-1)^{m-1}(s - \sqrt{+}) + iyx}{(s - (-1)^{m-1}i(x,y))^{m/2}}, \end{aligned}$$

164 *where $\sqrt{+} = \sqrt{s^2 + |x|^2|y|^2}$.*

165 When $m = 2$,

$$\mathcal{L}(e^{i\frac{\pi}{2}\Gamma_y^2}e^{-it(x,y)}) = \frac{1}{2\sqrt{+}}\left(\frac{\sqrt{+}}{s - i(x,y)} + \frac{s - iyx}{s + i(x,y)}\right).$$

166 By formula (2), (5), and the convolution formula (4), the kernel equals, putting $t = 1$,

$$K_{2,\Gamma^2}(x,y) = e^{i(x,y)} + J_0(|x||y|) + ix \wedge y \int_0^1 e^{-i(x,y)(1-\tau)} J_0(|x||y|\tau) d\tau.$$

167 In the following, we analyze each term in Theorem 3 in detail. By formula (3), (4) and (5), letting

168 $t = 1$, we get $U_m^1, U_m^2, U_m^3, U_m^4$ in the time domain as

$$\begin{aligned}
K_{U_m^1} &= \frac{e^{-i(x,y)}}{\Gamma(m/2)} + \frac{1}{\Gamma(m/2-1)} \int_0^1 \tau^{m/2-2} e^{-i(x,y)\tau} J_0(|x||y|(1-\tau)) d\tau \\
&\quad + \frac{ix \wedge y}{\Gamma(m/2)} \int_0^1 e^{-i(x,y)} J_0(|x||y|(1-\tau)) d\tau, \\
K_{U_m^2} &= \frac{e^{i(x,y)}}{\Gamma(m/2)} + \frac{1}{\Gamma(m/2-1)} \int_0^1 \tau^{m/2-2} e^{i(x,y)\tau} J_0(|x||y|(1-\tau)) d\tau \\
&\quad - \frac{ix \wedge y}{\Gamma(m/2)} \int_0^1 e^{i(x,y)} J_0(|x||y|(1-\tau)) d\tau, \\
K_{U_m^3} &= (-1)^{m-1} \left(\frac{1}{\Gamma(m/2)} e^{i(-1)^m(x,y)} \right. \\
&\quad \left. - \frac{1}{\Gamma(m/2-1)} \int_0^1 \tau^{m/2-2} e^{i(-1)^m(x,y)\tau} J_0(|x||y|(1-\tau)) d\tau \right) \\
&\quad - \frac{ix \wedge y}{\Gamma(m/2)} \int_0^1 e^{i(-1)^m(x,y)} J_0(|x||y|(1-\tau)) d\tau,
\end{aligned}$$

169

$$\begin{aligned}
K_{U_m^4} &= (-1)^{m-1} \left(-\frac{1}{\Gamma(m/2)} e^{i(-1)^{m-1}(x,y)} \right. \\
&\quad \left. + \frac{1}{\Gamma(m/2-1)} \int_0^1 \tau^{m/2-2} e^{i(-1)^{m-1}(x,y)\tau} J_0(|x||y|(1-\tau)) d\tau \right) \\
&\quad - \frac{ix \wedge y}{\Gamma(m/2)} \int_0^1 e^{i(-1)^{m-1}(x,y)} J_0(|x||y|(1-\tau)) d\tau.
\end{aligned}$$

170 **Theorem 4.** Let $m \geq 2$. For $x, y \in \mathbb{R}^m$, the generalized Fourier kernel is given by

$$\begin{aligned}
K_{m,\Gamma^2}(x, y) &= \frac{\Gamma(m/2)}{4} \left((1+i)K_{U_m^1} + (1-i)K_{U_m^2} \right. \\
&\quad \left. + e^{i\frac{\pi}{2}(m-1)^2} ((1+i)K_{U_m^3} + (1-i)K_{U_m^4}) \right).
\end{aligned}$$

171 There exists a constant c such that

$$|K_{m,\Gamma^2}(x, y)| \leq c(1 + |x||y|).$$

172 *Proof.* This follows from the fact that $J_0(y)$ and $e^{i(x,y)}$ are bounded functions and $|x \wedge y| \leq |x||y|$. \square

173 3.2 Closed expression for $e^{i\frac{\pi}{2}G(\Gamma_y)}e^{-i(x,y)}$

174 In this subsection, we consider the more general case. We act with $G(\Gamma_y)$ on the Fourier kernel. Here
175 $G(x)$ is a polynomial with integer coefficients,

$$G(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0, \quad a_k \in \mathbb{Z}.$$

176 Using the fact that $e^{i\frac{\pi}{2}j}$ is 4-periodic in j ,

$$e^{i\frac{\pi}{2}G(\Gamma_y)} M_k(y) = e^{i\frac{\pi}{2}G(-k)} M_k(y)$$

177 and

$$G(4n+k) \equiv G(k) \pmod{4},$$

178 we have

$$\begin{aligned} e^{i\frac{\pi}{2}G(\Gamma_y)}f(y) &= e^{i\frac{\pi}{2}G(0)}f_0 + e^{i\frac{\pi}{2}G(-1)}f_1 + e^{i\frac{\pi}{2}G(-2)}f_2 + e^{i\frac{\pi}{2}G(-3)}f_3 \\ &= i^{G(0)}f_0 + i^{G(-1)}f_1 + i^{G(-2)}f_2 + i^{G(-3)}f_3, \end{aligned}$$

179 with each f_k defined in (11). By

$$e^{i\frac{\pi}{2}G(\Gamma_y)}(yM_k(y)) = e^{i\frac{\pi}{2}G(m-1+k)}(yM_k)$$

180 and

$$G(4n + k + m - 1) \equiv G(k + m - 1)(\text{mod}4),$$

181 we have

$$\begin{aligned} &e^{i\frac{\pi}{2}G(\Gamma_y)}g(y) \\ &= \frac{iy}{s + \sqrt{+}} \left(e^{i\frac{\pi}{2}G(m-1)}f_0 + e^{i\frac{\pi}{2}G(m)}f_1 + e^{i\frac{\pi}{2}G(m+1)}f_2 + e^{i\frac{\pi}{2}G(m+2)}f_3 \right) x \\ &= \frac{iy}{s + \sqrt{+}} \left(i^{G(m-1)}f_0 + i^{G(m)}f_1 + i^{G(m+1)}f_2 + i^{G(m+2)}f_3 \right) x. \end{aligned}$$

182 Collecting everything and applying (13), we get

183 **Theorem 5.** For $G(x) \in \mathbb{Z}[x]$, the Laplace transform of $t^{m/2-1}e^{i\frac{\pi}{2}G(\Gamma_y)}e^{-it(x,y)}$ is given by

$$\mathcal{L}(t^{m/2-1}e^{i\frac{\pi}{2}G(\Gamma_y)}e^{-it(x,y)}) = \frac{\Gamma(m/2)}{8} \left(A_m^1 B C_m^T + \frac{iy}{s + \sqrt{+}} A_m^2 B C_m^T \right)$$

184 with A_m^1, A_m^2, B, C_m the matrices given by

$$\begin{aligned} A_m^1 &= (i^{G(0)} \quad i^{G(-1)} \quad i^{G(-2)} \quad i^{G(-3)}), \\ A_m^2 &= (i^{G(m-1)} \quad i^{G(m)} \quad i^{G(m+1)} \quad i^{G(m+2)}), \\ B &= \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -i & -1 & i \\ 1 & -1 & 1 & -1 \\ 1 & i & -1 & -i \end{pmatrix}, \\ C_m &= (f(y) \quad f_\alpha(y) \quad f_\beta(y) \quad f_\gamma(y)). \end{aligned}$$

185 **Remark 4.** We could get the regular Fourier kernel $e^{-i(x,y)}$ by setting $G(x) = 0$ or $4x$ for dimension
186 $m \geq 2$. When $G = 2x^2$, we get the inverse Fourier kernel $e^{i(x,y)}$ for even dimension. When $G(x) = \pm x$,
187 it is the Clifford-Fourier transform [8].

188 As the constant term of the polynomial will only contribute a constant factor to the integral kernel,
189 in the following we only consider polynomials without constant term

$$G(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x, \quad a_k \in \mathbb{Z}.$$

190 By

$$G(4n + k) \equiv G(k)(\text{mod}4),$$

191 it reduces to four cases $G(k)(\text{mod}4)$, $k = 0, 1, 2, 3$. The set $\{x^m\} \cup \{1\}$, $m \in \mathbb{N}$ is a basis for polynomials
192 over the ring of integers. We consider the four cases on this basis

$$\begin{aligned} x^j &= 0, & \text{when } x &= 0; \\ x^j &= 1, & \text{when } x &= 1; \\ x^j &\equiv \begin{cases} 2(\text{mod}4), & \text{when } j = 1 \text{ and } x = 2; \\ 0(\text{mod}4), & \text{when } j \geq 2 \text{ and } x = 2; \end{cases} \\ x^j &\equiv \begin{cases} 1(\text{mod}4), & \text{when } j \text{ is even and } x = 3; \\ 3(\text{mod}4), & \text{when } j \text{ is odd and } x = 3. \end{cases} \end{aligned}$$

193 For each $G(x)$, we denote $\frac{G(1)+G(-1)}{2} = s_0 = \sum_{j=0}^{\lfloor n/2 \rfloor} a_{2j}$ and $\frac{G(1)-G(-1)}{2} = s_1 = \sum_{j=0}^{\lfloor n/2 \rfloor} a_{2j+1}$ with n the
 194 degree of $G(x)$. We have

$$\begin{aligned} G(0) &= 0, \\ G(1) &= s_0 + s_1, \\ G(2) &\equiv 2a_1 \pmod{4}, \\ G(3) &\equiv G(-1) \equiv s_0 - s_1 \pmod{4}. \end{aligned}$$

195 Therefore

$$\begin{aligned} i^{G(0)} &= 1, & i^{G(-1)} &= i^{G(3)} = i^{s_0+3s_1}, \\ i^{G(-2)} &= i^{G(2)} = (-1)^{a_1}, & i^{G(-3)} &= i^{G(1)} = i^{s_0+s_1}. \end{aligned}$$

196 The class of integral transforms with polynomially bounded kernel is of great interest. For example,
 197 new uncertainty principles have been given for this kind of integral transforms in [13]. As we can see in
 198 Theorem 5, the generalized Fourier kernel is a linear combination of $f_\alpha, f_\beta, f_\gamma, f, g_\alpha, g_\beta, g_\gamma, g$. At present,
 199 very few of $f_\alpha, f_\gamma, g_\alpha, g_\gamma$ are known explicitly. The integral representations of $f_\alpha, f_\gamma, g_\alpha, g_\gamma$ are obtained
 200 in [5] but without the bound. Only in even dimensions, special linear combinations of $f_\alpha, f_\gamma, g_\alpha, g_\gamma$ are
 201 known to be polynomially bounded which is exactly the Clifford-Fourier kernel [8].

202 We have showed in Theorem 4 that f, f_β, g, g_β with polynomial bounds behaves better than $f_\alpha, f_\gamma, g_\alpha, g_\gamma$.
 203 So it is interesting to consider the generalized Fourier transform whose kernel only consists of f, f_β, g, g_β .
 204 **It also provides ways to define hypercomplex Fourier transforms with polynomially bounded kernel in odd**
 205 **dimensions.** We will hence characterize polynomials such that $e^{i\frac{\pi}{2}G(\Gamma_y)}e^{-i(x,y)}$ are only linear combination
 206 of f, f_β, g, g_β .

207 For fixed m , the kernel is a linear sum of f, f_β, g, g_β when the polynomial $G(x) \in \mathbb{Z}[x]$ satisfies the
 208 following conditions

$$\begin{cases} i^{G(0)} - ii^{G(-1)} - i^{G(-2)} + ii^{G(-3)} = 0, \\ i^{G(0)} + ii^{G(-1)} - i^{G(-2)} - ii^{G(-3)} = 0, \\ i^{G(m-1)} - ii^{G(m)} - i^{G(m+1)} + ii^{G(m+2)} = 0, \\ i^{G(m-1)} + ii^{G(m)} - ii^{G(m+1)} - ii^{G(m+2)} = 0. \end{cases} \quad (14)$$

209 We find that (14) is equivalent with

$$\begin{cases} G(0) \equiv G(-2) \pmod{4}, \\ G(-1) \equiv G(-3) \pmod{4}, \\ G(m-1) \equiv G(m+1) \pmod{4}, \\ G(m) \equiv G(m+2) \pmod{4}. \end{cases} \quad (15)$$

210 As $G(k) \pmod{4}$ is uniquely determined by $G(0), G(-1), G(-2)$ and $G(-3)$, the first two formulas in (15)
 211 imply the last two formulas for all $m \geq 2$ automatically. Now (15) becomes

$$\begin{cases} i^{G(0)} = 1 = i^{G(-2)} = (-1)^{a_1}, \\ i^{G(-1)} = i^{s_0+3s_1} = i^{G(-3)} = i^{s_0+s_1}. \end{cases}$$

212 It follows that the kernel only consists of f, f_β, g, g_β if and only if a_1 and s_1 are even. We have the
 213 following

214 **Theorem 6.** *Let $m \geq 2$. For $x, y \in \mathbb{R}^m$ and a polynomial $G(x)$ with integer coefficients, the ker-*
 215 *nel $e^{i\frac{\pi}{2}G(\Gamma_y)}e^{-i(x,y)}$ is a linear combination of f, f_β, g, g_β in the Laplace domain if and only if a_1 and*
 216 *$\frac{G(1)-G(-1)}{2}$ are even. Furthermore, the generalized Fourier kernel is bounded and equals*

$$\frac{1 + i^{G(1)}}{2} e^{-i(x,y)} + \frac{1 - i^{G(1)}}{2} K^\pi(x, y),$$

217 *with $K^\pi(x, y)$ the fractional Clifford-Fourier kernel in [5]. When $m \geq 2$ is even, the kernel is*

$$\frac{1 + i^{G(1)}}{2} e^{-i(x,y)} + \frac{1 - i^{G(1)}}{2} e^{i(x,y)}.$$

218 When $m \geq 2$ is odd, there exists a constant c which is independent of m such that

$$|e^{i\frac{\pi}{2}G(\Gamma_y)}e^{-i(x,y)}| \leq c(1 + |x||y|). \quad (16)$$

219 *Proof.* We only need to prove the generalized Fourier kernel is

$$\frac{1 + i^{s_0+s_1}}{2}e^{-i(x,y)} + \frac{1 - i^{s_0+s_1}}{2}K^\pi(x,y).$$

220 In fact, by verification, we have,

$$(e^{i0})^{m-1}A_m^1 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = A_m^2 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}; \quad (e^{i\pi})^{m-1}A_m^1 \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix} = A_m^2 \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix},$$

221 and

$$A_m^1 \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = 2 + 2i^{s_0+s_1}; \quad A_m^1 \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix} = 2 - 2i^{s_0+s_1}.$$

222 By Remark 3, $f + (e^{i0})^{m-1}g$ is the kernel K_0 and $f_\beta + (e^{i\pi})^{m-1}g_\beta$ is the fractional Clifford-Fourier kernel
223 K^π . The bound (16) follows from the integral expression of f, f_β, g, g_β in the time domain. \square

224 **Remark 5.** The case $G(x) = x^2$ is a special case of this theorem.

225 In the following, we consider the generalized Fourier kernel which has polynomial bound and con-
226 sists of $f_\alpha, f_\beta, f_\gamma, f, g_\alpha, g_\beta, g_\gamma, g$. For even dimension, we already know the Clifford-Fourier kernel has a
227 polynomial bound. If the polynomial $G(x)$ satisfies

$$(-i)^{m-1}A_m^1 \begin{pmatrix} 1 \\ -i \\ -1 \\ i \end{pmatrix} = A_m^2 \begin{pmatrix} 1 \\ i \\ -1 \\ -i \end{pmatrix}; \quad i^{m-1}A_m^1 \begin{pmatrix} 1 \\ i \\ -1 \\ -i \end{pmatrix} = A_m^2 \begin{pmatrix} 1 \\ -i \\ -1 \\ i \end{pmatrix}, \quad (17)$$

228 by Remark 3, $e^{i\frac{\pi}{2}G(\Gamma_y)}e^{-i(x,y)}$ is a linear combination of the Clifford-Fourier kernel and some function
229 bounded by $c(1 + |x||y|)$. Hence it has a polynomial bound as well. When $m = 4j$, (17) becomes

$$i(1 - i^{s_0+3s_1+1} - (-1)^{a_1} + i^{s_0+s_1+1}) = i^{s_0+3s_1} + i - i^{s_0+s_1} - i(-1)^{a_1}$$

230 and

$$-i(1 + i^{s_0+3s_1+1} - (-1)^{a_1} - i^{s_0+s_1+1}) = i^{s_0+3s_1} - i - i^{s_0+s_1} + i(-1)^{a_1}.$$

231 It shows that (17) is true for any $G(x) \in \mathbb{Z}[x]$ when $m = 4j$. When $m = 4j + 2$, (17) becomes

$$-i(1 - i^{s_0+3s_1+1} - (-1)^{a_1} + i^{s_0+s_1+1}) = i^{s_0+s_1} + i(-1)^{a_1} - i^{s_0+3s_1} - i$$

232 and

$$i(1 + i^{s_0+3s_1+1} - (-1)^{a_1} - i^{s_0+s_1+1}) = i^{s_0+s_1} - i(-1)^{a_1} - i^{s_0+3s_1} + i.$$

233 It also shows that (17) is true for any $G(x) \in \mathbb{Z}[x]$ when $m = 4j + 2$. Now we have

234 **Theorem 7.** Let $m \geq 2$ be even. For $x, y \in \mathbb{R}^m$ and any polynomial $G(x)$ with integer coefficients, the
235 kernel $e^{i\frac{\pi}{2}G(\Gamma_y)}e^{-i(x,y)}$ has a polynomial bound, i.e. there exists a constant c which is independent of
236 $G(x)$ such that

$$|e^{i\frac{\pi}{2}G(\Gamma_y)}e^{-i(x,y)}| \leq c(1 + |x||y|)^{\frac{m-2}{2}}.$$

237 At the end of this section, we give the formal generating function of the even dimensional generalized
 238 Fourier kernels for a class of polynomials. We define

$$H(x, y, a, G) = \sum_{m=2,4,6,\dots} \frac{K_{m,G}(x, y) a^{m/2-1}}{\Gamma(m/2)}.$$

239 **Theorem 8.** *Let $m \geq 2$ be even. For $x, y \in \mathbb{R}^m$ and any polynomial $G(x)$ with integer coefficients, the*
 240 *formal generating function of the even dimensional generalized Fourier kernel is given by*

$$\begin{aligned} & H(x, y, a, G) \\ = & \frac{1 - i^{G(-1)+1} - (-1)^{G'(0)} + i^{G(1)+1}}{2} \left(\cos(\sqrt{|x|^2|y|^2 - ((x, y) + a)^2}) - (x \wedge y - a) \frac{\sin \sqrt{|x|^2|y|^2 - ((x, y) + a)^2}}{\sqrt{|x|^2|y|^2 - ((x, y) + a)^2}} \right) \\ & + \frac{1 + i^{G(-1)+1} - (-1)^{G'(0)} - i^{G(1)+1}}{2} \left(\cos(\sqrt{|x|^2|y|^2 - ((x, y) - a)^2}) + (x \wedge y + a) \frac{\sin \sqrt{|x|^2|y|^2 - ((x, y) - a)^2}}{\sqrt{|x|^2|y|^2 - ((x, y) - a)^2}} \right) \\ & + \frac{1 + i^{G(-1)} + (-1)^{G'(0)} + i^{G(1)}}{2} e^{-i(x,y)-a} + \frac{1 - i^{G(-1)} + (-1)^{G'(0)} - i^{G(1)}}{2} e^{i(x,y)+a}. \end{aligned}$$

241 *Proof.* When m is even, the generalized Fourier kernel is

$$\begin{aligned} e^{i\frac{\pi}{2}G(\Gamma_y)} e^{-i(x,y)} &= \frac{1}{2} \left((1 - i^{s_0+3s_1+1} - (-1)^{a_1} + i^{s_0+s_1+1})(f_\alpha + e^{i\frac{-\pi}{2}(m-1)}g_\gamma) \right. \\ & \quad + (1 + i^{s_0+3s_1+1} - (-1)^{a_1} - i^{s_0+s_1+1})(f_\gamma + e^{i\frac{\pi}{2}(m-1)}g_\alpha) \\ & \quad \left. + (1 + i^{s_0+3s_1} + (-1)^{a_1} + i^{s_0+s_1})e^{-i(x,y)} + (1 - i^{s_0+3s_1} + (-1)^{a_1} - i^{s_0+s_1})e^{i(x,y)} \right), \end{aligned}$$

242 with $s_0 = \sum_{j=0}^{\lfloor n/2 \rfloor} a_{2j}$ and $s_1 = \sum_{j=0}^{\lfloor n/2 \rfloor} a_{2j+1}$.

243 By $s_0 + 3s_1 \equiv s_0 - s_1 \equiv G(-1) \pmod{4}$, $s_0 + s_1 = G(1)$, $a_1 = G'(0)$ and because $\frac{\Gamma(m/2)}{2}(f_\alpha +$
 244 $e^{i\frac{-\pi}{2}(m-1)}g_\gamma)$ and $\frac{\Gamma(m/2)}{2}(f_\gamma + e^{i\frac{\pi}{2}(m-1)}g_\alpha)$ are the Clifford-Fourier kernel $K^{\frac{-\pi}{2}}$ and $K^{\frac{\pi}{2}}$ in the Laplace
 245 domain, the result follows from the generating function of Clifford-Fourier kernel, see [5] Theorem 8. \square

246 **Remark 6.** *When $G(x) = x$, we get the generating function of the Clifford-Fourier kernel.*

247 For the case that the coefficients of $G(x)$ are not integers but fractions, we write $G_1(x) = cG(x)$ in
 248 which c is the least common multiple of each denominator of $G(x)$. So $G_1(x)$ is a polynomial with integer
 249 coefficients. We only need to compute $e^{i\frac{\pi}{2c}G_1(\Gamma_y)}f(y)$ and $e^{i\frac{\pi}{2c}G_1(\Gamma_y)}g(y)$. The same method will also
 250 work but f and g split into $4c$ parts.

251 4 Conclusion

252 By working on the Laplace domain, we found explicit expressions for the generalized kernel. For even
 253 dimension, we obtained the closed expression. **As the bound of the kernel is important to see in which**
 254 **function space the transform is well defined, we moreover determined which polynomials G give rise**
 255 **to polynomially bounded kernels for all dimensions and only even dimensions. When the kernel has**
 256 **polynomial bound, the transform can be proved to be well defined on the Schwartz function space and**
 257 **the transform is a continuous operator. Following a similar discussion in Section 6 of [7], we can get the**
 258 **existence of the inversion formula for this kind transform.** Also we determined the generating function
 259 corresponding to a fixed polynomial G . **We point out that the closed kernel of odd dimension is still**
 260 **an open problem since 2005 and deserves more study.** We think it is interesting to develop the Laplace
 261 method on the hyperbolic space or sphere to compute the closed kernel of the hypercomplex Fourier
 262 transform.

263 **First application of a quaternion Fourier transform to color images was reported in 1996 by Sangwine**
 264 **et al. The advantage of Fourier-type transforms of quaternionic signals over the classical Fourier transform**

265 is that the kernel through which they act are quaternion-valued and the transforms therefore "mix the
266 channels" rather than acting on each channel separately. The generalized transform we studied here will
267 also "mix the channels" and yet still has a simple inversion and Plancherel theorem.

268 **Acknowledgments** H. De Bie is supported by the UGent BOF starting grant 01N01513. P. Lian is supported by
269 a scholarship from Chinese Scholarship Council (CSC) under the CSC No. 201406120169.
270

271 References

- 272 [1] Brackx F, De Schepper N, Kou K I, Sommen F. The Mehler formula for the generalized CliffordHermite polynomials.
273 *Acta Math. Sin. (Engl. Ser.)* 2007, 23(4): 697-704.
- 274 [2] Brackx F, De Schepper N, Sommen F. The Clifford-Fourier transform. *J. Fourier Anal. Appl.* 2005, 11(6): 669-681.
- 275 [3] Brackx F, De Schepper N, Sommen F. The two-dimensional Clifford-Fourier transform. *J. Math. Imaging Vision*, 2006,
276 26(1-2): 5-18.
- 277 [4] Brackx F, Hitzer E, Sangwine S. History of quaternion and Clifford-fourier transforms and wavelets. *Quaternion and*
278 *Clifford fourier transforms and wavelets. In Trends in Mathematics 27.* 2013, XI-XXVII.
- 279 [5] Constales D, De Bie H, Lian P. A new construction of the Clifford-Fourier kernel. To appear in *J. Fourier Anal. Appl.*
- 280 [6] Craddock M J, Hogan J A. The Fractional Clifford-Fourier Kernel. *J. Fourier Anal. Appl.* 2013, 19(4): 683-711.
- 281 [7] De Bie H, Oste R, Van der Jeugt J. Generalized Fourier transforms arising from the enveloping algebras of $\mathfrak{sl}(2)$ and
282 $\mathfrak{osp}(1|2)$. *Int. Math. Res. Not. IMRN*, 2015: rnv293.
- 283 [8] De Bie H, Xu Y. On the Clifford-Fourier transform. *Int. Math. Res. Not. IMRN*, 2011, no.22, 5123-5163.
- 284 [9] Delanghe R, Sommen F, Soucek V. *Clifford algebra and spinor-valued functions, volume 53 of Mathematics and its*
285 *Applications*. Kluwer Academic Publishers Group, Dordrecht 1992.
- 286 [10] Eelbode D, Hitzer E. Operator Exponentials for the Clifford Fourier Transform on Multivector Fields in Detail. *Adv.*
287 *Appl. Clifford Al*, 2015, 1-16.
- 288 [11] Erdélyi, Arthur, ed. *Tables of integral transforms*. Vol. 1. New York: McGraw-Hill, 1954.
- 289 [12] Frappat L, Sciarrino A, Sorba P. *Dictionary on Lie algebras and superalgebras*. San Diego: Academic Press, 2000.
- 290 [13] Ghobber S, Jaming P. Uncertainty principles for integral operators. *Studia Math.* 2014, 220, 197-220.
- 291 [14] Howe R. The oscillator semigroup. *Proc. Symp. Pure Math.* 1988, 48: 61-132.
- 292 [15] Saïd S B, Kobayashi T, Ørsted B. Laguerre semigroup and Dunkl operators. *Compos. Math.* 2012, 148(04): 1265-1336.
- 293 [16] Schiff J L. *The Laplace transform: theory and applications*. Springer, 1999.