

26 **Abstract**

27 Methicillin-resistant *Staphylococcus aureus* (MRSA) is infrequently reported in mastitis. Yet,
28 as in many other countries, the prevalence of methicillin resistance among *S. aureus* from
29 mastitis is currently unknown in Belgium.

30 To elucidate this, the presence of *mecA* was investigated in 118 *S. aureus* strains originating
31 from diagnostic mastitis milk samples from 118 different farms experiencing *S. aureus*
32 mastitis. MRSA strains were characterized by disk diffusion susceptibility testing, *spa*-typing,
33 MLST and SCC*mec*-typing. In an additional study, four MRSA-positive farms were selected
34 to assess the in-herd prevalence of MRSA, by sampling all cows in lactation. Isolated MRSA
35 strains were similarly characterized.

36 The *mecA* gene was detected in eleven (9.3%) of the 118 *S. aureus* isolates, indicating that
37 nearly 10% of the Belgian farms suffering from *S. aureus* mastitis have an MRSA problem.

38 The in-herd prevalence varied between 0% and 7.4%. Characterization of the MRSA strains
39 showed that they were all resistant to tetracycline. Additional resistances to macrolides,
40 lincosamides and aminoglycosides were frequently detected. The strains were ST398, *spa*-
41 types t011 or t567 and had SCC*mec*-types IVa or V, proving they belong to the emerging
42 livestock-associated MRSA (LA-MRSA) strains of CC398.

43 Our study shows that after detection in Belgian pigs, horses and poultry, LA-MRSA has also
44 attained Belgian cattle. It is the first report on frequent isolation of LA-MRSA from bovine
45 infections. As the in-herd isolation rate resembles that of regular *S. aureus* in farms
46 experiencing *S. aureus* mastitis, the multi-resistance of LA-MRSA strains may cause future
47 treatment problems.

48 **Keywords**

49 methicillin-resistant *Staphylococcus aureus*, MRSA, mastitis, Belgium, ST398, multidrug
50 resistance

51 **Introduction**

52 *Staphylococcus aureus* is a major pathogen in dairy cattle mastitis (Waage et al., 1998;
53 Tenhagen et al., 2006; Piepers et al., 2007). Resistance of *S. aureus* to antimicrobial agents
54 can complicate treatment of its infections (Lowy, 2003). For treatment of mastitis, methicillin
55 resistance, which is caused by the expression of the *mecA* gene, is of particular interest.
56 Indeed, this mechanism confers resistance to almost all types of β -lactam antibiotics active
57 against *S. aureus*, and these antibiotics are still frequently used in mastitis treatment (Sawant
58 et al., 2005). However, methicillin-resistant *Staphylococcus aureus* (MRSA) has never been
59 important in mastitis. After the very first report of MRSA in mastitis in 1972 (Devriese et al.,
60 1972), MRSA has been described in mastitis only occasionally (Lee, 2003; Kwon et al., 2005;
61 Lee, 2006; Juhász-Kaszanyitzky et al., 2007; Moon et al., 2007; Hendriksen et al., 2008).
62 From such studies, it seems that the prevalence of MRSA in mastitis is generally low. Yet,
63 data on MRSA in mastitis need to be assessed carefully, as there are often ambiguities on
64 presence of *mecA*, level of investigation and origin of the detected MRSA strains.
65 Recently, a specific MRSA clone, CC398, has been found associated with pigs, veal calves,
66 broiler chickens, companion animals and people in close contact with livestock. MRSA of this
67 type, called Livestock-Associated MRSA (LA-MRSA), typically has closely related *spa*-types
68 (de Neeling et al., 2007; Denis et al., 2009), carries mostly SCC*mec*-types IVa and V (Witte et
69 al., 2007; Van den Eede et al., 2009) and cannot be typed with PFGE using *Sma*I digestion
70 (Bens et al., 2006). In addition, LA-MRSA shows resistance against tetracycline and, to a
71 lesser extent, macrolides, lincosamides, aminoglycosides and fluoroquinolones (Witte et al.,
72 2007). Generally LA-MRSA lacks common virulence factors found in other MRSA
73 (Monecke et al., 2007; Walther et al., 2009). This is remarkable because, although
74 infrequently compared to colonization, LA-MRSA has been isolated from infections, of both
75 animals and humans (e.g. Hermans et al., 2008; Krziwanek et al., 2009). To our knowledge,

76 so far only one study has reported on the isolation of MRSA ST398 from a case of mastitis
77 (Monecke et al., 2007).

78 We performed two studies to assess the role of MRSA in Belgian *S. aureus* mastitis. In a first
79 study we investigated how many *S. aureus* isolated from mastitis were resistant to methicillin.
80 Second, we investigated the in-herd prevalence of MRSA in Belgian herds where cows were
81 previously shown to suffer from MRSA mastitis.

82 **Methods**

83 *1. Methicillin resistance in S. aureus isolated from mastitis*

84 **Strains**

85 From November 2006 through April 2007, the regional veterinary laboratories were asked to
86 send us a representative isolate from all farms on which an *S. aureus*-mastitis problem was
87 detected. Care was taken to include only one strain per visited farm. As such, a collection of
88 118 non duplicate isolates of *S. aureus*, originating from cases of subclinical or clinical
89 mastitis from different farms was obtained.

90 **DNA extraction**

91 An Eppendorf cup (Eppendorf, Germany) containing a 500 µl Brain Heart Infusion broth
92 (BHI) (BioRad, France) overnight pure culture was centrifuged for 3.0 min at approx.
93 20,000 x g, at room temperature. After removal of the supernatant, 45 µL of sterile, distilled
94 water and 5 µL of a 1 mg/mL lysostaphin (Sigma-Aldrich, USA) solution at 4°C were
95 thoroughly mixed with the pellet of cells. After incubation for 10 min at 37°C, 45 µL of
96 sterile, distilled water, 5 µL of a 2 mg/mL proteinase K (Merck, Germany) solution at 4°C
97 and 150 µL of tris-HCl of 0.1 M at pH 8.0 were added. The resulting solution was incubated
98 for 10 min at 60°C, followed by 5 min at 100°C and then centrifuged for 5 min at approx.
99 20,000 x g, at room temperature. DNA was stored at -20°C until use.

100 **Identification of MRSA**

101 A triplex PCR, targeting a *Staphylococcus*-specific 16S rRNA sequence, the *mecA* gene and
102 the *S. aureus* specific region of the thermonuclease gene (*nuc*), was performed as previously
103 described (Maes et al., 2002). The amplified DNA fragments were separated by
104 electrophoresis on a 2% agarose (Sigma-Aldrich, USA) gel stained with SYBR Safe DNA gel
105 stain (Invitrogen, USA), for 2 h at 80V, using an O'RangeRuler 100bp DNA ladder
106 (Fermentas, Germany).

107 **Characterization of MRSA**

108 *Susceptibility testing*

109 Strains proven to be MRSA were tested for susceptibility to non β -lactam antimicrobial
110 agents, by using the disk diffusion method. A panel of 16 antimicrobial agents was used:
111 chloramphenicol, gentamicin, kanamycin, tobramycin, fucidic acid, erythromycin, tylosin,
112 lincomycin, linezolid, quinupristin + dalfopristin, mupirocin, ciprofloxacin, tetracycline,
113 rifampicin, sulfonamides and trimethoprim (NeoSensitabs, Rosco, Denmark). Results were
114 recorded after 24h incubation at 37°C and interpreted according to the directions for use of
115 Rosco with the method described by the CLSI guidelines (document M31-A3).

116 *Spa-typing*

117 Of all MRSA strains, the polymorphic X-region of the *Staphylococcus* protein A (*spa*) gene
118 was amplified according to the Ridom StaphType standard protocol
119 (www.ridom.de/staphtype). Amplicons were purified with a Nucleospin Extract II kit
120 (Macherey-Nagel, Germany) and then sequenced using the same primers. The sequenced
121 DNA was then run on a CEQ 8000 Genetic Analysis System (Beckman Coulter, United
122 Kingdom) according to the manufacturer's instructions. The resulting *spa*-types were assigned
123 by using the Ridom StaphType software package (Ridom GmbH, Germany).

124 *MLST*

125 Multi Locus Sequence Typing was performed on all MRSA strains. In short, seven household
126 genes of *S. aureus* were amplified using primers previously described (Enright et al., 2000).

127 Amplicons were purified with a Nucleospin Extract II kit (Macherey-Nagel, Germany) and
128 then sequenced using the same primers. The sequenced DNA was then run on a CEQ 8000
129 Genetic Analysis System (Beckman Coulter, United Kingdom) according to the
130 manufacturer's instructions. Allele numbers and sequence type (ST) were assigned by using
131 the *S. aureus* MLST website (<http://saureus.mlst.net>).

132 *SCCmec*-typing

133 The *SCCmec* type was determined using three different sets of primers (Oliveira and de
134 Lencastre, 2002; Zhang et al., 2005; Milheiriço et al., 2007). For differentiation among
135 *SCCmec* types I–IV we used all the primers described by Oliveira and de Lencastre (2002).
136 The PCR mix consisted of 25 µL of *Taq* PCR Master Mix (Qiagen GmbH, Germany), 4 µL of
137 H₂O and 16 µL of the primers, in the reported concentration. To this mix 5 µL DNA was
138 added.

139 For subtyping *SCCmec* of type IV, we used the primers described by Milheiriço et al. (2007).
140 The PCR-mix consisted of 25 µL of *Taq* PCR Master Mix (Qiagen GmbH, Germany), 6.4 µL
141 of H₂O, 0.2 µM of primers J IVa forward (F) and reverse (R), 0.2 µM of J IVb F and R, 0.4
142 µM of *ccr* B2 F and J IVc F and R, 0.8 µM of *ccr*B2 R, J IVd F and R, 0.9 µM of J IVg F and
143 R, and 0.9 µM of J IVh F and R. To the mix 5 µL DNA was added.

144 A third set, meant to detect *SCCmec*-type V and to have a control for *SCCmec*-types IVb,
145 IVc, IVe and IVf, was based on the method described by Zhang et al. (2005). The PCR mix
146 consisted of 25 µL of *Taq* PCR Master Mix (Qiagen GmbH, Germany), 10.4 µL of H₂O, 0.6
147 µM of primers Type V F and R, 0.8 µM of Type IVc F and R, and 1.0 µM of Type IVb F and
148 R. To the mix 5 µL DNA was added.

149 We used the same PCR program for all three sets: an initial denaturation of 4 min at 94°C, 35
150 cycles of denaturation at 94°C for 30 s, annealing at 53°C for 30 s and extension at 72°C for
151 1 min, followed by a final extension for 4 min at 72°C.

152 **2. MRSA in-herd prevalence**

153 From the results of the first study, four MRSA positive farms were selected for investigation
154 of the in-herd prevalence of MRSA, defined as the number of MRSA-positive cows relative to
155 the total number of lactating cows present in the specific farm. A randomly chosen fifth farm
156 volunteered to serve as control (Table 2). Milk samples were taken from each quarter. All
157 sampling was done by the same person, from February 2008 through April 2008.

158 Samples were immediately transported to the Veterinary and Agrochemical Research Center
159 (VAR), where each sample was plated on Columbia Colistine Aztreonam Plates (CAP)
160 supplemented with 5% sheep blood (Oxoid, Germany) and on chromID MRSA plates
161 (Biomérieux, France). Suspected *S. aureus* or MRSA colonies were purified. Pure colonies
162 were then subjected to the MRSA triplex PCR, as described above. Strains identified as
163 MRSA were characterized by susceptibility testing, *spa*-typing, MLST and SCC*mec*-typing,
164 as described above.

165 **Results**

166 **1 – Methicillin resistance in *S. aureus* isolated from mastitis**

167 **Detection of MRSA**

168 All 118 isolates phenotypically identified as *S. aureus* were confirmed to be *S. aureus* by the
169 triplex PCR. A total of 11 isolates (9.3%) contained *mecA* (Table 1). Two MRSA originated
170 from clinical mastitis, the other nine from subclinical mastitis (Table 1).

171 **Antimicrobial susceptibility testing**

172 Antibiotic resistance patterns of the 11 MRSA strains are shown in Table 1. Nine of them
173 showed additional resistance to at least two different antibiotics. All strains were resistant to
174 tetracycline; nine were resistant to trimethoprim, seven to aminoglycosides and lincomycin,
175 five to macrolides and two to ciprofloxacin (Table 1). No resistance was detected to the other
176 antimicrobial agents tested.

177 **MLST, *spa*- and SCC*mec*-typing**

178 Ten strains were *spa*-type t011. One strain had a different yet related *spa*-type, t567 (Table 1).
179 All MRSA strains were ST398 (Table 1). Five strains had *SCCmec*-type IVa, and five had
180 *SCCmec*-type V. The *SCCmec*-type of one strain could not be determined with the different
181 sets of primers we used (Table 1).

182 **2 – MRSA in-herd prevalence**

183 **Identification of MRSA**

184 The percentage of cows carrying MRSA in their milk varied between 0% and 7.4% (Table 2).
185 Quarter level prevalence ranged from 0% to 1.98% (Table 2). Three of the four selected farms
186 were positive. MRSA could not be detected in one farm previously found positive nor in the
187 control farm (Table 2).

188 One cow from the first farm carried MRSA in three of her quarters. In all other positive cows,
189 MRSA was found in only one quarter, resulting in 14 isolates in total (Table 3). Most isolates
190 were found in the right-hind (6 isolates) and right-front (4 isolates) quarter (Table 3). Of the
191 11 cows that had MRSA in only one quarter, nine of the MRSA isolates were found in one of
192 the hind-quarters.

193 **Antimicrobial susceptibility testing**

194 All six strains isolated from the first farm had the same susceptibility profile (Table 3). They
195 were all resistant to tetracycline, the tested macrolides and lincomycin, and were susceptible
196 to all other antimicrobial agents tested.

197 In the second farm, two out of five strains were resistant to trimethoprim, tetracycline and the
198 tested aminoglycosides, and susceptible to all other antimicrobial agents tested. The three
199 other strains had additional resistances to the tested macrolides and lincomycin (Table 3).

200 The three strains from the third farm were resistant to the tested aminoglycosides, macrolides,
201 lincomycin, tetracycline and trimethoprim (Table 3). They were susceptible to the other
202 antimicrobial agents tested.

203 **MLST, *spa*- and *SCCmec*-typing**

204 All MRSA strains showed *spa*-type t011 (Table 3). The strains originating from the first farm
205 had *SCCmec* type V, while all strains isolated from the cows of the other two farms had
206 *SCCmec* type IVa (Table 3). MLST was performed on one representative MRSA strain per
207 farm. As strains from farm 2 showed two different resistance profiles, one representative
208 strain from each profile was tested. The four strains tested all were ST398 (Table 3).

209 **Discussion**

210 The prevalence of methicillin resistance in *S. aureus* isolated from mastitis in our first study is
211 unexpectedly high. In the abundance of studies investigating the antibiotic resistance of
212 mastitis pathogens, few reports have noted a substantial occurrence of methicillin resistance,
213 meaning MRSA is usually negligible as a mastitis pathogen (Hendriksen et al., 2008).
214 However, we found nearly 10% of our 118 *S. aureus* strains to be MRSA. This means that
215 nearly 10% of the Belgian farms experiencing *S. aureus* mastitis is affected by MRSA.
216 Reports can be found in which a higher prevalence of MRSA among *S. aureus* isolated from
217 mastitis cases is described. In Turkey, Turutoglu et al. (2006) found 18 out of 103 (17.5%) *S.*
218 *aureus* isolates from mastitis milk samples to be MRSA. However, they did not mention
219 whether all strains were collected from different farms experiencing *S. aureus* mastitis. In
220 addition, their detection method was limited to phenotypic disk diffusion testing. Performing
221 only phenotypic tests has previously been shown to lead to false positive or false negative
222 results (Murakami et al., 1991; De Oliveira et al., 1999). Generally it is now accepted that
223 checking for the presence of *mecA* is the most reliable method for detection of methicillin
224 resistance, and staphylococci carrying *mecA* should be regarded as resistant to almost all types
225 of β -lactam antibiotics (CLSI guidelines, M31-A3). Consequently, to accurately assess our
226 results, only other reports in which *mecA* was proven to be present should be considered. Still,
227 even then, it remains difficult to make viable comparisons, due to differences in sampling
228 methodology or a lack of information on the source of the strains. For example, two South

229 Korean studies did not mention exactly how many of their samples originated from mastitis
230 (Lee, 2003; Lee, 2006). A Hungarian study sampled only a single farm (Juhász-Kaszanyitzky
231 et al., 2007). In two other studies from South Korea, the data involved quarter-level results
232 (Kwon et al., 2005; Moon et al., 2007).

233 Despite these difficulties to fully assess our results, it must be acknowledged that the MRSA
234 prevalence we found is quite high. However, some other remarks should be made. First, the
235 burden of MRSA for Belgian milk production cannot be assessed, because we have no data on
236 the total number of farms that were visited during the sampling period. Also, while our study
237 allows us to estimate the importance of methicillin resistance in Belgian *S. aureus* mastitis,
238 we cannot judge the importance of MRSA for mastitis as a whole. A hint to address the latter
239 can be found in a recent study that investigated the importance of *S. aureus* in Belgian
240 mastitis. It was found that *S. aureus* was the most prevalent species in Belgian quarter milk
241 samples from subclinical mastitis, with 25% of culture-positive quarter samples with a
242 geometric mean composite somatic cell count of $\geq 250\ 000$ cells/ml harboring *S. aureus*
243 (Piepers et al., 2007). Regarding this, our result is certainly quite worrying.

244 Another important fact is presented by our typing data. All our strains had characteristics
245 typical for the emerging livestock-associated MRSA CC398 strains. Consequently, it seems
246 that our findings should rather be regarded as a further expansion of the host range of the
247 CC398 MRSA clone than as an indication of a generally increasing incidence of methicillin
248 resistance in mastitis-associated *S. aureus*. This should however not be less worrying.

249 In addition to its resistance against all β -lactam antibiotics, which are still the most used
250 antimicrobial agents in the treatment of mastitis, the typical antibiotic resistances of LA-
251 MRSA also include some other antibiotics used to treat or prevent mastitis, such as
252 aminoglycosides and macrolides (Sawant et al., 2005). This could lead to serious treatment
253 problems. Moreover, in our second study we found that the in-herd prevalence of LA-MRSA

254 ranged between 0% and 7.4%. In the farms where MRSA was found, it varied from 3.9% to
255 7.4%, with a corresponding quarter level prevalence of 0.97% to 1.98%. This resembles the
256 in-herd quarter level prevalence of *S. aureus* described earlier in a cross-sectional collection
257 of Belgian milk samples (Piepers et al., 2007), suggesting that, considering its spread in
258 farms, LA-MRSA behaves similar to regular mastitis causing *S. aureus*. The possibility that
259 LA-MRSA could become equally important in mastitis as normal *S. aureus* should thus be
260 thoroughly investigated. Unfortunately, we have no data on the individual health status of the
261 cows from which MRSA was isolated in our second study, so we cannot state that the LA-
262 MRSA strains we found were actually involved in mastitis. As it was shown that within-cow
263 transmission between quarters likely occurs in *S. aureus* mastitis (Barkema et al., 1997), the
264 fact that 11 of the 12 cows carried LA-MRSA in only one quarter could mean that the isolates
265 concerned only contaminants. However, *S. aureus* infection of only one quarter also certainly
266 exists (Barkema et al., 1997). Moreover, *S. aureus* was shown to more frequently infect the
267 right and hind quarters (Barkema et al., 1997; Barkema et al., 2006). Of the 11 single-quarter
268 LA-MRSA isolates we found, 10 originated from right quarters and nine from hind quarters.
269 Considering also our first study, which clearly showed the capacity of LA-MRSA to cause
270 mastitis, the actual presence of LA-MRSA in Belgian mastitis should urgently be studied in
271 more depth, in order to profoundly assess its possible burden.

272 LA-MRSA has been reported only once before in mastitis in cows, one LA-MRSA strain that
273 was found among 128 *S. aureus* isolated from German mastitis cases (Monecke et al., 2007).
274 While this strain was *spa*-type t034, our strains were *spa*-types t011 and t567. It thus seems
275 unlikely that a specific subclone of LA-MRSA is associated with mastitis, but more research
276 is required to confirm this. Until now, it is also unclear whether LA-MRSA has an actual
277 reservoir in dairy cattle. Whereas veal calves have been found carrying LA-MRSA in the

278 Netherlands (Mooij et al., 2007) and Belgium (unpublished data), the colonization capacity of
279 LA-MRSA in milking cows has not yet been investigated.

280 The presence of LA-MRSA in infections has been reported substantially less frequent than
281 carriage, and has only been described occasionally in pigs (van Duijkeren et al., 2007), horses
282 (Hermans et al., 2008; Loeffler et al., 2009), humans (e.g. Krziwanek et al., 2009) and a dog
283 (Witte et al., 2007). Our findings thus seem to add new proof of a certain pathogenic potential
284 of LA-MRSA. Remarkably, many common virulence factors, including those considered to
285 be involved in mastitis, such as toxic shock syndrome toxin-1 (tsst-1), haemolysins and
286 enterotoxins (Matsunaga et al., 1993), have been shown to be largely absent in LA-MRSA
287 (Monecke et al., 2007; Walther et al., 2009). However, as we did not check for the presence of
288 virulence factors in our strains, the significance of our data regarding the pathogenic potential
289 of LA-MRSA is hard to assess. Yet, in addition to the other reports on LA-MRSA associated
290 with infections, our findings urge for further research into the virulence capacities of LA-
291 MRSA.

292 **Conclusions**

293 We found an unusual high prevalence of MRSA in Belgian cases of subclinical and clinical *S.*
294 *aureus* mastitis in cows. All strains belonged to the CC398 clone, which, seen its multi-
295 resistance, may lead to treatment problems. Future research is warranted to assess the actual
296 spread and corresponding burden that LA-MRSA may pose for dairy cattle farming and to
297 elucidate which virulence factors are involved.

298 **Acknowledgments**

299 DGZ and ARSIA are acknowledged for sending us the strains for analysis. Dr. Luc
300 Demeulemeester is acknowledged for helping us contacting the MRSA-positive farms. We
301 also thank Danielle Vanderghenst for her skilled technical assistance. These studies were
302 funded by the Belgian Antibiotic Policy Coordination Committee (BAPCOC) and by the

303 Belgian Federal Public Service of Public Health, Food Chain Safety and Environment, project
304 number RF-6189 MRSA.

305 **References**

- 306 1. Barkema, H.W., Schukken, Y.H., Lam, T.J., Galligan, D.T., Beiboer, M.L., Brand, A.,
307 1997. Estimation of interdependence among quarters of the bovine udder with subclinical
308 mastitis and implications for analysis. *J. Dairy Sci.* 80, 1592-1599.
- 309 2. Barkema, H.W., Schukken, Y.H., Zadoks, R.N., 2006. Invited review: The role of cow,
310 pathogen, and treatment regimen in the therapeutic success of bovine *Staphylococcus*
311 *aureus* mastitis. *J. Dairy Sci.* 89, 1877-1895.
- 312 3. Bens, C.C.P.M., Voss, A., Klaassen, C.H.W., 2006. Presence of a novel DNA methylation
313 enzyme in methicillin-resistant *Staphylococcus aureus* isolates associated with pig
314 farming leads to uninterpretable results in standard pulsed-field gel electrophoresis
315 analysis. *J. Clin. Microbiol.* 44, 1875-1876.
- 316 4. de Neeling, A.J., van den Broek, M.J.M., Spalburg, E.C., van Santen-Verheuevel, M.G.,
317 Dam-Deisz, W.D.C., Boshuizen, H.C., van de Giessen, A.W., van Duijkeren, E.,
318 Huijsdens, X.W., 2007. High prevalence of methicillin resistant *Staphylococcus aureus* in
319 pigs. *Vet. Microbiol.* 122, 366-372.
- 320 5. De Oliveira, A.P., Watts, J.L., Salmon, S.A., Aarestrup, F.M., 1999. Antimicrobial
321 susceptibility of *Staphylococcus aureus* isolated from bovine mastitis in Europe and the
322 United States. *J. Dairy Sci.* 83, 855-862.
- 323 6. Denis, O., Suetens, C., Hallin, M., Catry, B., Ramboer, I., Dispas, M., Willems, G., Gordts, B.,
324 Butaye, P., Struelens, M., 2009. Methicillin-resistant *Staphylococcus aureus* in swine farm
325 personnel, Belgium. *Emerg. Infect. Dis.* 15, 1098-1101.
- 326 7. Devriese, L.A., Van Damme, L.R., Fameree, L., 1972. Methicillin- (cloxacillin)-resistant
327 *Staphylococcus aureus* strains isolated from bovine mastitis cases. *Zentralbl.*
328 *Veterinarmed. B.* 19, 598-605.

- 329 8. Enright, M.C., Day, N.P.J., Davies, C.E., Peacock, S.J., Spratt, B.G., 2000. Multilocus
330 Sequence Typing for characterization of methicillin-resistant and methicillin-susceptible
331 clones of *Staphylococcus aureus*. J. Clin. Microbiol. 38, 1008-1015.
- 332 9. Hendriksen, R.S., Mevius, D.J., Schroeter, A., Teale, C., Meunier, D., Butaye, P., Franco,
333 A., Utinane, A., Amado, A., Moreno, M., Greko, C., Stärk, K., Berghold, C., Myllyniemi,
334 A.-L., Wasyl, D., Sunde, M., Aarestrup, F.M., 2008. Prevalence of antimicrobial
335 resistance among bacterial pathogens isolated from cattle in different European countries:
336 2002-2004. Acta Vet. Scan. 50, 28.
- 337 10. Hermans, K., Lipinska, U., Denis, O., Deplano, A., Struelens, M.J., Nemati, M., Pasmans,
338 F., Butaye, P., Martens, A., Deprez, P., Haesebrouck, F. 2008. MRSA clone ST398-
339 SCCmec IV as a cause of infections in an equine clinic. Vlaams Diergeneeskundig
340 Tijdschrift; 77, 429-433.
- 341 11. Juhász-Kaszanyitzky, É., Jánosi, S., Somogyi, P., Dán, Á., van der Graaf-van Bloois, L.,
342 van Duijkeren, E., Wagenaar, J.A., 2007. MRSA transmission between cows and humans.
343 Emerg. Infect. Dis. 13, 630-632.
- 344 12. Krziwanek, K., Metz-Gercek, S., Mittermayer H., 2009. Methicillin-resistant
345 *Staphylococcus aureus* ST398 from human patients, upper Australia. Emerg. Infect. Dis.
346 15, 766-769.
- 347 13. Kwon, N.H., Park, K.T., Moon, J.S., Jung, W.K., Kim, S.H., Kim, J.M., Hong, S.K., Koo,
348 H.C., Joo, Y.S., Park, Y.H., 2005. Staphylococcal cassette chromosome *mec* (SCCmec)
349 characterization and molecular analysis for methicillin-resistant *Staphylococcus aureus*
350 and novel SCCmec subtype IVg isolated from bovine milk in Korea. J. Antimicrob.
351 Chemother. 56, 624-632.

- 352 14. Lee, J.H., 2003. Methicillin (oxacillin)-resistant *Staphylococcus aureus* strains isolated
353 from major food animals and their potential transmission to humans. *App. Environ.*
354 *Microbiol.* 69, 6489-6494.
- 355 15. Lee, J.H., 2006. Occurrence of methicillin-resistant *Staphylococcus aureus* strains from
356 cattle and chicken, and analysis of their *mecA*, *mecRI* and *mecI* genes. *Vet. Microbiol.*
357 114, 155-159.
- 358 16. Loeffler, A., Kearns, A.M., Ellington, M.J., Smith, L.J., Unt, V.E., Lindsay, J.A., Pfeiffer,
359 D.U., Lloyd, D.H., 2009. First isolation of MRSA ST398 from UK animals: a new
360 challenge for infection control teams? *J. Hosp. Infect.* 72, 269-271.
- 361 17. Lowy, F.D., 2003. Antimicrobial resistance: the example of *Staphylococcus aureus*. *J.*
362 *Clin. Invest.* 111, 1265-1273.
- 363 18. Maes, N., Magdalena, J., Rottiers, S., De Gheldre, Y., Struelens, M.J., 2002. Evaluation of
364 a triplex PCR assay to discriminate *Staphylococcus aureus* from coagulase-negative
365 staphylococci and determine methicillin-resistance from blood cultures. *J. Clin. Microbiol.*
366 40, 1514-1517.
- 367 19. Matsunaga, T., Kamata, S., Kakiichi, N., Uchida, K., 1993. Characteristics of
368 *Staphylococcus aureus* isolated from peracute, acute and chronic bovine mastitis. *J. Vet.*
369 *Med. Sci.* 55, 297-300.
- 370 20. Milheiriço, C., Oliveira, D.C., de Lencastre, H., 2007. Multiplex PCR strategy for
371 subtyping the Staphylococcal Cassette Chromosome *mec* type IV in methicillin-resistant
372 *Staphylococcus aureus*: 'SCC*mec* IV multiplex'. *J. Antimicrob. Chemother.* 60, 42-48.
- 373 21. Monecke, S., Kuhnert, P., Hotzel, H., Slickers, P., Ehricht, R., 2007. Microarray based
374 study on virulence-associated genes and resistance determinants of *Staphylococcus aureus*
375 isolates from cattle. *Vet. Microbiol.* 125, 128-140.

- 376 22. Mooij, R., Toelstra, A., Jenkins, J.M., Thijsen, S.F.T., 2007. MRSA in calves. *English*
377 *abstract Infectieziekten bulletin* 18, 234-236.
- 378 23. Moon, J.-S., Lee, A.-R., Kang, H.-M, Lee, E.-S., Kim, M.-N., Paik, Y.H., Park, Y.H., Joo,
379 Y.S., Koo, H.C., 2007. Phenotypic and genetic antibiogram of methicillin-resistant
380 Staphylococci isolated from bovine mastitis in Korea. *J. Dairy Sci.* 90, 1176-1185.
- 381 24. Murakami, K., Minamide, W., Wada, K., Nakamura, E., Teraoka, H., Watanabe, S., 1991.
382 Identification of methicillin-resistant strains of staphylococci by polymerase chain
383 reaction. *J. Clin. Microbiol.* 29, 2240-2244.
- 384 25. Oliveira, D.C., de Lencastre, H., 2002. Multiplex PCR strategy for rapid identification of
385 structural types and variants of the *mec*-element in methicillin-resistant *Staphylococcus*
386 *aureus*. *Antimicrob. Agents Chemother.* 46, 2155-2161.
- 387 26. Piepers, S., De Meulemeester, L., de Kruif, A., Opsomer, G., Barkema, H.W., De
388 Vlieghe, S., 2007. Prevalence and distribution of mastitis pathogens in subclinically
389 infected dairy cows in Flanders, Belgium. *J. Dairy Res.* 74, 478-483.
- 390 27. Sawant, A.A., Sordillo, L.M., Jayarao, B.M., 2005. A survey on antibiotic usage in dairy
391 herds in Pennsylvania. *J. Dairy Sci.* 88, 2991-2999.
- 392 28. Tenhagen, B.-A., Köster, G., Wallmann, J., Heuwieser, W, 2006. Prevalence of mastitis
393 pathogens and their resistance against antimicrobial agents in dairy cows in Brandenburg,
394 Germany. *J. Dairy Sci.* 89, 2542-2551.
- 395 29. Turutoglu, H., Ercelik, S., Ozturk, D., 2006. Antibiotic resistance of *Staphylococcus*
396 *aureus* and coagulase-negative staphylococci isolated from bovine mastitis. *Bull. Vet. Inst.*
397 *Pulawy* 50, 41-45.
- 398 30. Van Den Eede, A., Martens, A., Lipinska, U., Struelens, M., Deplano, A., Denis, O.,
399 Haesebrouck, F., Gasthuys, F., Hermans, K., 2009. High occurrence of methicillin-

- 400 resistant *Staphylococcus aureus* ST398 in equine nasal samples. Vet. Microbiol. 133, 138-
401 144.
- 402 31. van Duijkeren, E., Jansen, M.D., Flemming, S.C., de Neeling, H., Wagenaar, J.A.,
403 Schoormans, A.H.W., van Nes, A., Fluit, A.C., 2007. Methicillin-resistant *Staphylococcus*
404 *aureus* in pigs with exudative epidermitis. Emerg. Infect. Dis. 13, 1408-1410.
- 405 32. Waage, S., Mørk, T., Røros, A., Aasland, D., Hunshamar, A., Odegaard, A.S., 1998.
406 Bacteria associated with clinical mastitis in dairy heifers. J. Dairy Sci. 82, 712-719.
- 407 33. Walther, B., Monecke, S., Ruscher, C., Friedrich, A.W., Ehricht, R., Slickers, P., Soba, A.,
408 Wleklinski, C.-G., Wieler, L.H., Lübke-Becker, A., 2009. Comparative molecular analysis
409 substantiates zoonotic potential of equine methicillin-resistant *Staphylococcus aureus*. J.
410 Clin. Microbiol. 47, 704-710.
- 411 34. Witte., W., Strommenger, B., Stanek, C., Cuny, C., 2007. Methicillin-resistant
412 *Staphylococcus aureus* ST398 in humans and animals, Central Europe. Emerg. Infect. Dis.
413 13, 255-258.
- 414 35. Zhang, K., McClure, J.-A., Elsayed, S., Louie, T., Conly, J.M., 2005. Novel multiplex
415 PCR assay for characterization and concomitant subtyping of Staphylococcal Cassette
416 Chromosome *mec* types I to V in methicillin-resistant *Staphylococcus aureus*. J. Clin.
417 Microbiol. 43, 5026-5033.

418 Table 1. Resistance profile, Multi-Locus Sequence Type (MLST), *spa*- and Staphylococcal
 419 Cassette Chromosome (SCC)*mec*-type per MRSA strain in study 1.

Strain	Type of mastitis	Resistance profile ^a	<i>spa</i>	MLST	SCC <i>mec</i> ^b
1	Subclinical	AG, TET, TMP	t011	398	IVa
2	Clinical	AG, TET, TMP	t011	398	IVa
3	Subclinical	AG, ML, LM, TET, TMP	t011	398	IVa
4	Subclinical	LM, CIP, TET, TMP	t011	398	V
5	Subclinical	TET	t567	398	NT
6	Subclinical	AG, ML, LM, TET, TMP	t011	398	IVa
7	Subclinical	LM, CIP, TET, TMP	t011	398	V
8	Clinical	AG, ML, LM, TET, TMP	t011	398	IVa
9	Subclinical	KAN, TOB, ML, LM, TET, TMP	t011	398	V
10	Subclinical	AG, ML, LM, TET, TMP	t011	398	V
11	Subclinical	TET	t011	398	V

420 ^a AG: all aminoglycosides tested; KAN: kanamycin; TOB: tobramycin; ML: all macrolides
 421 tested; LM: lincomycin; CIP: ciprofloxacin; TET: tetracycline; TMP: trimethoprim

422 ^b NT: not typeable with the primers used

423 Table 2. In-herd prevalence of MRSA • no. 5: control farm

Farm	Herd size	Herd size	Positive cows		Positive quarters	
	(n cows)	(n quarters)	n	%	n	%
1	63	252	4	6.3	6	1.98
2	68	272	5	7.4	5	1.83
3	77	308	3	3.9	3	0.97
4	51	204	0	0	0	0
5	69	276	0	0	0	0

424

425 Table 3. Resistance profile, Multi-Locus Sequence Type (MLST), *spa*- and Staphylococcal
 426 Cassette Chromosome (SCC)*mec*-type per MRSA strain in study 2.

Farm	Quarter ^a	Strain	Resistance profile ^b	<i>Spa</i>	MLST ^c	SCC _{<i>mec</i>}
1	LH	1	ML, LM, TET	t011	ND	V
	RH	2	ML, LM, TET	t011	ND	V
	RF	3	ML, LM, TET	t011	ND	V
	LF	4	ML, LM, TET	t011	398	V
	RH	5	ML, LM, TET	t011	ND	V
	RF	6	ML, LM, TET	t011	ND	V
2	LH	7	AG, TET, TMP	t011	398	IVa
	LF	8	AG, ML, LM, TET, TMP	t011	ND	IVa
	RH	9	AG, TET, TMP	t011	ND	IVa
	RH	10	AG, ML, LM, TET, TMP	t011	398	IVa
	RH	11	AG, ML, LM, TET, TMP	t011	ND	IVa
3	RH	12	AG, ML, LM, TET, TMP	t011	ND	IVa
	RH	13	AG, ML, LM, TET, TMP	t011	ND	IVa
	RF	14	AG, ML, LM, TET, TMP	t011	398	IVa

427 ^a LF: left-front; LH: left-hind; RH: right-hind; RF: right-front

428 ^b AG: all aminoglycosides tested; LM: lincomycin; ML: all macrolides tested; TET:
 429 tetracycline; TMP: trimethoprim

430 ^c ND: not determined

431