



biblio.ugent.be

The UGent Institutional Repository is the electronic archiving and dissemination platform for all UGent research publications. Ghent University has implemented a mandate stipulating that all academic publications of UGent researchers should be deposited and archived in this repository. Except for items where current copyright restrictions apply, these papers are available in Open Access.

This item is the archived peer-reviewed author-version of:

Title: An elementary approach to asymptotic behavior in the Cesaro sense and applications to Stieltjes and Laplace transforms

Authors: Dennis Nemzer and Jasson Vindas

In: Novi Sad Journal of Mathematics, Vol. 45 (1), 111-124, 2015.

Optional: http://www.dmi.uns.ac.rs/nsjom/Papers/45_1/NSJOM_45_1_111_124.pdf

To refer to or to cite this work, please use the citation to the published version:

D. Nemzer and J. Vindas (2015). An elementary approach to asymptotic behavior in the Cesaro sense and applications to Stieltjes and Laplace transforms. *Novi Sad J. Math.* 45(1) 111-124.

AN ELEMENTARY APPROACH TO ASYMPTOTIC BEHAVIOR IN THE CESÀRO SENSE AND APPLICATIONS TO THE LAPLACE AND STIELTJES TRANSFORMS

DENNIS NEMZER AND JASSON VINDAS

ABSTRACT. We present an elementary approach to asymptotic behavior of generalized functions in the Cesàro sense. Our approach is based on Yosida's subspace of Mikusiński operators. Applications to Laplace and Stieltjes transforms are given.

1. INTRODUCTION

Asymptotic analysis is a very much studied topic within generalized function theory and has shown to be quite useful for the understanding of structural properties of a generalized function in connection with its local behavior as well as its growth properties at infinity. Several applications have been developed in diverse areas such as Tauberian and Abelian theory for integral transforms, differential equations, number theory, and mathematical physics. There is a vast literature on the subject, see the monographs [6, 9, 11, 12, 14] and references therein.

The purpose of this paper is to present an elementary approach to asymptotic behavior in the Cesàro sense. The Cesàro behavior for Schwartz distributions was introduced by Estrada in [5] (see also [6]). The approach we develop in this article uses Yosida's algebra \mathcal{M} of operators [13], which provides a simplified but useful version of Mikusiński's operational calculus [8]. While Schwartz distribution theory is based on the duality theory of topological vector spaces, the construction of Yosida's space \mathcal{M} is merely algebraic, making only use of elementary notions from calculus. That is why we call our approach to Cesàro asymptotics *elementary*.

The plan of the article is as follows. In Section 2, we recall the construction of Yosida's space \mathcal{M} . We study some useful localization properties of elements of \mathcal{M} in Section 3. The asymptotics in the Cesàro sense is defined in Section 4 and its properties are investigated. As an application, we conclude the article with some Abelian and Tauberian theorems for Stieltjes and Laplace transforms in Section 5. It should be mentioned that Abelian and Tauberian theorems for Stieltjes and Laplace transforms of generalized functions have been extensively investigated by several authors, see, e.g., [7, 9, 11, 12, 14].

2010 *Mathematics Subject Classification*. Primary 41A60, 44A40. Secondary 44A10, 44A15, 46F10, 46F12.

Key words and phrases. Asymptotic behavior of generalized functions; asymptotics in the Cesàro sense; Laplace transform; Stieltjes transform.

Dedicated to Professor Bogoljub Stanković on the occasion of his 90th birthday.

J. Vindas gratefully acknowledges support by Ghent University, through the BOF-grant 01N01014.

2. PRELIMINARIES

We recall in this section the construction of Yosida's space of operators \mathcal{M} and explain some of its properties. See [13] for more details about \mathcal{M} .

Let $C_+^n(\mathbb{R})$ denote the space of all n -times continuously differentiable functions on \mathbb{R} which vanish on the interval $(-\infty, 0)$. We write $C_+(\mathbb{R}) = C_+^0(\mathbb{R})$. For $f, g \in C_+(\mathbb{R})$, the convolution is given by

$$(f * g)(x) = \int_0^x f(x-t)g(t)dt.$$

Let H denote the Heaviside function. That is, $H(x) = 1$ for $x \geq 0$ and zero otherwise. For each $n \in \mathbb{N}$, we denote by H^n the function $H * \dots * H$ where H is repeated n times. One has $H^n \in C_+^{n-2}(\mathbb{R})$, $n \geq 2$. Note that if $f \in C_+(\mathbb{R})$, then

$$(H^m * f)(x) = \frac{1}{(m-1)!} \int_0^x (x-t)^{m-1} f(t)dt.$$

The space \mathcal{M} is defined as follows,

$$\mathcal{M} = \left\{ \frac{f}{H^k} : f \in C_+(\mathbb{R}), k \in \mathbb{N} \right\}.$$

Two elements of \mathcal{M} are equal, denoted $\frac{f}{H^n} = \frac{g}{H^m}$, if and only if $H^m * f = H^n * g$. Addition, multiplication (using convolution), and scalar multiplication are defined in the natural way, and \mathcal{M} with these operations is a commutative algebra with identity $\delta = \frac{H^2}{H^2}$, the Dirac delta. We can embed $C_+(\mathbb{R})$ into \mathcal{M} . Indeed, for $f \in C_+(\mathbb{R})$, we set $W_f = \frac{H * f}{H}$. Obviously, $f \mapsto W_f$ is injective.

Let $W = \frac{f}{H^k} \in \mathcal{M}$. The generalized derivative of W is defined as $DW = \frac{f}{H^{k+1}}$. The product of x and W is given by

$$xW = \frac{xf - kH * f}{H^k} \quad (k \geq 2).$$

In the last formula $k \geq 2$ is no restriction because $\frac{f}{H^k} = \frac{H * f}{H^{k+1}}$. Clearly, these definitions do not depend on the representative of W .

Proposition 2.1. *The generalized derivative and product by x satisfy:*

- (a) $xW_f = W_{xf}$, $f \in C_+(\mathbb{R})$.
- (b) $DW_f = W_{f'}$, $f \in C_+^1(\mathbb{R})$.
- (c) $D(xW) = W + xDW$, $W \in \mathcal{M}$.

Proof. For (a),

$$\begin{aligned} xW_f &= \frac{x(H^2 * f)}{H^2} - \frac{2H^2 * f}{H} = \frac{xH^2 * f}{H^2} + \frac{H^2 * xf}{H^2} - \frac{2H^2 * f}{H} \\ &= \frac{2H^3 * f}{H^2} + \frac{H^2 * xf}{H^2} - \frac{2H^2 * f}{H} \\ &= \frac{H * xf}{H} = W_{xf}. \end{aligned}$$

If $f' \in C_+(\mathbb{R})$, we have $DW_f = \frac{H * f}{H^2} = \frac{f}{H} = \frac{H * f'}{H} = W_{f'}$, which shows (b). Next, let $W = \frac{f}{H^k} \in \mathcal{M}$. Then,

$$W + xDW = \frac{f}{H^k} + \left(\frac{xf}{H^{k+1}} - \frac{(k+1)f}{H^k} \right) = \frac{xf}{H^{k+1}} - \frac{kf}{H^k} = D(xW).$$

□

Remark 2.2. Notice by identifying $f \in L^1_{loc}(\mathbb{R}^+)$ with $\frac{H * f}{H} \in \mathcal{M}$, the space $L^1_{loc}(\mathbb{R}^+)$ can be considered a subspace of \mathcal{M} . Also, for the construction of \mathcal{M} , the space of locally integrable functions which vanish on $(-\infty, 0)$ could have been used instead of $C_+(\mathbb{R})$.

3. LOCALIZATION

We discuss in this section localization properties of elements of \mathcal{M} .

Definition 3.1. Let $W = \frac{f}{H^k} \in \mathcal{M}$. W is said to vanish on an open interval (a, b) , denoted $W(x) = 0$ on (a, b) , provided there exists a polynomial p with degree at most $k - 1$ such that $p(x) = f(x)$ for $a < x < b$. Two elements $W, V \in \mathcal{M}$ are said to be equal on (a, b) , denoted $W(x) = V(x)$ on (a, b) , provided $W - V$ vanishes on (a, b) .

The support of $W \in \mathcal{M}$, denoted $\text{supp } W$, is the complement of the largest open set on which W vanishes. The degree of a polynomial p will be denoted by $\text{deg } p$ in the sequel.

Example 3.2. Recall $\delta = \frac{H^2}{H^2}$. Notice that $H^2(x) = x$ on the open interval $(0, \infty)$. Thus, $\delta(x) = 0$ on $(0, \infty)$. Also, $H^2(x) = 0$ on $(-\infty, 0)$. So, $\delta(x) = 0$ on $(-\infty, 0)$. Therefore, $\text{supp } \delta = \{0\}$.

Proposition 3.3. Let $W \in \mathcal{M}$.

- (a) If $W(x) = 0$ on (a, b) , then $DW(x) = 0$ on (a, b) .
- (b) If $DW(x) = 0$ on (a, b) , then W is constant on (a, b) .

Proof. Part (a) follows immediately from definitions. See [10, Thm. 4.1] for (b). □

Proposition 3.4. Let $W \in \mathcal{M}$.

- (a) If $W(x) = 0$ on (a, b) , then $xW(x) = 0$ on (a, b) .

- (b) Suppose $xW(x) = 0$ on (a, b) . Then,
 (i) $W(x) = 0$ on (a, b) , provided $0 \notin (a, b)$.
 (ii) $W(x) = 0$ on $(a, 0) \cup (0, b)$, provided $0 \in (a, b)$.

Proof. Let $W = \frac{f}{H^k} \in \mathcal{M}$ ($k \geq 2$).

Part (a). Since $W(x) = 0$ on (a, b) , there exist $a_0, a_1, \dots, a_{k-1} \in \mathbb{C}$ such that $f(x) = a_0 + a_1x + \dots + a_{k-1}x^{k-1}$ for $a < x < b$. Now, there exists $A \in \mathbb{C}$ such that

$$xf(x) - k(H * f)(x) = A + a_0(1 - k)x + a_1 \left(1 - \frac{k}{2}\right)x^2 + \dots + a_{k-2} \left(1 - \frac{k}{k-1}\right)x^{k-1},$$

for $a < x < b$. Since $xW = \frac{xf - kH * f}{H^k}$, the above yields $xW(x) = 0$ on (a, b) .

Part (b). Suppose $xW(x) = 0$ on (a, b) .

- (i) If $a < 0$, then the conclusion is clearly true. So assume $a \geq 0$. Now, there exists a polynomial p with $\deg p \leq k - 1$ such that

$$xf(x) - k \int_0^x f(t)dt = p(x) \quad \text{for } a < x < b.$$

Thus, $f \in C^1(a, b)$ and $xf'(x) + (1 - k)f(x) = p'(x)$ for $a < x < b$. Solving this differential equation, it follows that $f(x) = q(x)$ for $a < x < b$, where q is a polynomial with $\deg q \leq k - 1$. Therefore, $W(x) = 0$ on (a, b) .

- (ii) Since for all $W \in \mathcal{M}$, we have $W(x) = 0$ on $(-\infty, 0)$, we only need to show that $W(x) = 0$ on $(0, b)$. Since $0 \in (a, b)$, one has

$$(3.1) \quad xf(x) - k \int_0^x f(t)dt = 0$$

for $a < x < b$, in particular, for $0 < x < b$. Similarly as in the proof of part (i), we obtain $f(x) = Ax^{k-1}$ for $0 < x < b$, where $A \in \mathbb{C}$. Thus, $W(x) = 0$ on $(0, b)$. Therefore, $W(x) = 0$ on $(a, 0) \cup (0, b)$. □

Proposition 3.5. *Let $W \in \mathcal{M}$.*

- (a) *If $xW(x) = 0$ on $(-\infty, \infty)$, then $W = \alpha\delta$ for some $\alpha \in \mathbb{C}$.*
 (b) *If $W = \frac{f}{H^k}$ ($k \geq 2$) and $W(x) = 0$ on $(0, \infty)$, then $W = \sum_{n=0}^{k-2} \alpha_n \delta^{(n)}$, for some $\alpha_n \in \mathbb{C}$, $n = 0, 1, 2, \dots, k - 2$.*

Proof. Let $W = \frac{f}{H^k}$ ($k \geq 2$).

Part (a). Suppose $xW(x) = 0$ on $(-\infty, \infty)$. Similarly as in the proof of Proposition 3.4(b), we obtain (3.1) on $(-\infty, \infty)$. It follows that $f(x) = \beta x^{k-1}$ on $(0, \infty)$, for some $\beta \in \mathbb{C}$. Since f is continuous on \mathbb{R} and vanishes on $(-\infty, 0)$, we have $f(x) = \beta x^{k-1}$ on $[0, \infty)$. Therefore,

$$W = \frac{f}{H^k} = (k - 1)! \beta \frac{H^k}{H^k} = \alpha \delta \quad \text{where } \alpha = (k - 1)! \beta.$$

Part (b). Suppose $W(x) = 0$ on $(0, \infty)$. Then, $f(x) = a_0 + a_1x + \cdots + a_{k-1}x^{k-1}$ on $(0, \infty)$, where $a_0, a_1, \dots, a_{k-1} \in \mathbb{C}$. Since f is continuous and vanishes on $(-\infty, 0)$, we obtain $a_0 = 0$. Thus,

$$\begin{aligned} W &= a_1 \frac{H^2}{H^k} + 2!a_2 \frac{H^3}{H^k} + \cdots + (k-1)! a_{k-1} \frac{H^k}{H^k} \\ &= a_1 \delta^{(k-2)} + 2! a_2 \delta^{(k-1)} + \cdots + (k-1)! a_{k-1} \delta. \end{aligned}$$

□

4. ASYMPTOTICS IN THE CESÀRO SENSE

We now introduce and study asymptotics in the Cesàro sense for elements of \mathcal{M} . As usual, Γ stands for the Euler Gamma function and o stands for the little o growth order symbol of Landau [6, Chap. 1].

Definition 4.1. Let $W \in \mathcal{M}$. For $\alpha \in \mathbb{R} \setminus \{-1, -2, \dots\}$, define

$$W(x) \sim \frac{\gamma x^\alpha}{\Gamma(\alpha + 1)} \quad (C),$$

if and only if there is $k \in \mathbb{N}$ such that $W = \frac{f}{H^k}$ with

$$(4.1) \quad \frac{\Gamma(\alpha + k + 1)f(x) - p(x)}{x^{\alpha+k}} \rightarrow \gamma, \quad x \rightarrow \infty,$$

where p is some polynomial with $\deg p \leq k - 1$. That is,

$$\Gamma(\alpha + k + 1)f(x) = p(x) + \gamma x^{\alpha+k} + o(x^{\alpha+k}), \quad \text{as } x \rightarrow \infty.$$

Remark 4.2. If $\alpha > -1$, then the polynomial p is not needed.

The next theorem tells us that Definition 4.1 is consistent with the choice of representatives.

Theorem 4.3. *If $W = \frac{f}{H^k} \in \mathcal{M}$ is such that (4.1) holds, then the continuous function $H^m * f$ in the representation $W = \frac{H^m * f}{H^{k+m}}$ ($m \in \mathbb{N}$) satisfies*

$$\frac{\Gamma(\alpha + k + m + 1)(H^m * f)(x) - q(x)}{x^{\alpha+k+m}} \rightarrow \gamma, \quad x \rightarrow \infty,$$

for some polynomial q of degree at most $k + m - 1$.

Proof. Suppose that (4.1) holds for some polynomial p with $\deg p \leq k - 1$. Consider

$$\Gamma(\alpha + k + 2) \int_0^x \left(f(t) - \frac{p(t)}{\Gamma(\alpha + k + 1)} \right) dt + K,$$

with K a constant to be determined.

Case 1. Suppose $\alpha + k < -1 \Rightarrow x^{\alpha+k} \in L^1(1, \infty)$. So, (4.1) implies $f - p/\Gamma(\alpha + k + 1) \in L^1(0, \infty)$. Therefore there exists a constant K such that

$$\Gamma(\alpha + k + 2) \int_0^x \left(f(t) - \frac{p(t)}{\Gamma(\alpha + k + 1)} \right) dt + K \rightarrow 0, \quad \text{as } x \rightarrow \infty.$$

Thus, using L'Hospital's rule and (4.1)

$$\lim_{x \rightarrow \infty} \frac{\Gamma(\alpha + k + 2) \int_0^x \left(f(t) - \frac{p(t)}{\Gamma(\alpha + k + 1)} \right) dt + K}{x^{\alpha + k + 1}} = \lim_{x \rightarrow \infty} \frac{\Gamma(\alpha + k + 1)f(x) - p(x)}{x^{\alpha + k}} = \gamma.$$

Let $q(x) = (\alpha + k + 1) \int_0^x p(t)dt - K$. Then q is a polynomial with $\deg q \leq k$. Moreover,

$$\frac{\Gamma(\alpha + k + 2)(H * f)(x) - q(x)}{x^{\alpha + k + 1}} \rightarrow \gamma, \quad \text{as } x \rightarrow \infty.$$

Case 2. Suppose $\alpha + k > -1$. Assume $\gamma \neq 0$. Then (4.1) implies

$$\int_0^x (\Gamma(\alpha + k + 1)f(t) - p(t)) dt \rightarrow \pm\infty, \quad \text{as } x \rightarrow \infty.$$

Using L'Hospital's Rule and (4.1),

$$\frac{\Gamma(\alpha + k + 2) \int_0^x \left(f(t) - \frac{p(t)}{\Gamma(\alpha + k + 1)} \right) dt}{x^{\alpha + k + 1}} = \frac{(\alpha + k + 1) \int_0^x (\Gamma(\alpha + k + 1)f(t) - p(t))}{x^{\alpha + k + 1}} \rightarrow \gamma,$$

as $x \rightarrow \infty$. For the case $\gamma = 0$ (and $\alpha + k > -1$), we set $g(x) = \Gamma(\alpha + k + 1)f(x) - p(x)$. It is well known that if $x^{-\alpha - k}g(x) \rightarrow 0$, then $x^{-\alpha - k - 1} \int_0^x g(t)dt \rightarrow 0$ (see e.g. [6, Chap. 1]).

The above shows that the claim is true for $m = 1$. By using induction, the result follows. \square

Unless otherwise stated, we assume from now on $\alpha \in \mathbb{R} \setminus \{-1, -2, \dots\}$.

The following provides an alternative to Definition 4.1.

Let $W \in \mathcal{M}$ and $\gamma \in \mathbb{C}$. Then

$$(4.2) \quad W(x) \sim \frac{\gamma x^\alpha}{\Gamma(\alpha + 1)} \quad (C), \quad x \rightarrow \infty$$

if and only if there exist $n \in \mathbb{N}$ with $\alpha + n > 0$, $g \in C_+(\mathbb{R})$, and $b > 0$ such that $W(x) = D^n g(x)$ on (b, ∞) and $g(x)/x^{\alpha + n} \rightarrow \gamma/\Gamma(\alpha + n + 1)$ as $x \rightarrow \infty$. We leave the verification of this fact to the reader.

Theorem 4.4. *We have:*

(a) *If $f \in C_+(\mathbb{R})$ such that $f(x) \sim \gamma x^\alpha$ as $x \rightarrow \infty$, then $W_f(x) \sim \gamma x^\alpha$ (C), $x \rightarrow \infty$.*

(b) *Let $W \in \mathcal{M}$ satisfy (4.2). Then,*

$$(4.3) \quad DW(x) \sim \frac{\alpha \gamma x^{\alpha - 1}}{\Gamma(\alpha + 1)} \quad (C), \quad x \rightarrow \infty.$$

and

$$(4.4) \quad xW(x) \sim \frac{\gamma x^{\alpha + 1}}{\Gamma(\alpha + 1)} \quad (C), \quad x \rightarrow \infty.$$

Proof. We only prove (4.4) since the other parts of the theorem follow from the definitions. We first assume a stronger condition on the polynomial p . Suppose $V = \frac{g}{H^n} \in \mathcal{M}$ ($n \geq 2$) with

$$(4.5) \quad \frac{\Gamma(\alpha + n + 1)g(x) - p(x)}{x^{\alpha+n}} \rightarrow \gamma \quad \text{as } x \rightarrow \infty,$$

for some polynomial p with $\deg p \leq n - 2$. Then there exists a polynomial q with $\deg q \leq n - 1$ such that

$$\frac{\Gamma(\alpha + n + 2)xg(x) - q(x)}{x^{\alpha+n+1}} \rightarrow (\alpha + n + 1)\gamma \quad \text{as } x \rightarrow \infty.$$

Therefore,

$$\frac{xg}{H^n} \sim \frac{(\alpha + n + 1)\gamma x^{\alpha+1}}{\Gamma(\alpha + 2)} \quad (C), \quad x \rightarrow \infty.$$

Also, from (4.5) it follows that

$$\frac{-ng}{H^{n-1}} \sim \frac{-n\gamma x^{\alpha+1}}{\Gamma(\alpha + 2)} \quad (C), \quad x \rightarrow \infty.$$

Thus,

$$xV(x) \sim \frac{\gamma x^{\alpha+1}}{\Gamma(\alpha + 1)} \quad (C), \quad x \rightarrow \infty.$$

We now remove the stronger condition that $\deg p \leq n - 2$ and complete the proof of the theorem. Let $W = \frac{f}{H^k} \in \mathcal{M}$ such that (4.2) holds. That is, there exists $a_0, a_1, \dots, a_{k-1} \in \mathbb{C}$ such that

$$(4.6) \quad \frac{\Gamma(\alpha + k + 1)f(x) - (a_0 + a_1x + \dots + a_{k-1}x^{k-1})}{x^{\alpha+k}} \rightarrow \gamma \quad \text{as } x \rightarrow \infty$$

Let $V = \frac{f(x) - \beta x^{k-1}}{H^k} \in \mathcal{M}$, where $\beta = a_{k-1}/\Gamma(\alpha + k + 1)$. Then, from (4.6) it follows that,

$$V(x) \sim \frac{\gamma x^\alpha}{\Gamma(\alpha + 1)} \quad (C), \quad x \rightarrow \infty.$$

And, by the first part of the proof,

$$xV(x) \sim \frac{\gamma x^{\alpha+1}}{\Gamma(\alpha + 1)} \quad (C), \quad x \rightarrow \infty.$$

Since, $xW = xV + \beta(k-1)!x\delta = xV$, it follows that

$$xW(x) \sim \frac{\gamma x^{\alpha+1}}{\Gamma(\alpha + 1)} \quad (C), \quad x \rightarrow \infty.$$

□

By Theorem 4.4, we obtain the following theorem.

Theorem 4.5. *If $(H * W)(x) \sim \frac{\gamma x^\alpha}{\Gamma(\alpha + 1)} \quad (C)$, then $(xW)(x) \sim \frac{\alpha\gamma x^\alpha}{\Gamma(\alpha + 1)} \quad (C)$, $x \rightarrow \infty$.*

The proofs of the next proposition and corollary follow from the definitions.

Proposition 4.6. *If V has compact support, then $V(x) \sim \frac{0}{\Gamma(\alpha + 1)}x^\alpha$ (C), $x \rightarrow \infty$.*

Asymptotics in the Cesàro sense is a local property.

Corollary 4.7. *Let $W, V \in \mathcal{M}$. Suppose that W has Cesàro asymptotics (4.2) and $W(x) = V(x)$ on (a, ∞) . Then, $V(x) \sim \frac{\gamma x^\alpha}{\Gamma(\alpha + 1)}$ (C), $x \rightarrow \infty$.*

5. APPLICATIONS

In this last section we give some Abelian and Tauberian theorems for Stieltjes and Laplace transforms of elements of \mathcal{M} .

We start by defining the Stieltjes transform [10]. Let $r > -1$ and suppose $W = \frac{f}{H^k} \in \mathcal{M}$, where $x^{-r-k+\sigma}f(x)$ is bounded as $x \rightarrow \infty$ for some $\sigma > 0$. The Stieltjes transform of W of index r is given by

$$\Lambda_r W(z) = (r+1)_k \int_0^\infty \frac{f(x)}{(x+z)^{r+k+1}} dx, \quad z \in \mathbb{C} \setminus (-\infty, 0],$$

where $(r+1)_k = \frac{\Gamma(r+k+1)}{\Gamma(r+1)} = (r+1)(r+2)\dots(r+k)$. Notice that $\Lambda_r W(z)$ is holomorphic in the variable z , as one readily verifies.

The following is a classical Abelian theorem for the Stieltjes transform.

Theorem 5.1 ([3]). *If $f \in C_+(\mathbb{R})$ such that $x^{-\nu}f(x) \rightarrow A$ as $x \rightarrow \infty$, with $\nu > -1$, then for $\rho > \nu$,*

$$\lim_{\substack{z \rightarrow \infty \\ |\arg z| \leq \theta < \pi/2}} \frac{z^{\rho-\nu} \Gamma(\rho+1) S_\rho f(z)}{\Gamma(\rho-\nu) \Gamma(\nu+1)} = A,$$

where $S_\rho f(z) = \int_0^\infty \frac{f(x)}{(x+z)^{\rho+1}} dx$.

Theorem 5.2. *Let $W \in \mathcal{M}$ and $r > -1$. Suppose that $W(x) \sim \frac{\gamma x^\alpha}{\Gamma(\alpha + 1)}$ (C), $x \rightarrow \infty$. Then:*

(i) *If $r > \alpha > -1$, then $\Lambda_r W(z)$ is well-defined and has asymptotic behavior*

$$\lim_{\substack{z \rightarrow \infty \\ |\arg z| \leq \theta < \pi/2}} \frac{z^{r-\alpha} \Gamma(r+1) \Lambda_r W(z)}{\Gamma(r-\alpha)} = \gamma,$$

(ii) *If $\alpha < -1$, $\alpha \notin \{-2, -3, -4, \dots\}$, then $\Lambda_r W(z)$ is well-defined and there are constants A_1, \dots, A_k such that*

$$(5.1) \quad \lim_{\substack{z \rightarrow \infty \\ |\arg z| \leq \theta < \pi/2}} \frac{z^{r-\alpha} \Gamma(r+1)}{\Gamma(r-\alpha)} \left[\Lambda_r W(z) - \sum_{j=1}^k \frac{A_j}{z^{r+j}} \right] = \gamma.$$

Proof. (i) Let $\alpha > -1$ and $W = \frac{f}{H^k} \in \mathcal{M}$ such that

$$\frac{\Gamma(\alpha + k + 1)f(x)}{x^{\alpha+k}} \rightarrow \gamma \quad \text{as } x \rightarrow \infty.$$

It follows that for $r > \alpha$, $f(x)x^{-r-k+\sigma}$ is bounded as $x \rightarrow \infty$ for some $\sigma > 0$. Now, by substituting $r + k$ for ρ , $\alpha + k$ for ν , and $\frac{\gamma}{\Gamma(\alpha+k+1)}$ for A in the above classical Abelian theorem, we obtain

$$\lim_{\substack{z \rightarrow \infty \\ |\arg z| \leq \theta < \pi/2}} \frac{z^{r-\alpha}\Gamma(r+k+1)S_{r+k}f(z)}{\Gamma(r-\alpha)} = \gamma.$$

Now, using the fact that $\Lambda_r W(z) = (r+1)_k S_{r+k}f(z)$, the result follows.

(ii) Suppose that $W = \frac{f}{H^k} \in \mathcal{M}$ with $k + \alpha > -1$ and that f can be written as $f(x) = p(x) + g(x)$, where $g \in C_+(\mathbb{R})$ satisfies $\lim_{x \rightarrow \infty} x^{-\alpha-k}g(x) = \gamma/\Gamma(\alpha + k + 1)$ and $p(x) = \sum_{j=0}^{k-1} a_j x^j$. It follows that $|f(x)| \leq Cx^{k-1}$ for some constant C and thus $f(x)x^{-r-k+\sigma}$ is bounded for any $0 < \sigma \leq 1 + r$. Observe that

$$\begin{aligned} S_{r+k}f(z) &= S_{r+k}g(z) + \sum_{j=0}^{k-1} a_j \int_0^\infty \frac{x^j}{(x+z)^{r+k+1}} dx \\ &= S_{r+k}g(z) + \sum_{j=0}^{k-1} \frac{j!\Gamma(r+k-j)}{\Gamma(r+k+1)z^{j+k+r}} a_j. \end{aligned}$$

By Theorem 5.1, we have that $z^{r-\alpha}S_{r+k}g(z) \rightarrow \frac{\gamma\Gamma(r-\alpha)}{\Gamma(r+k+1)}$ as $z \rightarrow \infty$ on sectors $|\arg z| \leq \theta < \frac{\pi}{2}$. Since $\Lambda_r W(z) = (r+1)_k S_{r+k}f(z)$, we obtain (5.1) with $A_j = (k-j)!(r+1)_{j-1}a_{k-j}$. \square

We illustrate our ideas with the ensuing example, a deduction of Stirling's formula for the Gamma function.

Example 5.3. (Stirling's formula) Recall that the digamma function ψ is defined as the logarithmic derivative of Γ . By using the product formula for Γ , namely,

$$\Gamma(z) = \frac{e^{-\gamma z}}{z} \prod_{n=1}^{\infty} \left(1 + \frac{z}{n}\right)^{-1} e^{\frac{z}{n}},$$

one has

$$\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)} = -\gamma + \sum_{n=0}^{\infty} \left(\frac{1}{n+1} - \frac{1}{n+z} \right), \quad z \in \mathbb{C} \setminus \{0, -1, -2, \dots\},$$

where γ is the Euler-Mascheroni constant. We define $W = \frac{f}{H^2}$, where $f(x) = \int_0^x ([t] - t + \frac{1}{2}) dt$ (here $[x]$ stands for the integer part of x).

Set $g(x) = [x] - x + \frac{1}{2}$ and note that g is periodic with period 1, $|g(x)| \leq \frac{1}{2}$ for all $x \in \mathbb{R}$, and $\int_n^{n+1} g(x)dx = 0$ for all $n \in \mathbb{N}$. This implies that $|\int_0^x g(t)dt| \leq \frac{1}{2}$ for all $x \geq 0$. Consequently, $W(x) \sim 0 \cdot x^{-\sigma}$ (C), $x \rightarrow \infty$ for any $0 < \sigma < 2$. Theorem

4 now yields (from the proof it is clear that the constants $A_1 = A_2 = 0$ in this case because f is bounded)

$$(5.2) \quad \lim_{\substack{z \rightarrow \infty \\ |\arg z| \leq \theta < \pi/2}} z^\sigma \Lambda_0 W(z) = 0,$$

for any $0 < \sigma < 2$. From now on we will work with $1 < \sigma < 2$. We compute an explicit expression for $\Lambda_0 W(z)$,

$$\begin{aligned} \Lambda_0 W(z) &= 2 \int_0^\infty \frac{f(x)}{(x+z)^3} dx = \int_0^\infty \frac{(\lfloor x \rfloor - x + \frac{1}{2})}{(x+z)^2} dx = \frac{1}{2z} + \lim_{N \rightarrow \infty} \int_0^N \frac{\lfloor x \rfloor - x}{(x+z)^2} dx \\ &= \frac{1}{2z} + \lim_{N \rightarrow \infty} \left(\int_0^N \frac{d\lfloor x \rfloor}{x+z} - \frac{N}{N+z} + \frac{N}{N+z} - \int_0^N \frac{dx}{x+z} \right) \\ &= \ln z + \frac{1}{2z} + \lim_{N \rightarrow \infty} \sum_{n=1}^N \frac{1}{n+z} - \ln N \\ &= \ln z + \frac{1}{2z} - \psi(z). \end{aligned}$$

The limit (5.2) then yields

$$(5.3) \quad \psi(z) = \ln z + \frac{1}{2z} + o\left(\frac{1}{z^\sigma}\right), \quad z \rightarrow \infty,$$

for z in the sectors $|\arg z| \leq \theta < \frac{\pi}{2}$. Note that integration of (5.3) implies for any $0 < \tau < 1$,

$$\ln \Gamma(z) = z(\ln z - 1) + \frac{1}{2} \ln z + C + o\left(\frac{1}{z^\tau}\right), \quad z \rightarrow \infty$$

on $|\arg z| \leq \theta < \frac{\pi}{2}$, which is Stirling's asymptotic formula for the Gamma function except for the evaluation of the constant C . The constant is of course well known to be $C = \sqrt{2\pi}$. We refer to [6, p. 43] for an elementary proof of the latter fact.

We now consider the Laplace transform [1]. If $W = \frac{f}{H^k} \in \mathcal{M}$, where $f(x)e^{-\sigma x}$ is bounded as $x \rightarrow \infty$ for some $\sigma \in \mathbb{R}$, then the Laplace transform of W is given by

$$\mathcal{L}W(z) = z^k \int_0^\infty e^{-zx} f(x) dx, \quad \Re z > \sigma.$$

Theorem 5.4. *Let $W \in \mathcal{M}$. Assume that $W(x) \sim \frac{\gamma x^\alpha}{\Gamma(\alpha+1)}$ (C), $x \rightarrow \infty$. Then W is Laplace transformable and*

(i) *If $\alpha > -1$, then*

$$\lim_{\substack{z \rightarrow 0 \\ |\arg z| \leq \theta < \pi/2}} z^{\alpha+1} \mathcal{L}W(z) = \gamma.$$

(ii) If $\alpha < -1$, $\alpha \notin \{-2, -3, \dots\}$, then there are constants A_0, \dots, A_{k-1} such that

$$(5.4) \quad \lim_{\substack{z \rightarrow 0 \\ |\arg z| \leq \theta < \pi/2}} z^{\alpha+1} \left(\mathcal{L}W(z) - \sum_{j=0}^{k-1} A_j z^j \right) = \gamma$$

Remark 5.5. In the terminology of finite part limits ([6, Sect. 2.4]), the limit (5.4) might be rewritten as

$$\text{F.p.} \quad \lim_{\substack{z \rightarrow 0 \\ |\arg z| \leq \theta < \pi/2}} z^{\alpha+1} \mathcal{L}W(z) = \gamma.$$

Proof. (i) Let $\alpha > -1$ and $W = \frac{f}{H^k} \in \mathcal{M}$ such that

$$\frac{\Gamma(\alpha + k + 1)f(x)}{x^{\alpha+k}} \rightarrow \gamma \quad \text{as } x \rightarrow \infty.$$

It follows that there exists $\sigma > 0$ such that $f(x)e^{-\sigma x}$ is bounded as $x \rightarrow \infty$. Now, by a well known classical Abelian theorem for the Laplace transform [4], we obtain

$$\lim_{\substack{z \rightarrow 0 \\ |\arg z| \leq \theta < \pi/2}} \frac{z^{\alpha+k+1} \mathcal{L}f(z)}{\Gamma(\alpha + k + 1)} = \frac{\gamma}{\Gamma(\alpha + k + 1)}.$$

So,

$$\lim_{\substack{z \rightarrow 0 \\ |\arg z| \leq \theta < \pi/2}} z^{\alpha+1} \mathcal{L}W(z) = \lim_{\substack{z \rightarrow 0 \\ |\arg z| \leq \theta < \pi/2}} z^{\alpha+k+1} \mathcal{L}f(z) = \gamma.$$

(ii) Suppose $W = \frac{f}{H^k} \in \mathcal{M}$ with $k + \alpha > -1$ and that f can be written as $f(x) = p(x) + g(x)$, where $g \in C_+(\mathbb{R})$ has asymptotic behavior $\lim_{x \rightarrow \infty} x^{-\alpha-k} g(x) = \gamma/\Gamma(\alpha + k + 1)$ and $p(x) = \sum_{j=0}^{k-1} \alpha_j x^j$. Note that $W = W_1 + W_2$, where $W_1 = \frac{g}{H^k}$ and $W_2 = \frac{p}{H^k}$. Exactly as above, one verifies that

$$\lim_{\substack{z \rightarrow 0 \\ |\arg z| \leq \theta < \pi/2}} z^{\alpha+1} \mathcal{L}W_1(z) = \gamma.$$

It remains to observe that $\mathcal{L}W_2(z) = \sum_{j=0}^{k-1} A_j z^j$, with $A_j = \alpha_{k-j-1}/(k-j-1)!$. \square

Theorem 5.6. (Tauberian Theorem) Let $W = \frac{f}{H^k} \in \mathcal{M}$, where in addition to $f \in C_+(\mathbb{R})$, f is real-valued and nonnegative. If $\mathcal{L}W(s) \sim \gamma s^{-\alpha-1}$, $s \rightarrow 0^+$ (for some $\alpha > -1$), then $W(x) \sim \frac{\gamma x^\alpha}{\Gamma(\alpha + 1)}$ (C), $x \rightarrow \infty$.

Proof. It follows that

$$(5.5) \quad \mathcal{L}f(s) \sim \gamma s^{-(\alpha+k+1)}, \quad s \rightarrow 0^+.$$

Also,

$$(5.6) \quad \mathcal{L}f(s) = \int_0^\infty e^{-st} d(H * f)(t), \quad s > 0.$$

Since $H * f$ is nondecreasing, by (5.5) and (5.6) and Hardy-Littlewood-Karamata Tauberian Theorem [2, 14], it follows that

$$(H * f)(x) \sim \frac{\gamma}{\Gamma(\alpha + k + 2)} x^{\alpha+k+1}, \quad x \rightarrow \infty.$$

That is,

$$\frac{\Gamma(\alpha + (k + 1) + 1)(H * f)(x)}{x^{\alpha+(k+1)}} \rightarrow \gamma, \quad x \rightarrow \infty.$$

Since $W = \frac{H * f}{H^{k+1}}$, the above yields $W(x) \sim \frac{\gamma x^\alpha}{\Gamma(\alpha + 1)} \quad (C), \quad x \rightarrow \infty. \quad \square$

REFERENCES

- [1] D. Atanasiu, D. Nemzer, *Extending the Laplace transform*, Math. Student **77** (2008), 203–212.
- [2] N. Bingham, C. Goldie, J. Teugels, *Regular variation*, Cambridge University Press, Cambridge, 1987.
- [3] R. D. Carmichael, *Abelian theorems for the Stieltjes transform of functions*, Bull. Cal. Math. Soc. **68** (1976), 49–52.
- [4] G. Doetsch, *Theorie der Laplace transformation*, Band I, Verlag Birkhäuser, Basel, 1950.
- [5] R. Estrada, *The Cesàro behaviour of distributions*, Proc. R. Soc. Lond. A **454** (1998), 2425–2443.
- [6] R. Estrada, R. P. Kanwal, *A distributional approach to asymptotics. Theory and applications*, Birkhäuser Boston, Inc., Boston, MA, 2002.
- [7] R. Estrada, J. Vindas, *On Tauber's second Tauberian theorem*, Tohoku Math. J. (2) **64** (2012), 539–560.
- [8] J. Mikusiński, *Operational calculus*, Vol. 1, International Series of Monographs in Pure and Applied Mathematics, 109, Pergamon Press, Oxford; PWN—Polish Scientific Publishers, Warsaw, 1983.
- [9] O. P. Misra, J. L. Lavoine, *Transform analysis of generalized functions*, North-Holland Publishing Co., Amsterdam, 1986.
- [10] D. Nemzer, *Mikusiński's operational calculus approach to the distributional Stieltjes transform*, Adv. Math., Sci. J. **2** (2013), 35–42.
- [11] S. Pilipović, B. Stanković, A. Takači, *Asymptotic behaviour and Stieltjes transformation of distributions*, Teubner-Texte zur Mathematik, Leipzig, 1990.
- [12] S. Pilipović, B. Stanković, J. Vindas, *Asymptotic behavior of generalized functions*, Series on analysis, applications and computation – Vol. 5, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2012.
- [13] K. Yosida, *Operational calculus. A theory of hyperfunctions*, Springer-Verlag, New York, 1984.
- [14] V. S. Vladimirov, Yu. N. Drozzinov, B. I. Zavalov, *Tauberian theorems for generalized functions*, Kluwer Academic Publishers Group, Dordrecht, 1988.

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY, STANISLAUS, ONE UNIVERSITY CIRCLE, TURLOCK, CA 95382, USA

E-mail address: nemzer@comcast.net

DEPARTMENT OF MATHEMATICS, GHENT UNIVERSITY, KRIJGSLAAN 281 GEBOUW S22, 9000 GENT, BELGIUM

E-mail address: jvindas@cage.UGent.be