



**[biblio.ugent.be](http://biblio.ugent.be)**

The UGent Institutional Repository is the electronic archiving and dissemination platform for all UGent research publications. Ghent University has implemented a mandate stipulating that all academic publications of UGent researchers should be deposited and archived in this repository. Except for items where current copyright restrictions apply, these papers are available in Open Access.

This item is the archived peer-reviewed author-version of:

**Title: Chebyshev upper estimates for Beurling's generalized prime numbers**

**Authors: Jasson Vindas**

**In: Bulletin of the Belgian Mathematical Society – Simon Stevin , 20(1), 175-180, 2013.**

**Optional: <http://projecteuclid.org/euclid.bbms/1366306723>**

**To refer to or to cite this work, please use the citation to the published version:**

**Jasson Vindas (2013). Chebyshev upper estimates for Beurling's generalized prime numbers. *Bull. Belg. Math. Soc. Simon Stevin* 20(1), 175-180.**

# CHEBYSHEV UPPER ESTIMATES FOR BEURLING'S GENERALIZED PRIME NUMBERS

JASSON VINDAS

ABSTRACT. Let  $N$  be the counting function of a Beurling generalized number system and let  $\pi$  be the counting function of its primes. We show that the  $L^1$ -condition

$$\int_1^\infty \left| \frac{N(x) - ax}{x} \right| \frac{dx}{x} < \infty$$

and the asymptotic behavior

$$N(x) = ax + O\left(\frac{x}{\log x}\right),$$

for some  $a > 0$ , suffice for a Chebyshev upper estimate

$$\frac{\pi(x) \log x}{x} \leq B < \infty.$$

## 1. INTRODUCTION

Let  $P = \{p_k\}_{k=1}^\infty$  be a set of Beurling generalized primes, namely, a non-decreasing sequence of real numbers  $1 < p_1 \leq p_2 \leq \dots \leq p_k \rightarrow \infty$ . The sequence  $\{n_k\}_{k=1}^\infty$  denotes its associated set of generalized integers [2, 3]. Consider the counting functions of generalized integers and primes

$$N(x) = N_P(x) = \sum_{n_k < x} 1 \quad \text{and} \quad \pi(x) = \pi_P(x) = \sum_{p_k < x} 1.$$

Beurling's problem consists in finding mild conditions over  $N$  that ensure a certain asymptotic behavior for  $\pi$ . This problem has been extensively investigated in connection with the prime number theorem (PNT), i.e.,

$$(1) \quad \pi(x) \sim \frac{x}{\log x}, \quad x \rightarrow \infty,$$

and Chebyshev two-sided estimates, that is,

$$(2) \quad 0 < \liminf_{x \rightarrow \infty} \frac{\pi(x) \log x}{x} \quad \text{and} \quad \limsup_{x \rightarrow \infty} \frac{\pi(x) \log x}{x} < \infty.$$

---

2000 *Mathematics Subject Classification*. Primary 11N80. Secondary 11N05, 11M41.

*Key words and phrases*. Chebyshev upper estimates; Beurling generalized primes.

The author gratefully acknowledges support by a Postdoctoral Fellowship of the Research Foundation–Flanders (FWO, Belgium).

On the other hand, there are no mild hypotheses in the literature for Chebyshev upper estimates,

$$(3) \quad \limsup_{x \rightarrow \infty} \frac{\pi(x) \log x}{x} < \infty .$$

The purpose of this article is to study asymptotic requirements over  $N$  that imply the Chebyshev upper estimate (3).

Beurling [3] proved that

$$(4) \quad N(x) = ax + O\left(\frac{x}{\log^\gamma x}\right), \quad x \rightarrow \infty \quad (a > 0),$$

where  $\gamma > 3/2$ , suffices for the PNT (1) to hold. See [3, 10, 13] for more general PNT. Beurling's condition is sharp, because when  $\gamma = 3/2$  there are generalized number systems for which the PNT fails [3, 5]. For  $\gamma < 1$ , not even Chebyshev estimates need to hold, as follows from an example of Hall [9] (see also [1]). Diamond has shown [6] that (4) with  $\gamma > 1$  is enough to obtain Chebyshev two-sided estimates (2). Furthermore, he conjectured [7] that the weaker hypothesis

$$(5) \quad \int_1^\infty \left| \frac{N(x) - ax}{x} \right| \frac{dx}{x} < \infty, \quad \text{with } a > 0,$$

would be enough for (2). His conjecture was shown to be false by Kahane [11]. Nevertheless, the author has recently shown [15] that if one adds to (5) the condition

$$(6) \quad N(x) = ax + o\left(\frac{x}{\log x}\right), \quad x \rightarrow \infty,$$

then (2) is fulfilled, extending thus earlier results from [6, 18].

It is natural to replace the little  $o$  symbol in (6) by an  $O$  growth estimate and investigate the effect of this new condition on the asymptotic distribution of the generalized primes. It turns out that one gets a Chebyshev upper estimate in this case. Our main goal is to give a proof of the following theorem.

**Theorem 1.** *Diamond's  $L^1$ -condition (5) and the asymptotic behavior*

$$(7) \quad N(x) = ax + O\left(\frac{x}{\log x}\right), \quad x \rightarrow \infty,$$

*suffice for the Chebyshev upper estimate (3).*

## 2. NOTATION

We will give an analytic proof of Theorem 1. Our technique follows distributional ideas already used in [13, 15, 16]. It employs the Wiener division theorem [12, Chap. 2] and the operational calculus for the Laplace transform of Schwartz distributions [4, 17]. The Schwartz spaces of test functions and distributions are denoted as  $\mathcal{D}(\mathbb{R})$ ,  $\mathcal{S}(\mathbb{R})$ ,  $\mathcal{D}'(\mathbb{R})$  and  $\mathcal{S}'(\mathbb{R})$ , see [8, 14, 17] for

their properties. If  $f \in \mathcal{S}'(\mathbb{R})$  has support in  $[0, \infty)$ , its Laplace transform is well defined as

$$\mathcal{L}\{f; s\} = \langle f(u), e^{-su} \rangle, \quad \Re s > 0,$$

and the Fourier transform  $\hat{f}$  is the distributional boundary value [4] of  $\mathcal{L}\{f; s\}$  on  $\Re s = 0$ . We use the notation  $H$  for the Heaviside function, it is simply the characteristic function of  $(0, \infty)$ .

Observe that (3) is equivalent to

$$(8) \quad \limsup_{x \rightarrow \infty} \frac{\psi(x)}{x} < \infty,$$

where  $\psi$  is the Chebyshev function

$$\psi(x) = \psi_P(x) = \sum_{n_k < x} \Lambda(n_k),$$

as follows from [2, Lem. 2E].

### 3. PROOF OF THEOREM 1

Assume (5) and (7). Set  $T(u) = e^{-u}\psi(e^u)$ . We must show (8), that is,

$$(9) \quad \limsup_{u \rightarrow \infty} T(u) < \infty.$$

The crude inequality  $T(u) \leq ue^{-u}N(e^u) = O(u)$  implies that  $T \in \mathcal{S}'(\mathbb{R})$ . The proof of (9) depends upon estimates on convolution averages of  $T$ :

**Lemma 1.** *There exists  $c > 0$  such that*

$$(10) \quad \int_{-\infty}^{\infty} T(u)\hat{\phi}(u-h)du = O(1),$$

whenever  $\phi \in \mathcal{D}(-c, c)$ .

Indeed, suppose that Lemma 1 has been already established. Choose then in (10) a test function  $\phi \in \mathcal{D}(-c, c)$  such that  $\hat{\phi}$  is non-negative. Since  $\psi(e^u)$  is non-decreasing, we have  $e^{-u}T(h) \leq T(u+h)$  whenever  $u$  and  $h$  are positive. Setting  $C = \int_0^{\infty} e^{-u}\hat{\phi}(u)du > 0$ , we obtain that

$$T(h) \leq \frac{1}{C} \int_0^{\infty} T(u+h)\hat{\phi}(u)du = O(1),$$

and Theorem 1 follows at once. It remains to prove the lemma.

*Proof of Lemma 1.* Set  $E_1(u) := e^{-u}N(e^u) - aH(u)$  and  $E_2(u) = uE_1(u)$ . The assumptions (5) and (7) take the form  $E_1 \in L^1(\mathbb{R})$  and  $E_2 \in L^\infty(\mathbb{R})$ . Consider

$$G(s) = \zeta(s) - \frac{a}{s-1} = s\mathcal{L}\{E_1; s-1\} + a.$$

Taking  $\Re s \rightarrow 1^+$ , in the distributional sense, we obtain  $G(1+it) = (1+it)\hat{E}_1(t) + a$ . Since  $E_1 \in L^1(\mathbb{R})$ ,  $\hat{E}_1$  is continuous; therefore  $G(s)$  extends to a continuous function on  $\Re s = 1$ . Consequently,  $(s-1)\zeta(s)$  is continuous on

$\Re s = 1$  and there exists  $c > 0$  such that  $it\zeta(1+it) \neq 0$  for all  $t \in (-3c, 3c)$ . Next, we study the boundary values, on the line segment  $1 + i(-c, c)$ , of

$$\mathcal{L}\{T(u); s-1\} = \mathcal{L}\{\psi(e^u); s\} = -\frac{\zeta'(s)}{s\zeta(s)}.$$

A quick calculation shows that

$$(11) \quad -\frac{\zeta'(s)}{s\zeta(s)} = \frac{\mathcal{L}\{E'_2; s-1\}}{(s-1)\zeta(s)} - \frac{(2s-1)\mathcal{L}\{E_1; s-1\} + a}{s(s-1)\zeta(s)} - \frac{1}{s} + \frac{1}{s-1},$$

Consider the boundary distributions

$$g_1(t) = \lim_{\sigma \rightarrow 1^+} \frac{\mathcal{L}\{E'_2; \sigma-1+it\}}{(\sigma-1+it)\zeta(\sigma+it)} \quad \text{in } \mathcal{S}'(\mathbb{R}),$$

and

$$g_2(t) = -\lim_{\sigma \rightarrow 1^+} \left( \frac{(2\sigma-1+2it)\mathcal{L}\{E_1; \sigma-1+it\} + a}{(\sigma+it)(\sigma-1+it)\zeta(\sigma+it)} + \frac{1}{\sigma+it} \right) \quad \text{in } \mathcal{S}'(\mathbb{R}).$$

Taking boundary values in (11), we have  $\hat{T}(t) = g_1(t) + g_2(t) + \hat{H}(t)$ , where  $H$  is the Heaviside function. Fix  $\phi \in \mathcal{D}(-c, c)$ . Notice that  $g_2$  is actually a continuous function on  $(-3c, 3c)$ , thus,

$$\begin{aligned} \int_{-\infty}^{\infty} T(u)\hat{\phi}(u-h)du &= \left\langle g_1(t), e^{iht}\phi(t) \right\rangle + \int_{-c}^c e^{iht}g_2(t)\phi(t)dt + \int_{-h}^{\infty} \hat{\phi}(u)du \\ &= \left\langle g_1(t), e^{iht}\phi(t) \right\rangle + o(1) + O(1). \end{aligned}$$

Our task is then to demonstrate that  $\langle g_1(t), e^{iht}\phi(t) \rangle = O(1)$ . Let  $M \in \mathcal{S}'(\mathbb{R})$  be the distribution supported in the interval  $[0, \infty)$  that satisfies  $\mathcal{L}\{M; s-1\} = ((s-1)\zeta(s))^{-1}$ . Notice that  $g_1 = \widehat{(E'_2 * M)}$ . Fix an even function  $\eta \in \mathcal{D}(-3c, 3c)$  such that  $\eta(t) = 1$  for all  $t \in (-2c, 2c)$ . Then,  $\eta(t)it\zeta(1+it) \neq 0$  for all  $t \in (-2c, 2c)$ ; moreover, it is the Fourier transform of the  $L^1$ -function  $\chi_1 * E_1 + \chi_2$ , where  $\hat{\chi}_1(t) = it(1+it)\eta(t)$  and  $\hat{\chi}_2(t) = a(1+it)\eta(t)$ . We can therefore apply the Wiener division theorem [12, p. 88] to  $\eta(t)it\zeta(1+it)$  and  $\phi(t)$ . So we find  $f \in L^1(\mathbb{R})$  such that

$$\hat{f}(t) = \frac{\phi(t)}{\eta(t)it\zeta(1+it)}.$$

Hence,

$$\left\langle g_1(t), e^{iht}\phi(t) \right\rangle = \left\langle (E'_2 * M)(u), \hat{\phi}(u-h) \right\rangle = (E_2 * (\hat{\eta})' * f)(h) = O(1),$$

because  $E_2 \in L^\infty(\mathbb{R})$  and  $(\hat{\eta})' * f \in L^1(\mathbb{R})$ , whence (10) follows.  $\square$

## REFERENCES

- [1] E. P. Balanzario, *On Chebyshev's inequalities for Beurling's generalized primes*, Math. Slovaca **50** (2000), 415–436.
- [2] P. T. Bateman, H. G. Diamond, *Asymptotic distribution of Beurling's generalized prime numbers*, Studies in Number Theory, pp. 152–210, Math. Assoc. Amer., Prentice-Hall, Englewood Cliffs, N.J., 1969.
- [3] A. Beurling, *Analyse de la loi asymptotique de la distribution des nombres premiers généralisés*, Acta Math. **68** (1937), 255–291.
- [4] H. Bremermann, *Distributions, complex variables and Fourier transforms*, Addison-Wesley, Reading, Massachusetts, 1965.
- [5] H. G. Diamond, *A set of generalized numbers showing Beurling's theorem to be sharp*, Illinois J. Math. **14** (1970), 29–34.
- [6] H. G. Diamond, *Chebyshev estimates for Beurling generalized prime numbers*, Proc. Amer. Math. Soc. **39** (1973), 503–508.
- [7] H. G. Diamond, *Chebyshev type estimates in prime number theory*, in: Sémin. Théor. Nombres, 1973–1974, Univ. Bordeaux, Exposé **24**, (1974).
- [8] R. Estrada, R. P. Kanwal, *A distributional approach to asymptotics. Theory and applications*, Second edition, Birkhäuser, Boston, 2002.
- [9] R. S. Hall, *Beurling generalized prime number systems in which the Chebyshev inequalities fail*, Proc. Amer. Math. Soc. **40** (1973), 79–82.
- [10] J.-P. Kahane, *Sur les nombres premiers généralisés de Beurling. Preuve d'une conjecture de Bateman et Diamond*, J. Théor. Nombres Bordeaux **9** (1997), 251–266.
- [11] J.-P. Kahane, *Le rôle des algèbres  $A$  de Wiener,  $A^\infty$  de Beurling et  $H^1$  de Sobolev dans la théorie des nombres premiers généralisés de Beurling*, Ann. Inst. Fourier (Grenoble) **48** (1998), 611–648.
- [12] J. Korevaar, *Tauberian theory. A century of developments*, Grundlehren der Mathematischen Wissenschaften, 329, Springer-Verlag, Berlin, 2004.
- [13] J.-C. Schlage-Puchta, J. Vindas, *The prime number theorem for Beurling's generalized numbers. New cases*, Acta Arith. **153** (2012), 299–324.
- [14] L. Schwartz, *Théorie des Distributions*, Hermann, Paris, 1966.
- [15] J. Vindas, *Chebyshev estimates for Beurling generalized prime numbers. I*, preprint (arXiv:1201.1405v1).
- [16] J. Vindas, R. Estrada, *A quick distributional way to the prime number theorem*, Indag. Math. (N.S.) **20** (2009), 159–165.
- [17] V. S. Vladimirov, *Methods of the theory of generalized functions*, Analytical Methods and Special Functions, 6, Taylor & Francis, London, 2002.
- [18] W.-B. Zhang, *Chebyshev type estimates for Beurling generalized prime numbers*, Proc. Amer. Math. Soc. **101** (1987), 205–212.

DEPARTMENT OF MATHEMATICS, GHENT UNIVERSITY, KRIJGSLAAN 281 GEBOUW S22, B 9000 GENT, BELGIUM

*E-mail address:* jvindas@cage.Ugent.be