1	Published as: Veraverbeke, S., Somers, B., Gitas, I., Katagis, T., Polychronaki,
2	A., Goossens, R. (2012), Spectral mixture analysis to assess post-fire vegetation
3	regeneration using Landsat Thematic Mapper imagery: accounting for soil
4	brightness variation. International Journal of Applied Earth Observation and
5	Geoinformation, vol. 14(1): 1-11.
6	
7	
8	
9	Spectral mixture analysis to assess post-fire vegetation regeneration using Landsat
10	Thematic Mapper imagery: accounting for soil brightness variation
11	S. VERAVERBEKE* ^{1,2} , B. SOMERS ³ , I. GITAS ⁴ , T. KATAGIS ⁴ , A. POLYCHRONAKI ⁴
12	AND R. GOOSSENS ¹
13	¹ Department of Geography, Ghent University, Krijgslaan 281 S8, BE-9000 Ghent, Belgium,
14	rudi.goossens@ugent.be
15	² Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, US-
16	91109 Pasadena, USA, Sander.S.Veraverbeke@jpl.nasa.gov
17	³ Centre for Remote Sensing and Earth Observation Processes (TAP), Flemish Institute for
18	Technological Research (VITO), Boeretang 200, BE-2400 Mol, Belgium,
19	ben.somers@vito.be
20	⁴ Laboratory of Forest Management and Remote Sensing, Aristotle University of
21	Thessaloniki, PO BOX 248, GR-54124, Greece, igitas@for.auth.gr thkatag@for.auth.gr
22	anpolych@for.auth.gr
23	*Corresponding author, phone 18183540278, fax 18183545148
24	Abstract

25 Post-fire vegetation cover is a crucial parameter in rangeland management. This study aims to 26 assess the post-fire vegetation recovery three years after the large 2007 Peloponnese (Greece) 27 wildfires. Post-fire recovery landscapes typically are mixed vegetation-substrate environments 28 which makes Spectral Mixture Analysis (SMA) a very effective tool to derive fractional 29 vegetation cover maps. Using a combination of field and simulation techniques this study 30 aimed to account for the impact of background brightness variability on SMA model 31 performance. The field data consisted out of a spectral library of in situ measured reflectance 32 signals of vegetation and substrate and 78 line transect plots. In addition, a Landsat Thematic 33 Mapper (TM) scene was employed in the study. A simple SMA, in which each constituting 34 terrain feature is represented by its mean spectral signature, a multiple endmember SMA 35 (MESMA) and a segmented SMA, which accounts for soil brightness variations by forcing 36 the substrate endmember choice based on ancillary data (lithological map), were applied. In 37 the study area two main spectrally different lithological units were present: relatively bright 38 limestone and relatively dark flysch (sand-siltstone). Although the simple SMA model 39 resulted in reasonable regression fits for the flysch and limestones subsets separately 40 (coefficient of determination R^2 of respectively 0.67 and 0.72 between field and TM data), the performance of the regression model on the pooled dataset was considerably weaker (R^2 = 41 42 (0.65). Moreover, the regression lines significantly diverged among the different subsets 43 leading to systematic over-or underestimations of the vegetative fraction depending on the 44 substrate type. MESMA did not solve the endmember variability issue. The MESMA model 45 did not manage to select the proper substrate spectrum on a reliable basis due to the lack of 46 shape differences between the flysch and limestone spectra,. The segmented SMA model 47 which accounts for soil brightness variations minimized the variability problems. Compared 48 to the simple SMA and MESMA models, the segmented SMA resulted in a higher overall correlation ($R^2 = 0.70$), its regression slope and intercept were more similar among the 49

different substrate types and its resulting regression lines more closely resembled the expected one-one line. This paper demonstrates the improvement of a segmented approach in accounting for soil brightness variations in estimating vegetative cover using SMA. However, further research is required to evaluate the model's performance for other soil types, with other image data and at different post-fire timings.

55 Keywords: fire; vegetation recovery; Landsat Thematic Mapper; Spectral Mixture Analysis;
56 MESMA; segmentation

57 1 Introduction

58 Wildfires are a determining disturbance in almost all terrestrial ecosystems (Dwyer et al., 59 1999; Bond and Keeley, 2005; Riaño et al., 2007). They partially or completely consume the 60 protective vegetation and organic litter cover, which can destabilize surface soils on steep 61 slopes (Shakesby and Doerr, 2006). Shortly after the fire, infiltration significantly decreases 62 whereas surface erosion increases due the bares soil's elevated exposure to raindrop impact 63 and surface run-off. What is more, biomass burning instigates abrupt changes in ecological 64 processes and carbon fluxes (Epting and Verbyla, 2005; Amiro et al., 2006). After the fire 65 event a more gradual regeneration process is generally initiated (Viedma et al., 1997; van 66 Leeuwen, 2008). Post-fire recovery rates depend on fire severity (Díaz-Delgado et al., 2003), 67 soil properties (Bisson et al., 2008), post-fire meteorological conditions (Henry and Hope, 68 1998; van Leeuwen et al., 2010) and ecotype (Viedma et al., 1998; Veraverbeke 2010a, 2011, 69 Lhermitte et al. 2011). In fire-adapted sclerophyllous shrub lands, for example, recovery only 70 takes a few years (Viedma et al., 1997; Pausas and Verdu, 2005) whereas in boreal forests 71 recovery lasts several decades (Nepstad et al., 1999). The carbon sequestration by 72 regenerating plants partly compensates the fire's emissions and thus importantly influences the 73 net changes caused by fire (Amiro et al., 2006; Randerson et al., 2006). Vegetation recovery is 74 thus the main factor in limiting the damage of fire and its consequences. The assessment of post-fire vegetation regeneration is of crucial importance for the understanding of the environmental impacts of fire and to support sustainable rangeland management after fire. In comparison with labor-intensive field work, the synoptic nature of remote sensing systems offers a time-and cost-effective means to fulfill this duty.

79 The remote sensing of post-fire vegetation recovery has a long tradition in the use of the 80 Normalized Difference Vegetation Index (NDVI) (a.o. Viedma et al., 1997; Díaz-Delgado et 81 al., 2003; van Leeuwen 2008; Clemente et al., 2009; Lhermitte et al., 2011) because of the 82 well established relationship between the index and above-ground biomass in a wide range of 83 ecosystems (Carlson and Ripley, 1997; Henry and Hope, 1998; Cuevas-González et al., 84 2009). At moderate resolution scale Landsat data typically are the standard of choice. A 85 plethora of studies demonstrated the utility of Landsat NDVI to assess post-fire vegetation 86 dynamics (a.o. Viedma et al., 1997; Díaz-Delgado et al., 2003; McMichael et al., 2004; Malak 87 and Pausas, 2006; Clemente et al., 2009). These studies were restricted to a limited number of 88 images. Some other studies, however, used low resolution time series to monitor regeneration 89 processes. Cuevas-González et al. (2009), for example, monitored post-fire forest recovery in 90 Siberia using Moderate Resolution Imaging Spectroradiometer (MODIS)-derived NDVI data, 91 while van Leeuwen et al. (2010) conducted a similar study in three different study areas 92 (Spain, Israel and USA). At the expense of spatial detail, these studies offer the advantage of 93 image acquisition with high temporal frequency (van Leeuwen et al., 2010; Veraverbeke et 94 al., 2011). Including the temporal dimension, however, often hampers the differentiation 95 between post-fire effects and seasonal dynamics (Veraverbeke et al., 2010a, Lhermitte et al., 96 2011).

97 The post-fire environment typically consists of a mixture of vegetation and substrate. Thus, 98 monitoring post-fire regeneration processes essentially poses a sub-pixel issue at the 99 resolution of most operational satellite systems such as Landsat. A number of image analysis

100 techniques accommodating mixing problems exist (Atkinson et al., 1997; Arai, 2008) with 101 Spectral Mixture Analysis (SMA) being the most common technique utilized in many 102 applications (a.o. Roberts et al., 1998; Asner and Lobell, 2000; Riaño et al., 2002; Roder et 103 al., 2008; Somers et al. 2010ab). SMA effectively addresses this issue by quantifying the sub-104 pixel fraction of cover of different endmembers, which are assumed to represent the spectral 105 variability among the dominant terrain features. A major advantage of SMA is its ability to 106 detect low cover fractions, something which remains difficult with the traditional vegetation 107 indices (VIs) approach (Henry and Hope, 1998; Elmore et al., 2000; Rogan and Franklin, 108 2001). Moreover, SMA directly results in quantitative abundance maps, without the need of 109 an initial calibration based on field data as with VIs (Somers et al. 2010a, Vila and Barbosa, 110 2010). With regards to post-fire effects, rather few studies employed SMA to monitor post-111 fire vegetation responses (Riaño et al., 2002; Roder et al., 2008; Sankey et al., 2008; Vila and 112 Barbosa, 2010). Although results of these studies were consistent, they were all restricted to 113 simple linear SMA models in which only one spectrum was allowed for each endmember. As 114 a consequence, the performance of these SMA models often appeared to be suboptimal 115 (Roder et al., 2008; Vila and Barbosa, 2010) because these models did not incorporate the 116 natural variability in scene conditions of terrain features inherent in remote sensing data 117 (Asner, 1998). To overcome this variability effect a number of solutions have been presented 118 (Asner and Lobell, 2000; Zhang et al. 2004, 2006; Somers et al. 2010b). Multiple endmember 119 SMA (MESMA), as presented by Roberts et al. (1998), probably is the most widely used 120 technique to reduce the variability effects. In this model natural variability is included by 121 allowing multiple endmembers for each constituting terrain feature. These endmember sets 122 represent the within-class variability (Somers et al., 2009a) and MESMA models search for 123 the most optimal endmember combination by reducing the residual error when estimating 124 fractional covers (Asner and Lobell, 2000). Rogge et al. (2006), however, clearly

demonstrated that reducing the residual error by applying MESMA not always results in the selection of the most appropriate endmember spectrum. An initial segmentation of the area prior to the unmixing process in order to retain areas which reveal a high similarity in the spectral properties of a certain endmember has been presented as a sound and computationally efficient solution to address this issue (Rogge et al., 2006).

130 In this context, we aim to map vegetation abundance three year after the large 2007 131 Peloponnese (Greece) wildfires using Landsat Thematic Mapper (TM) imagery. We contrast 132 traditional simple SMA with one spectrum for each endmember with two approaches who 133 account for the natural variability in substrates. The first approach is MESMA while the 134 second method is a segmented SMA in which ancillary information (lithological map) is used 135 to force the endmember selection. Using a combination of field and simulation techniques the 136 accuracy of the MESMA and segmented SMA is assessed and compared to the traditional 137 simple SMA.

138 **2 Methodology**

139 **2.1 Study area**

140 The study focuses on the recovery of several large burned areas situated at the Peloponnese 141 peninsula, in southern Greece (36°50'-37°40' N, 21°30'-22°20' E) (Figure 1). The fire scars 142 date from the 2007 summer. These fires were the worst natural disaster of the last decades in 143 Greece, both in terms of human losses and the extent of the burned area. Elevations range 144 between 0 and 2404 m above sea level. Limestone sediments cover most of the mountainous 145 inland. Also significant outcrops of flysch sediments occur (Institute of Geology and Mineral 146 Exploration, 1983; Higgins and Higgins, 1996). Flysch sediments are dominated by sandstone 147 with finer siltstone and clay (Institute of Geology and Mineral Exploration, 1983; Higgins and 148 Higgins, 1996). The hilly and mountainous inland is covered with shallow and gravely soils 149 (European Commission, 2005). The climate is typically Mediterranean with hot, dry summers

150 and mild, wet winters. For the Kalamata meteorological station $(37^{\circ}4' \text{ N}, 22^{\circ}1' \text{ E})$ the average 151 annual temperature is 17.8 °C and the mean annual precipitation equals 780 mm (Hellenic 152 National Meteorological Service, www.hnms.gr, accessed 20 December 2010). The fires 153 consumed more than 175 000 ha, which merely consisted of shrub land and pine forest 154 (Veraverbeke et al., 2010a). Black pine (Pinus nigra) is the dominant conifer species. The 155 shrub layer is mainly characterised by Quercus coccifera, Q. frainetto, Erica arborea and 156 Arbutus unedo. Perennial grasses cover significant parts of the ground. These grasses reveal 157 summer-dormancy and are not photosynthetically active during the Mediterranean summer 158 drought (Volaire and Lelievre, 2010). Mediterranean-type shrub lands are highly resilient to 159 burning due to both obligate seeder and resprouter fire-adapted strategies. They regenerate in 160 a couple of years (Trabaud, 1981; Capitaino and Carcaillet, 2008) in a so called 161 autosuccession process (Hanes, 1971). Conversely, the recovery of the forests is considerably 162 slower and can take up to several decades (Viedma et al., 1997; van Leeuwen et al, 2010).

163 FIGURE 1 HERE

164 **2.2 Field data**

165 **2.2.1 Spectral library**

166 In September 2010, field spectrometry measurements of the dominant terrain features (i.e. 167 endmembers) were collected in the burned areas three years after the fire. Measurements were 168 obtained within one hour of local solar noon on clear-sky days with a single channel 169 spectroradiometer (UniSpec-SC) covering the 300-1100 nm spectral domain with a 3.7 nm 170 resolution (PP Systems, 2006). 59 top-of-canopy (TOC) measurements of regenerating 171 vegetation were recorded. Canopy height ranged between 0.5 and 2 m which made it possible 172 to collect TOC signatures without the need of a measurement platform. In addition, 39 spectra 173 of non-photosynthetic (i.e. brown) vegetation and 29 spectra of shallow and gravelly soils of 174 both flysch and limestone sediments were also obtained: 15 above flysch substrate and 14

175 above limestone substrate. The spectra of each class were collected from various locations 176 throughout the study area. All measurements were obtained while holding the sensor 0.3 to 177 0.5 m above the sample. The shadow endmember was assumed to be a uniformly dark 178 component and was modeled as a flat 1 % reflectance (Lelong et al., 1998; Somers et al., 179 2009a, 2010ab). The spectra were resampled to the TM wavebands to facilitate further 180 analysis. Figure 2A shows the mean spectral signatures of each constituting endmember. The 181 TM visual and near-infrared (VNIR) band passes are indicated in the figure. In the area two 182 main substrate classes appear: limestone and flysch sediments. Corresponding spectral 183 signatures are plotted with dashed lines in figure 2A, whereas figure 2B shows the occurrence 184 of these two substrate classes in the 2007 burned areas. This classification was obtained after 185 interpreting and digitizing the geological map of the area (Institute for Geology and Mineral 186 Exploitation 1983). The difference in the substrates' optical properties is clear from the figure. 187 The limestone substrate is relatively bright compared to the darker flysch substrate.

188 FIGURE 2 HERE

189 2.2.2 Line transect data

190 78 line transect plots were sampled to estimate the abundance of regenerating vegetation in 191 the 2007 burned areas three years post-fire, in September 2010. 46 plots were measured in 192 flysch areas whereas the remaining 32 samples were obtained on a limestone substrate. The 193 sample scheme was designed for the 30 m Landsat resolution. The plots were selected during 194 several one-day hikes based on a stratified sampling approach taking into account the 195 constraints on mainly accessibility and time, encompassing the range of variability in 196 recovery rates in the study area. The plot's centre coordinates were recorded with a handheld 197 Garmin eTrex Visa Global Positioning System (GPS, 15 m error in x and y: Garmin, 2005). 198 To minimize the influence of spatial autocorrelation plots were located at least 500 m apart. 199 They consist of two perpendicular 60 m line transects, of which the first was directed northsouth. Using the point-intercept method (Bonham, 1989; Clemente et al., 2009; van Leeuwen
et al., 2010; Vila and Barbosa, 2010) at a 1 m interval along the line transect, vegetation
abundance was estimated. The fraction of vegetation cover equals the total number of
vegetation interception points divided by the total number of interception points (Bonham,
1989) (Figure 3). 60 m linear transects were preferred to 30 m transects to anticipate potential
satellite misregistration. Moreover, samples were located in relatively homogeneous areas of
regrowth. Figure 4 shows example plot photographs of shrub land at different recovery rates.

207 FIGURE 3 HERE

208 FIGURE 4 HERE

209 **2.3 Satellite data and preprocessing**

210 One 30 m resolution Landsat TM image (path/row 184/34, acquired on July 18, 2010) was 211 used in this study. We have tried to minimize the difference in phenology between the image 212 and field data acquisition, however, small differences in phenology and resulting reflectance 213 might influence the analysis. The image of July 18, 2010 was the acquisition that most closely 214 resembled the ecosystem status as measured during the field campaign in September 2010. 215 Analysis was restricted to wavebands between 400 and 1100 nm because of the consistency 216 with the field spectral library. In addition, these wavebands show a higher signal-to-noise 217 ratio than other spectral regions (Chen and Vierling, 2006). For TM imagery this results in 218 four spectral bands: blue (B, 450-520 nm), green (G, 520-600 nm), red (R, 630-690 nm) and 219 near infrared (NIR, 760-900).

The TM image was geometrically corrected using a set of homologous points of a previously georeferenced TM image of the study area (Veraverbeke et al., 2010ab, 2011). The resulting Root Mean Squared Error (RMSE) was lower than 0.5 pixels. The image was registered in Universal Transverse Mercator (UTM, zone 34S), with ED 50 (European Datum 1950) as geodetic datum. 225 Raw digital numbers (DNs) were scaled to at-sensor radiance values (L_s) (Chander et al., 226 2007). The radiance to reflectance conversion was performed using the COST method 227 (Chavez 1996):

228
$$r_a = \frac{\pi (L_s - L_d)}{(E_o / d^2) (\cos \theta_z)^2}$$
(1)

where r_a is the atmospherically corrected reflectance at the surface; L_s is the at-sensor radiance (Wm⁻²sr⁻¹); L_d is the path radiance (Wm⁻²sr⁻¹); E_o is the solar spectral irradiance (Wm⁻²); d is the earth-sun distance (astronomical units); and θ_z is the solar zenith angle. The COST method is a dark object subtraction (DOS) approach that assumes 1 % surface reflectance for dark objects (e.g. deep water).

Additionally, it was necessary to correct for differing illumination effects due to topography. This was done based on the modified c-correction method (Veraverbeke et al., 2010c), a modification of the original c-correction approach (Teillet et al., 1982), using a digital elevation model (DEM) and knowledge of the solar zenith and azimuth angle at the moment of image acquisition. Topographical slope and aspect data were derived from a 30 m DEM (Hellenic Military Geographical Service, HMGS) resampled and co-registered with the TM images. The illumination is modeled as:

241
$$\cos \gamma_i = \cos \theta_p \cos \theta_z + \sin \theta_p \sin \theta_z \cos(\varphi_a - \phi_o)$$
 (2)

where γ_i is the incident angle (angle between the normal to the ground and the sun rays); θ_p is the slope angle; θ_z is the solar zenith angle; ϕ_a is the solar azimuth angle; and ϕ_o is the aspect angle. Then terrain corrected reflectance r_i is defined as:

245
$$r_{t} = r_{a} \left(\frac{1 + c_{k}}{\cos \gamma_{i} + c_{k}} \right)$$
(3)

where c_k is a band specific parameter $c_k = b_k/m_k$ where b_k and m_k are the respective intercept and slope of the regression equation $r_a = b_k + m_k \cos \gamma_i$. Since topographic normalization works better when applied separately for specific land cover types (Bishop and Colby, 2002) specific c-values for the recovering 2007 scars were calculated by masking the unburned areas (Veraverbeke et al., 2010c).

251 2.4 SMA

SMA is a commonly used image analysis technique to derive abundance estimates of dominant ground components (e.g. green vegetation, substrates, etc.). Although some authors recognize the occurrence of multiple photon scattering (Ray and Murray, 1996; Somers et al. 2009b), most vegetation monitoring studies consider a mixed pixel spectrum (r) as a linear combination of pure spectral signals of its constituent components or endmembers, weighted by their corresponding sub-pixel fractional covers (Adams et al., 1986):

$$258 r = Mf + \varepsilon (4)$$

where *M* is a matrix in which each column corresponds with the pure spectral signal of a specific endmember, *f* is a column vector $[f_1,...,f_m]^T$ denoting the cover fractions occupied by each of the endmembers in the pixel. In this study, green vegetation, brown vegetation, substrate and shadow are the endmembers of interest. ε represents the residual error.

Equation 4 is often solved by estimating abundance fractions using least squares error estimates. Once the pure spectral signals of the endmembers are known, the fraction vector f is calculated by minimizing the following equation:

266
$$\sum_{i=1}^{n} \varepsilon_{i}^{2} = \sum_{i=1}^{n} \left(\sum_{j=1}^{n} (M_{i,j} \times f_{j}) - r_{i} \right)^{2}$$
(5)

where *n* is the number of spectral bands (Barducci and Mecocci, 2005). Generally, physically meaningful abundance estimates are obtained by constraining the cover fraction to sum to unity and to be positive (Roberts et al., 1993). 270 Endmembers may be derived from spectral libraries built from field or laboratory 271 measurements (Roberts et al., 1998). Yet, endmember reference spectra can also be derived 272 directly from the image data themselves (Bateson et al., 2000). Even in quickly recovering 273 ecosystems, the diameter of woody individuals seldom exceeds 2 m in a medium-term 274 perspective (3 years post-fire) (Keeley and Keeley, 1981; Malanson and Trabaud, 1988; 275 Clemente et al., 1996). As a result, the occurrence of pure image pixels in the post-fire 276 recovery areas is very rare at the Landsat 30 m resolution. As a consequence we acquired pure 277 field spectra as described in section 2.2.1. To account for endmember variability, several 278 authors suggest to evaluate multiple endmember combinations from the spectral library 279 instead of using a fixed mean signature per endmember (Roberts et al., 1998; Asner and 280 Lobell, 2000). Then, pixels are iteratively decomposed using different sets of endmember 281 combinations and ultimately these fractional covers corresponding with the iteration that 282 revealed the lowest least squares error are selected. This method is widely known as MESMA 283 (Roberts et al., 1998). MESMA, however, does not always select the most appropriate 284 endmember spectrum (Rogge et al., 2006). A prior segmentation of the imagery in zones that 285 reveal a high similarity in the spectral properties of a certain endmember has been presented 286 as a sound and computationally efficient solution for this issue (Rogge et al., 2006).

We executed three linear unmixing models. Each model used the mean spectrum as endmember for green and brown vegetation. The difference between the different models, however, is the definition of the substrate endmember:

- 290 291
- The first model using the mean substrate spectrum as soil endmember and is referred to as simple SMA.
- 292 293
- The second model is a simple MESMA in which two different soil spectra are incorporated (the mean flysch and limestone spectra).

The third model forces the choice between the mean limestone or flysch spectrum
 based on ancillary data. We used a generalized lithological map (Figure 2B) to ensure
 the proper substrate endmember selection. This technique is referred to a segmented
 SMA.

298 Preliminary experiments indicated that it was impossible to discriminate between the brown 299 vegetation and substrate endmembers. This is explained by their high spectral similarity 300 (Figure 2A) and corroborates with previous findings of Goodwin et al. (2005), Gill and Phinn 301 (2009) and Somers et al. (2010b). As such, the best characterization of image variance was 302 achieved with a three-endmember (green vegetation, substrate and shadow) model. To obtain 303 ecologically meaningful estimates, the shadow cover fraction cover was distributed over the 304 green vegetation and substrate components, proportionally to the estimated fractional cover of 305 these components (Roder et al., 2008).

306 2.5 Analysis method

307 2.5.1 Simulated data

308 The analysis is twofold. Firstly, we used the spectral library with pure substrate (29) and 309 vegetation signals (59) to create simulated mixed pixels. According to equation 4, a total of 310 1000 mixed vegetation-substrate spectra were calculated. 500 of them were constituted with a 311 limestone spectrum while for the other half a flysch endmember was used. Pure pixel spectra 312 combinations and fractional covers were randomly assigned to each pixel. To account for 313 ambient and instrumental error, normally distributed noise was added to the signal (with a 314 mean of zero and standard deviation ranging from 0 % to 15 % of the mixed signal, Asner and 315 Lobell, 2000). Subsequently, each mixed spectrum was unmixed using the three different 316 models. The first model, traditional simple SMA, uses one spectrum for each endmember. 317 The second model, MESMA, chooses the substrate endmember (flysch or limestone) 318 corresponding with the lowest residual error. Finally, the segmented SMA model forces the

319 choice between the limestone or flysch endmember based on ancillary knowledge. Simulated 320 data supply a reliable means to evaluate the performance of the various models as it inherently 321 provides correct validation data (Rogge et al., 2006). The performance of each model was 322 expressed in the coefficient of determination (R^2) of the linear regression with the estimated 323 vegetation fractions as independent variable and the modeled fractional vegetation covers a 324 dependent variable. Separate regression models were performed for the limestone mixtures, 325 the flysch mixtures and the pooled dataset comining limestone and flysch mixtures. In 326 addition, the selection of the proper substrate endmember by the MESMA model was 327 evaluated using the knowledge of the set-up of the simulation experiment as reference data.

328 2.5.2 Landsat imagery

329 The second part of the analysis focused on the Landsat TM data. The same three unmixing 330 models were applied and vegetation fractional covers of the line transect locations were 331 extracted by calculating the mean index value of a 3-by-3 pixels matrix. It is widely accepted 332 that using the mean of a pixel matrix minimizes the effect of potential misregistration (Ahern 333 et al. 1991). Linear regressions were performed to correlate the TM fractional covers 334 (independent variables) and line transect field data of vegetation recovery (dependent variables). Regression model results were compared using the R² statistic. Again, separate 335 336 regression models were performed for the 32 limestone plots, for the 46 flysch samples and 337 for the 78 field ratings together. The ancillary knowledge of the constituting substrate 338 endmember was also used to assess the performance of the MESMA model's endmember 339 spectrum selection. The best method was used to map the vegetation abundance three years 340 after the large 2007 Peloponnese wildfires.

341 3 Results

342 **3.1 Simulated data**

343 Figure 5 displays the scatter plots and regression lines of the simulation experiments. In figure 344 5A the results of the traditional SMA model are visualized, while figure 5C and 5E 345 respectively depict the outcomes of the MESMA and segmented SMA models. A comparison between the simple SMA and MESMA model learns that the R^2 between modeled and 346 347 estimated fraction covers was higher for MESMA compared to simple SMA for the flysch 348 subset, limestone subset and the whole dataset (respectively 0.75, 0.75 and 0.68 for simple 349 SMA and 0.79, 0.79 and 0.77 for the three datasets for MESMA). However, the goodness-offit of the segmented SMA for the pooled dataset was yet higher ($R^2 = 0.79$), whereas R^2 350 values of the substrate subsets were equal to the MESMA model. Moreover, for the 351 352 segmented SMA the regression parameters of the flysch subset, limestone subset and pooled 353 dataset closely resembled each other (slope respectively 0.77, 0.78 and 0.78 and intercept 0.12 354 for three datasets) whereas with simple unmixing regression slope (0.90 for the flysch subset, 355 0.70 for the limestone subset and 0.74 overall) and intercept (-0.04 for the flysch subset, 0.21 356 for the limestone subset and 0.12 pooled) significantly diverged. Also for MESMA a similar 357 divergence was present in the data: the regression slope equalled respectively 1, 0.79 and 0.86 358 for the flysch, limestone and pooled data, whereas the intercept was respectively -0.08, 0.12 359 and 0.04. The divergence of the different regression lines as observed with simple SMA and 360 MESMA was especially obvious for low vegetation cover estimates. Figures 5B, 5D and 5F 361 respectively show the same model as presented in figures 5A, 5C and 5E, however, in these 362 models randomly distributed noise was added. This did not impact the trends described above, however, R^2 values revealed a small drop compared to their noise-free counterpart. The only 363 364 exception against this drop was the traditional SMA model of the pooled dataset which retained its $R^2 = 0.68$. 365

366 FIGURE 5 HERE

The error matrix of the selection of the substrate spectrum by MESMA based on simulated data is tabulated in Table 1. The overall accuracy equalled 61 % and a relatively low Kappa coefficient of 0.21 was obtained. The MESMA model's substrate spectrum selection revealed a high omission error for the flysch class (producer's accuracy of 29 %) and relatively high commission error for the limestone class (user's accuracy of 57 %).

372 TABLE 1 HERE

373 **3.2 Landsat imagery**

374 Figure 6 presents scatter plots and regression line between the line transect field ratings and 375 the vegetation fractional covers retrieved from the Landsat imagery. In corroboration with the 376 results from the simulations (Figure 5), the regression parameters of the segmented SMA 377 model were very similar for the flysch subset, limestone subset and pooled dataset (slope 378 respectively 1.02, 0.99 and 1.03 and intercept respectively -0.06, -0.08 and -0.08). This 379 contrasts with the more differing regression slope and intercept of the simple SMA (slope 380 respectively 1.07, 0.93 and 0.83 for the flysch subset, limestone subset and pooled dataset and 381 intercept respectively -0.15, -0.01 and 0) and MESMA models (slope respectively 0.71, 0.89) 382 and 0.86 for the flysch subset, limestone subset and pooled dataset and intercept respectively 383 0.10, -0.01 and 0.06). For the simple SMA, this did not result in less optimal regression models for the substrate subsets, however, the overall R^2 was clearly higher for the model that 384 forced the flysch-limestone endmember choice ($R^2=0.70$ versus $R^2=0.65$ for simple SMA). 385 For MESMA, the goodness-of-fit was lower for both subset and pooled data (e.g. $R^2=0.63$ for 386 387 the pooled dataset). In addition, the regression lines of the segmented SMA more closely 388 resembled the expected one-one line compared to the other models.

389 FIGURE 6 HERE

390 The error matrix of the selection of the substrate spectrum by MESMA based on field data is

391 listed in Table 2. Similar to the results of table 1, the overall accuracy equalled 62 % and a

relatively low Kappa coefficient of 0.18 was obtained. The MESMA model's substrate spectrum selection revealed a high omission and commission error for the limestone class which resulted in a relatively low producer's accuracy (41 %) and user's accuracy (54 %) for this class. Producer's and user's accuracy for the flysch category were slightly higher (respectively 76 % and 65 %).

397 The ancillary information of figure 2B was used to differentiate between relatively bright 398 (limestone) and dark (flysch) substrates when mapping the post-fire vegetation cover while 399 accounting for background variability using the segmented SMA model (Figure 7).

400 4 Discussion

401 Post-fire recovery landscapes essentially are mixed vegetation-substrate environments. A 402 plethora of studies made use of this feature to map post-fire vegetation cover with the NDVI 403 (a.o. Viedma et al., 1997; Díaz-Delgado et al., 2003; McMichael et al., 2004; Malak and 404 Pausas, 2006; Clemente et al., 2009). To obtain qualitative fractional cover maps, these index 405 values require a prior calibration with field estimates of vegetation cover (Clemente et al., 406 2009). In this study, SMA demonstrated to be a strong alternative for the spectral indices 407 approach, as SMA outputs fraction images without an initial regression fit between remotely 408 sensed data and field ratings.

409 The regression fit between the line transect field estimates of recovery and the most optimal SMA resulted in moderate-high $R^2 = 0.70$. The residual variation can be explained by the fact 410 411 that both field and remotely sensed estimates are imperfect proxies for vegetation cover. The 412 line transect method is a relatively rough approach to estimate fractional vegetation cover 413 while several noise factors hamper satellite image analysis. Inaccurate atmospheric correction 414 (Gong et al., 2008), suboptimal illumination correction (Veraverbeke et al. 2010c), sensor 415 noise (Plaza et al., 2004), slight differences in acquisition timing between field and image data 416 or the unmixing model structure itself (e.g. non-linear mixing due to multiple photon scattering among different ground components, Borel and Gerstl, 1994; Somers et al., 2009b)
are all known to create noise in image analyses. The influence of soil brightness variation,
however, was a very important factor impacting model performance.

420 Both the simulation experiment and Landsat application demonstrated that accounting for soil 421 brightness variations by the segmented approach significantly improved the SMA model. The 422 simple SMA with one single spectrum for each endmember provided reasonable regression 423 models for each substrate class separately, however, model performance of the pooled dataset 424 was considerably weaker. This is explained by the fact that traditional SMA resulted in clearly 425 different regression lines depending on substrate class (Figure 5A, 5B and 6A). In other 426 words, the relationship between the observed (field or modeled) vegetative fraction and the 427 estimated fraction from the simple SMA model was determined by the brightness of the 428 background. Thus, neglecting this background brightness difference produced a weaker 429 overall fit. Moreover, the simple SMA model underestimates the vegetative fraction in 430 limestone areas while in flysch areas the opposite is true. As shown in figure 2A the optical 431 properties of these two substrate types are clearly different. They represent a relatively bright 432 (limestone) and dark (flysch) background. MESMA is the most widely used technique to 433 include endmember variability in a SMA model (Roberts et al., 1998). Table 1 and 2, 434 however, clearly indicated that MESMA did not manage to select the appropriate substrate 435 spectrum in this case study. The Kappa coefficients of 0.18 an 0.21 for respectively the 436 simulation experiment and the Landsat application revealed that the substrate spectrum 437 selection was only slightly better than an agreement by chance. As a consequence, MESMA 438 did not solve the substrate variability issue in this application. This can be explained by the 439 fact that the spectral signatures of limestone and flysch are almost linear translations of each 440 other (Figure 2A). Due to the lack of shape differences between these two substrate spectra, 441 MESMA did not demonstrate a strong tendency to select the appropriate soil endmember. In

442 contrary, the ultimate selection of the substrate endmember appeared to be rather arbitrarily. It
443 is recognized that when different substrate endmember spectra reveal clear shape differences,
444 MESMA can be a very straightforward solution to find the proper substrate spectrum based on
445 an iterative process (Roberts et al., 1998).

446 Because of the failure of the MESMA model in this case study, we applied a segmented 447 approach in which the substrate endmember choice was based on ancillary knowledge (i.e. the 448 simulation set up in the case of the simulations and a generalized lithological map for the 449 Landsat application). For this model, regression slope and intercept did not depend on 450 substrate class (Figures 5E, 5F and 6C). So irrespective which substrate type, the regression 451 lines were similar. As a consequence, potential over- or underestimation of vegetative cover 452 was eliminated and the performance of the pooled regression model was equally high. The 453 SMA model that accounts for soil brightness variations also produced regression fits very 454 close to the expected one-one line, which proves its consistency. These beneficial results of 455 the segmentation approach corroborate with Rogge et al. (2006) who demonstrated the 456 effectiveness of prior segmentation to overcome poor endmember spectrum selection by 457 MESMA. In addition, limiting the number of the potential endmember spectra favors the 458 computational efficiency compared to MESMA models (Rogge et al., 2006).

459 In post-fire recovery studies using SMA, Riaño et al. (2002), Roder et al. (2008) and Vila and 460 Barbosa (2010) all employed one single substrate endmember. Disregarding soil brightness 461 variations potentially adds an explanation to the observed suboptimality of the SMA outcomes 462 observed by Roder et al. (2008) and Vila and Barbosa (2010). We want to remark that in the 463 simulation model vegetation cover was slightly overestimated for very low vegetative covers, 464 while the model slightly underestimated the vegetative fraction for mixtures in which the 465 vegetation component dominates (Figure 5). For extreme fractional vegetation covers (close 466 to zero and one) the SMA simulation models showed a tendency to estimate vegetative

467 fractional cover as respectively zero and one. This explains the slight over- and 468 underestimation observed in the simulation experiment. Due to the fact that most field ratings 469 range between 20-70 % vegetative coverage, this behavior is not present in the regression fit 470 between Landsat and line transect data. In contrast, the overall regression intercept of the 471 modified SMA regression model is slightly negative (-0.08). However, the general SMA 472 constraint that fraction estimates have to be positive (Roberts et al., 1998), prevents the 473 occurrence of negative fractional covers without biophysical meaning. In the field, the 474 presence of extreme fractional covers (close to zero or one) was extremely rare, so these cases 475 do not nullify the performance of the model. In this respect, a totally different scenario would 476 emerge when one would aim to estimate the post-fire vegetation regrowth very shortly after 477 the fire, e.g. one year after the fire. Then, it would be wise to additionally evaluate the model 478 performance for very low vegetation covers. However, in contrast with our study, a one year 479 post-fire assessment would also need to include a char endmember in the model (Lewis et al., 480 2007; Robichaud et al., 2007).

481 A drawback of the proposed method is the need of ancillary data. With a combination of field 482 knowledge and lithological maps it is relatively easy to construct spectrally similar 483 lithological units, however, this possibility depends on the availability of such data layers. 484 Besides among substrates, endmember variability is also present among vegetation species. In 485 our case study, however, the variability in the spectral response of different vegetation species 486 was very small compared to large spectral differences between the substrate classes. For this 487 reason and because of the small sensitivity of broadband sensors to discriminate between 488 different vegetation types (Somers et al. 2010a), we disregarded vegetation variability in our 489 analyses. Other pathways to improve the accuracy of the recovery assessment are multiple. A 490 possible amelioration could be the inclusion of the short-wave infrared (SWIR: 1300-1700 491 nm) and mid infrared (MIR: 1700-2400 nm) spectral regions in the unmixing process. These

492 spectral regions have proven to be very effective in discriminating soil and vegetation (Drake 493 et al., 1999; Asner and Lobell, 2000). Moreover, the SWIR-MIR spectrum is very sensitive to 494 moisture content (Hunt and Rock, 1989; Zarco-Tejada et al., 2003) and are consequently 495 strongly related to plant water content. Carreiras et al. (2006) demonstrated that adding the 496 SWIR-MIR Landsat bands resulted in better estimates of tree canopy cover in Mediterranean 497 shrublands. To retain consistency with the field spectral library these wavebands were not 498 included in our study (Somers et al., 2010a). Additionally, enhancing the spectral resolution 499 by employing hyperspectral data would increase the amount of spectral detail which would 500 benefit the differentiation between spectra. By including more and other spectral wavebands 501 the unmixing model could gain discriminative power. Potentially, this would make it even 502 possible to distinguish between non-photosynthetic vegetation and substrate (Asner and 503 Lobell, 2000; Somers et al; 2010a), which appeared to be impossible based on the Landsat 504 VNIR bands.

505 5 Conclusions

506 Using a combination of field and simulation techniques, the importance of accounting for 507 background brightness variability in estimating fractional vegetation cover using SMA was 508 highlighted. Although the traditional SMA model in which the substrate endmember was 509 defined as the arithmetic mean of two flysch and limestone substrates subclasses resulted in 510 reasonable regression fits for the flysch and limestone datasets separately, the regression fit 511 performed on the pooled dataset was considerable weaker. The regression lines of the 512 different datasets (only limestone, only flysch and pooled) significantly diverged and as such 513 vegetative cover estimations depended on substrate type. The use of a single spectrum 514 substrate endmember thus resulted in an over- or underestimation of the vegetative cover 515 fraction related to background brightness differences. Traditionally, MESMA is applied to 516 address the endmember variability issue, however, in this case study MESMA did not manage

517 to select the appropriate substrate endmember due to the lack of shape difference between the 518 flysch and limestone spectra. Therefore, a prior segmentation based on ancillary information 519 (lithological map) was executed to incorporate soil color variation in a segmented SMA 520 model. This model forces the proper substrate endmember spectrum choice. The overall 521 regression fit of the segmented approach significantly improved and the discrepancy between 522 the regression of the different subsets significantly reduced. Moreover, the resulting 523 regression line very closely resembled the expected one-one line between observed and 524 estimated fractional vegetation covers.

This paper demonstrated the utility of SMA for monitoring post-fire vegetation regeneration three year after the 2007 Peloponnese wildfires. Although a segmented approach to account for soil brightness variations significantly improved the model, further research is required to evaluate the model's performance for other soil types, with other image data and at different post-fire timings.

530 Acknowledgements

The study was financed by the Ghent University special research funds (BOF: Bijzonder
Onderzoeksfonds). Part of the work was carried out at the Jet Propulsion Laboratory,
California Institute of Technology, under a contract with the National Aeronautics and Space
Administration. The contribution of Dr. Ben Somers is funded by the Belgian Science Policy
Office in the frame of the STEREO II programme – project VEGEMIX (SR/67/146).

536 **References**

Adams, J., Smith, M. & Johnson, P., 1986. Spectral mixture modelling: A new analysis of
rock and soil types at Viking Lander. J. of Geophys. Res., 91 8113–8125

- 539 Ahern, F., Erdle, T., Maclean, D., Kneppeck, I., 1991. A quantitative relationship between
- 540 forest growth rates and Thematic Mapper reflectance measurements. Int. J. Remote Sens. 12,
- 541 387-400

- 542 Amiro, B., Orchansky, A., Barr, A., Black, T., Chambers, S., Chapin, F., Goulden, M., Litvak,
- 543 M., Liu, H., McCaughey, J., McMillan, A., Randerson, J., 2006. The effect of post-fire stand
- age on the boreal forest energy balance. Agric. For. Meteorology 140, 41-50.
- 545 Arai, K., 2008. Nonlinear mixture model of mixed pixels in remote sensing satellite images
- 546 based on Monte Carlo simulation. Adv. Sp. Res. 41, 1725-1743.
- 547 Asner, G., 1998. Biophysical and biochemical sources of variability in canopy reflectance.
- 548 Remote Sens. of Environ. 64, 234-253
- 549 Asner, G., Lobell, D., 2000. A biogeophysical approach for automated SWIR unmixing of
- soils and vegetation. Remote Sens. Environ. 74, 99-112
- 551 Atkinson, P., Cutler, M., Lewis, H., 1997. Mapping sub-pixel proportional land-cover with
- 552 AVHRR imagery. Int. J. Remote Sens. 18, 917-935
- 553 Barducci, A., Mecocci, A., 2005. Theoretical and experimental assessment of noise effects on
- least-squares spectral unmixing of hyperspectral images. Opt. Eng. 44, 087008
- 555 Bateson, C., Asner, G., Wessman, C., 2000. Endmember bundles: a new approach to
- 556 incorporating endmember variability in spectral mixture analysis. IEEE Trans. Geosc. Remote
- 557 Sens. 38, 1083-1094
- 558 Bishop, M., Colby, J., 2002. Anisotropic reflectance correction of SPOT-3 HRV imagery. Int.
- 559 J. Remote Sens. 23, 2125–2131
- 560 Bisson, M., Fornaciai, A., Coli, A., Mazzarini, F., Pareschi, M., 2008. The Vegetation
- 561 Resilience After Fire (VRAF) index: development, implementation and an illustration from
- 562 central Italy. Int. J. Appl. Earth Observation Geoinf. 10, 312–329
- 563 Bonham, C., 1989. Measurements for terrestrial vegetation, New York, Wiley
- 564 Bond, W., Keeley, J., 2005. Fire as a global 'herbivore': the ecology and evolution of
- 565 flammable ecosystems. Trends Ecol. Evol. 20, 387-394

- 566 Borel, C., Gerstl, S., 1994. Nonlinear spectral mixing models for vegetative and soil surfaces.
- 567 Remote Sens. Enviorn. 47, 403-416
- 568 Capitanio, R., Carcaillet, C., 2008. Post-fire Mediterranean vegetation dynamics and
- 569 diversity: a discussion of succession models. For. Ecol. Manage. 255, 431–439
- 570 Carreiras, J., Pereira, J., 2006. Estimation of tree canopy cover in evergreen oak
- 571 woodlands using remote sensing. Forest Ecol. and Manage. 23, 45-53
- 572 Carlson, T., Ripley, T., 1997. On the relation between NDVI, fractional vegetation cover and
- 573 leaf area index. Remote Sens. Environ. 62, 241-252
- 574 Chander G., Markham, L., Barsi, J., 2007. Revised Landsat-5 Thematic Mapper radiometric
- 575 calibration. IEEE Geosc. Remote Sens. Lett. 4, 490–494
- 576 Chavez, P., 1996. Image-based atmospheric corrections revisited and improved.
 577 Photogramm. Eng. Remote Sens., 6, 1025–1036
- 578 Chen, X., Vierling, L., 2006. Spectral mixture analysis of hyperspectral data acquired using a
- tethered balloon. Remote Sens.Environ. 103, 338–350
- 580 Clemente, A., Rego, F., Correia, O., 1996. Demographic patterns and productivity of post-fire
- 581 regeneration in Portuguese Mediterranean maquis. Int. J. Wildland Fire 6, 5–12
- 582 Clemente, R., Navarro Cerrillo, R., Gitas, I., 2009. Monitoring post-fire regeneration in
- 583 Mediterranean ecosystems by employing multitemporal satellite imagery. Int. J. Wildland Fire
- 584 18, 648–658
- 585 Cuevas-González, M., Gerard, F., Baltzer, H., Riaño, D., 2009. Analysing forest recovery
- 586 after wildfire disturbance in boreal Siberia using remotely sensed vegetation indices. Glob.
- 587 Chang. Biol. 15, 561-577
- 588 Díaz-Delgado, R., Lloret, F., Pons, X., 2003. Influence of fire severity on plant regeneration
- 589 by means of remote sensing. Int. J. Remote Sens. 24, 1751-1763

- 590 Drake, N., Mackin, S., Settle, J., 1999. Mapping vegetation, soils and geology in semiarid
- 591 shrublands using spectral matching and mixture modeling of SWIR AVIRIS imagery. Remote
- 592 Sens. Environ. 68, 12-25
- 593 Dwyer, E., Perreira, J., Grégoire, J., DaCamara, C., 1999. Characterization of the spatio-
- 594 temporal patterns of global fire activity using satellite imagery for the period April 1992 to
- 595 March 1993. J. Biogeogr. 27, 57-69
- 596 Elmore, A., Mustard, J., Manning, S., Lobell, D., 2000. Quantifying vegetation change in
- 597 semiarid environments: precision and accuracy of spectral mixture analysis and the 598
- normalized difference vegetation index. Remote Sens. Environ. 73, 87-102
- 599 Epting, J., Verbyla, D., 2005. Landscape-level interactions of prefire vegetation, burn
- 600 severity, and postfire vegetation over a 16-year period in interior Alaska. Can. J. For. Res., 35,
- 601 1367-1377
- 602 European Commission, 2005. Soil atlas of Europe, Office for Official Publications of the 603 European Communities, Luxembourg
- 604 Garmin, 2005. Garmin eTrex Vista personal navigator. Owner's manual and reference guide.
- 605 https://buy.garmin.com/shop/store/manual.jsp?product=010-00243-Available from:
- 606 00&cID=167&pID=163 (Last visited on 21/01/2011).
- 607 Gill, T., Phinn, S., 2009. Improvements to ASTER-derived fractional estimates of bare ground
- 608 in as savanna rangeland. IEEE Trans. Geosci. Remote Sens. 46, 662-670
- 609 Gong, S., Huang, J., Li, Y., Wang, H., 2008. Comparison of atmospheric correction
- 610 algorithms for TM image in inland waters. Int. J. Remote Sens. 29, 2199-2210
- Goodwin, N., Coops, N., Stone, C., 2005. Assessing plantation canopy condition from 611
- 612 airborne imagery using spectral mixture analysis and fractional abundances. Int. J. Appl. Earth
- 613 Observation Geoinf. 7, 11–28

- Hanes, T., 1971. Succession after fire in the chaparral of southern California. Ecol. Monogr.
 41, 27-52
- 616 Henry, M., Hope, A., 1998. Monitoring post-burn recovery of chaparral vegetation in
- 617 southern California using multi-temporal satellite data. Int. J. Remote Sens., 19, 3097-3107
- 618 Higgins, M., Higgins, R., 1996. A geological companion to Greece and the Aegean, Cornell
- 619 University Press, Cornell
- 620 Hunt, E., Rock, B., 1989. Detection of changes in leaf water content using near- and middle-
- 621 infrared reflectance. Remote Sens. Environ. 30, 43-54
- 622 Institute for Geology and Mineral Exploration, 1983. Geological map of Greece 1:500 000,
- 623 Ordnance Survey, Southampton
- Keeley, J., Keeley, S., 1981. Post-fire regeneration of southern California chaparral. Am. J.
 Bot. 68, 524-530
- 626 Lelong, C., Pinet, P., Poilvé, H. (1998). Hyperspectral imaging and stress mapping in
- 627 agriculture: a case study on wheat in Beauce (France). Remote Sens. Environ. 66, 179–191
- 628 Lewis, S., Lentile, L., Hudak, A., Robichaud, P., Morgan, P., Bobbitt, M., 2007. Mapping
- 629 ground cover using hyperspectral remote sensing after the 2003 Simi and Old wildfires in
- 630 Southern California, Fire Ecol. 3, 109-128
- 631 Lhermitte, S., Verbesselt, J., Verstraeten, W.W., Veraverbeke, S., Coppin, P., 2011. Assessing
- 632 intra-annual vegetation regrowth after fire using the pixel based regeneration index. ISPRS J.
- 633 Photogramm. Remote Sens. 66, 17-27
- 634 Malak, D., Pausas, J., 2006. Fire regime and post-fire Normalized Difference Vegetation
- 635 Index changes in the eastern Iberian peninsula. Int. J. Wildland Fire 15, 407–413
- 636 Malanson, G., Trabaud, L., 1988. Vigour of post-fire resprouting by *Quercus coccifera* L. J.
- 637 Ecol. 76, 351–365

- 638 McMichael, C., Hope, A., Roberts, D., Anaya, M., 2004. Post-fire recovery of leaf area index
- 639 in California chaparral: a remote sensing-chronosequence approach. Int. J. Remote Sens. 25,
 640 4743–4760.
- 641 Nepstad, D., Verssimo, A., Alencar, A., Nobre, C., Lima, E., Lefebvre, P., Schlesinger, P.,
- 642 Potter, C., Moutinho, P., Mendoza, E., Cochrane, M., Brooks, V., 1999. Large-scale
- 643 impoverishment of Amazonian forest by logging and fire. Nat. 398, 505–508
- Pausas, J., Verdu, M., 2005. Plant persistence traits in the fire-prone ecosystems of the
 Mediterranean basin: a phylogenetic approach. Oikos 109, 196-202
- 646 Plaza, A., Martínez, P., Pérez, R., Plaza, J., 2004. A quantitative and comparative analysis of
- 647 endmember extraction algorithms from hyperspectral data. IEEE Trans. Geosci. Remote Sens.
- 648 42, 650–663
- PP Systems, 2006. Unispec-SC (single channel) spectral analysis system operation manual
 version 2.02, Hertfordshire
- 651 Randerson, J., Liu, H., Flanner, M., Chamber, S., Jin, Y., Hess, P., Pfister, G., Mack, M.,
- 652 Treseder, K., Welp, L., Chapin, F., Harden, J., Goulden, M., Lyons, E., Neff, J., Schuur, E.,
- EXAMPLE 2006. The impact of boreal forest fire on climate warming. Sci. 314, 1130-1132.
- Ray, T., Murray, B., 1996. Nonlinear spectral mixing in desert vegetation. Remote Sens.
- 655 Environ. 55, 59–64
- 656 Riaño, D., Chuvieco, E., Ustin, S., Zomer, R., Dennison, P., Roberts, D., Salas, J., 2002.
- 657 Assessment of vegetation regeneration after fire through multitemporal analysis of AVIRIS
- 658 images in the Santa Monica mountains. Remote Sens. Environ. 79, 60-71
- 659 Riaño, D., Moreno-Ruiz, J., Isidoros, D., Ustin, S., 2007. Global spatial patterns and temporal
- trends of burned area between 1981 and 2000 using NOAA-NASA Pathfinder. Glob. Chang.
- 661 Biol. 13, 40-50

- 662 Roberts, D., Smith, M., Adams, J., 1993. Green vegetation, nonphotosynthetic vegetation, and
- soils in AVIRIS data. Remote Sens. Environ. 44, 255–269
- 664 Roberts, D., Gardner, M., Church, R., Ustin, S., Scheer, G., Green, R., 1998. Mapping
- 665 chaparral in the Santa Monica mountains using multiple endmember spectral mixture models.
- 666 Remote Sens. Environ. 65, 267-279
- 667 Robichaud, P., Lewis, S., Laes, D., Hudak, A., Kokaly, R., Zamudio, R., 2007. Post-fire soil
- burn severity mapping with hyperpspectral image unmixing. Remote Sens. Environ. 108, 467-480
- 670 Roder, A., Hill, J., Duguy, B., Alloza, J., Vallejo, R., 2008. Using long time series of Landsat
- data to monitor fire events and post-fire dynamics and identify driving factors. A case study in
- the Ayora region (eastern Spain). Remote Sens. Environ. 112, 259-273
- 673 Rogan, J., Franklin, J., 2001. Mapping wildfire burn severity in southern California forests and
- 674 shrublands using Enhanced Thematic Mapper imagery. Geocarto Int., 16, 1–11
- 675 Rogge, D., Rivard, B., Zhang, J., Feng, J., 2006. Iterative spectral unmixing for optimizing
- 676 per-pixel endmember sets. IEEE Trans. Geosci. Remote Sens. 44, 3725–3736
- 677 Sankey, T., Moffet, C., Weber, K., 2008. Postfire recovery of sagebrush communities:
- 678 assessment using SPOT-5 and very large-scale aerial imagery. Rangel. Ecol. Manage. 61,
- 679 598–604
- Shakesby, R., Doerr, S., 2006. Wildfire as hydrological and geomorphological agent. Earth
 Sci. Rev. 74, 269-307
- 682 Somers, B., Delalieux, S., Stuckens, J., Verstraeten, W.W., Coppin, P. 2009a. A weighted
- 683 linear spectral mixture analysis to address endmember variability in agricultural production
- 684 systems., Int. J. Remote Sens. 30, 139–147

- 685 Somers, B., Cools, K., Delalieux, S., Stuckens, J., Van der Zande, D., Verstraeten, W.W. &
- 686 Coppin, P., 2009b. Nonlinear hyperspectral mixture analysis for tree cover estimates in
- orchards. Remote Sens. of Environ. 113, 1183–1193
- 688 Somers, B., Verbesselt, J., Ampe, E., Sims, N., Verstraeten, W.W., Coppin, P. 2010a. Spectral
- 689 mixture analysis to monitor defoliation in mixed-age Eucalyptus globulus Labill plantations in
- 690 southern Australia using Landsat 5-TM and EO-A Hyperion data. Int. J. Appl. Earth
- 691 Observation Geoinf. 12, 270–277
- 692 Somers, B., Delalieux, S., Verstraeten, W.W., van Aardt, J., Albrigo, J., Coppin, P. 2010b. An
- 693 automated waveband selection technique for optimized hyperspectral mixture analysis. Int. J.
- 694 Remote Sens. 31, 5549-5568
- Teillet, P., Guindon, B., Goodenough, D., 1982. On the slope-aspect correction of
 multispectral scanner data. Can. J. Remote Sens. 8, 84–106
- 697 Trabaud, L., 1981. Man and fire: impacts on Mediterranean vegetation, in di Castri, F.,
- Goodall, D., Specht, R. (Eds.), Mediterranean-type shrublands. Elsevier, Amsterdam, pp. 523537
- van Leeuwen, W., 2008. Monitoring the effects of forest restoration treatments on post-fire
- vegetation recovery with MODIS multitemporal data. Sens. 8, 2017-2042
- van Leeuwen, W., Casady, G., Neary, D., Bautista, S., Alloza, J., Carmel, J., Wittenberg, L.,
- 703 Malkinson, D., Orr, B., 2010. Monitoring post-wildfire vegetation response with remotely
- sensed time series data in Spain, USA and Israel. Int. J. Wildland Fire 19, 75-93
- 705 Veraverbeke, S., Lhermitte, S., Verstraeten, W.W., Goossens, R., 2010a. The temporal
- dimension of differenced Normalized Burn Ratio (dNBR) fire/burn severity studies: the case
- 707 of the large 2007 Peloponnese wildfires in Greece. Remote Sens. Environ. 114, 2548-2563

- 708 Veraverbeke, S., Verstraeten, W., Lhermitte, S., Goossens, R., 2010b. Evaluation Landsat
- 709 Thematic Mapper spectral indices for estimating burn severity of the 2007 Peloponnese
- 710 wildfires in Greece. Int. J. Wildland Fire 19, 558-569
- 711 Veraverbeke, S., Verstraeten, W.W., Lhermitte, S., Goossens, R., 2010c. Illumination effects
- on the differenced Normalized Burn Ratio's optimality for assessing fire severity. Int. J. Appl.
- 713 Earth Observation Geoinf. 12, 60-70
- 714 Veraverbeke, S., Lhermitte, S., Verstraeten, W.W., Goossens, R., 2011. A time-integrated
- 715 MODIS burn severity assessment using the multi-temporal differenced Normalized Burn
- 716 Ratio (dNBR_{MT}). Int. J. Appl. Earth Observation Geoinf.13, 52-58
- 717 Viedma, O., Melia, J., Segarra, D., Garcia-Haro, J., 1997. Modeling rates of ecosystem
- recovery after fires by using Landsat TM data. Remote Sens. Environ. 61, 383-398
- 719 Vila, G., Barbosa, P., 2010. Post-fire vegetation regrowth detection in the Deiva Marina
- region (Liguria-Italy) using Landsat TM and ETM+ data. Ecol. Model. 221, 75-84
- 721 Volaire, F., Lelievre, F., 2010. Role of summer dormant perennial grasses as intercrops in
- rainfed Mediterranean vineyards. Crop Sci. 50, 2046-2054
- 723 Zarco-Tejada, P., Rueda, C., Ustin, S., 2003. Water content estimation in vegetation with
- MODIS reflectance data and model inversion methods. Remote Sens. Environ. 85, 109-124
- 725 Zhang, J., Rivard, B., Sanchez-Azofeifa, A., 2004. Derivative spectral unmixing of
- 726 hyperspectral data applied to mixtures of lichen and rock. IEEE Trans. Geosc. Remote Sens.
- 727 42, 1934-1940
- 728 Zhang, J., Rivard, B., Sanchez-Azofeifa, A., Castro-Esau, K., 2006. Intra- and inter-class
- 729 spectral variability of tropical tree species at La Selva, Costa Rica: Implications for species
- 730 identification using HYDICE imagery. Remote Sens. Environ. 105, 129-141
- 731

Figure 1. Location of the study area (the areas encircled with black represent the 2007 burned areas) and
distribution of the field plots (marked with white crosses) (Landsat Thematic Mapper image July 18, 2010 RGB-

734 432).

Figure 2. Mean spectral signatures of green vegetation, brown vegetation, and substrate acquired in the field with a Unispec single channel field spectroradiometer (A). The shadow endmember is modeled as a flat 1 % reflectance (Lelong et al., 1998). Specific spectra for limestone and flysch substrate are indicated by the dashed lines. The Thematic Mapper (TM) visual and near infrared bandpasses are also shown. B shows the presence of flysch and limestone substrates in the 2007 burned areas (based on Institute for Geology and Mineral Exploration, 1983).

741 Figure 3. Line transect plot design (Bonham 1989)

Figure 4. Example plot photographs of shrub land with a high (A), moderate (B) and low (C) recovery rate.

743 Figure 5. Scatter plots and regression lines of modeled versus estimated fractional vegetation cover of the

simulation experiments for the noise-free simple Spectral Mixture Analysis (SMA) (A), the noise-free multiple

rd45 endmember SMA (MESMA) (C) and the noise-free segmented SMA (E) and the equivalent models with noise

746 (Asner and Lobell, 2000) (respectively B, D and F). Separate scatter plots and regression lines are displayed for

747 the flysch subset (n = 500) and limestone subset (n = 500). Regression lines of the pooled dataset (n = 1000) are 748 also indicated.

749 Figure 6. Scatter plots and regression lines of line transect ratings versus fractional vegetation cover derived from

750 Landsat imagery for the simple Spectral Mixture Analysis (SMA) (A), the multiple endmember SMA (MESMA)

(B) and segmented SMA (C). Separate scatter plots and regression lines are displayed for the flysch subset (n =

46) and limestone subset (n = 32). Regression lines of the pooled dataset (n = 78) are also indicated.

Figure 7. Fractional vegetation cover map three years after the fires based on the segmented SMA model.

754

755 Table 1. Error matrix of the substrate spectrum selection by the multiple endmember Spectral Mixture Analysis

756 (MESMA) model for the simulation experiment. The reference data were retrieved from the experimental set-up.

757 Table 2. Error matrix of the substrate spectrum selection by the multiple endmember Spectral Mixture Analysis

758 (MESMA) model for the Landsat application. The reference data are the line transect field plots.