

RNA Interference Links Oxidative Stress to the Inhibition of Heat Stress Adaptation

Zoltán Spiró,^{1,*} Mehmet Alper Arslan,¹ Milán Somogyvári,¹ Minh Tu Nguyen,¹ Arne Smolders,² Balázs Dancsó,^{1,#} Nóra Németh,¹ Zsuzsanna Elek,¹ Bart P. Braeckman,² Péter Csermely,¹ and Csaba Solti¹

Abstract

Increased oxidative stress is associated with various diseases and aging, while adaptation to heat stress is an important determinant of survival and contributes to longevity. However, the impact of oxidative stress on heat resistance remains largely unclear. **Aim:** In this study we investigated how oxidative stress impinges on heat stress responses. **Results:** We report that hydrogen-peroxide (H₂O₂) pretreatment inhibits both acquired thermotolerance and heat-induced Hsp70 expression in mammalian cells, as well as acquired thermotolerance in the nematode *Caenorhabditis elegans*, via RNA interference. Moreover, we demonstrate that elimination of RNA interference by silencing key enzymes in microRNA biogenesis, *dcr-1* or *pash-1*, restores the diminished intrinsic thermotolerance of aged and H₂O₂-elimination compromised (catalase-2 and peroxiredoxin-2 deficient) worms. **Innovation and Conclusion:** These results uncover a novel post-transcriptional element in the regulation of heat stress adaptation under oxidative conditions that may have implications in disease susceptibility and aging. *Antioxid. Redox Signal.* 00, 000–000.

Introduction

BASIC PHYSIOLOGICAL PROCESSES such as metabolism, cellular signaling, and immunity are associated with the production of reactive oxygen species (ROS) (16). An accumulation of ROS, called oxidative stress, plays a critical role in various diseases and in aging (13, 16, 34). Although an excess of ROS generates diverse molecular and cellular damages and evokes a plethora of signaling events, how it is involved in the induction or aggravation of these pathological states is not entirely understood.

Increased resistance to heat stress protects against degenerative diseases in mammals (9, 32) and associates with longevity in *Caenorhabditis elegans* (10, 26). Intrinsic thermotolerance is maintained by multiple mechanisms. A preconditioning (*i.e.*, heat) stress induces acquired thermotolerance, mediated by the heat shock response via heat shock factor (HSF1)-dependent induction of heat shock proteins (Hsp-s) (30, 47). Previous studies reported contrasting results of oxidative stress on HSF1 activation (2, 28) and Hsp70 levels (14, 22, 43). However, the effect of oxidative stress on thermotolerance remains largely unexplored.

RNA interference is a powerful post-transcriptional regulator of gene expression that operates via ~22 nt microRNAs

(miRNAs) (27). Genomic miRNA precursors are processed by highly specific RNases: the nuclear Droscha/PASH-1 produces hairpin pre-miRNAs, which are transported to the cytoplasm and cleaved to mature miRNAs by Dicer/DCR-1 (capital names indicate the respective nematode orthologs). Hence, Dicer/Droscha knockout is a reliable tool to investigate the general role of miRNAs (5, 41, 44). miRNAs bind to the mRNA 3' untranslated region (3'UTR), repress translation, or promote mRNA degradation (27). miRNAs modulate diverse

Innovation

Oxidative stress is a serious cause of cell and tissue damage associated with many human diseases. Our observations beyond demonstrating a novel crosstalk between various types of stresses via RNA interference extend our understanding on how oxidative stress may debilitate physiological function. As RNA interference exhibits a significant functional conservation from nematodes to humans, we anticipate that the mechanism identified herein may be involved in human diseases and aging.

¹Department of Medical Chemistry, Semmelweis University, Budapest, Hungary.

²Laboratory for Aging Physiology and Molecular Evolution, Department of Biology, Ghent University, Ghent, Belgium.

*Current affiliation: ISREC, Ecole Polytechnique Federal de Lausanne, Lausanne, Switzerland.

#Current affiliation: Emergency Department, Doncaster and Bassetlaw Hospitals, NHS Foundation Trust, Worksop, United Kingdom.

biological processes. Their connection with stress is exemplified by imparting robustness to gene expression networks in response to environmental change (24) and by the profound alterations of miRNA expression upon heat and oxidative stresses (25, 42, 49) [reviewed in (23)]. Heat and ischemic preconditioning-induced miRNAs induce Hsp70 and are cardioprotective during ischemia-reperfusion in mice (48, 49). Moreover, miRNAs modulate the life span and stress resistance of *C. elegans* involving DAF-16 and HSF1 (6, 11), underscoring a vital role of RNA interference in stress responses.

In this study we focused on the impact of oxidative stress on heat stress adaptation and found that hydrogen-peroxide (H_2O_2) pretreatment inhibited acquired thermotolerance in both COS-7 mammalian cells and in *C. elegans*. As an underlying mechanism, H_2O_2 inhibited the heat-induction of Hsp70 in cells, consistent with a recent study (1). Moreover, H_2O_2 prevented the heat-induction of an Hsp70 3'UTR reporter. H_2O_2 -induced effects required Dicer, a key enzyme in miRNA biogenesis, in both cells and worms. We further found that RNAi against Dicer and Drosha orthologs restored the compromised thermotolerance of two worm strains deficient in H_2O_2 disposal. Finally, Dicer silencing delayed the decline of thermotolerance in aging worms and phenocopied the effect

of the antioxidant *N*-acetyl-L-cysteine (NAC). Our results reveal RNA interference as a mediator of oxidative stress-induced inhibition of heat stress responses.

Results

H₂O₂ inhibits acquired thermotolerance and Hsp70 induction at the post-transcriptional level in COS-7 cells

The effect of a transient H_2O_2 exposure on thermotolerance of COS-7 cells was determined by subjecting cells to a lethal heat stress 24h after H_2O_2 and/or preconditioning heat treatments. Heat preconditioning elicited a large increase in survival (acquired thermotolerance, Fig. 1A). A prior H_2O_2 treatment slightly increased intrinsic thermotolerance. Importantly, it potently inhibited acquired thermotolerance in a concentration-dependent manner (Fig. 1A).

To examine whether the decrease in acquired thermotolerance is due to the inhibition of the heat shock response, we pretreated COS-7 cells with a series of H_2O_2 concentrations and monitored the heat induction of Hsp70 by flow cytometry (Fig. 1B). Cells, exposed to heat shock, exhibited an ~10-fold induction of Hsp70, concordant with the induction of thermotolerance (cf. Fig. 1A). H_2O_2 treatment did not affect basal




FIG. 1. Hydrogen-peroxide (H_2O_2) impairs heat-preconditioned thermotolerance and Hsp70 heat-induction in COS-7 cells. (A) Effect of H_2O_2 on thermotolerance. Cells were treated by the indicated concentrations of H_2O_2 for 2 h, then kept at 37°C (control) or at 43°C for 30 min (heat shock). About 24 h later cells were subjected to a lethal heat stress (45°C, 60 min). Cell survival was analyzed 24 h later by Trypan blue exclusion. Values are means \pm standard deviations (SDs) of three experiments. (B, C) Effect of H_2O_2 on Hsp70 and Hsp90 protein levels. Cells were treated by H_2O_2 and heat shock as in panel A. Five hours later Hsp70 levels were analyzed by flow cytometry using a monoclonal antibody (B) or by Western blot using a polyclonal anti-Hsp70 and monoclonal anti-Hsp90 and anti-actin antibodies, respectively (C). Values are means \pm SDs of five experiments compared to their respective controls, and image is a representative of three experiments. (D) Densitometric analysis of relative Hsp70 and Hsp90 levels from (C). (E) H_2O_2 titration curve of cytosolic HyPer-C in COS-7 cells. The 490/420-nm fluorescence excitation ratio of HyPer was calculated after background fluorescence subtraction from two experiments. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Hsp70 level, but inhibited Hsp70 heat induction in a concentration-dependent manner (Fig. 1B). Western blots using a polyclonal anti-Hsp70 antibody showed a similar inhibition of Hsp70, but not of the specific chaperone Hsp90 (Fig. 1C, D). These results exclude an H₂O₂-induced modification or degradation of Hsp70 as well as a general, stress-induced transcriptional or translational block. The efficacy of H₂O₂ was verified by cells expressing the H₂O₂-sensor Hyper-C (Fig. 1E) (12). Thus, H₂O₂ pretreatment compromises both acquired thermotolerance and Hsp70 heat-induction in COS-7 cells.

Next, we investigated the site of action of H₂O₂ along the heat shock regulon. Upon heat, misfolded proteins activate HSF1, which binds to heat shock promoter elements and induces *hsp* gene transcription (32). To assess the level of HSF1-dependent transactivation, we transfected COS-7 cells with a *hsp70pr*/luciferase vector and performed reporter gene assays after cells had either been oxidatively stressed and/or heat-shocked. Heat shock markedly induced reporter activity, while H₂O₂ treatment significantly affected neither basal nor heat-induced transactivation (Fig. 2A). Likewise, H₂O₂ treatment did not decrease *hsp70* mRNA level (Fig. 2B). Thus, a

transcriptional inhibition does not seem to underlie the H₂O₂-induced decrease in Hsp70 protein expression.

To assess, if H₂O₂ could down-regulate Hsp70 post-translationally, we changed the order of stresses (*i.e.*, employed H₂O₂ after heat shock) and followed the Hsp70 protein level by flow cytometry (Fig. 2C). H₂O₂ did not change the heat shock-induced sustained elevation of Hsp70, which excluded the possibility of an accelerated Hsp70 turnover.

The 3'UTR is intimately connected with the post-transcriptional regulation of mRNAs. To investigate the molecular events at the Hsp70 3'UTR, we took use of a reporter harboring the mouse *hsp70.1* 3'UTR fused to Firefly luciferase (18). Monitoring luciferase activity provided an estimate of the impact of the *hsp70* 3'UTR on the translation of luciferase mRNA following H₂O₂ and/or heat shock treatments. 3'UTR reporter activity displayed a time-dependent increase after heat-shock peaking at 6 h (Fig. 2C). This finding is consistent with early reports in *Drosophila* and mammalian cells on the role of the 3'UTR in the regulation of Hsp70 protein synthesis during heat shock (33, 38). Neither H₂O₂ nor the combination of H₂O₂ and heat shock increased luciferase activity above the




FIG. 2. H₂O₂ inhibition of Hsp70 involves a potential post-transcriptional regulation. (A) H₂O₂ does not affect *hsp70* promoter activation. Cells transfected with the *hsp70.1pr*/luc and control plasmids were treated as in Figure 1. Enzyme activities were measured 18 h later, and their ratios were expressed. (B) H₂O₂ does not diminish *hsp70* (HSPA1A) mRNA expression. Cells were treated as in Figure 1. mRNA levels were determined 1-h after treatments by quantitative reverse transcriptase–polymerase chain reaction and expressed relative to β -actin. (C) H₂O₂ does not affect Hsp70 protein turnover. Cells were heat shocked as above, after 2 h at 37°C cells were incubated in the absence (control), or presence of 800 μ M of H₂O₂ for 2 h, and then harvested at the indicated timepoints, and analyzed by flow cytometry. (D) H₂O₂ inhibits the heat-induced luciferase reporter translation mediated by the Hsp70 3' untranslated region (3'UTR). Cells transfected with a pGL3/luc/*hsp70.1* 3'-UTR and control plasmids were treated by 650 μ M H₂O₂ for 2 h, and then kept at 37°C or heat shocked. At the indicated timepoints enzyme activities were determined, and expressed as a ratio. Values are means \pm SDs of three experiments. n.s., non-significant, * p < 0.05, ** p < 0.01, *** p < 0.001.




FIG. 3. RNA interference mediates H₂O₂-induced inhibition of Hsp70 induction and acquired thermotolerance in COS-7 cells. (A) Effect of H₂O₂ treatment and anti-Dicer siRNA on Dicer protein level. Two days after transfection by anti-Dicer or control siRNA, respectively, cells were treated with 800 μ M H₂O₂ for 2 h. Protein levels were analyzed by Western blot. Image is a representative of three experiments. (B) Effect of Dicer siRNA and H₂O₂ on the Hsp70 3'UTR activation. Cells undergoing a 2-day co-transfection with a control/Dicer siRNA and the 3'UTR reporter plasmids were treated by 650 μ M H₂O₂ for 2 h, and then heat shocked. About 6 h later enzyme activities were determined and expressed as a ratio. (C) Effect of Dicer siRNA and H₂O₂ on Hsp70 protein expression. Cells transfected with a control/Dicer siRNA were treated by 800 μ M H₂O₂ for 2 h, and then kept at 37°C or heat shocked. Five hours later Hsp70 levels were analyzed by flow cytometry. (D) Effect of Dicer siRNA and H₂O₂ on heat preconditioned thermotolerance. Cells transfected with a control/Dicer siRNA were treated by 800 μ M H₂O₂ for 2 h, and then kept at 37°C or heat shocked. Lethal heat stress and survival assay was performed as in Figure 1A. Values are means \pm SDs of three experiments. n.s., non-significant, * p < 0.05, ** p < 0.01, *** p < 0.001.

baseline demonstrating that H₂O₂ entirely prevented the heat-induced activation by the *hsp70* 3'UTR (Fig. 2C).

RNA interference mediates H₂O₂-induced inhibition of Hsp70 induction and acquired thermotolerance in COS-7 cells

RNA interference is a powerful modulator of stress responses (23). To address whether RNA interference may mediate the events involving the Hsp70 3'UTR, we blocked miRNA maturation by anti-Dicer siRNA transfection. Only the siRNA, but not H₂O₂ led to a knock-down of Dicer (Fig. 3A). Intriguingly, anti-Dicer siRNA led to a large decrease in heat-induced 3'UTR reporter activity, suggesting that Dicer was necessary for the 3'UTR-mediated translational activation of the luciferase mRNA upon heat shock. This inhibition was comparable to that induced by H₂O₂, and a combination of anti-Dicer siRNA and H₂O₂ was not additive (Fig. 3B). Thus, H₂O₂ prevents the heat-induced Hsp70 3'UTR activation primarily *via* RNA interference.

To investigate how the inhibition of the 3'UTR by RNA interference is reflected in Hsp70 translation, we determined Hsp70 protein levels in anti-Dicer siRNA-transfected cells

undergoing H₂O₂ and heat shock treatments. Dicer silencing inhibited Hsp70 heat-induction to approximately two-thirds of the control siRNA transfected value, comparable to the effect of Dicer silencing (Fig. 3C). Remarkably, H₂O₂ could not further reduce Hsp70 expression in Dicer-silenced cells, suggesting that the effect of H₂O₂ required an intact RNA interference.

These results suggested that RNA interference might play a role in the H₂O₂-induced inhibition of acquired thermotolerance. Indeed, we found that Dicer siRNA reduced acquired thermotolerance in heat-preconditioned cells, which was similar to the effect of H₂O₂ (Fig. 3D). Moreover, H₂O₂ did not further diminish thermotolerance in Dicer-silenced cells, in agreement with our observations on Hsp70 induction (cf. Fig. 3C). Hence, we conclude that RNA interference mediates the H₂O₂-induced inhibition of heat stress adaptation in COS-7 cells.

H₂O₂ inhibits acquired thermotolerance through DCR-1 in *C. elegans*

To address if the effect of H₂O₂ on heat stress adaptation was conserved during evolution, we used *C. elegans*, a




FIG. 4. A prior H₂O₂ treatment inhibits acquired thermotolerance in an HSF1-dependent manner in *Caenorhabditis elegans*. (A) Effect of preconditioning H₂O₂ treatments on oxidative tolerance. Oxidative stress was applied in the liquid nematode growth medium (NGM) for 1 h at 20°C, 12–14 h before a lethal oxidative challenge. Data are means ± SDs of two separate experiments. (B) Intensity-normalized ratio image demonstrating a rapid rise of oxidized/reduced HyPer ratio in *jrIs[Prpl-17::HyPer]* worms in response to a 1-min challenge by 100 μM H₂O₂ in liquid NGM. Representative image from five independent experiments. (C) Effect of H₂O₂ (100 μM for 1 h) on intrinsic and acquired thermotolerance induced by a preconditioning heat shock (heat shock, 30° for 2 h). Lethal heat stress was employed 12 h later. Only heat shock induces a significant difference in survival ($p < 0.0001$). (D) No change in thermotolerance by a preconditioning heat shock and/or an H₂O₂ treatment in *hsf-1(sy441)* mutant worms ($p > 0.1$). (E) H₂O₂ employed after the preconditioning heat shock does not abrogate acquired thermotolerance ($p > 0.1$ compared to heat shock). (To see this illustration in color the reader is referred to the web version of this article at www.liebertonline.com/ars).

powerful model system exhibiting an organismal complexity. In search of an H₂O₂-exposure that did not cause significant damage in nematodes, we found that a treatment by 100 μM for 1 h was below the threshold to induce oxidative tolerance to a lethal H₂O₂ challenge (Fig. 4A). This concentration induced a rapid signal elevation in the pharynx and intestine of worms ubiquitously expressing HyPer (Fig. 4B) (3). We used the 100-μM pretreatment to investigate its effect on nematodal

thermotolerance. A preconditioning heat shock at 30°C for 2 h resulted in a 20%–40% increase in thermotolerance (Fig. 4C). A prior H₂O₂ treatment did not affect intrinsic thermotolerance of worms; however, it entirely abolished acquisition of thermotolerance by the preconditioning heat shock.

To address whether H₂O₂ would affect an HSF1-dependent process, we employed the *hsf-1(sy441)* point mutant strain

harboring a truncated transactivation domain that prevented the heat-induction of HSF1-target genes (15). In line with recently published data of McColl and colleagues (29), HSF1 was required for acquired, but not for intrinsic thermotolerance (Fig. 4D). H₂O₂ treatment was not additive to the *hsf-1(sy441)* background; it affected neither basal nor heat-preconditioned survival. Moreover, in wild-type worms, H₂O₂, if applied after the preconditioning heat shock, was unable to inhibit acquired thermotolerance (Fig. 4E), suggesting that H₂O₂ needs to precede heat preconditioning. Thus, H₂O₂ specifically inhibits the acquisition of HSF1-dependent thermotolerance in *C. elegans*.

If, similarly to mammalian cells, H₂O₂ inhibited the heat shock response *via* RNA interference in *C. elegans*, then worms deficient in miRNA synthesis would escape from the H₂O₂-dependent inhibition of thermotolerance. Investigating this hypothesis we found that silencing the Dicer ortholog by *dcr-1(RNAi)* restored the acquired thermotolerance of H₂O₂-treated worms to levels comparable to heat shock alone (Fig. 5A, B). *dcr-1(RNAi)* *per se* did not affect thermotolerance (Fig. 5B). We obtained similar results using loss-of-function *dcr-1* mutant nematodes (Fig. 5C). The efficiency of *dcr-1* silencing and the lack of a

general disruption of RNA interference by H₂O₂, respectively, were demonstrated by an RNA interference reporter strain (20) (Fig. 5D). We made attempts to investigate an analogous involvement of Hsp70 regulation. Unfortunately, a number of antibodies were unable to detect nematode Hsp70. Quantitative polymerase chain reaction (PCR) measurements revealed a tendency of H₂O₂ preconditioning to augment heat-induced *hsp-70* mRNA expression. However, *dcr-1(RNAi)* significantly altered neither heat-induced mRNA level nor the H₂O₂-induced elevation (Supplementary Fig. S1; Supplementary Data are available online at www.liebertonline.com/ars). Despite the unclear involvement of Hsp70, RNA interference is required for H₂O₂ to inhibit acquired thermotolerance in worms.

Inhibition of RNA interference restores thermotolerance in endogenous models of oxidative stress

Next we asked how thermotolerance might be affected by chronic genetic disturbances in antioxidant defense. Antioxidant enzymes provide protection against oxidative stress




FIG. 5. DCR-1 mediates the H₂O₂-induced inhibition of thermotolerance in *C. elegans*. Effect of H₂O₂ on intrinsic and acquired thermotolerance in worms fed by empty (EV, **A**) or *dcr-1(RNAi)* (**B**) vectors, respectively. Treatments were as in Figure 4. Note that the activatory effect of preconditioning heat shock was less pronounced on RNAi plates. In EV-fed worms (**A**) only heat shock, while in *dcr-1(RNAi)*-fed worms (**B**) both heat shock as well as H₂O₂ + heat shock induced a significantly higher survival compared with controls ($p < 0.001$). (**C**) Both heat shock and H₂O₂ + heat shock induces a significant increase in thermotolerance in *dcr-1(ok247);unc-32(e189)* nematodes ($p < 0.0001$ vs. control). Survival curves are representatives of three experiments yielding similar results. (**D**) H₂O₂ does not compromise RNA interference. Epifluorescence image demonstrating increased expression of *pajm::GFP* (harboring an anti-GFP hairpin siRNA in addition to the GFP sequence) in the GR1401 RNA interference reporter strain fed by *dcr-1(RNAi)*. Arrows point to specific dots localized to epithelial seam cells. In contrast, H₂O₂ (100 μ M for 1 h) treatment followed by a 12- or 24-h recovery did not inhibit GFP silencing. Please note the autofluorescence of oxidatively stressed worms. Representative image from three independent experiments. (To see this illustration in color the reader is referred to the web version of this article at www.liebertonline.com/ars).




FIG. 6. Loss of RNA interference rescues thermotolerance in nematodes with genetic defects of H₂O₂ disposal. (A) Effect of *dcr-1(RNAi)* on thermotolerance of N2 and *ctl-2(ok1137)* worms. *ctl-2(ok1137)* worms exhibited significantly shorter survival ($p < 0.001$), while other survivals were not significantly different ($p > 0.2$), compared to N2 control. (B) *pash-1(RNAi)* phenocopies *dcr-1(RNAi)* by inducing a significant increase in thermotolerance of *ctl-2(ok1137)* ($p < 0.0001$) compared to that of the EV control. (C) *pash-1(RNAi)* does not change thermotolerance of wild-type worms ($p > 0.1$) compared to that of the EV control. (D) *pash-1(RNAi)* extends thermotolerance independently of *hsf-1* in *ctl-2(ok1137)* worms [$p < 0.01$ vs. *pash-1/hsf-1(RNAi)*]. (E) Effect of *dcr-1(RNAi)* on thermotolerance of N2 and *prdx-2(gk169)* worms. *prdx-2(gk169)* worms fed by EV exhibited significantly shorter ($p < 0.0001$), while those fed by *dcr-1(RNAi)* exhibited slightly longer survival ($p < 0.05$) compared to N2 control. Survival curves are representatives of three independent experiments giving similar results.

by removing ROS. Catalase-2 is a peroxisomal enzyme involved in H₂O₂ elimination accounting for ~80% of total catalase activity in the worm (39). *ctl-2* loss of function elevates endogenous H₂O₂ levels (3), decreases oxidative tolerance and shortens lifespan (39). We observed that *ctl-2(ok1137)* animals exhibited impaired intrinsic thermotolerance compared to wild type, which was completely restored by *dcr-1(RNAi)* (Fig. 6A). Silencing the Drosha ortholog PASH-1, the other key enzyme in miRNA biogenesis, phenocopied the effect of *dcr-1(RNAi)* in the *ctl-2(ok1137)* strain (Fig. 6B) without affecting wild-type thermotolerance (Fig. 6C). Neither the survival decrease in *ctl-2(ok1137)* nor the amelioration by *pash-1(RNAi)* was prevented by *hsf-1(RNAi)* (Fig. 6D). Hence, *ctl-2*

loss of function modulates intrinsic thermotolerance, not involving the HSF1-Hsp axis.

To test whether the observed phenomena might be attributed to the general impairment of H₂O₂ elimination, we examined the lack of peroxiredoxin-2, involved in H₂O₂ reduction in the cytosol. *prdx-2(gk169)* worms, similarly to the *ctl-2(ok1137)* strain, are susceptible to H₂O₂ injury, and display a shortened lifespan (37). We found that *prdx-2* knockout also markedly decreased *C. elegans* thermotolerance (Fig. 6E). Importantly, *dcr-1(RNAi)* prevented thermotolerance inhibition in *prdx-2(gk169)* worms. Together these data suggest that genetic defects in H₂O₂ elimination compromise heat stress adaptation via RNA interference.




FIG. 7. Loss of RNA interference and the antioxidant *N*-acetyl-L-cysteine ameliorate age-associated decline of thermotolerance in *C. elegans*. Thermotolerance of 2-day (A) and 10-day (B) old nematodes treated by *dcr-1(RNAi)* and/or 5 mM *N*-acetyl-L-cysteine (NAC; from day 1 of adulthood). There was no significant difference in survival between treatments at day 2 ($p > 0.1$). The 10-day old control (EV) exhibited a significantly shorter survival ($p < 0.001$ vs. 2-day EV). All *dcr-1(RNAi)* and/or 5 mM NAC induced a significant increase in survival ($p < 0.001$ vs. 10-day old EV), which approached ($p = 0.028$ 10-day EV + NAC vs. 2-day EV), and became non-significant ($p > 0.05$ 10-day *dcr-1(RNAi)* and *dcr-1(RNAi)* + NAC vs. 2-day EV) compared with the survival of the 2-day old EV control. Survivals of 10-day *dcr-1(RNAi)* and *dcr-1(RNAi)* + NAC strains were not significantly different ($p > 0.4$). (C) Mean thermotolerance of nematodes treated by *dcr-1(RNAi)* and/or 5 mM NAC as a function of age. Panels are representatives of two independent experiments yielding similar results. ** $p < 0.01$, *** $p < 0.001$.

Inhibition of RNA interference delays age-dependent decline of thermotolerance

Aging is characterized by a collapse of proteostasis and an impairment of the heat shock response in *C. elegans* (4). Consistent with this, we observed a decline in *C. elegans* thermotolerance during aging (Fig. 7A–C). Oxidative stress and H_2O_2 increases during aging and ROS are considered a major cause of aging (3, 34). To address if oxidative stress affected thermotolerance during aging, we treated worms with the small molecular antioxidant, NAC. Intriguingly, NAC was able to reduce the decline of thermotolerance during aging resulting in a milder slope and a significant difference at the old worms at day 10 of age (Fig. 7A–C).

Then we asked, whether RNA interference was involved in the age-associated decline of thermotolerance of worms. *dcr-1(RNAi)* did not significantly influence the thermotolerance of young animals at day 1 (Fig. 5), but efficiently suppressed the age-induced decline of thermotolerance similarly to NAC treatment (Fig. 7A–C). Moreover, the combination of *dcr-1(RNAi)* with NAC was not significantly different from the effect of *dcr-1(RNAi)* at any time points tested. These findings indicate that RNA interference is involved in the oxidative stress-induced age-dependent decline of heat stress adaptation in *C. elegans*.

Discussion

In this study, we have presented evidence that oxidative stress inhibits the adaptive responses to heat stress in both mammalian cells and *C. elegans*. Silencing Dicer and Drosha orthologs, key enzymes specific to miRNA maturation reveals a conserved role for RNA interference. In mammalian cells H_2O_2 abolishes a positive action of RNA interference on acquired thermotolerance. Inhibition of RNA interference does not alter thermotolerance in young nematodes, suggesting that H_2O_2 may induce miRNA(s) that inhibit the acquisition of thermotolerance. Intrinsic thermotolerance decrease of *prdx-2* and *ctl-2* knockouts and aged worms might require accumulation of miRNA(s) inhibiting HSF1-independent processes. Despite species-specific and context-dependent mechanisms, our results provide support to the modulation of stress responses by RNA interference (Fig. 8) (23).

Our findings on the post-transcriptional inhibition of Hsp70 expression offer a potential molecular mechanism underlying the H_2O_2 -induced compromise of acquired thermotolerance. Early reports demonstrating a heat-induced stabilization of *hsp70* mRNA by its 3'UTR (33, 38) and the decrease in heat-induced *hsp70* mRNA by H_2O_2 in glioma cells (1) suggested that H_2O_2 may prevent mRNA stabilization. However, our results showing no impact of H_2O_2 on




FIG. 8. Model for the role of RNA interference in the modulation of heat stress responses by oxidative stress. Oxidative stress may be induced by various sources, such as increased production/extrinsic factor (H_2O_2), decreased elimination (impaired defense, endogenous mutants), or a combination of the two (aging). RNA interference mediates an inhibitory action of oxidative stress and reduces heat resistance.

hsp70 mRNA and inhibition of 3'UTR reporter, respectively, are consistent with a compromised translation by H_2O_2 . Interestingly, inflammatory cytokines inhibit colonic Hsp70 translation by recruiting its mRNA to stress granules (17, 18). Although it may be one plausible mechanism, our results using Dicer knockdown suggest the involvement of miRNA(s). Possible scenarios include an H_2O_2 -induced decrease of activatory miRNA(s), or displacement/domination of heat-induced activatory miRNA(s) by H_2O_2 -induced inhibitory/neutral miRNA(s) from the *hsp70* mRNA. Both mechanisms are generally employed by RNA interference (23, 27). Moreover, recent articles provide evidence on miRNAs either inhibiting (miR-378*, miR-711, miR-146a, miR-146b-5b) (35, 46), or ischemic preconditioning-induced miRNAs (miR-1, miR-21, miR-24 or others) (48, 49) activating Hsp70 expression. Identification of the exact mechanism(s) and miRNA(s), as well as an analogous Hsp70 regulation in nematodes requires further studies. Nevertheless, our study raises the idea that pathophysiological oxidative conditions (inflammation, wound healing, aging) might employ RNA interference to post-transcriptionally regulate Hsp70 in various tissues (7, 18, 36).

Our use of mutants deficient in H_2O_2 elimination demonstrates that a chronic disturbance in ROS metabolism impairs intrinsic thermotolerance, independently of HSF1 (Fig. 6). This defect can entirely be reversed by blocking miRNA maturation, suggesting a profound post-transcriptional remodeling of heat stress adaptation by RNA interference in response to oxidative stress. McColl *et al.* elegantly showed that increased intrinsic thermotolerance in *daf-2* mutant worms is mediated by a *daf-16*-dependent translational response (29). A common motif in the two studies is that RNA interference or translation do not limit survival in young

worms; however, they differentially condition heat resistance in both long-lived insulin-like signaling mutants and short-lived oxidative defense-deficient mutants and aged worms, respectively [(29) and our study]. It is tempting to speculate that the *daf-16*-regulated response of insulin signaling mutants might involve miRNAs. The clarification of a possible interaction of the translational response and RNA interference in the regulation of stress resistance remains the task of future studies.

Aging in the worm is characterized by an increased accumulation of ROS as well as a collapse of protein homeostasis (4). Our results on the age-induced decline of intrinsic thermotolerance support these observations, and use of the antioxidant NAC demonstrates a progressive causal role for ROS in decline of stress resistance during aging (Fig. 7). Importantly, both the comparable pattern of NAC and *dcr-1(RNAi)* protection and the lack of significant synergism imply a substantially overlapping mode of action. Moreover, increased protection by *dcr-1(RNAi)* suggests that RNA interference adversely affects heat resistance with aging. Single miRNAs do not seem to play an essential role in *C. elegans* development and growth, but both RNA interference and single miRNAs are indispensable to ensure proper development during environmental stress (20, 24, 31). Likewise, there is an extensive change in miRNA expression during *C. elegans* aging (11, 19) and several individual miRNAs similarly modulate longevity and stress resistance in *C. elegans* (11). Inhibition of the entire RNA interference in adulthood provides strong evidence to the general dysregulation of miRNAs in aging and in oxidative stress with a negative impact on stress resistance (Figs. 6 and 7). It remains to be seen whether RNA interference would pose a trade-off between fine-tuning developmental programs and growth during stress in exchange for a self-maintenance later in life. Our results imply that beyond well-characterized stress-responsive HSF1 and DAF-16 pathways, RNA interference may offer a novel target to alleviate decline of stress responses during aging.

Materials and Methods

Materials

Reagents for cell culture were from Invitrogen. Solutions for flow cytometry were from BD Biosciences. Electrophoresis and blotting reagents were from Bio-Rad. *N*-acetyl-L-cysteine and H_2O_2 were from Sigma. All other reagents were from either from Sigma or Fluka.

Cell culture and survival

COS-7 cells were obtained from the ATCC. Cells were cultured as described (40). Cell survival was analyzed by Trypan Blue exclusion 24 h after challenge.

Determination of protein levels

Flow cytometry using a fluorescein-isothiocyanate-conjugated monoclonal anti-Hsp70 antibody (StressGen), cell lysis, and Western blotting using a polyclonal anti-Hsp70 antibody (21), or antibodies against Hsp90 (Stressgen), Dicer (CST), actin (Sigma) was carried out as previously described (40).

Transfection and reporter gene assays

Cells were transfected at a density of 40% using Lipofectamine (Invitrogen). Control/anti-Dicer siRNA (Quiagen)

was introduced at 100 nM. Further treatments were applied at 48 h post-transfection. For the hsp70-promoter reporter gene assay, cells were transfected with 0.35 μ g *hsp70.1pr*/Firefly luciferase plasmid (Rick Morimoto, Northwestern University) and cytomegalovirus/ β -galactosidase plasmids, while for the 3'UTR reporter assay 0.35 μ g pGL3 basic or pGL3/*hsp70.1 3'-UTR* plasmid (Eugene Chang, University of Chicago) and thymidine kinase/Renilla luciferase plasmids were employed, respectively. Treatments were performed 24 h post-transfection. About 18 h post-treatment reporter activities were measured using commercial assay kits (Promega) and expressed.

C. elegans strains and RNA interference

Strains were obtained from the CGC, if not otherwise specified. The following strains were used in this study: wild type (N2), jrIs[Prpl-17::HyPer], PS3551 *hsf-1(sy441)*, BB1 *dcr-1(ok247);unc-32(e189)* III, VC289 *prdx-2(gk169)* II, VC574 *ctl-2(ok1137)* II, and the GFP RNAi-reporter GR1401 (Gary Ruvkun, Harvard University). Strains were backcrossed to the wild type at least three times to clear potential background mutations, and were maintained as described (8). RNAi was performed as described by feeding worms with HT115(DE3) bacteria transformed with empty vector, *dcr-1(RNAi)* Gary Ruvkun (Harvard University) or *pash-1(RNAi)* (Source BioScience) vectors, respectively (45). Experiments were carried out in the second generation. Experiments were performed in the second generation with synchronized young 1-day-old adults, except for age-related thermotolerance.

Thermotolerance assay

Thermotolerance was performed on nematode growth medium plates at 35°C till complete extinction of the population using 25 animals per condition in at least two independent trials. Viability was determined hourly by assaying for movement in response to gentle prodding.

mRNA expression analysis

mRNA was prepared using the GeneJET RNA Purification Kit (Fermentas). mRNA was reverse transcribed using the RevertAid™ cDNA Synthesis Kit (Fermentas). Quantitative PCR was performed in an ABI 7300 System by Taqman Gene Expression Assays: HSPA1A: Hs_00359147_s1; β -actin: Hs_99999903_m1 (Applied Biosystems). Relative amounts of *hsp70* mRNA were determined using the Comparative Cycle Threshold Method for quantitation and normalized to actin mRNA levels. Please see Supplementary Materials and Methods for the analysis of *hsp70* mRNA expression in nematodes.

Analysis of H₂O₂ levels and fluorescence microscopy

Fluorescence measurements in COS-7 cells transfected by HyPer-C (Miklós Geiszt, Semmelweis University) were performed as described (12). HyPer titration was achieved by sequential addition of increasing concentrations of H₂O₂. Mean fluorescence intensities over individual cells were calculated from 3-min recordings. H₂O₂ in worms was monitored using the jrIs[Prpl-17::HyPer] strain, ubiquitously

expressing the H₂O₂-biosensor HyPer, was used. Worms were immobilized and imaged as described (3).

Statistical analysis

Data were analyzed using SPSS software 15.0 (SPSS, Inc.). Survival curves were compared by the log-rank test. If not stated otherwise, all experiments were repeated at least three times. Variables were expressed as mean \pm standard deviation. Statistical significance was indicated as follows: * p < 0.05, ** p < 0.01, *** p < 0.001.

Acknowledgments

We thank Eugene Chang, Miklós Geiszt, Gary Ruvkun, and Rick Morimoto for reagents; Tibor Vellai for help in setting up our worm lab; Melinda Zana and Balázs Enyedi (cellular HyPer measurements), Zsolt Rónai (qPCR), Bea Gilányi, and Ákos Putics for technical help; Eszter Daubner and members of the Sóti Group for discussions; the anonymous reviewers for their helpful advice; and the *Caenorhabditis* Genetics Center for nematode strains. This work was supported by grants from the EU (FP6-518230, FP7-200970, TÁMOP-4.2.2/B-10/1-2010-0013), a grant of the Hungarian Science Foundation/Norway Grants (NNF-78794), the Hungarian Science Foundation (OTKA-K69105 and OTKA-K83314). During the completion of this study, C.S. was a Bolyai Research Scholar of the Hungarian Academy of Sciences.

Author Disclosure Statement

No competing financial interests exist.

References

- Adachi M, Liu Y, Fujii K, Calderwood SK, Nakai A, Imai K, and Shinomura Y. Oxidative stress impairs the heat stress response and delays unfolded protein recovery. *PLoS One* 4: e7719, 2009.
- Ahn SG and Thiele DJ. Redox regulation of mammalian heat shock factor 1 is essential for Hsp gene activation and protection from stress. *Genes Dev* 17: 516–528, 2003.
- Back P, De Vos WH, Depuydt GG, Matthijssens F, Vanfleteren JR, and Braeckman BP. Exploring real-time *in vivo* redox biology of developing and aging *Caenorhabditis elegans*. *Free Rad Biol Med* 52: 850–859, 2012. <http://dx.doi.org/10.1016/j.freeradbiomed.2011.11.037> 2011.
- Ben-Zvi A, Miller EA, and Morimoto RI. Collapse of proteostasis represents an early molecular event in *Caenorhabditis elegans* aging. *Proc Natl Acad Sci U S A* 106: 14914–14919, 2009.
- Bernstein E, Kim SY, Carmell MA, Murchison EP, Alcorn H, Li MZ, Mills AA, Elledge SJ, Anderson KV, and Hannon GJ. Dicer is essential for mouse development. *Nat Genet* 35: 215–217, 2003.
- Boehm M and Slack F. A developmental timing microRNA and its target regulate life span in *C. elegans*. *Science* 310: 1954–1957, 2005.
- Bonelli MA, Alfieri RR, Petronini PG, Brigotti M, Campanini C, and Borghetti AF. Attenuated expression of 70-kDa heat shock protein in WI-38 human fibroblasts during aging *in vitro*. *Exp Cell Res* 252: 20–32, 1999.

8. Brenner S. The genetics of *Caenorhabditis elegans*. *Genetics* 77: 71–94, 1974.
9. Currie RW, Karmazyn M, Kloc M, and Mailer K. Heat-shock response is associated with enhanced postischemic ventricular recovery. *Circ Res* 63: 543–549, 1988.
10. Cypser JR and Johnson TE. Multiple stressors in *Caenorhabditis elegans* induce stress hormesis and extended longevity. *J Gerontol A Biol Sci Med Sci* 57: B109–B114, 2002.
11. de Lencastre A, Pincus Z, Zhou K, Kato M, Lee SS, and Slack FJ. MicroRNAs both promote and antagonize longevity in *C. elegans*. *Curr Biol* 20: 2159–2168, 2010.
12. Enyedi B, Varnai P, and Geiszt M. Redox state of the endoplasmic reticulum is controlled by Ero1L-alpha and intraluminal calcium. *Antioxid Redox Signal* 13: 721–729, 2010.
13. Finkel T. Radical medicine: treating ageing to cure disease. *Nat Rev Mol Cell Biol* 6: 971–976, 2005.
14. Gosslau A, Ruoff P, Mohsenzadeh S, Hobohm U, and Rensing L. Heat shock and oxidative stress-induced exposure of hydrophobic protein domains as common signal in the induction of hsp68. *J Biol Chem* 276: 1814–1821, 2001.
15. Hajdu-Cronin YM, Chen WJ, and Sternberg PW. The L-type cyclin CYL-1 and the heat-shock-factor HSF-1 are required for heat-shock-induced protein expression in *Caenorhabditis elegans*. *Genetics* 168: 1937–1949, 2004.
16. Halliwell B and Gutteridge JMC. *Free Radicals in Biology and Medicine*. Oxford; New York: Oxford University Press; 2007. xxxvi, 851 p., [8] p. of plates p.
17. Hu S, Claud EC, Musch MW, Chang EB. Stress granule formation mediates the inhibition of colonic Hsp70 translation by interferon-gamma and tumor necrosis factor-alpha. *Am J Physiol Gastrointest Liver Physiol* 298: G481–G492, 2010.
18. Hu S, Zhu X, Triggs JR, Tao Y, Wang Y, Lichtenstein L, Bissonnette M, Musch MW, and Chang EB. Inflammation-induced, 3'UTR-dependent translational inhibition of Hsp70 mRNA impairs intestinal homeostasis. *Am J Physiol Gastrointest Liver Physiol* 296: G1003–G1011, 2009.
19. Ibanez-Ventoso C, Yang M, Guo S, Robins H, Padgett RW, and Driscoll M. Modulated microRNA expression during adult lifespan in *Caenorhabditis elegans*. *Aging Cell* 5: 235–246, 2006.
20. Kim JK, Gabel HW, Kamath RS, Tewari M, Pasquinelli A, Rual JF, Kennedy S, Dybbs M, Bertin N, Kaplan JM, Vidal M, and Ruvkun G. Functional genomic analysis of RNA interference in *C. elegans*. *Science* 308: 1164–1167, 2005.
21. Kurucz I, Tombor B, Prechl J, Erdo F, Hegedus E, Nagy Z, Vitai M, Koranyi L, and Laszlo L. Ultrastructural localization of Hsp-72 examined with a new polyclonal antibody raised against the truncated variable domain of the heat shock protein. *Cell Stress Chaperones* 4: 139–152, 1999.
22. Lee JS and Seo JS. Differential expression of two stress-inducible hsp70 genes by various stressors. *Exp Mol Med* 34: 131–136, 2002.
23. Leung AK and Sharp PA. MicroRNA functions in stress responses. *Mol Cell* 40: 205–215, 2010.
24. Li X, Cassidy JJ, Reinke CA, Fischboeck S, and Carthew RW. A microRNA imparts robustness against environmental fluctuation during development. *Cell* 137: 273–282, 2009.
25. Lin Y, Liu X, Cheng Y, Yang J, Huo Y, and Zhang C. Involvement of MicroRNAs in hydrogen peroxide-mediated gene regulation and cellular injury response in vascular smooth muscle cells. *J Biol Chem* 284: 7903–7913, 2009.
26. Lithgow GJ, White TM, Melov S, and Johnson TE. Thermotolerance and extended life-span conferred by single-gene mutations and induced by thermal stress. *Proc Natl Acad Sci U S A* 92: 7540–7544, 1995.
27. Liu Q and Paroo Z. Biochemical principles of small RNA pathways. *Ann Rev Biochem* 79: 295–319, 2010.
28. Manalo DJ, Lin Z, and Liu AY. Redox-dependent regulation of the conformation and function of human heat shock factor 1. *Biochemistry* 41: 2580–2588, 2002.
29. McColl G, Rogers AN, Alavez S, Hubbard AE, Melov S, Link CD, Bush AI, Kapahi P, and Lithgow GJ. Insulin-like signaling determines survival during stress via posttranscriptional mechanisms in *C. elegans*. *Cell Metab* 12: 260–272.
30. McMillan DR, Xiao X, Shao L, Graves K, and Benjamin IJ. Targeted disruption of heat shock transcription factor 1 abolishes thermotolerance and protection against heat-inducible apoptosis. *J Biol Chem* 273: 7523–7528, 1998.
31. Miska EA, Alvarez-Saavedra E, Abbott AL, Lau NC, Hellman AB, McGonagle SM, Bartel DP, Ambros VR, and Horvitz HR. Most *Caenorhabditis elegans* microRNAs Are Individually Not Essential for Development or Viability. *PLoS Genet* 3: e215, 2007.
32. Morimoto RI. Proteotoxic stress and inducible chaperone networks in neurodegenerative disease and aging. *Genes Dev* 22: 1427–1438, 2008.
33. Moseley PL, Wallen ES, McCafferty JD, Flanagan S, and Kern JA. Heat stress regulates the human 70-kDa heat-shock gene through the 3'-untranslated region. *Am J Physiol* 264: L533–L537, 1993.
34. Muller FL, Lustgarten MS, Jang Y, Richardson A, and Van Remmen H. Trends in oxidative aging theories. *Free Radic Biol Med* 43: 477–503, 2007.
35. Namba T, Tanaka K, Hoshino T, Azuma A, and Mizushima T. Suppression of expression of heat shock protein 70 by gefitinib and its contribution to pulmonary fibrosis. *PLoS ONE* 6: e27296, 2011.
36. Oberringer M, Baum HP, Jung V, Welter C, Frank J, Kuhlmann M, Mutschler W, and Hanselmann RG. Differential expression of heat shock protein 70 in well healing and chronic human wound tissue. *Biochem Biophys Res Commun* 214: 1009–1014, 1995.
37. Olahova M, Taylor SR, Khazaipoul S, Wang J, Morgan BA, Matsumoto K, Blackwell TK, and Veal EA. A redox-sensitive peroxiredoxin that is important for longevity has tissue- and stress-specific roles in stress resistance. *Proc Natl Acad Sci U S A* 105: 19839–19844, 2008.
38. Petersen RB and Lindquist S. Regulation of HSP70 synthesis by messenger RNA degradation. *Cell Regul* 1: 135–149, 1989.
39. Petriv OI and Rachubinski RA. Lack of peroxisomal catalase causes a progeric phenotype in *Caenorhabditis elegans*. *J Biol Chem* 279: 19996–20001, 2004.
40. Putics A, Vegh EM, Csermely P, and Soti C. Resveratrol induces the heat-shock response and protects human cells from severe heat stress. *Antioxid Redox Signal* 10: 65–75, 2008.
41. Rehwinkel J, Natalin P, Stark A, Brennecke J, Cohen SM, and Izaurralde E. Genome-wide analysis of mRNAs regulated by Drosophila and Argonaute proteins in *Drosophila melanogaster*. *Mol Cell Biol* 26: 2965–2975, 2006.
42. Simone NL, Soule BP, Ly D, Saleh AD, Savage JE, Degraff W, Cook J, Harris CC, Gius D, and Mitchell JB. Ionizing radiation-induced oxidative stress alters miRNA expression. *PLoS One* 4: e6377, 2009.

43. Steels EL, Watson K, and Parsons PG. Relationships between thermotolerance, oxidative stress responses and induction of stress proteins in human tumour cell lines. *Biochem Pharmacol* 44: 2123–2129, 1992.
44. Tang F, Kaneda M, O'Carroll D, Hajkova P, Barton SC, Sun YA, Lee C, Tarakhovsky A, Lao K, and Surani MA. Maternal microRNAs are essential for mouse zygotic development. *Genes Dev* 21: 644–648, 2007.
45. Timmons L, Court DL, and Fire A. Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in *Caenorhabditis elegans*. *Gene* 263: 103–112, 2001.
46. Tranter M, Helsley RN, Paulding WR, McGuinness M, Brokamp C, Haar L, Liu Y, Ren X, and Jones WK. Coordinated post-transcriptional regulation of Hsp70.3 gene expression by microRNA and alternative polyadenylation. *J Biol Chem* 286: 29828–29837, 2011.
47. Walker GA and Lithgow GJ. Lifespan extension in *C. elegans* by a molecular chaperone dependent upon insulin-like signals. *Aging Cell* 2: 131–139, 2003.
48. Yin C, Salloum FN, and Kukreja RC. A novel role of microRNA in late preconditioning: upregulation of endothelial nitric oxide synthase and heat shock protein 70. *Circ Res* 104: 572–575, 2009.
49. Yin C, Wang X, and Kukreja RC. Endogenous microRNAs induced by heat-shock reduce myocardial infarction following ischemia-reperfusion in mice. *FEBS Lett* 582: 4137–4142, 2008.

Address correspondence to:
 Dr. Csaba Söti
 Department of Medical Chemistry
 Semmelweis University
 Tüzoltó u. 37-47
 Budapest H-1094
 Hungary

E-mail: soti.csaba@med.semmelweis-univ.hu

Date of first submission to ARS Central, July 14, 2011; date of final revised submission, February 5, 2012; date of acceptance, February 23, 2012.

Abbreviations Used

3'UTR = 3' untranslated region
 H₂O₂ = hydrogen-peroxide
 HSF1 = heat shock transcription factor 1
 Hsp = heat shock protein
 miRNA = microRNA
 NAC = N-acetyl-L-cysteine
 NGM = nematode growth medium
 PCR = polymerase chain reaction
 ROS = reactive oxygen species
 MFI = mean fluorescence intensity