

Phosphodiesterase 8 governs cAMP/PKA-dependent reduction of L-type calcium current in human atrial fibrillation: a novel arrhythmogenic mechanism

Nefeli Grammatika Pavlidou^{1,2}, Shokoufeh Dobrev³, Kira Beneke^{1,2}, Franziska Reinhardt^{1,2,4}, Simon Pecha ^{2,4}, Eric Jacquet ⁵, Issam H. Abu-Taha³, Constanze Schmidt ⁶, Niels Voigt ^{7,8,9}, Markus Kamler ¹⁰, Renate B. Schnabel^{2,11}, Istvan Baczkó ¹², Anne Garnier¹³, Hermann Reichenspurner^{2,4}, Viacheslav O. Nikolaev ^{1,2}, Dobromir Dobrev ^{3,14,15}, and Cristina E. Molina ^{1,2,3*}

¹Institute of Experimental Cardiovascular Research and University Center of Cardiovascular Sciences, University Medical Center Hamburg Eppendorf (UKE), Martinistrasse 52, W23, 20246 Hamburg, Germany; ²German Centre for Cardiovascular Research (DZHK), partner site Hamburg/Kiel/Lübeck, Potsdamer Strasse 58, 10785 Berlin, Germany; ³Institute of Pharmacology, West German Heart and Vascular Center, University Duisburg-Essen, Hufelandstrasse 55, 45147 Essen, Germany; ⁴Department of Cardiovascular Surgery, University Heart Center Hamburg, Martinistrasse 52, O70, 20246 Hamburg, Germany; ⁵Université Paris-Saclay, CNRS, Institut de Chimie des Substances Naturelles, UPR 2301, 1 avenue de la Terrasse, 91198 Gif-sur-Yvette, France; ⁶Department of Cardiology, University Medical Center Heidelberg, Im Neuenheimer Feld 672, 69120 Heidelberg, Germany; ⁷Institute of Pharmacology and Toxicology, University Medical Center Göttingen, Robert-Koch-Strasse 40, 37075 Göttingen, Germany; ⁸German Centre for Cardiovascular Research (DZHK), partner Site Göttingen, Potsdamer Strasse 58, 10785 Berlin, Germany; ⁹Cluster of Excellence 'Multiscale Bioimaging: from Molecular Machines to Networks of Excitable Cells' (MBExC), University of Göttingen, Robert-Koch-Strasse 40, 37075 Göttingen, Germany; ¹⁰Department of Thoracic and Cardiovascular Surgery, West German Heart and Vascular Center, University Duisburg-Essen, Hufelandstrasse 55, 45147 Essen, Germany; ¹¹Cardiology Department, University Heart and Vascular Centre Hamburg-Eppendorf, Martinistrasse 52, O70, 20246 Hamburg, Germany; ¹²Department of Pharmacology and Pharmacotherapy, University of Szeged, H-6721, Szeged, Dóm tér 12, Szeged, Hungary; ¹³Inserm, UMR-S 1180, Université Paris-Saclay, Faculté de pharmacie, 17 avenue des Sciences, 91400 Orsay, France; ¹⁴Department of Medicine and Research Center, Montreal Heart Institute and Université de Montréal, 4100 Molson Street, Suite 340, H1Y 3N1 Montréal, Canada; and ¹⁵Department of Molecular Physiology & Biophysics, Baylor College of Medicine, One Baylor Plaza, 77030 Houston, USA

Received 1 June 2022; revised 22 December 2022; accepted 3 February 2023

Abstract

Aims

Atrial fibrillation (AF) is associated with altered cAMP/PKA signaling and an AF-promoting reduction of L-type Ca^{2+} -current ($I_{\text{Ca,L}}$), the mechanisms of which are poorly understood. Cyclic-nucleotide phosphodiesterases (PDEs) degrade cAMP and regulate PKA-dependent phosphorylation of key calcium-handling proteins, including the $I_{\text{Ca,L}}$ -carrying Cav1.2 _{α 1C} subunit. The aim was to assess whether altered function of PDE type-8 (PDE8) isoforms contributes to the reduction of $I_{\text{Ca,L}}$ in persistent (chronic) AF (cAF) patients.

Methods and results

mRNA, protein levels, and localization of PDE8A and PDE8B isoforms were measured by RT-qPCR, western blot, co-immunoprecipitation and immunofluorescence. PDE8 function was assessed by FRET, patch-clamp and sharp-electrode recordings. PDE8A gene and protein levels were higher in paroxysmal AF (pAF) vs. sinus rhythm (SR) patients, whereas PDE8B was upregulated in cAF only. Cytosolic abundance of PDE8A was higher in atrial pAF myocytes, whereas PDE8B tended to be more abundant at the plasmalemma in cAF myocytes. In co-immunoprecipitation, only PDE8B2 showed binding to Cav1.2 _{α 1C} subunit which was strongly increased in cAF. Accordingly, Cav1.2 _{α 1C} showed a lower phosphorylation at Ser1928 in association with decreased $I_{\text{Ca,L}}$ in cAF. Selective PDE8 inhibition increased Ser1928 phosphorylation of Cav1.2 _{α 1C}, enhanced cAMP at the subsarcolemma and rescued the lower $I_{\text{Ca,L}}$ in cAF, which was accompanied by a prolongation of action potential duration at 50% of repolarization.

* Corresponding author. Tel: +49 40741057095, Fax: +49 40741040180, Email: c.molina@uke.de

© The Author(s) 2023. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Conclusion

Both PDE8A and PDE8B are expressed in human heart. Upregulation of PDE8B isoforms in cAF reduces $I_{Ca,L}$ via direct interaction of PDE8B2 with the Cav1.2 α_{1C} subunit. Thus, upregulated PDE8B2 might serve as a novel molecular mechanism of the proarrhythmic reduction of $I_{Ca,L}$ in cAF.

Structured Graphical Abstract**Key Question**


What is the molecular mechanism of reduced L-type calcium current that promotes persistent atrial fibrillation?

Key Finding

Cyclic-nucleotide phosphodiesterase 8B (PDE8B) was expressed in the human atria and upregulated in persistent atrial fibrillation, contributing to the proarrhythmic reduction of L-type calcium current by decreasing sarcolemmal cAMP levels.

Take Home Message

Targeting PDE8B may represent a new atrial-selective therapeutic strategy for atrial fibrillation.



The reduction of L-type Ca^{2+} -current in atrial myocytes promotes atrial fibrillation. Phosphodiesterase type 8B binds L-type Ca^{2+} channels and reduces local cAMP levels and PKA-dependent channel phosphorylation, thereby decreasing L-type Ca^{2+} -current and abbreviating atrial action potential that promotes atrial fibrillation persistence. ADP, adenosine diphosphate; cAF, persistent (chronic) atrial fibrillation; cAMP, cyclic adenosine monophosphate; DAD, delayed afterdepolarization; $I_{Ca,L}$, L-type Ca^{2+} current; LTCC, L-type Ca^{2+} channel; PDE, phosphodiesterase; PLB, phospholamban; RyR2, ryanodine receptor type 2; SERCA2a, sarcoplasmic reticulum Ca^{2+} ATPase type 2a; SLN, sarcolipin; SR, sinus rhythm.

Keywords

Phosphodiesterases • PDE8 • cAMP • Atrial fibrillation • Calcium handling • $I_{Ca,L}$

Translational perspective

The mechanism of the proarrhythmic reduction of L-type calcium current ($I_{Ca,L}$) in patients with persistent (chronic) atrial fibrillation (cAF) remains elusive. Here we show that patients with cAF exhibit a stronger binding of phosphodiesterase 8B2 (PDE8B2) to the pore-forming Cav1.2 $_{\alpha1C}$ subunit of $I_{Ca,L}$. Selective PDE8 inhibition enhanced Ser1928 phosphorylation of Cav1.2 $_{\alpha1C}$, increased subsarcolemmal cAMP, and rescued the reduced $I_{Ca,L}$ leading to action potential prolongation in cAF atria. We thus discovered a new molecular mechanism for the proarrhythmic reduction of $I_{Ca,L}$ in cAF and established PDE8B2 as a novel atrial-selective target for cAF treatment.

Introduction

Atrial fibrillation (AF) is a common disease with poorly understood pathophysiology.^{1,2} Remodeling of membrane receptors and alterations in 3',5'-cyclic adenosine monophosphate (cAMP)-dependent regulation of Ca^{2+} handling mechanisms^{3,4} are considered important contributors to AF promotion. For instance, upregulated sarcoplasmic reticulum Ca^{2+} ATPase type 2a (SERCA2a)^{3,5} and PKA-dependent hyperphosphorylation of ryanodine receptor type 2 (RyR2),^{6–8} are hallmarks of AF-related remodeling. Indeed, cAMP levels are increased in long-standing persistent (chronic) AF (cAF),³ probably due to a reduction of cAMP-hydrolyzing phosphodiesterases (PDEs).⁹ Decreased L-type Ca^{2+} current ($I_{Ca,L}$) density^{3,10} linked to PKA-dependent dephosphorylation of L-type Ca^{2+} channel (LTCC) macromolecular complex¹¹ is another feature of AF. Although type 2A protein phosphatase (PP2A) activity in AF is globally increased,¹² the inhomogeneous AF-associated changes of protein phosphorylation point to a local regulation of PKA within distinct subcellular compartments.

cAMP levels, and thus the degree of PKA activation, are tightly regulated by PDEs that degrade cAMP in subcellular nanodomains.¹³ So far, most of our knowledge of cAMP compartmentation was predominantly obtained in the ventricular myocytes where PDE3 and PDE4 are considered to be the main contributors to total cAMP-hydrolysis.^{14,15} PDE3 is predominant in microsomal fractions in large mammals,¹⁶ with PDE4 accounting for only $\approx 20\%$ of the total PDE activity in canine ventricles.¹⁷ Our previous studies demonstrated that although all the classical cardiac PDEs (PDE1–4) are expressed in human atria,^{9,18} PDE3 and PDE4 account for only 45% and 15% of total cAMP-hydrolysis, respectively, with a large portion of cAMP-hydrolysis remaining unexplained.⁹ In rodent^{19–21} and human^{20,22} ventricles, PDE4 was found to be associated with β -adrenoceptors, LTCC, RyR2 and phospholamban (PLN)-SERCA2 complexes. Previously, we could show that pharmacological inhibition of PDE4 increased the amplitude of $I_{Ca,L}$ and frequency of potentially proarrhythmic Ca^{2+} -sparks and Ca^{2+} -waves in human atrial myocytes (HAMs) of sinus rhythm (SR) patients.⁷ PDE3 and PDE2 may also contribute to $I_{Ca,L}$ regulation of HAMs.²³ The total cAMP-hydrolytic activity in response to the unselective PDE inhibitor IBMX, which inhibits PDE1–4, is reduced in AF compared to SR patients,⁹ suggesting that a reduction in PDE1–4 activity may contribute to RyR2 hyperphosphorylation and the related RyR2-mediated Ca^{2+} -dependent triggered activity in AF.^{6,7} However, this global reduction of PDE activity is inconsistent with the postulated cAF-associated reduction of LTCC,^{11,12} pointing to a contribution of additional PDE families. Since PDE8 is not inhibited by IBMX,²⁴ an increase in PDE8 activity could underlie the decrease of $I_{Ca,L}$ in cAF. The PDE8 family is encoded by the two genes, PDE8A and PDE8B, of which only PDE8A was found in human ventricles.^{25,26} We recently showed that PDE8A is also expressed in human atria.¹⁸ A previous study demonstrated that PDE8A is involved in the regulation of Ca^{2+} homeostasis in mouse ventricular myocytes.²⁷

Whether PDE8B is expressed in human atria and how PDE8 isoforms may contribute to the proarrhythmic reduction of $I_{Ca,L}$ in cAF is unknown and was the object of this study.

Here we show that both PDE8A and PDE8B subfamilies are present in HAMs. Patients with cAF exhibited a stronger binding of a specific PDE8B2 isoform to the pore-forming Cav1.2 $_{\alpha1C}$ subunit of LTCC. Selective PDE8 inhibition enhanced Ser1928 phosphorylation (pS1928) of Cav1.2 $_{\alpha1C}$, increased subsarcolemmal cAMP, and rescued the lower $I_{Ca,L}$, leading to action potential (AP) prolongation in cAF atria. This provides a new molecular mechanism for the proarrhythmic reduction of $I_{Ca,L}$ in cAF with PDE8B2 as a potential novel atrial-selective target for cAF treatment.

Methods

Study design and human atrial samples

Right atrial (RA) tissues were obtained from patients in SR, with paroxysmal AF (pAF), or with cAF who had undergone open-heart surgery. All available RA appendages were used consecutively. Of the 165 RA samples obtained, 32 were used for qPCR, 69 for biochemical experiments, 47 for HAM isolation (thereof 19 for immunofluorescence, 18 for live cell imaging, and 10 for patch-clamp), and 17 for AP recordings (see [Supplementary data online, Tables S1–S8](#)). Experimental protocols were approved by ethical boards of University Hospital Essen (#12-5268-BO) and Medical Chamber Hamburg (WF-088/18) and were conducted in accordance with the Declaration of Helsinki. Each individual provided written informed consent. In addition, we included RA and left ventricular (LV) tissues from eight healthy (Ctl) and eight heart failure (HF) patients undergoing heart transplantation. This study was approved by the Scientific Board at the Hungarian Ministry of Health (ETT-TUKEB:4991-0/2010-1018EKU). Patient characteristics are provided for all samples, samples used for qPCR, immunoblotting, immunostaining, FRET, patch-clamp, and AP recordings (see [Supplementary data online, Tables S1–S8](#)).

RNA isolation, cDNA synthesis, real-time qPCR and quantification

Frozen tissue samples from 48 patients (16 SR, 8 pAF, 8 cAF, 8 Ctl and 8 HF patients) were used for real-time PCR assays performed as previously described¹⁸ to study the expression of 2 target genes (*PDE8A*, *PDE8B*) related to specifically chosen²⁸ reference genes (*POLR2A*, *YWHAZ*, *GAPDH*, *IPO8*, *PPIA*). Further experimental details are available in the online Supplement.

Western blot analysis and co-immunoprecipitation

54 snap-frozen atrial samples were used to perform western blot analysis. From those, 12 samples were, initially, incubated for 5 min with control buffer (CON), 30 nM PF-04957325 (PF), or 100 nM β -adrenergic agonist isoprenaline (ISO), and then snap-frozen. Another 15 snap-frozen samples were homogenized in immunoprecipitation buffer and used for co-immunoprecipitation experiments. Further experimental details are available in the online Supplement.

Isolation and culture of HAMs

After surgical excision, tissue samples were placed into Custodiol® solution (Dr. Franz Köhler Chemie GmbH, Bensheim, Germany) for the transport to the laboratory. Cell isolations from 47 patients were carried out as previously described.⁷ Please see also [Supplementary material online](#).

Perforated patch-clamp in freshly-isolated and cultured HAMs

Whole-cell perforated patch-clamp configuration was used to record $I_{Ca,L}$ in 16 HAMs from 10 patients as previously described.^{4,11}

Live cell imaging of sarcolemmal cAMP in HAMs

Förster resonance energy transfer (FRET) based measurements of cAMP were performed on 38 living HAMs isolated from 18 patients and transfected with an adenovirus encoding for the pmEpac1-camps biosensor to measure cAMP at the membrane.²⁹ See online Supplement.

Immunocytochemistry and confocal imaging

HAMs from 7 SR patients, 5 pAF patients and 7 cAF patients were used for immunostaining as described in online Supplement.

Sharp-electrode AP recordings

APs were recorded as previously described using standard intracellular microelectrodes in atrial trabeculae from 5 SR, 5 pAF and 7 cAF patients.³⁰ Further experimental details are available in the online Supplement.

Data analysis and statistics

Results are presented as mean \pm SEM. $P < 0.05$ was considered statistically significant. Normal distribution was tested by Kolmogorov–Smirnov or by Shapiro–Wilk tests. Statistical differences between groups with normally distributed data were analyzed using mixed ANOVA followed by Wald χ^2 test, Student's *t*-test for two-group comparison, or one-way ANOVA and Sidak multiple comparison test for multiple groups. Non-normally distributed continuous data or data for which normality could not be assessed were compared using Mann–Whitney tests or one-way ANOVA for multiple groups with a Kruskal–Wallis test. To investigate potential influences of individual clinical variables on specific parameters, two-way ANOVA sub-analyses with factors rhythm and either age, sex, bypass surgery, hyperlipidemia, use of AT1-blockers, ACE inhibitors, diuretics or amiodarone were performed (see [Supplementary data online, Figures S4–S8](#)).

Results

Patients

Patients with pAF had more often hyperlipidemia, diabetes and cardiopulmonary bypass surgery and were younger in some subgroups. pAF and cAF patients were more often taking digitalis, beta-blockers and diuretics, with cAF patients being more frequently treated with amiodarone (see [Supplementary data online, Table S1](#)). There were no significant differences between groups in sex, body mass index, comorbidities, degree of valvular regurgitation, left ventricular ejection fraction. Multivariate analysis for hyperlipidemia, cardiopulmonary bypass, digitalis, beta-blockers, and amiodarone showed no alterations in the results (see [Supplementary data online, Figures S4–S8](#)). Although use of diuretics was associated with higher mRNA levels of PDE8B in SR (see [Supplementary data online, Figure S5](#)), this association was not observed in any of the other data sets. Other medication was not different between groups.

Expression and localization of PDE8 isoforms and Cav1.2 in human atrium

To determine if both PDE8 isoforms are present in human atria and altered in AF, we assessed gene expression ([Figure 1A](#)) and protein levels ([Figure 1B](#)) of PDE8A and PDE8B in atria from SR, pAF and cAF patients. PDE8A mRNA was present in human atria and increased in both AF groups, whereas PDE8B was increased in cAF only ([Figure 1A](#)).

The full-length human PDE8A1 isoform has 829 amino acids with a calculated molecular weight of 93 kDa, whereas the PDE8A2 isoform consists of 783 amino acids with a calculated molecular weight of 88 kDa.³¹ Similarly, the full-length human PDE8B1 isoform has 885 amino acids and an expected molecular mass of approximately 99 kDa, whereas the PDE8B2 variant has 838 amino acids and an expected molecular mass of 87 kDa.³² Previous work could detect differential protein expression of different PDE8A and PDE8B isoforms in rat brain, mouse ovary and bovine testis.^{33,34} In human atria of SR, pAF, and cAF patients we could detect two specific bands between 75 and 100 kDa, which were in reasonable agreement with the expected molecular masses predicted for PDE8A1, PDE8A2, PDE8B1, and PDE8B2 ([Figure 1B](#)). The protein levels of PDE8A1 and PDE8A2 were significantly higher in pAF but unaltered in cAF. Conversely, PDE8B1 and PDE8B2 were up-regulated in cAF only ([Figure 1B](#)). In contrast, human LV do not express PDE8B and expression of PDE8A was similar between control (Ctl) and HF patients ([Figure 1C](#)). Most important, although human atria express both PDE8A and PDE8B, their mRNA levels were not changed during HF, indicating that the PDE8B increases were AF-specific ([Figure 1C](#)).

In addition, PDE8A was more abundant in the cytosol in HAMs and its abundance was increased in pAF vs. SR and cAF ([Figure 2A and C](#) and [Supplementary data online, Figure S1](#)). In contrast, PDE8B was preferentially localized at the plasma membrane, especially in cAF where its expression was increased compared to SR and pAF ([Figure 2B and C](#) and [Supplementary data online, Figure S1](#)).

Since PDEs degrade cAMP^{13,15,22} and impaired PKA-dependent LTCC regulation was suggested to be the main mechanism for $I_{Ca,L}$ reduction in cAF,^{11,35} we examined whether PDE8B, which we found up-regulated in cAF, co-localizes with LTCC in the macromolecular complex ([Figure 3A](#)). In co-immunoprecipitations PDE8B was the only partner of Cav1.2, with PDE8B2 being the major isoform ([Figure 3A](#)). PDE8B2 showed a stronger binding to Cav1.2 α_{1C} subunit in cAF ([Figure 3A](#)).

Subsequently, we studied the consequences of selective PDE8 inhibition on cAMP/PKA-regulation of LTCC by assessing the phosphorylation state of Cav1.2 α_{1C} at Ser1928 (Cav1.2-pS1928), a validated PKA phosphorylation site of Cav1.2 α_{1C} ,³⁶ in atrial tissue lysates incubated with control buffer (CON), the selective PDE8 inhibitor PF-04957325 (PF) or the β -adrenoceptor agonist isoprenaline (ISO) from SR and cAF patients. Although phosphorylation of Cav1.2-pS1928 is not required for β -adrenoceptor stimulation of LTCC in mouse ventricular myocytes and HEK293 cells,^{37–39} it is a well-established Cav1.2 PKA phosphorylation site³⁶ and phosphorylation of its α_{1C} subunit is relevant for channel function by increasing trafficking to the plasma membrane.⁴⁰ In CON, the relative Cav1.2-pS1928 phosphorylation was significantly lower in cAF compared to SR and increased significantly after incubation with ISO or PF to levels seen in corresponding CONs ([Figure 3C](#) and [Supplementary data online, Figure S2](#)), pointing to significant contribution of PDE8 to LTCC dephosphorylation. In agreement with previous studies,^{12,41} no significant difference in total Cav1.2 α_{1C} protein expression was observed in cAF compared to SR (see [Supplementary data online, Figure S2](#)).




Figure 1 PDE8A and PDE8B expression in human atrium. (A) Plot of the individual and mean \pm SEM gene expression ratio of PDE8A and PDE8B normalized to a set of five housekeeping genes (POLR2A, YWHAZ, GAPDH, IPO8, PPIA) in atrial tissue homogenates from 16 sinus rhythm (SR), 8 paroxysmal atrial fibrillation (pAF) and 8 persistent (chronic) atrial fibrillation (cAF) patients. $*P < 0.05$ vs. sinus rhythm SR based on analysis of variance (ANOVA with a Kruskal–Wallis test). (B) Representative western blot (top) and protein expression quantification (bottom, mean \pm SEM) of PDE8A and PDE8B in atrial tissue homogenates from 16 SR, 10 pAF and 16 cAF patients. GAPDH was used as loading control. $*P < 0.05$ vs. SR based on unpaired Student's *t*-test analysis. (C) Plot of the individual and mean \pm SEM gene expression ratio of PDE8A and PDE8B normalized to the set of five housekeeping genes in right atrial and left ventricular tissue homogenates from eight healthy control (Ctl) and eight end-stage heart failure (HF) patients. $*P < 0.05$ vs. Ctl based on ANOVA and Mann–Whitney test.




Figure 2 PDE8A and PDE8B localization in human atrial myocytes. (A) Representative immunocytochemistry confocal images of PDE8A showing its cytosolic distribution in 36 isolated human atrial myocytes (HAMs) from 7 sinus rhythm (SR), 25 HAMs from 5 paroxysmal atrial fibrillation (pAF) and 29 HAMs from 7 persistent (chronic) atrial fibrillation (cAF) patients. Lower panels show transversal line profiles of fluorescence intensity (F.I.) of PDE8A staining for all the myocytes analyzed (thin colored lines) and the average within all cells (thick black lines). (B) Similar representative immunocytochemistry confocal images and fluorescence intensity profiles of PDE8B showing mainly plasma membrane localization of this PDE isoform, in 38 HAMs from 7 SR, 25 HAMs from 5 pAF and 31 HAMs from 7 cAF patients. (C) Average F.I. of PDE8A and PDE8B at the plasma membrane (first and last 10% of the cell width) related to the F.I. in the cytosol (middle), in SR, pAF, and cAF. * $P < 0.05$ between groups of patients, # $P < 0.05$ vs. PDE8A, by mixed ANOVA followed by Wald χ^2 test and Sidak multiple comparison test. All images were captured under the same conditions in order to obtain comparable fluorescence intensities.




Figure 3 PDE8 co-localizes with LTCC. (A) Co-immunoprecipitation (Co-IP) assay followed by immunoblotting (IB) confirmed that with L-type Ca²⁺ channel (Cav1.2) binds to PDE8B, PKA_{RII} and PKA_c in human atria from patients with persistent (chronic) atrial fibrillation (cAF). NSB, non-specific binding. (B) Top, representative Co-IP and IB of Cav1.2 binding to PDE8B in human atrial samples from patients in sinus rhythm (SR), paroxysmal AF (pAF) and cAF. Bottom, mean data of PDE8B binding to Cav1.2 for all groups ($n = 6$ SR, $n = 4$ pAF, and 5 cAF patients). Lys, Lysate. Vertical white lines separate non-adjacent lanes on the same blot. (C) Representative western blot (left) and quantification (right, mean \pm SEM) of relative steady-state LTCC phosphorylation (pSer1928/total Cav1.2) in atrial tissue homogenates from 6 SR and 6 cAF patients, at baseline and after 5 min stimulation with the selective PDE8 inhibitor PF-04957325 (30 nM) or the β -adrenoceptor agonist isoprenaline (ISO, 100 nM). GAPDH was used as loading control. * $P < 0.05$ based on ANOVA with a Kruskal–Wallis test.




Figure 4 Functional consequences of selective PDE8 inhibition in human atrial myocytes (HAMs). (A) Left panel, representative experiments showing the time course of the FRET signals indicating cAMP changes in HAMs from sinus rhythm (SR) and in persistent (chronic) atrial fibrillation (cAF) patients exposed to selective PDE8 inhibition with PF-04957325 (30 nM). Right panel, effects of the selective PDE8 inhibitor and the non-selective PDE blocker IBMX (100 mM) on the individual and mean values of cAMP levels expressed as percentage of increase of CFP/YFP over its control value measured in 14 HAMs from 6 SR, 12 HAMs from 6 paroxysmal AF (pAF), and 12 HAMs from 6 cAF patients. The changes in FRET ratio were expressed as a percentage change vs. basal. * $P < 0.05$ compared with SR, # $P < 0.05$ compared with PF. (B) Left panel, current-voltage (I - V) relationship for peak inward $I_{Ca,L}$ density in HAMs from SR and cAF patients before (CON) and after exposure to PF. Middle panels, representative L-type Ca^{2+} current ($I_{Ca,L}$) recordings measured at 0 mV in HAMs from SR and cAF patients before (CON) and after exposure to PF. Right panel, average current densities before and after exposure to PF ($n = 8$ cells/5 patients SR and 8/5 cAF). * $P < 0.05$ vs. sinus rhythm SR, # $P < 0.05$ vs. CON. Statistical differences analyzed by mixed ANOVA followed by Wald χ^2 test and Sidak multiple comparison test.

Altogether, these results show that PDE8B localizes at the plasma membrane of HAMs and accumulates in the LTCC complex of cAF patients, whereby it causes LTCC dephosphorylation. This positions PDE8B as a potential molecular mechanism of the reduced $I_{Ca,L}$ that promotes AF.

PDE8 controls subsarcolemmal cAMP concentration in HAMs

To directly test whether PDE8 controls cAMP levels near the LTCC, cAMP concentration at the plasma membrane was measured in living HAMs from SR, pAF and cAF patients using the FRET-based targeted biosensor pmEpac1-camps.^{9,29} Application of PF strongly increased basal levels of cAMP in the subsarcolemmal compartment in cAF compared to SR (Figure 4A), reaching approximately 74% of the respective maximal response elicited by the non-selective PDE inhibitor IBMX (100 μ M). These results indicate that PDE8 controls basal cAMP

levels at the plasma membrane of HAMs in cAF, proving that the observed increase of PDE8B2 in the LTCC complex could contribute to the dephosphorylation of LTCC and the reported reduction in $I_{Ca,L}$ in cAF.

No changes in $I_{Ca,L}$ density or cell capacitance were observed when comparing freshly-isolated and 48-h cultured HAMs from the same patients at baseline and after ISO stimulation (see [Supplementary data online, Figure S3](#)).

PDE8 controls $I_{Ca,L}$ function

We then studied the direct consequences of increased PDE8B2 for $I_{Ca,L}$ function with perforated patch-clamp recordings. $I_{Ca,L}$ amplitude was lower in cAF, but was rescued by selective PDE8 inhibition with PF (Figure 4B) and this effect was fully reversible. These findings validate the novel concept that enhanced PDE8B is a major molecular mechanism for the reduction of $I_{Ca,L}$ in cAF.




Figure 5 Effect of selective PDE8 inhibition on action potential (AP) properties of human atrial trabeculae. (A) Representative traces of AP at baseline (CON) and after 20 min of exposure to the selective PDE8 inhibitor PF-04957325 (PF, 100 nM) recorded in atrial trabeculae from patients in sinus rhythm (SR) and in persistent (chronic) atrial fibrillation (cAF). (B) Left panel, average resting membrane potentials in trabeculae from five SR, five paroxysmal AF (pAF), and seven cAF patients. Middle panel, individual and mean values of the effects of the selective PDE8 inhibitor (PF) on AP duration at 50% of repolarization (APD₅₀) in the three groups of patients. Right panel, plateau potential before (CON) and after exposure to the selective PDE8 inhibitor (PF). * $P < 0.05$ vs. SR, # $P < 0.05$ vs. CON based on ANOVA with a Kruskal–Wallis test.

Effects of PDE8 inhibition on AP in cAF

To assess the consequences of PDE8 inhibition and the related upregulation of $I_{Ca,L}$ for AP properties, we performed sharp-electrode AP recordings in atrial trabeculae from SR, pAF and cAF patients. Resting membrane potentials were similar between all groups. In cAF PDE8 inhibition shifted the plateau potential, calculated as the average voltage during a time window of 20 to 90 ms after AP upstroke, to more positive values. This shift leads to a slight prolongation of the AP duration at 50% of repolarization (APD₅₀) in cAF vs. SR and pAF (Figure 5), which is consistent with a stimulation of $I_{Ca,L}$. The PF effect was reversible on washout.

Discussion

Our study demonstrates that PDE8A and PDE8B are expressed in human atria and HAMs. PDE8B preferentially localizes at the plasma membrane in close proximity to Cav1.2_{α1C}, and the PDE8B2 isoform is enriched in the LTCC complex in cAF. Furthermore, PDE8B is central in controlling cAMP at the plasma membrane which provides a new molecular mechanism of the reduced $I_{Ca,L}$ in cAF (Structured Graphical Abstract). These novel insights suggest that selective PDE8B inhibition may constitute a novel atrial-selective anti-AF strategy.

Although a reduction in $I_{Ca,L}$ amplitude has been consistently reported in HAMs from cAF patients,^{3,10,11,42–44} the underlying molecular mechanism has been poorly understood. Reduced $I_{Ca,L}$ is considered a main cause of the reentry-promoting APD shortening,⁴⁵ and atrial contractile dysfunction in AF.⁴⁶ Different mechanisms have been suggested to contribute to the reduced $I_{Ca,L}$ in AF. Although expression of the accessory subunit $\alpha 2\delta$ seems to be reduced,⁴⁷ protein expression of the pore-forming Cav1.2 subunits $\alpha 1C$ or $\beta 2$ are unchanged in AF,⁴¹ suggesting that the reduced $I_{Ca,L}$ might be due to altered regulation of single-channel properties. In addition, we and others have demonstrated a higher sensitivity of $I_{Ca,L}$ to β -adrenoceptor stimulation in AF compared to SR^{10,11,44} suggesting that, opposite to the hyperphosphorylation of RyR2 or PLN,^{3,7,8} basal phosphorylation of the LTCC complex appears reduced.¹² The stronger stimulatory effect of isoprenaline on $I_{Ca,L}$ in AF¹¹ indicates that the alterations in Ca^{2+} -handling in AF are likely due to a specific alteration of cAMP-dependent protein phosphorylation in discrete subcellular nanodomains. These results are in agreement with recent studies reporting atrial-specific cAMP nanodomains in the sarcolemmal axial tubule dyadic junctions with the SR in association with RyR2 hyperphosphorylation and clustering,³⁵ and in the caveolar structures linked to extradyadic LTCC subpopulations.⁴⁸ Since protein phosphorylation states depend on the dynamic balance between protein kinases and protein phosphatases, it has been suggested

that a local regulation of PP2A activity may contribute to these compartmental differences in cAMP/PKA-dependent phosphorylation.¹² However, although increased PP2A may contribute to the reduced $I_{Ca,L}$, the strong decrease in $I_{Ca,L}$ in the face of hyperphosphorylated RyR2 and PLN in AF^{3,7,8} likely results from different molecular mechanisms. Since in ventricular cardiomyocytes local PKA activity highly depends on the local availability of cAMP, which in turn depends on its degradation by PDEs at each discrete nanodomain,^{13,15,22} PDEs could contribute to the unique LTCC complex dephosphorylation in AF. Although the classical cardiac PDE2–4 were associated with $I_{Ca,L}$,^{9,23} their activity decreases in AF,⁹ suggesting that the decreased $I_{Ca,L}$ in AF cannot result from altered PDE2–4 activities. Here we show that a novel PDE8 isoform, PDE8B, is present in human atria, but unlike PDE8A,²⁵ PDE8B is not present in human ventricles,²⁶ and constitutes an atrial-selective PDE isoform. Importantly, our results demonstrate that PDE8B1 and PDE8B2 proteins are upregulated in cAF. Furthermore, PDE8B localizes at the sarcolemma of HAMs, with PDE8B2 strongly binding to Cav1.2_{α1C} in cAF. Functionally, PDE8B inhibition increases Cav1.2-pS1928 phosphorylation, cAMP levels in the subsarcolemma, and rescues $I_{Ca,L}$, thereby prolonging APD₅₀ in cAF.

We speculate that the PDE8B2 upregulation in the LTCC complex, the subsequent Cav1.2_{α1C} dephosphorylation at pS1928 and the related $I_{Ca,L}$ reduction in cAF may be a protective mechanism to counteract the cytotoxic Ca^{2+} overload and related negative consequences for cardiac metabolic activity due to the high atrial rate and energy demand, although this occurs at the expense of reentry-promoting APD shortening.⁴⁹ The PDE8B upregulation could promote Rad-mediated inhibition of LTCC^{37,50} and consequently reduce Ca^{2+} influx to compensate the increase in diastolic intracellular $[Ca^{2+}]$ that causes AF-promoting remodeling.¹ Although Ca^{2+} -channel blockers can be used to control the ventricular rate in AF, they could promote AF and cause ventricular dysfunction.^{51,52} Since PDE8B and particularly PDE8B2 occurrence appear atrial-selective, selective inhibition of PDE8B2 could constitute a viable approach to normalize the abnormal LTCC function and may constitute a novel atrial-selective anti-AF target. However, $I_{Ca,L}$ is not downregulated in patients with pAF.⁵³ Thus, the reduction of $I_{Ca,L}$ in cAF could be a consequence of the persistency of the arrhythmia and/or the sustained high atrial rate during AF. The related shortening of ERP/APD, which promotes AF-maintaining reentry, promotes the progression and the stabilization of AF. Thus, restoration of physiological $I_{Ca,L}$ by selective inhibition of PDE8B is expected to prevent the progression of AF to persistent forms rather than the initiation of AF and might also constitute a rhythm control strategy after cardioversion. These possibilities require direct verification in large animal models and suitable patient populations.

Our results clearly show that both PDE8A and PDE8B are expressed in human atrial tissue and myocytes. However, the immunoblot experiments showed multiple bands (Figure 1B and Figure 3A and B) and although the molecular weights of these bands were comparable to specific PDE8 splicing variants in previous studies,^{31–34} their identities as PDE8A and PDE8B isoforms should be further validated in future work.

Because PDE8 isoforms are important for the regulation of steroidogenesis,^{54–56} it remains to be determined whether selective PDE8B inhibition could be used for AF treatment without serious effects on adrenal gland or other organs.

Because of limited accessibility to human left atria at our centers, experiments were restricted to right atrial appendages. Different atrial regions may undergo distinct remodeling and further work should be directed to assess whether the observed changes in PDE8 in the right atria also apply to left atrial cells.

pAF patients had similar clinical parameters to SR patients except that they had more often hyperlipidemia and cardiopulmonary bypass surgery, were younger in some subgroups, and were more often taking digitalis, beta-blockers, and diuretics. cAF patients were more frequently treated with AT1-blockers, diuretics and amiodarone. The levels of PDE8B were higher in treated vs. non-treated SR patients with diuretics (see Supplementary data online, Figure S5). However, this was not observed at the protein level. Overall, controlling for individual comorbidities and medication did not alter the results of the individual sets of experiments.

A large number of atrial samples were collected, but each sample could only be used for a limited subset of experiments. Therefore, we performed extensive analyses to assess the influence of factors like patient age, sex, the influence of comorbidities and medications (see Supplementary data online, Figures S4–S8), we cannot fully exclude contributions from other unrecognized factors. The clinical profile of the patient subgroups used for biochemical experiments, FRET/patch-clamp, and multicellular AP recordings were comparable. The patient cohort used for qPCR and immunofluorescence showed differences in distribution of some clinical parameters; this needs to be considered when relating the differences that we noted in SR and AF patients to the other findings in our manuscript.

We observed an upregulation of PDE8A in pAF, but not cAF, and an upregulation of PDE8B in cAF, but not pAF. The underlying mechanisms for this differential regulation are still unclear. The increase of PDE8A in pAF could result from a specific atrial cardiomyopathy and might alter the regulation of targets other than $I_{Ca,L}$, although it might also constitute a transient event during the progression to persistent AF. Once persistent AF is established, the duration of the arrhythmia or/and the sustained high atrial rate could cause adaptive changes that may lead to a switch from PDE8A to PDE8B, along with a reduction in $I_{Ca,L}$, which is consistent with the selective increase in PDE8B in AF, but not in HF atria. Future work should directly address whether tachypacing upregulates atrial PDE8B.

Ser1928 is a well-established PKA phosphorylation site of Cav1.2_{α1C}³⁶ relevant for the trafficking of the channel to the membrane.⁴⁰ In agreement, we found that both ISO and PDE8 inhibition increased Ser1928 phosphorylation. However, the phosphorylation of this residue is not required for $I_{Ca,L}$ modulation in mouse ventricular myocytes and HEK293 cells.^{37–39} Previous works have identified Rad as the channel inhibitor responsible of the modulation of $I_{Ca,L}$ via PKA phosphorylation in mice,^{37,50} but the functional role of Ser1928 and Rad in HAMs and in AF has not been addressed here and warrants additional studies. Finally, we studied the effects of PDE8 on $I_{Ca,L}$ only. Whether and how PDE8 affects the function of other ionic currents that contribute to the atrial AP remains to be determined in future studies.

In summary, we establish PDE8B as a novel regulator of atrial Ca^{2+} handling and arrhythmogenesis and a new integral component of the LTCC macromolecular complex, which importantly contributes to the reentry-promoting reduction of $I_{Ca,L}$ in cAF.

Author contribution

Istvan Bacsko, Anne Garnier (Investigation: Supporting), Markus Kamler (Formal analysis: Supporting; Resources: Supporting), Renate B. Schnabel (Investigation: Supporting), Dobromir Dobrev (Formal analysis: Supporting; Funding acquisition: Supporting; Writing – original draft: Supporting), Cristina E. Molina, Ph.D. (Conceptualization: Lead; Data curation: Lead; Funding acquisition: Lead; Investigation:

Supporting; Methodology: Lead; Supervision: Lead; Writing – original draft: Lead), Hermann Reichenspurner (Resources: Supporting), Viacheslav O. Nikolaev (Conceptualization: Supporting; Data curation: Supporting), Niels Voigt (Conceptualization: Supporting), Kira Beneke (Investigation: Lead; Writing – original draft: Supporting), Franziska Reinhardt (Formal analysis: Supporting), Nefeli Grammatika Pavlidou (Investigation: Lead; Writing – original draft: Supporting), Shokoufeh Dobrev (Formal analysis: Supporting; Investigation: Supporting), Issam H. Abu-Taha, Constanze Schmidt (Investigation: Supporting), Simon Pecha (Resources: Equal), and Eric Jacquet (Investigation: Supporting)

Acknowledgements

We thank Claudine Deloménie (ACTAGen platform of UMS-IPSIT) for her expertise in molecular biology, Ramona Löcker and the Imaging Center Campus Essen—University of Duisburg-Essen for technical support.

Supplementary data

Supplementary data is available at *European Heart Journal* online.

Data and materials availability

All data are available in the manuscript or the [supplementary materials](#).

Conflict of interest

The authors have no competing interests. R.B.S. reports grants from European Union Horizon 2020, the German Ministry of Research and Education, ERACOSysMed3 and personal fees from BMS/Pfizer, outside this work.

Funding

This work was supported by the Deutsche Forschungsgemeinschaft (ES 569/2-1 to CM), the German Centre for Cardiovascular Research (DZHK), the National Institutes of Health (R01HL136389, R01HL131517, R01HL089598, and R01HL163277 to D.D.), the European Union (large-scale integrative project MEASTRIA, No. 965286 to D.D.), and the Hungarian National Research, Development and Innovation Office (TKP2021-EGA-32, K-128851 and GINOP-2.3.2-15-2016-00006).

References

- Larupa Santos J, Rodríguez I, Olesen MS, Hjorth Bentzen B, Schmitt N. Investigating gene-microRNA networks in atrial fibrillation patients with mitral valve regurgitation. *PLoS One* 2020;**15**:e0232719. <https://doi.org/10.1371/journal.pone.0232719>
- Nattel S, Heijman J, Zhou L, Dobrev D. Molecular basis of atrial fibrillation pathophysiology and therapy: a translational perspective. *Circ Res* 2020;**127**:51–72. <https://doi.org/10.1161/CIRCRESAHA.120.316363>
- Voigt N, Li N, Wang Q, Wang W, Trafford AW, Abu-Taha I, et al. Enhanced sarcoplasmic reticulum Ca²⁺ leak and increased Na⁺-Ca²⁺ exchanger function underlie delayed afterdepolarizations in patients with chronic atrial fibrillation. *Circulation* 2012;**125**:2059–2070. <https://doi.org/10.1161/CIRCULATIONAHA.111.067306>
- Llach A, Molina CE, Prat-Vidal C, Fernandes J, Casado V, Ciruela F, et al. Abnormal calcium handling in atrial fibrillation is linked to up-regulation of adenosine A_{2A} receptors. *Eur Heart J* 2011;**32**:721–729. <https://doi.org/10.1093/eurheartj/ehq464>
- Shanmugam M, Molina CE, Gao S, Severac-Bastide R, Fischmeister R, Babu GJ. Decreased sarcolipin protein expression and enhanced sarco(endo)plasmic reticulum Ca²⁺ uptake in human atrial fibrillation. *Biochem Biophys Res Commun* 2011;**410**:97–101. <https://doi.org/10.1016/j.bbrc.2011.05.113>
- Hove-Madsen L, Llach A, Bayes-Genis A, Roura S, Font ER, Aris A, et al. Atrial fibrillation is associated with increased spontaneous calcium release from the sarcoplasmic Reticulum in human atrial myocytes. *Circulation* 2004;**110**:1358–1363. <https://doi.org/10.1161/01.CIR.0000141296.59876.87>
- Vest JA, Wehrens XHT, Reiken SR, Lehnart SE, Dobrev D, Chandra P, et al. Defective cardiac ryanodine receptor regulation during atrial fibrillation. *Circulation* 2005;**111**:2025–2032. <https://doi.org/10.1161/01.CIR.0000162461.67140.4C>
- El-Armouche A, Boknik P, Eschenhagen T, Carrier L, Knaut M, Ravens U, et al. Molecular determinants of altered Ca²⁺ handling in human chronic atrial fibrillation. *Circulation* 2006;**114**:670–680. <https://doi.org/10.1161/CIRCULATIONAHA.106.636845>
- Molina CE, Leroy J, Richter W, Xie M, Scheitrum C, Lee IO, et al. Cyclic adenosine monophosphate phosphodiesterase type 4 protects against atrial arrhythmias. *J Am Coll Cardiol* 2012;**59**:2182–2190. <https://doi.org/10.1016/j.jacc.2012.01.060>
- Wagoner DRV, Pond AL, Lamorgese M, Rossie SS, McCarthy PM, Nerbonne JM. Atrial L-type Ca²⁺ currents and human atrial fibrillation. *Circ Res* 1999;**85**:428–436. <https://doi.org/10.1161/01.RES.85.5.428>
- Reinhardt F, Beneke K, Pavlidou NG, Conradi L, Reichenspurner H, Hove-Madsen L, et al. Abnormal calcium handling in atrial fibrillation is linked to changes in cyclic AMP dependent signaling. *Cells* 2021;**10**:3042. <https://doi.org/10.3390/cells10113042>
- Christ T, Boknik P, Wöhrl S, Wettwer E, Graf EM, Bosch RF, et al. L-Type Ca²⁺ current downregulation in chronic human atrial fibrillation is associated with increased activity of protein phosphatases. *Circulation* 2004;**110**:2651–2657. <https://doi.org/10.1161/01.CIR.0000145659.80212.6A>
- Mika D, Leroy J, Vandecasteele G, Fischmeister R. PDEs create local domains of cAMP signaling. *J Mol Cell Cardiol* 2012;**52**:323–329. <https://doi.org/10.1016/j.yjmcc.2011.08.016>
- Rochais F, Abi-Gerges A, Horner K, Lefebvre F, Cooper DMF, Conti M, et al. A specific pattern of phosphodiesterases controls the cAMP signals generated by different Gs-coupled receptors in adult rat ventricular myocytes. *Circ Res* 2006;**98**:1081–1088. <https://doi.org/10.1161/01.RES.0000218493.09370.8e>
- Molina CE, Johnson DM, Mehl H, Spätjens RLHMG, Mika D, Algalarrondo V, et al. Interventricular differences in β-adrenergic responses in the canine heart: role of phosphodiesterases. *J Am Heart Assoc* 2014;**3**:e000858. <https://doi.org/10.1161/JAHA.114.000858>
- Weishaar RE, Kobylarz-Singer DC, Steffen RP, Kaplan HR. Subclasses of cyclic AMP-specific phosphodiesterase in left ventricular muscle and their involvement in regulating myocardial contractility. *Circ Res* 1987;**61**:539–547. <https://doi.org/10.1161/01.RES.61.4.539>
- Lugnier C, Muller B, Bec AL, Beaudry C, Rousseau E. Characterization of indolandin- and rolipram-sensitive cyclic nucleotide phosphodiesterases in canine and human cardiac microsomal fractions. *J Pharmacol Exp Ther* 1993;**265**:1142–1151.
- Garnier A, Bork NI, Jacquet E, Zipfel S, Muñoz-Guijosa C, Baczkó I, et al. Mapping genetic changes in the cAMP-signaling cascade in human atria. *J Mol Cell Cardiol* 2021;**155**:10–20. <https://doi.org/10.1016/j.yjmcc.2021.02.006>
- Leroy J, Richter W, Mika D, Castro LRV, Abi-Gerges A, Xie M, et al. Phosphodiesterase 4B in the cardiac L-type Ca²⁺ channel complex regulates Ca²⁺ current and protects against ventricular arrhythmias in mice. *J Clin Invest* 2011;**121**:2651–2661. <https://doi.org/10.1172/JCI44747>
- Lehnart SE, Wehrens XHT, Reiken S, Warrior S, Belevych AE, Harvey RD, et al. Phosphodiesterase 4D deficiency in the ryanodine-receptor Complex promotes heart failure and arrhythmias. *Cell* 2005;**123**:25–35. <https://doi.org/10.1016/j.cell.2005.07.030>
- Sprenger JU, Perera RK, Steinbrecher JH, Lehnart SE, Maier LS, Hasenfuss G, et al. In vivo model with targeted cAMP biosensor reveals changes in receptor-microdomain communication in cardiac disease. *Nat Commun* 2015;**6**:6965. <https://doi.org/10.1038/ncomms7965>
- Berisha F, Götz KR, Wegener JW, Brandenburg S, Subramanian H, Molina CE, et al. cAMP imaging at ryanodine receptors reveals β₂-adrenoceptor driven arrhythmias. *Circ Res* 2021;**129**:81–94. <https://doi.org/10.1161/CIRCRESAHA.120.318234>
- Rozmaritsa N, Christ T, Wagoner DRV, Haase H, Stasch JP, Matschke K, et al. Attenuated response of L-type calcium current to nitric oxide in atrial fibrillation. *Cardiovasc Res* 2014;**101**:533–542. <https://doi.org/10.1093/cvr/cvt334>
- Fisher DA, Smith JF, Pillar JS, Denis SHS, Cheng JB. Isolation and characterization of PDE8A, a novel human cAMP-specific phosphodiesterase. *Biochem Biophys Res Commun* 1998;**246**:570–577. <https://doi.org/10.1006/bbrc.1998.8684>
- Soderling SH, Bayuga SJ, Beavo JA. Cloning and characterization of a cAMP-specific cyclic nucleotide phosphodiesterase. *Proc Natl Acad Sci U S A* 1998;**95**:8991–8996. <https://doi.org/10.1073/pnas.95.15.8991>
- Hayashi M, Matsushima K, Ohashi H, Tsunoda H, Murase S, Kawarada Y, et al. Molecular cloning and characterization of human PDE8B, a novel thyroid-specific isozyme of 3',5'-cyclic nucleotide phosphodiesterase. *Biochem Biophys Res Commun* 1998;**250**:751–756. <https://doi.org/10.1006/bbrc.1998.9379>
- Patrucco E, Albergine MS, Santana LF, Beavo JA. Phosphodiesterase 8A (PDE8A) regulates excitation-contraction coupling in ventricular myocytes. *J Mol Cell Cardiol* 2010;**49**:330–333. <https://doi.org/10.1016/j.yjmcc.2010.03.016>
- Molina CE, Jacquet E, Ponien P, Muñoz-Guijosa C, Baczkó I, Maier LS, et al. Identification of optimal reference genes for transcriptomic analyses in normal and diseased human heart. *Cardiovasc Res* 2018;**114**:247–258. <https://doi.org/10.1093/cvr/cvx182>
- Perera RK, Sprenger JU, Steinbrecher JH, Hübscher D, Lehnart SE, Abesser M, et al. Microdomain switch of cGMP-regulated phosphodiesterases leads to ANP-induced

- augmentation of β -adrenoceptor-stimulated contractility in early cardiac hypertrophy. *Circ Res* 2015;**116**:1304–1311. <https://doi.org/10.1161/CIRCRESAHA.116.306082>
30. Wettwer E, Håla O, Christ T, Heubach JF, Dobrev D, Knaut M, et al. Role of IKur in controlling action potential shape and contractility in the human atrium: influence of chronic atrial fibrillation. *Circulation* 2004;**110**:2299–2306. <https://doi.org/10.1161/01.CIR.0000145155.60288.71>
 31. Wang P, Wu P, Egan RW, Billah MM. Human phosphodiesterase 8A splice variants: cloning, gene organization, and tissue distribution. *Gene* 2001;**280**:183–194. [https://doi.org/10.1016/S0378-1119\(01\)00783-1](https://doi.org/10.1016/S0378-1119(01)00783-1)
 32. Hayashi M, Shimada Y, Nishimura Y, Hama T, Tanaka T. Genomic organization, chromosomal localization, and alternative splicing of the human phosphodiesterase 8B gene. *Biochem Biophys Res Commun* 2002;**297**:1253–1258. [https://doi.org/10.1016/S0006-291X\(02\)02371-9](https://doi.org/10.1016/S0006-291X(02)02371-9)
 33. Sasseville M, Albus FK, Côté N, Guillemette C, Gilchrist RB, Richard FJ. Characterization of novel phosphodiesterases in the bovine ovarian follicle1. *Biol Reprod* 2009;**81**:415–425. <https://doi.org/10.1095/biolreprod.108.074450>
 34. Kobayashi T, Gamanuma M, Sasaki T, Yamashita Y, Yuasa K, Kotera J, et al. Molecular comparison of rat cyclic nucleotide phosphodiesterase 8 family: unique expression of PDE8B in rat brain. *Gene* 2003;**319**:21–31. [https://doi.org/10.1016/S0378-1119\(03\)00809-6](https://doi.org/10.1016/S0378-1119(03)00809-6)
 35. Brandenburg S, Pawlowitz J, Steckmeister V, Subramanian H, Uhlenkamp D, Scardigli M, et al. A junctional cAMP compartment regulates rapid Ca²⁺ signaling in atrial myocytes. *J Mol Cell Cardiol* 2022;**165**:141–157. <https://doi.org/10.1016/j.yjmcc.2022.01.003>
 36. Hulme JT, Westenbroek RE, Scheuer T, Catterall WA. Phosphorylation of serine 1928 in the distal C-terminal domain of cardiac CaV1.2 channels during beta1-adrenergic regulation. *Proc Natl Acad Sci U S A* 2006;**103**:16574–16579. <https://doi.org/10.1073/pnas.0607294103>
 37. Liu G, Papa A, Katchman AN, Zakharov SI, Roybal D, Hennessey JA, et al. Mechanism of adrenergic CaV1.2 stimulation revealed by proximity proteomics. *Nature* 2020;**577**:695–700. <https://doi.org/10.1038/s41586-020-1947-z>
 38. Yang L, Katchman A, Kushner J, Kushnir A, Zakharov SI, Chen BX, et al. Cardiac CaV1.2 channels require β subunits for β -adrenergic-mediated modulation but not trafficking. *J Clin Invest* 2019;**129**:647–658. <https://doi.org/10.1172/JCI123878>
 39. Lemke T, Welling A, Christel CJ, Blaich A, Bernhard D, Lenhardt P, et al. Unchanged beta-adrenergic stimulation of cardiac L-type calcium channels in ca v 1.2 phosphorylation site S1928A mutant mice. *J Biol Chem* 2008;**283**:34738–34744. <https://doi.org/10.1074/jbc.M804981200>
 40. Ito DW, Hannigan KI, Ghosh D, Xu B, Villar SGD, Xiang YK, et al. β -adrenergic-mediated dynamic augmentation of sarcolemmal CaV 1.2 clustering and co-operativity in ventricular myocytes. *J Physiol* 2019;**597**:2139–2162. <https://doi.org/10.1113/jp277283>
 41. Schotten U, Haase H, Frechen D, Greiser M, Stellbrink C, Vazquez-Jimenez JF, et al. The L-type Ca²⁺-channel subunits alpha1C and beta2 are not downregulated in atrial myocardium of patients with chronic atrial fibrillation. *J Mol Cell Cardiol* 2003;**35**:437–443. [https://doi.org/10.1016/S0022-2828\(03\)00012-9](https://doi.org/10.1016/S0022-2828(03)00012-9)
 42. Molina CE, Abu-Taha IH, Wang Q, Roselló-Díez E, Kamler M, Nattel S, et al. Profibrotic, electrical, and calcium-handling remodeling of the atria in heart failure patients with and without atrial fibrillation. *Front Physiol* 2018;**9**:1383. <https://doi.org/10.3389/fphys.2018.01383>
 43. Llach A, Molina CE, Fernandes J, Padró J, Cinca J, Hove-Madsen L. Sarcoplasmic reticulum and L-type Ca²⁺ channel activity regulate the beat-to-beat stability of calcium handling in human atrial myocytes. *J Physiol* 2011;**589**:3247–3262. <https://doi.org/10.1113/jphysiol.2010.197715>
 44. Dinanian S, Boixel C, Juin C, Hulot JS, Coulombe A, Rücker-Martin C, et al. Downregulation of the calcium current in human right atrial myocytes from patients in sinus rhythm but with a high risk of atrial fibrillation. *Eu Heart J* 2008;**29**:1190–1197. <https://doi.org/10.1093/eurheartj/ehh140>
 45. Nattel S, Li D, Yue L. Basic mechanisms of atrial fibrillation—very new insights into very old ideas. *Annu Rev Physiol* 2000;**62**:51–77. <https://doi.org/10.1146/annurev.physiol.62.1.51>
 46. Schotten U, Ausma J, Stellbrink C, Sabatschus I, Vogel M, Frechen D, et al. Cellular mechanisms of depressed atrial contractility in patients with chronic atrial fibrillation. *Circulation* 2001;**103**:691–698. <https://doi.org/10.1161/01.CIR.103.5.691>
 47. Grammer JB, Zeng X, Bosch RF, Köhlkamp V. Atrial L-type Ca²⁺-channel, β -adrenoreceptor, and 5-hydroxytryptamine type 4 receptor mRNAs in human atrial fibrillation. *Basic Res Cardiol* 2001;**96**:82–90. <https://doi.org/10.1007/s003950170081>
 48. Glukhov AV, Balycheva M, Sanchez-Alonso JL, Ilkan Z, Alvarez-Laviada A, Bhogal N, et al. Direct evidence for microdomain-specific localization and remodeling of functional L-type calcium channels in rat and human atrial myocytes. *Circulation* 2015;**132**:2372–2384. <https://doi.org/10.1161/CIRCULATIONAHA.115.018131>
 49. Qi XY, Yeh YH, Xiao L, Burstein B, Maguy A, Chartier D, et al. Cellular signaling underlying atrial tachycardia remodeling of L-type calcium current. *Circ Res* 2008;**103**:845–854. <https://doi.org/10.1161/CIRCRESAHA.108.175463>
 50. Finlin BS, Crump SM, Satin J, Andres DA. Regulation of voltage-gated calcium channel activity by the rem and rad GTPases. *Proc Natl Acad Sci U S A* 2003;**100**:14469–14474. <https://doi.org/10.1073/pnas.2437756100>
 51. Furberg CD, Psaty BM, Meyer JV. Nifedipine. Dose-related increase in mortality in patients with coronary heart disease. *Circulation* 1995;**92**:1326–1331. <https://doi.org/10.1161/01.CIR.92.5.1326>
 52. Multicenter Diltiazem Postinfarction Trial Research Group. The effect of diltiazem on mortality and reinfarction after myocardial infarction. *N Engl J Med* 1988;**319**:385–392. <https://doi.org/10.1056/NEJM198808183190701>
 53. Voigt N, Heijman J, Wang Q, Chiang DY, Li N, Karck M, et al. Cellular and molecular mechanisms of atrial arrhythmogenesis in patients with paroxysmal atrial fibrillation. *Circulation* 2014;**129**:145–156. <https://doi.org/10.1161/CIRCULATIONAHA.113.006641>
 54. Shimizu-Albergine M, Tsai LCL, Patrucco E, Beavo JA. cAMP-specific phosphodiesterases 8A and 8B, essential regulators of Leydig cell steroidogenesis. *Mol Pharmacol* 2012;**81**:556–566. <https://doi.org/10.1124/mol.111.076125>
 55. Tsai LCL, Shimizu-Albergine M, Beavo JA. The high-affinity cAMP-specific phosphodiesterase 8B controls steroidogenesis in the mouse adrenal gland. *Mol Pharmacol* 2011;**79**:639–648. <https://doi.org/10.1124/mol.110.069104>
 56. Brown KM, Lee LCY, Findlay JE, Day JP, Baillie GS. Cyclic AMP-specific phosphodiesterase, PDE8A1, is activated by protein kinase A-mediated phosphorylation. *FEBS Lett* 2012;**586**:1631–1637. <https://doi.org/10.1016/j.febslet.2012.04.033>