

## RESISTING VERSUS ASSISTING THE ANKLE MUSCULATURE HAS AN UNIDIRECTIONAL EFFECT ON WALK-TO-RUN TRANSITION SPEED

Philippe Malcolm, Veerle Segers, Ine Van Caekenberghe, Dirk De Clercq

Department of Movement and Sports Sciences, Ghent University,

email: [philippe.malcolm@ugent.be](mailto:philippe.malcolm@ugent.be)

### INTRODUCTION

Various descriptive studies point towards the dorsiflexor or the plantarflexor muscles as determinants for the walk-to-run transition (WRT). It is generally accepted that an experimental design is the next step after descriptive analysis towards finding the underlying (pool of) determinant(s). Until now there is only one such study in which specifically the dorsiflexors were weakened by a fatigue protocol and indeed a lower WRT speed was found [1]. An even more critical test would be to check if there is an increase in WRT speed after experimentally assisting potential determinants. In order to accomplish this goal a powered exoskeleton was developed because this is the only method that allows to provide direct and focused assistance to the dorsiflexor or plantarflexor muscles during the WRT.

### METHODS

In two separate experiments respectively 8 and 11 young adult female subjects performed several WRT's on an accelerating treadmill while wearing a bilateral ankle-foot-exoskeleton. The ankle musculature was assisted or resisted by pneumatic muscles mounted on the exoskeleton (figure 1). In the dorsiflexor experiment the pneumatic muscles were activated around heel contact i.e. when the endogenous dorsiflexors are highly activated. The conditions were: dorsiflexion resist, control and dorsiflexion assist [2]. In the plantarflexor experiment the pneumatic muscles were activated during the push off phase. The conditions were: plantarflexion resist, control and plantarflexion assist. WRT-speed was measured as the average treadmill belt velocity during the transition step. Differences between conditions within each experiment were analyzed by means of a Wilcoxon non parametric analysis.

### RESULTS AND DISCUSSION

WRT speed was significantly lower in the resist condition than in the corresponding assist ( $p \leq 0.006$ ) and control condition ( $p \leq 0.002$ ) in the dorsiflexor experiment as well as in the plantarflexor experiment. Also in both experiments there was no significant difference in WRT-speed between the assist and control condition ( $p \geq 0.132$ ). (table 1)


	WRT speed (m.s <sup>-1</sup> )	p		test
<b>dorsiflexor experiment</b>				
resist	2.06 ± 0.09	0.001	▪▪	resist-control
control	2.10 ± 0.10	0.006	**	resist-assist
assist	2.12 ± 0.11	0.132		assist-control
<b>plantarflexor experiment</b>				
resist	1.93 ± 0.15	0.002	**	resist-control
control	2.11 ± 0.21	0.002	**	resist-assist
assist	2.09 ± 0.17	0.722		assist-control

**Table 1:** WRT speed in the resist, control and assist condition in the dorsiflexor and plantarflexor experiment.

\*\*= $p < 0.01$       ▪▪= $p < 0.01$  after elimination of one outlier

### CONCLUSIONS

The fact that the effect from the resist condition cannot be extrapolated in the opposite direction is a striking analogy between both experiments. A possible explanation is that it would be easy to lower WRT speed by sufficiently resisting specific muscle groups whereas when assisting the same muscle group some presently unknown weak link (figure 2) prevents a substantial increase in WRT speed. These results differ from a study from Bartlett et al. [3] who did observe significant increases in transition speed. This could be due to the fact that the experimental conditions had a more general effect whereas in the present study the effects would have been more focused in space and time.




**Figure 2:** Weak link concept: a=control, b=resist, c=assist  
D1= manipulated determinant    D2, D3=other determinants

### ACKNOWLEDGEMENTS

The authors wish to thank Technische Orthopedie België ([www.orthoshop.be](http://www.orthoshop.be)) and Festo for their technical support.

### REFERENCES

1. Segers V, et al., *Gait Posture*; **25**: 639-47, 2007.
2. Malcolm P, et al., *Gait Posture*; **29**: 6-10, 2009.
3. Bartlett JL, et al., *J Exp Biol*; **211**: 1281-8, 2008.



**Figure 1:** Experimental conditions (the difference between the dorsiflexor and plantarflexor experiment consists in the timing of the pneumatic muscle activation)