

Geophysical Research Abstracts
Vol. 14, EGU2012-4391-2, 2012
EGU General Assembly 2012
© Author(s) 2012



Transport of root-derived CO₂ via the transpiration stream affects aboveground tree physiology

J. Bloemen (1), M.A. McGuire (2), D.P. Aubrey (2), R.O. Teskey (2), and K. Steppe (1)

(1) Laboratory of Plant Ecology, Ghent University, Coupure links 653, 9000 Gent, Belgium (jasper.bloemen@ugent.be), (2) Warnell school of Forestry and Natural Resources, University of Georgia, 180 East Green St, Athens, GA, 30602-2152, USA

Recent research on soil CO₂ efflux has shown that belowground autotrophic respiration is largely underestimated using classical net CO₂ flux measurements. Aubrey & Teskey (2009) found that in forest ecosystems a substantial portion of the CO₂ released from root respiration remained within the root system and was transported aboveground in the stem via the transpiration stream. The magnitude of this upward movement of CO₂ from belowground tissues suggested important implications for how we measure above- and belowground respiration. If a considerable fraction of root-respired CO₂ is transported aboveground, where it might be fixed in woody and leaf tissues, then we are routinely underestimating the amount of C needed to sustain belowground tissues.

In this study, we infused ¹³C labeled water into the base of field-grown poplar trees as a surrogate for root-respired CO₂ to investigate the possible role of root-derived CO₂ as substrate for carbon fixation. The label was transported upwards from the base of the tree toward the top. During its ascent, the ¹³C label was removed from the transpiration stream and fixed by chlorophyll-containing woody (young bark and xylem) and leaf (petiole) tissues. Moreover, based on ¹³C analysis of gas samples, we observed that up to 88 ± 0.10 % of the label applied was lost to the atmosphere by stem and branch efflux higher in the trees. Given that one-half of root-respired CO₂ may follow this internal flux pathway (Aubrey & Teskey, 2009), we calculated that up to 44% of the root-respired CO₂ could diffuse to the atmosphere once transported to the stem and branches. Thus, a large portion of CO₂ that diffuses out of aboveground tissues may actually result from root respiration.

Our results show that CO₂ originating belowground can be transported internally to aboveground parts of trees, where it will have an important impact on tree physiology. Internal transport of CO₂ indicates that the gas exchange approach to estimating above- and belowground autotrophic respiration is inadequate. Accurate quantification of this internal carbon flux is necessary to understand plant physiological mechanisms and to explain variations in above- and belowground respiratory patterns, but these results do not imply the necessity for a reevaluation of net CO₂ flux at the ecosystem level.

Reference:

Aubrey DP, Teskey RO (2009) Root-derived CO₂ efflux via xylem stream rivals soil CO₂ efflux. *New Phytologist* 184: 35-40.