

First ihsig International symposium, March 28-29,2012 – Ghent, Belgium
**The mycotoxin deoxynivalenol predisposes for the development
of necrotic enteritis in broilers**

Gunther Antonissen^{1,2}, Siska Croubels², Frank Pasmans¹, Richard Ducatelle¹, Freddy Haesebrouck¹, Leen Timbermont¹, Marc Verlinden¹, Jeroen Dewulf³, Mia Eeckhout⁴, Sarah De Saeger⁵, Evelyne Delezie⁶, An Martel¹, Filip Van Immerseel¹

1) Department of Pathology, Bacteriology and Poultry Diseases, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, 9820 Merelbeke, Belgium 2) Department of Pharmacology, Toxicology and Biochemistry, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, 9820 Merelbeke, Belgium 3) Department of Reproduction, Obstetrics and Herd Health, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, 9820 Merelbeke 4) Department of Food Science and Technology, Faculty of Biosciences and Landscape Architecture, Ghent University, Schoonmeerstraat 52, 9000 Ghent, Belgium 5) Department of Bio-analysis, Faculty of Pharmaceutical Sciences, Ghent University, Harelbekestraat 72, 9000 Ghent, Belgium 6) Institute for Agricultural and Fisheries Research (ILVO) Animal Sciences Unit, Scheldeweg 68, 9090 Melle, Belgium

E-mail: Gunther.Antonissen@UGent.be

Background

Subclinical necrotic enteritis (NE) is an economically important enteric disease caused by a netB producing, Gram-positive, anaerobic bacterium, *Clostridium perfringens*. Massive intestinal proliferation of *C. perfringens* is promoted by increased availability of free amino acids. The *Fusarium* mycotoxin deoxynivalenol (DON) is a common feed contaminant with a maximum guidance level set at 5000 µg/kg feed (2006/576/EC) and may damage intestinal epithelial cells and/or their intercellular junctions, subsequently inducing protein leakage (Girish and Smith, 2008) and may thus predispose to the development of NE.

Methods

A highly reproducible *in vivo* infection model mimicking subclinical NE was used (Gholamiandehkordi et al., 2007). A total of 360 one-day-old Ross 308 broilers from a commercial hatchery were randomly divided into four groups of three replicates with 30 birds per replica. All birds were fed a starter diet during the first eight days of the experiment, subsequently a grower diet for eight days, followed by a finisher diet during the remaining days. Throughout the entire experiment, group 1 and 4 received a blank diet while group 2 and 3 received an experimentally contaminated diet with DON. All birds in group 1 and 2 were challenged orally with one ml of a culture of *C. perfringens* strain 56 containing approximately 4×10^8 cfu/ml for four consecutive days starting at day 17. The remaining groups received sterile medium orally. The contamination level of DON and other mycotoxins was assayed using a validated multi-mycotoxin LC-MS/MS method (Monbaliu et al., 2010).

Results

The blank feed contained DON at 75 ± 22 µg/kg (starter), 83 ± 24 µg/kg (grower) and 100 ± 29 µg/kg (finisher). The contaminated feed contained DON at 3761 ± 1100 µg/kg (starter), 4281 ± 1300 µg/kg (grower) and 4384 ± 1300 µg/kg (finisher). At 1, 2 or 3 days after the final challenge with *C. perfringens*, chickens were euthanized and scored macroscopically for intestinal NE lesions. Chickens that received DON and *C. perfringens* had significantly ($\alpha=0.05$, $P<0.001$) more lesions than chickens that received only *C. perfringens*, with 46.6% and 19.5% of chickens positive for NE lesions, respectively. No NE lesions were seen in the groups receiving no *C. perfringens* inoculations.

Conclusion

Feeding DON contaminated feed in concentrations lower than the maximum guidance contamination level of 5000 µg/kg to broilers is a predisposing factor for the development of NE.

References

- Gholamiandehkordi, A.R.; Timbermont, L.; Lanckriet, A.; Van Den Broeck, W.; Pedersen, K.; Dewulf, J.; Pasmans, F.; Haesebrouck, F.; Ducatelle, R.; Van Immerseel F., 2007. Quantification of gut lesions in a subclinical necrotic enteritis model. *Avian Pathology* 36, 375-382.
- Girish, C.K.; Smith, T.K., 2008. Impact of feed-borne mycotoxins on avian cell-mediated and humoral immune responses. *World Mycotoxin Journal*, 1, 105-121.
- Monbaliu S.; Van Poucke C.; Detavernier C.; Dumoulin F.; Van de Velde F.; Schoeters E.; Van Dyck S.; Averkiewa A.; Van Peteghem C.; De Saeger S., 2010. Occurrence of mycotoxins in feed as analyzed by a multi-mycotoxin LC-MS/MS method. *Journal of Agricultural and Food Chemistry* 58, 66-71.

Acknowledgements

This work was financially supported by BIOMIN GmbH, Austria.