

Speciation-analysis for Recycling of Critical Elements in Aqueous Waste Streams

Karel FOLENS¹, Frank VANHAECKE² and Gijs DU LAING¹

¹Laboratory of Analytical Chemistry and Applied Ecochemistry, Faculty of Bioscience Engineering, Ghent University, Coupure Links 653, 9000 Ghent, Belgium, URL: ecochem.ugent.be, E-mail: Karel.Folens@UGent.be

²Department of Analytical Chemistry, Faculty of Sciences, Ghent University, Krijgslaan 281 S12, 9000 Ghent, Belgium, URL: analchem.ugent.be





Presence of Critical Elements in Waste Streams

Limited availability of inorganic commodities is one of the largest challenges for a future sustainable material use as set by European Commission. Growing life standards, demand for mobility and electronic devices require more and more specialty materials. Combining this economic importance and their supply risk, a class of critical materials is defined.

Platinum Group Metals

PGM's include platinum (Pt), rhodium (Rh), palladium (Pd), iridium (Ir), ruthetium (Ru) and osmium (Os). They possess unique properties, making them indispensable in autocatalysts, jewelry, electrical and electronical applications and industrial catalysts. PGM prices, with Pt being the far most important, have seriously risen in the last years (*Table 1*).

Table 1. Overview of PGM prices and the current fracition of recycling. ^[1]

ELEMENT	PRICE (€ kg⁻¹)	CURRENT RECYCLED FRACTION (%)
Platinum	38,240	24
Rhodium	26,520	26
Palladium	21,500	26
Irridium	14,940	25-50
Osmium	9,790	0
Ruthenium	1,850	10-25

Presence

of Critical Elements in Waste Streams

Rare Earth Elements

REE's comprehend 17 elements with a large chemical similarity, essential in i.a. permanent magnets and related 'clean energy' applications such as wind turbines, rechargeable batteries and electric vehicles. Tight export quota set by the world's largest exporting country of REE's, China, hardly meet this rising demand.^[2]



Figure 1. Applications which involve REE's with their relative fraction of overall REE use. [3]

Tahlo 2	Main	elemental	com	nosition	of RE	F hase	dan	nlication	c
uble z	• IVIUIII	elementui	COM	position	UJ NE	E DUSE	ւ սր	pilcution	5.

APPLICATION	ELEMENTS
Permanent magnets	Nd, Pr, Dy, Gd
Catalysts	Ce, La, Nd, Pr
Polishing	Ce, La, Pr
Rechargeable batteries	La, Ce, Nd, Pr, Sm
Steel and metal alloys	Ce, La, Nd, Pr
Phosphors	Y, Ce, La, Eu, Tb, Gd
Ceramics	Y, La, Ce, Nd, Pr
Other	Ce, La, Y, Nd, Pr, Sm, Gd

Focus on Recycling Strategies

- + Along with political and strategic considerations, alternative sources for these materials are considered vital. Having a non-geographically bounded supply, recycling and "urban mining" have strong economic advantages besides primary mining.
- + Efforts and possibilities in processing solid waste streams of critical elements is growing. [4] However, scientific and technical expertise are missing effective strategies for liquid media. Although some PGM's and REE's are toxic, absence of discharge limits in the current legislation certainly contributes to this. ^[5]

Focus on Recycling Strategies

+ The ongoing research focusses on developing strategies to recover and recuperate valuable elements in wastewaters. The value of potential recuperation is considerable to overcome operational costs. (Bio)medical use of platinum for example, extends to 8290 ton of platinum demand, representing a worldwide (input) value of 255 million EUR.



Speciation-analysis as Analytical tool

Where quantification reveals information on the scale, chemical determination guides towards well-aimed recovery strategies. Characterisation by speciation-analysis is key to fully understand the target's nature. However strictly inorganic, these elements are associated with ligands and functionalities, especially considering their matrix.

- + A screening for platinum content in several waste water was performed. Microfiltration was used to distinguish between soluble or particulate matter.
- + In order to overcome challenging detection limits, Solid Phase Extraction was evaluated for the pre-concentration of species (*Figure 2*).
- + Further techniques, such as **Reversed Phase HPLC-ICP-QMS** differentiate between species of select elements. The originally applied versus degraded products could be revealed.



Figure 2. Recovery profile of platinum chloride species in Solid Phase Extraction (SPE) experiment. Initial loading of 27.1 ng Pt(IV) and 65.4 ng *Pt(II) were eluted with 0.1 M thiourea, 0.1 M HCl at 60 ° C. Adsorption* efficencies of 98% for Pt(IV) and 94% for Pt(II) were obtained.

Speciation as Analytical tool

Outlook

Based hereupon, different removal techniques such as adsorption were tested at lab scale.



Outlook

Clarify the current distribution of critical elements in the environment for Flanders.

- Study the influence of urine and wastewater matrix on the removal process.
- + Further **optimisation** of chromatographic conditions for speciation of platinum chloride complexes.

References

- 1. Johnson Matthey (2014) Platinum Today, *platinum.matthey.com*
- 2. K. Binnemans, P.T. Jones, B. Blanpain, T. Van Gerven, Y. Yang, A. Walton and M. Buchert (2013) Journal of Cleaner Production 51, 1-22.
- 3. L. Tercero Espinoza (2014) CRM_InnoNet Substitution of Critical Raw Materials, European Commission.
- 4. A. Dubiella-Jackowska, Z. Polkowska and J. Namiesnik (2013) Polish Journal of Enivironmental *Studies* **16**, 17.
- 5. F. A. Chmilenko and S. N. Khudyakova (2013) Journal of Analytical Chemistry 68, 409-416.

— Pt(IV) ··O·· Pt(II)







FACULTY OF BIOSCIENCE ENGINEERING