



biblio.ugent.be

The UGent Institutional Repository is the electronic archiving and dissemination platform for all UGent research publications. Ghent University has implemented a mandate stipulating that all academic publications of UGent researchers should be deposited and archived in this repository. Except for items where current copyright restrictions apply, these papers are available in Open Access.

This item is the archived peer-reviewed author-version of:

Title: Dietary mycotoxins, co-exposure, and carcinogenesis in humans: short review.

Authors: De Ruyck K., Boevre M., Huybrechts I., De Saeger S.

In: Mutation Research-Reviews in Mutation Research, 766, 32-41, 2015

To refer to or to cite this work, please use the citation to the published version:

De Ruyck K., Boevre M., Huybrechts I., De Saeger S. (2015). Dietary mycotoxins, co-exposure, and carcinogenesis in humans: short review. Mutation Research-Reviews in Mutation Research, 766, 32-41. DOI:10.1016/j.mrrev.2015.07.003

Dietary mycotoxins, co-exposure, and carcinogenesis in humans: short review.

Karl De Ruyck¹, Marthe De Boevre¹, Inge Huybrechts², and Sarah De Saeger¹

¹ *Laboratory of Food Analysis, Faculty of Pharmaceutical Sciences, Ghent University, Ottergemsesteenweg 460, 9000 Ghent, Belgium*

² *International Agency for Research on Cancer, Dietary Exposure Assessment Group, 150 Cours Albert Thomas, 69008 Lyon, France*

karl.deruyck@ugent.be; marthe.deboevre@ugent.be; huybrechtsi@iarch.fr; sarah.desaeger@ugent.be

Corresponding author: Karl De Ruyck; +32 4 97 128 377; karl.deruyck@ugent.be

Abstract

Mycotoxins, toxic secondary metabolites of fungi, affect global agriculture so prolifically that they are virtually ubiquitous at some concentration in the average human diet. Studies of *in vitro* and *in vivo* toxicity are discussed, leading to investigations of co-exposed mycotoxins, as well as carcinogenic effects. Some of the most common and toxicologically significant mycotoxins, such as the aflatoxins, ochratoxins, fumonisins, trichothecenes, and zearalenone, are outlined. The wide variety of pathogenic mechanisms these compounds employ are shown capable of inducing a complex set of interactions. Of particular note are potential synergisms between mycotoxins with regard to carcinogenic attributable risk, indicating an important field for future study.

Keywords: mycotoxins; carcinogenesis; mycotoxin co-exposure; mycotoxin carcinogenicity; colorectal cancer; hepatocellular cancer

Introduction

Fungi have been observed for millennia, and are found to be relatively ubiquitous in nature, with

spores able to travel vast distances across the surface of the planet (Hirst & Stedman 1967). Many important agricultural products, especially those rich in carbohydrates, are attractive colonization sites for fungi. Some toxic secondary metabolites of fungal growth are identified as mycotoxins, and may be found to contaminate agricultural products (Chelkowski 1998).

Many species from the *Alternaria*, *Aspergillus*, *Claviceps*, and *Fusarium* genera, as well as some *Penicillium* species, and several others are known to produce mycotoxins (Zain 2011). Mycotoxins are small organic molecules produced as secondary metabolites of fungal growth, observed as toxic to animals and humans who consume them. Mycotoxicosis is the term used for poisoning associated with exposures to mycotoxins. The symptoms of mycotoxicosis depend on the type of mycotoxin, the concentration and duration of exposure, as well as age, health, and sex of the exposed individual (Bennett & Klich 2003).

Like all toxicological syndromes, mycotoxicoses can be categorized as acute or chronic. Acute toxicity generally has a rapid onset and an obvious toxic response, while chronic toxicity is characterized by low dose exposure over a long time period, resulting in cancers and other generally irreversible effects (Williams et al. 2015). Although the main human and veterinary health burdens of mycotoxin exposure are related to chronic exposure (e.g., cancer, kidney damage, immune suppression), the best-known mycotoxin episodes are manifestations of acute effects (e.g., Turkey X-syndrome, human ergotism, stachybotryotoxicosis in livestock) (Bennett & Klich 2003).

Some of the most frequently encountered mycotoxins ochratoxin A (OTA) and deoxynivalenol (DON), are reported to interfere with mammalian cellular processes including DNA replication and protein synthesis (Bensassi et al. 2009; Pfohl-Leszkowicz & Manderville 2012). Other mycotoxins, particularly aflatoxin B₁ (AFB₁) and its metabolic precursor sterigmatocystin, have been identified as carcinogenic by the World Health Organization's (WHO) International Agency for Research on Cancer (IARC) Monographs Program (International Agency for Research on Cancer 2007). The IARC Monographs identify environmental factors associated by scientific literature with increased risk of carcinogenesis in humans.

As a result of crops affected by fungal infection being eaten, either directly by humans or as feed for livestock, mycotoxins are introduced to the food chain. Mycotoxins are able to resist decomposition or being broken down by mammalian digestion, even by ruminant livestock, allowing

these compounds to persist in meat and even dairy products (Kang'ethe & Lang'a 2009). This gives rise to certain partially metabolized mycotoxins, such as aflatoxin M₁ (AFM₁), present in milk from cows or humans that consumed feed or food contaminated by aflatoxins. Even temperature treatments, such as cooking and freezing, do not inactivate some mycotoxins.

Due to the broad, overlapping habitats of fungal species, it is observed that multiple species may affect a given region (Alkadri et al. 2014; Jackson et al. 2012). Furthermore, one fungal species may produce many different mycotoxins, and the same mycotoxin may also be produced by several different species. Since many species are each capable of producing multiple mycotoxin compounds, and agricultural products from many sources may be aggregated prior to processing *en masse*, there is a very high likelihood of multiple mycotoxins co-occurring in food and feed products (Mngadi et al. 2008). Consequently, in the richly varied modern diet, individuals undergo dietary exposure to a very wide variety of mycotoxins (Abia et al. 2013; Solfrizzo et al. 2014).

Various mycotoxins were investigated since early last century for individual toxicity, usually in regard to acute pathologies (Schreiner & Reed 1908). However, recent research is indicating complex interactions between chronic co-exposure to multiple mycotoxins possibly having synergistic, or even mitigating effects (Bensassi et al. 2014; Grenier & Oswald 2011). Some of the most common and toxicologically significant mycotoxins, such as the aflatoxins, ochratoxins, fumonisins, trichothecenes, and zearalenone, are outlined below, followed by a description of possible health burdens caused by co-occurrence and co-exposure of multiple mycotoxins, as well as their carcinogenic effects.

Mycotoxins

Aflatoxins

There are six predominant aflatoxins, named AFB₁, aflatoxin B₂ (AFB₂), aflatoxin G₁ (AFG₁), aflatoxin G₂ (AFG₂), AFM₁, and aflatoxin M₂ (AFM₂) (Reddy et al. 2010). In this nomenclature, 'B' and 'G' are used to denote compounds that fluoresce blue or green, respectively, under ultraviolet light. The 'M₁' and 'M₂' compounds are not found on cereal products themselves, but are metabolites expressed in milk of mammals whose diet was contaminated by aflatoxins B₁ and B₂, respectively (Garrido et al. 2003). Finally, the '2' numbered aflatoxins are structural isomers missing one double bond, as compared to the respective '1' numbered molecule, illustrated in **figure 1** below. Among the *Aspergillus* genus, species in the section *Flavi* are most frequently reported producers of AFB₁, while

A. flavus and *A. parasiticus* are commonly used as producers of AFB₁, AFB₂, AFG₁ and AFG₂ for morphological studies (Klich 2007; Varga et al. 2011).

AFB₁ is identified as the most potent naturally occurring carcinogen due to being metabolised in the liver to a reactive epoxide, which may form DNA adducts at some guanine residues (Gouas et al. 2009; Bedard & Massey 2006). The epoxide adduct is reported to catalyze a G → T mutation in the *p53* tumour suppressor protein gene, where it causes a missense mutation effectively inactivating the gene's protein product (Besaratina et al. 2009). This identified mechanism, as well as previous studies in animal models, has led the IARC to classify AFB₁, AFB₂, AFG₁, and AFG₂ as Group 1 carcinogens, denoting their explicit carcinogenicity to humans (World Health Organization & International Agency for Research on Cancer 2002). Developing nations, including most of Africa, Latin and South Americas, and Asia are identified as high risk areas for aflatoxin exposure, leading to aflatoxicosis (Williams et al. 2004).

An Italian study using lactating goats illustrated the relatively rapid absorption of AFB₁ and its subsequent metabolism and excretion as AFM₁, with peak concentrations of AFM₁ detected in expressed milk between 3 and 6 hours after oral AFB₁ administration, and non-detection of AFM₁ after 3.5 days post-dosage (Battacone et al. 2012). However, the amount of AFM₁ detected only accounted for less than 0.2% of the original dose, leaving a large majority as either degraded or excreted by other mechanisms. The rates of aflatoxin absorption and degradation were shown to be affected not only by species, but by diet, specifically through varying the roughage component of livestock feed (Upadhaya et al. 2009).

Ochratoxins

Ochratoxin is a mycotoxin that comes in three secondary metabolite forms, A, B, and C, all of which are produced by *Penicillium* and *Aspergillus* species contaminating in a wide range of commodities including staple food crops and beverages such as beer and wine (Bayman & Baker 2006). OTA is one of the most commonly encountered mycotoxins, and was reportedly detected in 60% of a healthy Moroccan study population of blood donors (Filali et al. 2002). The structure of OTA is depicted in **figure 2**; the related compound ochratoxin B differs only in having hydrogen in place of the chlorine atom, and ochratoxin C is an ethyl ester form of OTA produced in presence of rumen fluid (Fuchs et al. 1984).

OTA has been hypothesized to cause oxidative damage to DNA, leading to mutagenesis and potential carcinogenesis (Zepnik 2001). Recent papers also propose direct genotoxic mechanisms for OTA, describing a pathway that metabolizes OTA into an electrophilic species capable of directly binding to some nucleotide bases (Pfohl-Leszkowicz & Manderville 2012). Consequently, OTA is classified as a possible human carcinogen by IARC, citing sufficient evidence of carcinogenicity in animal models, but insufficient evidence from human studies (World Health Organization & International Agency for Research on Cancer 2002).

OTA is known to bind to blood plasma proteins, delaying its excretion in urine by up to 30 days, a fact greatly relevant to the development of methods for exposure detection (Kumagai 1985). Despite this temporary bioaccumulation, no significant relationship was found between age and plasma OTA levels in a British study population (Gilbert et al. 2001). However, a variable relationship between plasma and urine levels was hypothesized to result from decreased efficiency of OTA metabolic and excretion mechanisms with age.

Fumonisin

Since discovery of this group of mycotoxins in 1988, at least 28 compounds have been identified as fumonisins, with some of the most common forms illustrated in **figure 3** (Voss et al. 2007; Bolger et al. 2001; Tamura et al. 2014). These compounds are predominantly produced on maize and maize products by *F. proliferatum* and *F. verticillioides* (formerly *F. moniliforme*), but have also been reported in cultures of several other *Fusarium* spp., as well as the *Alternaria alternata* f. sp. *lycopersici* natively growing on tomatoes, and *Aspergillus niger* on coffee and grapes (Weidenbörner 2001; Scott 2012). Among the fumonisin compounds, fumonisin B₁ (FB₁) is produced most abundantly, and features most prominently in the literature (Mahmoodi et al. 2012). In an analysis of various foods and feeds from Burkina Faso, FB₁ was detected in over 80% of samples, with FB₂ and FB₃ also found in 69% and 46% of samples, respectively (Warth et al. 2012).

Dietary exposure to fumonisins was reported to have rapid but low absorption, based on oral administration of FB₁ in a rat model, with 3.5% of the dose observed in plasma, peaking in concentration after one hour (Voss et al. 2007). Another study in rats reported observations supporting a hypothesis of FB₁ genotoxicity mediated by oxidative stress mechanisms (Theumer et al. 2010). *Fusaria*-derived toxins such as FB₁ and FB₂ are listed as possible carcinogens in Group 2B of the

IARC classification (World Health Organization & International Agency for Research on Cancer 2002). A recent etiological study also found some positive correlation between prevalence of FB₁ contamination in rice and incidence of esophageal cancer, though this relationship was not seen in other staple foods, such as corn, in the same region of Iran (Alizadeh et al. 2012).

Due to a structural resemblance with ceramide, fumonisins are also reported to disrupt the *de novo* synthesis of ceramide and sphingolipid metabolism, potentially leading to broad impairment of cellular signalling mechanisms (Dutton 1996). FB₁ neurotoxicity was also observed in a carp model (Kovacic et al. 2009). However, formation of fumonisin A₁ (FA₁) through acetylation of the amine site in FB₁ was reported to block these effects (Stockmann-Juvala & Savolainen 2008). Additionally, types of maize processing that involved cooking in alkaline conditions were estimated to mitigate some adverse health effects through hydrolysis of the fumonisins (Voss et al. 2007).

T-2 Toxin and HT-2 Toxin

The T-2 and HT-2 toxins are produced by various *Fusarium* species, particularly *F. sporotrichioides* and *F. poae*, reportedly affecting several major cereal crops including oats, barley, corn, and wheat (Weidner et al. 2013). A European evaluation found oats and oat products contained the highest summed concentrations of these two contaminants (EFSA Panel on Contaminants in the Food Chain 2011).

These toxins are common representatives of the type A tricothecenes, a group identified by specific arrangements of ligands around a sesquiterpenoid cyclic ring (Cano-Sancho et al. 2012). As illustrated in **figure 4**, HT-2 is structurally differentiated from T-2 by oxidation of an acetate ligand at C-4.

Exposure to these compounds is associated with diverse pathologies including skin irritation, gastrointestinal issues (nausea, cramping, and vomiting), impaired mitochondrial function, and hypotrophy of the spleen and thymus (Ueno 1984). Recent research has discovered evidence for these toxins crossing the blood-brain barrier *in vitro*, linking them to neurotoxic effects (Weidner et al. 2013). However, the predominant human health effects reported in literature associate these toxins with inhibited protein synthesis and apoptosis in a wide variety of *in vitro* and *ex vitro* cells, as well as *in vivo* organs, including brain, gastrointestinal tract, skin, spleen, and thymus (Li et al. 2011; EFSA Panel on Contaminants in the Food Chain 2011). Though these tricothecenes were not observed to

produce genotoxic or directly mutagenic effects *in vitro*, the aforementioned widespread disruptive effects of exposure may potentially increase susceptibility to other carcinogenic factors (Rocha et al. 2005).

Deoxynivalenol

DON is produced by *F. graminearum* and *F. culmorum* on cereal crops, and is one of the most commonly encountered mycotoxins (Reddy et al. 2010). Acute dietary exposure to DON is characterized in animals by emesis and feed refusal. DON is rapidly metabolized for urinary excretion as glucuronide conjugates within 24hrs of dietary exposure (Lake et al. 1987).

As illustrated in **figure 5**, DON may be identified as a type-B trichothecene by the carbonyl group at C-8. Like all trichothecenes, DON is a strong inhibitor of protein synthesis (Ji et al. 2014). Literature reports DON preventing eukaryotic polypeptide assembly by interfering with activity of the 60S ribosomal subunit (Pestka 2007).

In 2002, an IARC review of available literature listed DON as a Group 3 (not classifiable) human carcinogen due to inadequate evidence of animal carcinogenicity, and lack of investigation in humans (World Health Organization & International Agency for Research on Cancer 2002). However, a more recent *in vivo* mouse study concluded oral exposure to DON was able to induce lung adenocarcinomas, with the carcinogenic effect synergized by co-exposure to sterigmatocystin (Huang et al. 2004). Unfortunately, the results of this study were argued to be invalid by a following review article, citing an inappropriate experimental design that did not represent real-world conditions of human exposure to DON (Ma & Guo 2008). A third study reported observing clear, dose-dependent induction of DNA strand breakage by DON treatment of *in vitro* liver cells (Zhang et al. 2009).

Zearalenone

Zearalenone (ZEN) is another common mycotoxin produced by various *Fusarium* fungi, detected in some 90% of sampled cereal crops that were grown in central European countries such as Austria, France, and Germany (Baliukoniene et al. 2003). ZEN shares a structural similarity with the human sex hormone 17 β -estradiol, and therefore also has a similar affinity to and effect on estrogen receptors, resulting in associations with fertility problems in both humans and livestock (Nordic Council of Ministers 1998).

IARC found limited evidence of ZEN carcinogenicity in animal models, classifying it together with DON in Group 3 (World Health Organization & International Agency for Research on Cancer 2002). Due to rapid metabolism, bioaccumulation is understood to be insignificant, though co-exposure with other mycotoxins is estimated to present a probable health risk due to inadequately researched interactions (Speijers & Speijers 2004).

Interestingly, ZEN is frequently reported to have antagonistic effects on the toxicity of some co-exposed mycotoxins. For example, OTA-induced kidney damage in rats was observed to be significantly mitigated in comparison cases where the animals were given both ZEN and OTA (Grenier & Oswald 2011). In another study, ZEN co-exposed with DON was also observed to produce sub-additive or antagonistic effect on levels of some serum immunoglobulins, as compared to individual ZEN or DON exposure cases (Forsell et al. 1986).

Co-occurrence and co-exposure of multiple mycotoxins

In a South African study, the co-occurrence of mycotoxin-producing fungi on crops was reported for almost half of contaminated samples (Mngadi et al. 2008). As mentioned before, many species of mycotoxin-producing fungi are known to be capable of producing more than one mycotoxin. A Polish study reported co-occurrence of mycotoxins in over half the rye samples analysed, with the most common combination being DON and ZEN (Błajet-Kosicka et al. 2014). This mycotoxin pair was also detected in a study on several Chinese wheat producing regions (Ji et al. 2014). Further, analysis of a variety of foods and feeds produced in Burkina Faso and Mozambique reported up to 28 different mycotoxins present and quantifiable in a single sample (Warth et al. 2012). Finally, even greater co-occurrence was reported after a recent analysis detecting between 5 and 41 of the previously discussed mycotoxins in individual samples of maize from various agricultural sites in Malawi (Matumba et al. 2014).

Government and industry regulations exist to minimize the concentrations of individual mycotoxins allowed into food and feed products (Zain 2011; The Commission of the European Communities 2006). However, these regulations are based on individual toxicities, and as such, do not take into account the complex dynamics of compounded risk from co-exposure to groups of mycotoxins, for even acute pathologies. Agricultural products are effectively aggregated and heterogeneously distributed for consumer convenience, leading to a highly unpredictable mixture of

possible mycotoxin contamination in food products (Turner et al. 2012). An analysis of 174 cereal-based food products from Belgian supermarkets reported a median of four mycotoxins present in these consumer-available products (De Boevre et al. 2012). Population studies of mycotoxin exposure therefore also frequently report detectable amounts of multiple mycotoxins in blood and urine samples from human populations all over the globe (Abia et al. 2013; Solfrizzo et al. 2014; Shirima et al. 2014; Njumbe-Ediage et al. 2012). Consequently, an important question remains posed by the current literature, to identify the chronic risks associated with repetitive exposure to low levels of multiple mycotoxins (Warth et al. 2013).

Several *in vivo* studies with various species of farm animals co-exposed to pairs of mycotoxins were compared in the literature, illustrating a complex set of possible synergistic, additive, sub-additive, or antagonistic effects on animal health and growth metrics, as compared with single mycotoxin exposures (Grenier & Oswald 2011; Speijers & Speijers 2004; Bensassi et al. 2014). These reports serve to highlight not only the complex and dynamic interaction between co-exposures, but also the great variety of possible co-exposure situations, and the dearth of studies representative of the real-world situation, particularly in humans (Streit et al. 2013).

Mycotoxins and carcinogenesis

Although mycotoxin exposure has been associated with several acute and chronic human health effects, one of the most important health burdens associated with mycotoxin exposure is the development of cancers. The possible causes of carcinogenesis are virtually innumerable, however, exposure to some chemical factors (*e.g.*, (multi-)mycotoxin exposure) is known to be a strong modulator of carcinogenic risk (Cohen & Arnold 2011). In the unfortunate case of such exposures, three stages are identified in the carcinogenic process: initiation, promotion, and progression (Oliveira et al. 2007). For example, AFB₁ is known to be a potent initiator of hepatocellular carcinoma, while fumonisins are reported to be strong tumour promoters after initiation of carcinogenesis by another agent, such as AFB₁ (Riley 1998). Although several cancers might be associated with multi-mycotoxin exposure (*e.g.* liver, colon, rectum, esophageal, breast cancer, etc.), direct evidence is still scarce for most of these cancer types, and multi-mycotoxin exposure may not yet have been investigated for causal relationship. Therefore, only the two most frequently studied cancer types in relation to mycotoxin exposure will be outlined in this review paper, namely colorectal and hepatocellular

carcinoma.

Colorectal carcinoma

Cancers of either the colon or rectum are the fourth most common causes of death by cancer worldwide (World Health Organization 2014). Rates of incidence rank colorectal cancer slightly higher, as the second and third most common site of cancer diagnosis in women and men, respectively (EpiCast Reports 2012). As with all cancers, the risk of incidence increases with age (Howlander et al. 2011). This is largely due to cumulative risk of carcinogenesis over the course of chronic exposure to potential carcinogens. Particularly, since the prevalent vector of human exposure to mycotoxins is through the diet, it is postulated that the colorectal region is itself highly exposed to these compounds, and therefore a prominent target for pathological developments (Pfohl-Leszkowicz et al. 1995; Bouhet & Oswald 2005).

The HT-29 human colonic cell line was used for *in vitro* studies of DON and FB₁, both of which were observed to induce apoptosis, though by different mechanisms (Schmelz et al. 1998; Ma et al. 2012). DON-induced apoptosis was separately reported to be caused by mitochondrial dysfunction, while treatment with FB₁ resulted in accumulation of endogenous free sphingoid bases, inhibiting growth and inducing apoptosis. A different colonic cell line, HCT-116, was used in an investigation of AFB₁, reporting induction of DNA damage and lesions, as well as inactivation of the ATR/Chk1 pathway that would normally help address an appropriate response to this damage (Gursoy-Yuzugullu et al. 2011). Further investigation of synergistic interactions between co-exposed mycotoxins is especially required in the colorectal context.

Hepatocellular carcinoma

Almost 700,000 individuals were recorded to have died of liver cancer in 2008, increasing to over 750,000 by 2010, making hepatocellular carcinoma the third most deadly form of cancer worldwide (Lozano et al. 2012; Ferlay et al. 2010). A slightly older report estimates the annual global diagnosis rate at over 560,000 individuals in 2000, though such estimates are reported elsewhere to experience growth at a rate of at least 3% annually (Bosch et al. 2004; Howlander et al. 2011).

There is a strong, established link between aflatoxin exposure and development of malignant hepatomas, with a speculated 5% to 28% of cases actually caused by aflatoxins (Gursoy-Yuzugullu et

al. 2011). The associated mechanism, as previously discussed, involves metabolism of AFB₁ in the liver to a highly reactive species capable of forming mutagenic DNA adducts. One well-known mutation results in the inactivation of p53 tumour suppressor gene, enabling initiation of carcinogenesis. Subsequently, a synergistic effect promoting tumour growth in the liver was reported for co-exposure to fumonisins (Carlson et al. 2001). Less pronounced effects were reported by multiple studies on co-exposed OTA also resulting in increased hepatic lipid levels and increased relative weight of the organ (Grenier & Oswald 2011).

Conclusion

Though literally innumerable fungal secondary metabolites exist, a few were identified as particularly harmful by centuries of experience and research. The acute and chronic health effects of exposure to these xenobiotics vary almost as widely as the mechanisms by which various mycotoxins produce these effects. Nevertheless, great efforts of research have helped in characterizing the hazards of mycotoxins. Unfortunately, the rate at which mycotoxins are produced and the ubiquity of their production throughout the world's agricultural industries, coupled with modern food processing and distribution techniques, indicates a deeper understanding of actual exposure is needed. In particular, both co-exposure and chronic exposure are increasingly identified as real-world factors, consequences of which are most relevant to inquiries regarding real-world health effects.

Acknowledgements

This work was written as part of a doctoral project, funded by a doctoral grant for strategic basic research (SB #141177) from the Flemish agency for Innovation by Science and Technology (IWT).

Conflict of Interest Statement

The authors declare that there are no conflicts of interest.

References

- Abia, W. a et al., 2013. Bio-monitoring of mycotoxin exposure in Cameroon using a urinary multi-biomarker approach. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*, 62, pp.927–34. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24128729> [Accessed October 31, 2014].
- Alizadeh, A.M. et al., 2012. Fumonisin B1 contamination of cereals and risk of esophageal cancer in a high risk area in northeastern Iran. *Asian Pacific journal of cancer prevention : APJCP*, 13, pp.2625–8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22938431>.
- Alkadri, D. et al., 2014. Natural co-occurrence of mycotoxins in wheat grains from Italy and Syria. *Food chemistry*, 157, pp.111–8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24679759> [Accessed October 18, 2014].
- Baliukoniene, V., Bakutis, B. & Stankevicius, H., 2003. Mycological and mycotoxicological evaluation of grain. *Annals of agricultural and environmental medicine : AAEM*, 10(2), pp.223–7. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14677916> [Accessed January 1, 2015].
- Battacone, G. et al., 2012. Excretion pattern of aflatoxin M1 in milk of goats fed a single dose of aflatoxin B1. *Journal of dairy science*, 95(5), pp.2656–61. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22541493> [Accessed December 4, 2014].
- Bayman, P. & Baker, J.L., 2006. Ochratoxins: A global perspective. *Mycopathologia*, 162(3), pp.215–223.
- Bedard, L.L. & Massey, T.E., 2006. Aflatoxin B1-induced DNA damage and its repair. *Cancer letters*, 241(2), pp.174–83. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16458422> [Accessed December 31, 2014].
- Bennett, J.W. & Klich, M., 2003. Mycotoxins. *Clinical Microbiology Reviews*, 16(3), pp.497–516. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=164220&tool=pmcentrez&rendertype=abstract> [Accessed January 14, 2015].
- Bensassi, F. et al., 2014. In vitro investigation of toxicological interactions between the fusariotoxins deoxynivalenol and zearalenone. *Toxicon : official journal of the International Society on Toxinology*, 84, pp.1–6. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24680766> [Accessed October 18, 2014].
- Bensassi, F. et al., 2009. Pathway of deoxynivalenol-induced apoptosis in human colon carcinoma cells. *Toxicology*, 264(1-2), pp.104–9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19664677> [Accessed October 18, 2014].
- Besaratinia, A. et al., 2009. In vitro recapitulating of TP53 mutagenesis in hepatocellular carcinoma associated with dietary aflatoxin B1 exposure. *Gastroenterology*, 137(3), pp.1127–37, 1137.e1–5. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2736365&tool=pmcentrez&rendertype=abstract> [Accessed December 31, 2014].
- Blajet-Kosicka, A. et al., 2014. Co-occurrence and evaluation of mycotoxins in organic and conventional rye grain and products. *Food Control*, 38, pp.61–66. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0956713513005252> [Accessed January 9, 2015].

- De Boevre, M. et al., 2012. Natural occurrence of mycotoxins and their masked forms in food and feed products. *World Mycotoxin Journal*, 5, pp.207–219.
- Bolger, M. et al., 2001. Fumonisin: Safety evaluations of certain mycotoxins in foods. *FAO Food and Nutrition Paper*, 74, pp.103–279. Available at: <http://www.inchem.org/documents/jecfa/jecmono/v47je03.htm> [Accessed January 6, 2015].
- Bosch, F.X. et al., 2004. Primary liver cancer: Worldwide incidence and trends. *Gastroenterology*, 127(5), pp.S5–S16. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0016508504015902> [Accessed January 8, 2015].
- Bouhet, S. & Oswald, I.P., 2005. The effects of mycotoxins, fungal food contaminants, on the intestinal epithelial cell-derived innate immune response. *Veterinary Immunology and Immunopathology*, 108, pp.199–209.
- Cano-Sancho, G. et al., 2012. Exposure assessment of T2 and HT2 toxins in Catalonia (Spain). *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*, 50(3-4), pp.511–7. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22261462> [Accessed December 5, 2014].
- Carlson, D.B. et al., 2001. Fumonisin B1 promotes aflatoxin B1 and N-methyl-N'-nitro-nitrosoguanidine-initiated liver tumors in rainbow trout. *Toxicology and applied pharmacology*, 172(1), pp.29–36. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11264020> [Accessed January 14, 2015].
- Chelkowski, J., 1998. Distribution of Fusarium species and their mycotoxins in cereal grains. In *Mycotoxins in Agriculture and Food Safety*. pp. 45–64.
- Cohen, S.M. & Arnold, L.L., 2011. Chemical carcinogenesis. *Toxicological sciences : an official journal of the Society of Toxicology*, 120 Suppl , pp.S76–92. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21147961> [Accessed January 9, 2015].
- Dutton, M.F., 1996. Fumonisin, mycotoxins of increasing importance: Their nature and their effects. *Pharmacology & Therapeutics*, 70(2), pp.137–161. Available at: <http://linkinghub.elsevier.com/retrieve/pii/016372589600006X> [Accessed December 5, 2014].
- EFSA Panel on Contaminants in the Food Chain, 2011. Scientific Opinion on the risks for animal and public health related to the presence of T-2 and HT-2 toxin in food and feed. *EFSA Journal*, 9(12), p.187. Available at: <http://www.efsa.europa.eu/en/efsajournal/doc/2481.pdf>.
- EpiCast Reports, 2012. *Colorectal Cancer: Epidemiology Forecast to 2022*, London.
- Ferlay, J. et al., 2010. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *International Journal of Cancer*, 127(12), pp.2893–2917.
- Filali, A. et al., 2002. Ochratoxin A in human plasma in Morocco: a preliminary survey. *Human & experimental toxicology*, 21(5), pp.241–5. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12141394> [Accessed December 4, 2014].
- Forsell, J.H. et al., 1986. Effects of 8-week exposure of the B6C3F1 mouse to dietary deoxynivalenol (vomitoxin) and zearalenone. *Food and Chemical Toxicology*, 24(3), pp.213–219. Available at: <http://linkinghub.elsevier.com/retrieve/pii/0278691586902310>.
- Fuchs, R. et al., 1984. Conversion of ochratoxin C into ochratoxin A in vivo. *Applied and Environmental Microbiology*, 48(1), pp.41–42.

- Garrido, N.S. et al., 2003. Occurrence of aflatoxins M(1) and M(2) in milk commercialized in Ribeirão Preto-SP, Brazil. *Food additives and contaminants*, 20(1), pp.70–3. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12519721> [Accessed December 4, 2014].
- Gilbert, J., Brereton, P. & MacDonald, S., 2001. Assessment of dietary exposure to ochratoxin A in the UK using a duplicate diet approach and analysis of urine and plasma samples. *Food additives and contaminants*, 18(12), pp.1088–93. Available at: <http://cat.inist.fr/?aModele=afficheN&cpsidt=14158551> [Accessed December 4, 2014].
- Gouas, D., Shi, H. & Hainaut, P., 2009. The aflatoxin-induced TP53 mutation at codon 249 (R249S): biomarker of exposure, early detection and target for therapy. *Cancer letters*, 286(1), pp.29–37. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19376640> [Accessed December 31, 2014].
- Grenier, B. & Oswald, I.P., 2011. Mycotoxin co-contamination of food and feed: meta-analysis of publications describing toxicological interactions. *World Mycotoxin Journal*, 4(3), pp.285–313. Available at: <http://wageningenacademic.metapress.com/openurl.asp?genre=article&id=doi:10.3920/WMJ2011.1.1281> [Accessed January 14, 2015].
- Gursoy-Yuzugullu, O. et al., 2011. Aflatoxin genotoxicity is associated with a defective DNA damage response bypassing p53 activation. *Liver international : official journal of the International Association for the Study of the Liver*, 31(4), pp.561–71. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21382167> [Accessed October 18, 2014].
- Hirst, J.M. & Stedman, O.J., 1967. Long-distance spore transport: methods of measurement, vertical spore profiles and the detection of immigrant spores. *Journal of general microbiology*, 48(3), pp.329–55. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6052628> [Accessed November 28, 2014].
- Howlader, N. et al., 2011. *SEER Cancer Statistics Review, 1975-2009 (Vintage 2009 Populations)*, Bethesda, MD. Available at: http://seer.cancer.gov/archive/csr/1975_2009_pops09/.
- Huang, X. hua et al., 2004. Carcinogenic effects of sterigmatocystin and deoxynivalenol in NIH mice. *Zhonghua zhong liu za zhi [Chinese journal of oncology]*, 26, pp.705–708.
- International Agency for Research on Cancer, 2007. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. *Monograph*, 89, pp.223 – 276.
- Jackson, L.C., Kudupoje, M.B. & Yiannikouris, A., 2012. Simultaneous multiple mycotoxin quantification in feed samples using three isotopically labeled internal standards applied for isotopic dilution and data normalization through ultra-performance liquid chromatography/electrospray ionization tandem mass spe. *Rapid communications in mass spectrometry : RCM*, 26(23), pp.2697–713. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23124660> [Accessed December 5, 2014].
- Ji, F. et al., 2014. Natural occurrence of deoxynivalenol and zearalenone in wheat from Jiangsu province, China. *Food chemistry*, 157(1881), pp.393–7. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24679796> [Accessed October 18, 2014].
- Kang'ethe, E.K. & Lang'a, K.A., 2009. Aflatoxin B1 and M1 contamination of animal feeds and milk from urban centers in Kenya. *African health sciences*, 9(4), pp.218–26. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3074399&tool=pmcentrez&rendertype=abstract> [Accessed November 28, 2014].
- Klich, M.A., 2007. *Aspergillus flavus*: the major producer of aflatoxin. *Molecular plant pathology*, 8(6), pp.713–22. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20507532> [Accessed December 2, 2014].

- Kovacić, S. et al., 2009. Fumonisin B1 neurotoxicity in young carp (*Cyprinus carpio* L.). *Arhiv za higijenu rada i toksikologiju*, 60, pp.419–426.
- Kumagai, S., 1985. Ochratoxin A: plasma concentration and excretion into bile and urine in albumin-deficient rats. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*, 23(10), pp.941–3. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/4065769> [Accessed December 4, 2014].
- Lake, B.G. et al., 1987. Studies on the metabolism of deoxynivalenol in the rat. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*, 25(8), pp.589–92. Available at: <http://linkinghub.elsevier.com/retrieve/pii/0278691587900196> [Accessed December 5, 2014].
- Li, Y. et al., 2011. T-2 toxin, a trichothecene mycotoxin: review of toxicity, metabolism, and analytical methods. *Journal of agricultural and food chemistry*, 59(8), pp.3441–53. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21417259> [Accessed January 8, 2015].
- Lozano, R. et al., 2012. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*, 380(9859), pp.2095–128. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23245604> [Accessed July 10, 2014].
- Ma, Y. et al., 2012. A mitochondria-mediated apoptotic pathway induced by deoxynivalenol in human colon cancer cells. *Toxicology in vitro : an international journal published in association with BIBRA*, 26(3), pp.414–20. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22269384> [Accessed October 18, 2014].
- Ma, Y.Y. & Guo, H.W., 2008. Mini-review of studies on the carcinogenicity of deoxynivalenol. *Environmental Toxicology and Pharmacology*, 25, pp.1–9.
- Mahmoodi, M. et al., 2012. Impact of fumonisin B1 on the production of inflammatory cytokines by gastric and colon cell lines. *Iranian Journal of Allergy, Asthma, and Immunology*, 11(2), pp.165–73. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22761190> [Accessed November 27, 2014].
- Matumba, L. et al., 2014. Fungal metabolites diversity in maize and associated human dietary exposures relate to micro-climatic patterns in Malawi. *World Mycotoxin Journal*, pp.1–14.
- Mngadi, P.T., Govinden, R. & Odhav, B., 2008. Co-occurring mycotoxins in animal feeds. *Journal of Biotechnology*, 7, pp.2239–2243.
- Njumbe-Ediage, E. et al., 2012. A direct assessment of mycotoxin biomarkers in human urine samples by liquid chromatography tandem mass spectrometry. *Analytica chimica acta*, 741, pp.58–69. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22840705> [Accessed November 5, 2014].
- Nordic Council of Ministers, 1998. *Fusarium Toxins in Cereals: A Risk Assessment* G. Eriksen & J. Alexander, eds., Copenhagen: Nordic Council of Ministers. Available at: <https://books.google.com/books?hl=en&lr=&id=jU5Wx8LkZHUC&pgis=1> [Accessed January 8, 2015].
- Oliveira, P.A. et al., 2007. Chemical carcinogenesis. *Anais da Academia Brasileira de Ciências*, 79(4), pp.593–616. Available at: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0001-37652007000400004&lng=en&nrm=iso&tlang=en [Accessed January 9, 2015].
- Pestka, J.J., 2007. Deoxynivalenol: Toxicity, mechanisms and animal health risks. *Animal Feed Science and Technology*, 137(3-4), pp.283–298. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0377840107002209> [Accessed December 5, 2014].

- Pfohl-Leszkowicz, a. et al., 1995. High levels of DNA adducts in human colon are associated with colorectal cancer. *Cancer Research*, 55, pp.5611–5616.
- Pfohl-Leszkowicz, A. & Manderville, R. a, 2012. An update on direct genotoxicity as a molecular mechanism of ochratoxin a carcinogenicity. *Chemical research in toxicology*, 25(2), pp.252–62. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22054007> [Accessed January 6, 2015].
- Reddy, K. et al., 2010. An overview of mycotoxin contamination in foods and its implications for human health. *Toxin Reviews*, 29(1), pp.3–26. Available at: <http://informahealthcare.com/doi/abs/10.3109/15569541003598553> [Accessed August 28, 2014].
- Riley, R.T., 1998. Mechanistic Interactions of Mycotoxins: Theoretical Considerations. In K. K. Shinha & D. Bhatnagar, eds. *Mycotoxins in Agriculture and Food Safety*. Athens, USA: CRC Press, pp. 227–256. Available at: <http://books.google.com/books?hl=en&lr=&id=FBDzbLAUYMcC&pgis=1> [Accessed January 14, 2015].
- Rocha, O., Ansari, K. & Doohan, F.M., 2005. Effects of trichothecene mycotoxins on eukaryotic cells: a review. *Food additives and contaminants*, 22(4), pp.369–78. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16019807> [Accessed December 27, 2014].
- Schmelz, E.M. et al., 1998. Induction of apoptosis by fumonisin B1 in HT29 cells is mediated by the accumulation of endogenous free sphingoid bases. *Toxicology and applied pharmacology*, 148(2), pp.252–60. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9473533> [Accessed January 15, 2015].
- Schreiner, O. & Reed, H., 1908. The toxic action of certain organic plant constituents. *Botanical gazette*, 45(2), pp.73–102. Available at: <http://www.jstor.org/stable/2467619> [Accessed November 28, 2014].
- Scott, P.M., 2012. Recent research on fumonisins: a review. *Food additives & contaminants. Part A, Chemistry, analysis, control, exposure & risk assessment*, 29(2), pp.242–8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21337235> [Accessed December 5, 2014].
- Shirima, C.P. et al., 2014. A Prospective Study of Growth and Biomarkers of Exposure to Aflatoxin and Fumonisin during Early Childhood in Tanzania. *Environmental health perspectives*. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25325363> [Accessed October 31, 2014].
- Solfrizzo, M., Gambacorta, L. & Visconti, A., 2014. Assessment of multi-mycotoxin exposure in southern Italy by urinary multi-biomarker determination. *Toxins*, 6(2), pp.523–38. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3942749&tool=pmcentrez&rendertype=abstract> [Accessed October 18, 2014].
- Speijers, G.J. a & Speijers, M.H.M., 2004. Combined toxic effects of mycotoxins. *Toxicology letters*, 153(1), pp.91–8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15342085> [Accessed January 14, 2015].
- Stockmann-Juvala, H. & Savolainen, K., 2008. A review of the toxic effects and mechanisms of action of fumonisin B1. *Human & experimental toxicology*, 27(11), pp.799–809. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19244287> [Accessed December 28, 2014].
- Streit, E. et al., 2013. Multi-mycotoxin screening reveals the occurrence of 139 different secondary metabolites in feed and feed ingredients. *Toxins*, 5(3), pp.504–23. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3705275&tool=pmcentrez&rendertype=abstract> [Accessed January 3, 2015].
- Tamura, M. et al., 2014. Characterization of fumonisin A-series by high-resolution liquid chromatography-orbitrap mass spectrometry. *Toxins*, 6(8), pp.2580–93. Available at:

<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4147598&tool=pmcentrez&rendertype=abstract> [Accessed January 6, 2015].

The Commission of the European Communities, 2006. Commission Regulation (EC) No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs. *Official Journal of the European Union*, L364(1881), pp.5–24.

Theumer, M.G. et al., 2010. Subchronic mycotoxicoses in Wistar rats: Assessment of the in vivo and in vitro genotoxicity induced by fumonisins and aflatoxin B1, and oxidative stress biomarkers status. *Toxicology*, 268, pp.104–110.

Turner, P.C. et al., 2012. The role of biomarkers in evaluating human health concerns from fungal contaminants in food. *Nutrition research reviews*, 25(1), pp.162–179. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22651937> [Accessed October 28, 2014].

Ueno, Y., 1984. Toxicological features of T-2 toxin and related trichothecenes. *Fundamental and applied toxicology : official journal of the Society of Toxicology*, 4(2 Pt 2), pp.S124–32. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6609858> [Accessed December 5, 2014].

Upadhaya, S.D. et al., 2009. Comparative study on the aflatoxin B1 degradation ability of rumen fluid from Holstein steers and Korean native goats. *Journal of Veterinary Science*, 10, pp.29–34.

Varga, J., Frisvad, J.C. & Samson, R.A., 2011. Two new aflatoxin producing species, and an overview of *Aspergillus* section *Flavi*. *Studies in Mycology*, 69, pp.57–80.

Voss, K. a., Smith, G.W. & Haschek, W.M., 2007. Fumonisins: Toxicokinetics, mechanism of action and toxicity. *Animal Feed Science and Technology*, 137(3-4), pp.299–325. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0377840107002210> [Accessed December 5, 2014].

Warth, B. et al., 2012. Quantitation of mycotoxins in food and feed from Burkina Faso and Mozambique using a modern LC-MS/MS multitoxin method. *Journal of agricultural and food chemistry*, 60(36), pp.9352–63. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22835072> [Accessed January 12, 2015].

Warth, B., Sulyok, M. & Krska, R., 2013. LC-MS/MS-based multibiomarker approaches for the assessment of human exposure to mycotoxins. *Analytical and bioanalytical chemistry*, 405(17), pp.5687–95. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3695324&tool=pmcentrez&rendertype=abstract> [Accessed November 4, 2014].

Weidenbörner, M., 2001. Foods and fumonisins. *European Food Research and Technology*, 212(3), pp.262–273. Available at: <http://link.springer.com/10.1007/s002170000259> [Accessed December 5, 2014].

Weidner, M. et al., 2013. Influence of T-2 and HT-2 toxin on the blood-brain barrier in vitro: new experimental hints for neurotoxic effects. *PloS one*, 8(3), p.e60484. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3609806&tool=pmcentrez&rendertype=abstract> [Accessed January 8, 2015].

Williams, J.H. et al., 2004. Human aflatoxicosis in developing countries: A review of toxicology, exposure, potential health consequences, and interventions. *American Journal of Clinical Nutrition*, 80(5), pp.1106–1122.

Williams, P.L., James, R.C. & Roberts, S.M., 2015. *Principles of Toxicology: Environmental and Industrial Applications*, John Wiley & Sons. Available at: <https://books.google.com/books?id=KrqnBQAAQBAJ&pgis=1> [Accessed January 27, 2015].

World Health Organization, 2014. Cancer. *Fact Sheet N°297*. Available at: <http://www.who.int/mediacentre/factsheets/fs297/en/> [Accessed January 15, 2015].

World Health Organization & International Agency for Research on Cancer, 2002. INTERNATIONAL AGENCY FOR RESEARCH ON CANCER IARC MONOGRAPHS ON THE EVALUATION OF CARCINOGENIC RISKS TO HUMANS. In *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*. Lyon: IARC Press, pp. 1–601.

Zain, M.E., 2011. Impact of mycotoxins on humans and animals. *Journal of Saudi Chemical Society*, 15(2), pp.129–144. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S1319610310000827> [Accessed October 18, 2014].

Zepnik, H., 2001. Ochratoxin A-Induced Tumor Formation: Is There a Role of Reactive Ochratoxin A Metabolites? *Toxicological Sciences*, 59(1), pp.59–67. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11134544> [Accessed December 4, 2014].

Zhang, X. et al., 2009. The role of oxidative stress in deoxynivalenol-induced DNA damage in HepG2 cells. *Toxicon*, 54(4), pp.513–518. Available at: <http://dx.doi.org/10.1016/j.toxicon.2009.05.021>.

List of Figures

Figure 1: Chemical structures of six different aflatoxins: aflatoxin B₁, aflatoxin B₂, aflatoxin G₁, aflatoxin G₂, aflatoxin M₁, and aflatoxin M₂.

Figure 2: Chemical structure of ochratoxin A.

Figure 3: Chemical structure and ligands identifying some common fumonisins.

Figure 4: Chemical structures of T-2 toxin and HT-2 toxin.

Figure 5: Chemical structure of deoxynivalenol.

Figure 6: Chemical structure of zearalenone.




Figure 1: Chemical structures of six different aflatoxins: aflatoxin B₁, aflatoxin B₂, aflatoxin G₁, aflatoxin G₂, aflatoxin M₁, and aflatoxin M₂.




Figure 2: Chemical structure of ochratoxin A.




Figure 3: Chemical structure and ligands identifying some common fumonisins (Bolger et al. 2001; Tamura et al. 2014).




Figure 4: Chemical structures of T-2 toxin and HT-2 toxin.




Figure 5: Chemical structure of deoxynivalenol.




Figure 6: Chemical structure of zearalenone.