

# Analysis of controlling factors on boulder production on the hillslopes of the upper Bhote Koshi catchment, Nepal

Dini, Benedetta; Bennett, Georgina L.

DOI:

[10.5194/egusphere-egu21-15605](https://doi.org/10.5194/egusphere-egu21-15605)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Dini, B & Bennett, GL 2021, Analysis of controlling factors on boulder production on the hillslopes of the upper Bhote Koshi catchment, Nepal. in *EGU General Assembly 2021.*, EGU21-15605, European Geosciences Union, EGU General Assembly 2021, 19/04/21. <https://doi.org/10.5194/egusphere-egu21-15605>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

EGU21-15605

<https://doi.org/10.5194/egusphere-egu21-15605>

EGU General Assembly 2021

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



## Analysis of controlling factors on boulder production on the hillslopes of the upper Bhote Koshi catchment, Nepal

**Benedetta Dini**<sup>1</sup> and Georgina L. Bennett<sup>2</sup>

<sup>1</sup>School of Environmental Sciences, University of East Anglia, UK

<sup>2</sup>College of Life and Environmental Sciences, University of Exeter, UK

Landslides from mountainous bedrock hillslopes often contain boulders, the presence of which has been shown to influence landscape evolution by altering hillslope geomorphic processes and river erosion. Furthermore, the presence in various proportions of large grain sizes on hillslopes can amplify both landslide and flood hazards in largely unquantified ways. Boulders can have an immediate destructive potential on properties and infrastructure and can hinder response and recovery by blocking access routes, posing a challenge for removal. On entering the river network, they might have far reaching effects if remobilised in high flows, damaging or destroying key infrastructure such as hydropower plants and inducing significant knock-on effects on local economies. A fundamental step towards quantification of increased hazard potential is the understanding environmental controls on boulder production. Despite their potential to enhance hazard, the probability of large boulders being produced within different landslide types has not been directly accounted for in landslide hazard mapping.

Our study focuses on the upper Bhote Koshi catchment, northeast of Kathmandu (Nepal), characterised by extreme topographic gradients, seismicity and monsoonal climate, and subjected to frequent landslides and floods. This, coupled with increased population pressure and infrastructure growth, makes the area prone to natural disasters.

We used high resolution optical imagery to map more than 11300 boulders and analysed this large dataset in combination with lithology and topography, and well as structural and landslide data, to investigate controls on boulder production and grain size distributions in different lithological, structural and geomorphic settings of the landscape.

Lithology appears to exert a significant influence on boulder sizes, with statistically significantly larger boulders observed in crystalline rocks of the Higher Himalaya Sequence than in metasedimentary rocks of the Lesser Himalaya Sequence. We also observe that the spacing of the most pervasive fracture set, parallel to foliation, influences boulder size distributions in some lithologies, whilst other dominant regional fracture sets appear not to strongly correlate with mapped boulder sizes.

Although recent studies have shown the importance of structural control on boulder sizes, our large dataset reveals that for complex, high-relief landscapes, with high erosion rates, fracture characteristics do not fully explain grain size distribution.

The type of processes involved in boulder production and transport on slopes, before reaching the river network, also appears to exert a control over grain size distributions and boulder density, with rockfall processes appearing to be responsible for producing boulders with largest sizes as opposed to rockslides, where the high energy and mode of transport is likely associated with increased fragmentation.

Analysing lithological and structural characteristics alone may not be sufficient to explain the observed distribution and would thus only give a limited insight in the enhanced hazard levels posed by boulders across different sectors of a landscape and other factors, such as distance from source and mode of transport at shorter temporal scales, must be taken into account.