

COMPUTERIZED TOMOGRAPHY (CT) OF JOINTS

Ingrid Gielen, DVM, PhD, MSc

Department of Medical Imaging & Small Animal Orthopaedics,
Faculty of Veterinary Medicine,
Ghent University, Belgium.
Email: ingrid.gielen@ugent.be

The history of musculoskeletal imaging begins with Roentgen's discovery of x rays in 1895. In small animal orthopaedics, plain radiography was the routine imaging technique for diagnostic purposes and follow up for decades, and is still routinely used in common practice.

Computerised tomography (CT) has been introduced in the seventies in human medicine and has been more readily available to veterinarians over the last decade. It is a cross-sectional imaging technique using x-rays and computers. Better soft-tissue differentiation and absence of superimposition are the major advantages of CT over conventional x-ray techniques. The first reports in small animals were published in the 1980s and concerned the normal CT brain anatomy and the CT aspect of various tumour types. Although the spatial resolution of CT images is poorer when compared with classical film-screen radiography, the cross-sectional image display and superior discrimination of tissue attenuation enables differentiation of soft tissue structures that cannot be perceived on conventional radiographs. Subtle new bone formation and bone lysis are better identified on CT images when compared with conventional radiography because of their greater physical density discrimination, the ability to manipulate the grey scale of the digital image, and the elimination of overlying structures. While a loss of 30% of bone density is often required for a lesion to be visible on conventional radiographs, CT is able to reliably detect density changes of only 0.5 – 2%.

Another advantage is that the transverse CT images can be reformatted in multiple anatomic planes.

CT enables more detailed and specific morphological diagnosis than radiography. CT greatly facilitates examining complex joint structures like the elbow and tarsus.

CT has been proved to be superior in the diagnosis of fragmented coronoid process of the **elbow joint**. Its use in the diagnosis of elbow incongruity has also been reported. With the use of CT, a clear distinction between kissing lesions and real OCD lesions is possible and is of help in treatment planning. On the transverse scans, OCD lesions present typical features being radiolucencies surrounded by a sclerotic rim. Kissing lesions on the other hand, are appearing as sclerotic stripe formed lesions without radiolucency. CT also helps in the difficult diagnosis of intercondylar fissures, which are difficult to pick up on conventional radiographs. CT can help in the diagnosis of flexor tendon enthesopathy, especially after iv. contrast resulting in enhancement in clinical lesions.

In the **stifle**, compared to radiographic examination, CT provides additional useful information in all processes where avulsions or fragmentation are involved. These disorders are not always visible on radiographs. CT proved to be extremely useful in the detection of avulsion fractures of intra-articular ligaments like the cranial cruciate ligament and the tendons of the extensor digitorum longus and the popliteus muscles. In cases of discrete OCD lesions, CT confirmed the diagnosis. Compared to radiography, the use of CT could detect many more intra-articular fragments, which provides important information to the surgeon, especially when arthroscopic treatment is envisaged. Degenerative changes can be identified in an earlier stage than on conventional radiographs.

In **tarsocrural** OCD, CT is superior in the diagnosis of lateral ridge involvement. In both, medial and lateral tarsocrural OCD, CT allows assessing the exact localisation, the size and number of the fragments. It helps in decision making when using minimal exposure

techniques to treat these lesions. In evaluating intra-articular fractures CT helps in complete evaluation of the fracture status.

In the treatment of **hip** dysplasia CT can be used to check the status of the dorsal acetabular rim which is an important criterion when triple pelvic osteotomy (TPO) is considered.

Within the **carpus**, CT can help in diagnosing radial carpal bone fractures or fissures.

In cases where treatment of **bone** tumours is considered, CT enables a more exact demarcation of the affected tissue and helps to decide to what extent the tumour has to be excised.

Arthro-CT

A rather recent new application of CT is arthro-CT, the intra-articular application of iodine containing contrast media. Before the application the contrast medium should be diluted till maximum 80-100 mg of iodine per ml. In severely inflamed joints the admixture of epinephrine can be useful to counteract the rapid absorption of the contrast medium by the inflamed, hyperaemic synovial membrane.

Within the **shoulder** joint intra-articular structures including the biceps tendon, glenohumeral ligaments and the joint cartilage can be visualised. Application of arthro-CT in the **elbow** joint is of use in appreciating the status of the articular cartilage in medial compartment syndrome. Within the **stifle** joint arthro-CT is helpful in evaluating meniscal damage and in evaluating fragment stability.

In **tarsal** OCD arthro-CT can be used to evaluate fragment stability.

The disadvantage of CT is the need for general anaesthesia and the cost for maintaining the equipment.

Suggested reading

Crawford JT, Manley PA, Adams WM. Comparison of computed tomography, tangential view radiography, and conventional radiography in evaluation of canine pelvic trauma. *Vet Radiol Ultrasound*. 2003, 44: 619-628.

Davis GJ, Kapatkin AS, Craig LE, et al. Comparison of radiography, computed tomography, and magnetic resonance imaging for evaluation of appendicular osteosarcoma in dogs. *Journal of the American Veterinary Medical Association* 2002, 220: 1171-1176.

de Bakker, Evelien, Ingrid Gielen, Annemie Van Caelenberg, Henri van Bree, and Bernadette Van Ryssen. 2014. Computed Tomographic Findings of Canine Elbow Joints Affected by Primary and Concomitant Flexor Enthesopathy. *Veterinary Radiology & Ultrasound* 55 (1): 45–55.

De Rycke LM, Gielen IM, van Bree H, et al. Computed tomography of the elbow joint in clinically normal dogs. *American Journal of Veterinary Research* 2002, 63: 1400-1407.

Fitch RB, Wilson ER, Hathcock JT, et al. Radiographic, computed tomographic and magnetic resonance imaging evaluation of a chronic long digital extensor tendon avulsion in a dog. *Vet Radiol Ultrasound*. 1997, 38: 177-181.

Gemmill T. Completing the picture: use of CT to investigate elbow dysplasia. *J Small Anim Pract*. 2004, 45: 429-430.

Gielen I, van Ryssen B, van Bree H. Computerized tomography compared with radiography in the diagnosis of lateral trochlear ridge talar osteochondritis dissecans in dogs. *Vet Comp Orthop Traumatol*. 2005, 18: 77-82.

Gielen I, Jimmy Saunders, Bernadette Van Ryssen, Henri van Bree. *Advances in the Canine Cranial Cruciate Ligament*, Editor: Peter Muir, Wiley-Blackwell, 2010. Computed tomography of the stifle, Chapter 20, p. 123-133.

Kramer A, Holsworth IG, Wisner ER, et al. Computed tomographic evaluation of canine radioulnar incongruence in vivo. *Vet Surg.* 2006, 35: 24-29.

Meyer-Lindenberg A., Heinen V., Fehr M., et al. Incomplete ossification of the humeral condyle as the cause of lameness in dogs. *Vet Comp Orthop Traumatol.* 2002, 15: 187-194.

Newberg A.H. Computed tomography of joint injuries. *Radiologic Clinics of North America* 1990, 28: 445-460.

Samii VF, Dyce J. Computed tomographic arthrography of the normal canine stifle. *Vet Radiol Ultrasound.* 2004, 45: 402-406.

Samoy, Yves, Ingrid Gielen, Annemie Van Caelenberg, Henri van Bree, Luc Duchateau, and Bernadette Van Ryssen. 2012. Computed Tomography Findings in 32 Joints Affected with Severe Elbow Incongruity and Fragmented Medial Coronoid Process. *Veterinary Surgery* 41 (4): 486–494.

van Bree, Henri, Ingrid Gielen, Jimmy Saunders, and Bernadette Van Ryssen. 2003. Computed Tomographic (CT) Features of Elbow OCD Compared with Radiography and Arthroscopy. *Veterinary Radiology & Ultrasound*, 44:371–371.