



Potential Yield of Cocksfoot (*Dactylis Glomerata*) Monocultures in Response to Irrigation and Nitrogen

A. Mills

Lincoln University, New Zealand

D. J. Moot

Lincoln University, New Zealand

R. L. Lucas

Lincoln University, New Zealand


P. D. Jamieson

Crop & Food Research, New Zealand

B. A. McKenzie

Lincoln University, New Zealand

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium4/28>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

Potential yield of cocksfoot (*Dactylis glomerata*) monocultures in response to irrigation and nitrogen

A. Mills¹, D.J. Moot¹, R.L. Lucas¹, P.D. Jamieson² and B.A. McKenzie¹

¹Agriculture and Life Sciences Division, PO Box 84, Lincoln University, Canterbury, New Zealand, Email: millsa@lincoln.ac.nz, ²Crop & Food Research, Private Bag 4704, Christchurch, New Zealand

Keywords: *Dactylis glomerata*, orchardgrass, irrigation, nitrogen, yield

Introduction Cocksfoot is a widely sown grass in temperate pastures. However, while potential yield of cocksfoot can exceed 28 t DM/ha per year, it is often restricted by water, temperature and nitrogen (N). Of these, Peri *et al.* (2002) showed that N was severely limiting in all seasons. The aim of this study was to confirm the potential yield of cocksfoot and quantify the extent of yield reductions due to environmental constraints.

Materials and methods The experiment was a split plot design with three replicates established in October 2003 onto an 8-year old 'Grasslands Wana' cocksfoot monoculture. Irrigation was the mainplot (irrigated (I) or dryland (D)) and N (0 (-N) or 800 kg/ha/y (+N)) the subplot. Irrigation maintained the soil moisture deficit above 50 mm in the top 0.5 m. Nitrogen was applied in eight split applications of 100 kg N/ha at the beginning of ~28-30 d regrowth periods during active growth. Dry matter production was determined from a 0.2 m² quadrat cut at the end of each regrowth period. The site was mown to 30 mm and herbage removed from the site.

Results Accumulated DM production was 22.6 t DM/ha per year for I+N (Figure 1a). In comparison, yield was 10.5, 15.1 and 7.5 t DM/ha/y for I+N, D+N and D-N treatments, respectively. The seasonal effect of different temperatures was accounted for by estimating yield against thermal time (Tt) (Figure 1b). Yield increased by 6.5 kg DM/°Cd above a base temperature (T_b) of 2.5°C under non-limiting N and moisture conditions. In the absence of N, yield consistently increased by 2.9 kg DM/°Cd whereas the D-N treatment produced 2.1 kg DM/°Cd. A broken stick model was used to determine slope for the D+N treatment. The first period was water stressed and D+N pastures grew at a similar rate (1.5 kg DM/°Cd) as the D-N treatments. In the second period water stress had been alleviated by rainfall and D+N treatments produced a similar rate (7.1 kg DM/°Cd) to the I+N treatment. The point of inflexion was 2015 Tt units (17/3/04) indicating an eight week lag phase after rainfall began. The broken stick model increased the R² from 88.6 to 99.3%. The LSD of the slope was 0.83.




Figure 1 Accumulated DM production of 'Grasslands Wana' cocksfoot, at Lincoln University, Canterbury, New Zealand, against time (a) and thermal time (b) Error bars are LSD ($p \leq 0.05$) for the I*N interactions

Conclusions Cocksfoot pastures grown in Canterbury have the potential to produce 23-28 t DM/ha per year. However, without irrigation or N, yield was 15.1 t DM/ha per year lower. The addition of N through legume fixation or strategic fertiliser use could double the current yield, particularly in periods of feed deficit in the autumn and spring. However, during drought, water stress may limit N uptake and yield will be proportional to total accumulated shoot N. Linear relationships between DM production and Tt provide a repeatable basis for extrapolating results for comparison of cocksfoot monoculture growth in other environments and interpreting seasonal temperature effects.

Acknowledgements A. Mills acknowledges AGMARDT, Lincoln University and the Sinclair Cummings Trust for financial assistance during her PhD study.

References

Peri, P. L., D. J. Moot, & R. J. Lucas (2002). Urine patches indicate yield potential of cocksfoot. *Proceedings of the New Zealand Grassland Association*, 64, 73-80.