



Cattle and Sheep Mixed Grazing 1: Species Equivalence

R. D. Améndola-Massiotti
Universidad Autónoma Chapingo, México

S. J. C. González-Montagna
Universidad Autónoma Chapingo, México

P. A. Martínez-Hernández
Universidad Autónoma Chapingo, México

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium4/92>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

Cattle and sheep mixed grazing 1: species equivalence

R.D. Améndola-Massiotti, S.J.C. González-Montagna and P.A. Martínez-Hernández

Universidad Autónoma Chapingo, Posgrado en Producción Animal, Programa de Investigación en Forrajes, Carretera México-Texcoco, Km 38.5, Chapingo, Edo. Méx., CP56230, México, Email: r_amendola@yahoo.com

Keywords: heifers, ewes, stocking rate, herbage intake

Introduction The effects of mixed grazing of cattle and sheep depend on stocking rate (SR) and species ratio (Nicol, 1997). Calculations of SR and species ratio require the use of species equivalence. Equivalents are often estimated in terms of intake requirements calculated on the basis of $LW^{0.75}$. Freer (1981) stated that $LW^{0.9}$ would be more appropriate for comparisons of intake requirements for maintenance of sheep and cattle. Nonetheless, Nolan & Connolly (1977) stated that the equivalence is system-specific and depends on the species being considered. The objective of this experiment was to estimate species equivalence for the evaluation of the introduction of sheep into a cattle dairy system based on grazing in temperate Mexico.

Materials and methods The experiment was carried out at Chapingo, Mexico (19°29' N, 98°54' W, 2240 m a.s.l.), between 15 March and 25 May 2000. Holstein heifers (initial LW 336±7 kg) and pregnant Suffolk ewes (initial LW 75.8±0.8 kg) were used; most ewes lambed during the experiment. There were 8 treatments, 3 heifers were used per mixture treatment and numbers of ewes varied to achieve the proportions of species shown in Table 1. Nine paddocks of *Medicago sativa* and *Dactylis glomerata* of 0.46 ha were strip-grazed for 7 days each. Three blocks of 3 paddocks were formed and treatments were randomly allotted to paddocks; only one paddock was grazed per week, the paddocks of the first and second week of each block were grazed by three treatments (each on separate areas), while the third-week paddock was grazed by the remaining two treatments. Fresh areas of pasture were allotted when already grazed areas reached 10 cm height (falling disc); no back fencing was used. The areas effectively grazed after 7 days were measured. Herbage dry matter (DM) intake was measured using chromium oxide as an external marker and *in situ* digestibility of hand plucked herbage samples.

Results Excluding the treatment of grazing by ewes only, a linear equation was calculated of area allotted (m^2/d) on number of ewes: $y = 93.4 + 4.43 x$, $R^2 = 0.94$, ($P < 0.05$) (Table 1); the intercept and the regression coefficient represent the area allotted to 3 heifers and the additional area allotted per ewe, respectively. The equivalence based on area allotted resulted in 7.0 ewes per heifer [(93.4/3)/4.43]. The equivalence based on average herbage DM intake in Table 1 (8.76 and 2.31 kg DM/animal/d) for heifers and ewes, respectively) resulted in 3.8 ewes per heifer (8.76/2.31), corresponding with the estimate of intake requirements using $LW^{0.9}$ ($336^{0.9}/75.8^{0.9}$), as recommended by Freer (1981). This difference between estimates of equivalence, expresses the higher efficiency of herbage utilisation under mixed grazing than under single species grazing by heifers, which was due to grazing by sheep below 10 cm height (data not shown here).

Table 1 Area allotted and herbage intake of heifers and ewes under single species grazing and mixed grazing with different ewes to heifer's ratios

Nr. of heifers	Nr. of ewes	Area allotted (m^2/d)	Intake per species		Total intake (kg DM/ m^2/d)
			(kg DM/heifer/d)	(kg DM/ewe/d)	
3	0	93.4	8.17 (0.49)		0.262
3	6	131.6	10.50 (0.33)	2.08 (0.15)	0.334
3	9	148.2	7.68 (0.49)	2.07 (0.15)	0.281
3	12	116.0	9.54 (0.33)	1.98 (0.15)	0.452
3	24	218.6	8.98 (0.33)	2.52 (0.15)	0.400
3	36	269.5	6.49 (0.49)	2.49 (0.15)	0.405
3	48	287.6	9.93 (0.33)	2.49 (0.15)	0.519
0	15	91.5		2.54 (0.15)	0.416

Conclusions The use of an equivalence based on DM intake (3.8 ewes/heifer), which concurs with intake requirements based on $LW^{0.9}$, would neglect benefits of more efficient pasture utilisation under mixed grazing; the use of the higher equivalent based on area allotted (7.0 ewes/heifer) should lead to better performance of the system.

References

- Freer, M. (1981). The control of food intake by grazing animals. In: F. H. W. Morley (ed.). *Grazing Animals*. Elsevier Scientific Publishing Company, Amsterdam, 105-124.
- Nicol A. M. (1997). The application of mixed grazing. In: J. G. Buchanan-Smith, L. D. Bailey and P. Mc Caughey (eds.) *Proceedings of the XVIII International Grassland Congress*, 525-534.
- Nolan, T. & J. Connolly. (1977). Mixed stocking by sheep and steers. A review. *Herbage Abstracts* 47,367-364.