



Effects of Particle Size in Forage Samples for Protein Breakdown Studies

G. Pichard

Pontificia Universidad Católica de Chile

C. Tapia

Pontificia Universidad Católica de Chile

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium4/113>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

Effects of particle size in forage samples for protein breakdown studies

G. Pichard and C. Tapia

Department of Animal Sciences, School of Agriculture, Pontificia Universidad Católica de Chile, Email: gpichard@puc.cl

Keywords: forages, particle size, proteolysis, rumen, nitrogen solubility

Introduction Coupling ruminal processes of hydrolysis and synthesis continues to be a research issue where more progress is needed. This requires the development of good protein assessment methods, particularly when representing the breakdown processes that occur in fresh pastures eaten by herbivores. Laboratory analyses need to deal with small and homogeneous samples, but the mechanical reduction of particle size may not reflect the actual digestion kinetics occurring when the original fresh forage is consumed. Such physical traits may alter the release of non-structural compounds and the penetration of microbial enzymes (Boudon *et al.*, 2002). The objective of this work was to assess in fresh samples the effect of reducing particle size upon the *in vitro* breakdown of proteins during the early rumen fermentation period.

Materials and methods Eight fresh forage samples with contrasting endopeptidase activities were subjected to different strategies for particle size reduction. Protein hydrolysis was assessed by measuring the residual neutral detergent insoluble nitrogen (NDIN) (Licitra *et al.*, 1996) and the accumulation of non-protein soluble nitrogen (NPSN) after 6 h *in vitro* rumen fermentation (IIV, Broderick, 1987). In fresh samples mastication-like damage was obtained with a device in which forage samples were pressed between two stony surfaces that simulated the animal molar surfaces. During the development of this method, microscopic observation was used in order to obtain a similar damage to that observed in samples obtained from the cardias of a fistulated adult cow fed the same type of fresh long forage. Three chopping sizes and two macerations were tested. Chopping was preceded by laboratory-mastication and further cutting to 3 cm, 1 cm or 0.25 cm; maceration was thoroughly done in a mortar with dry ice (CO₂) or liquid nitrogen. Treatments means were compared by Tukey-Kramer test at $P < 0.05$.

Results and discussion Sample size significantly affected ($P < 0.05$) the fractions of NDIN and NPSN (Figure 1), but the two macerates were essentially identical. As expected, the smaller the chopping the greater the solubility, with this effect being more pronounced in cultivars with lower endopeptidase activity. Mechanical particle comminution may facilitate access of external enzymes and activate the endogenous enzymatic system.




Figure 1 Effect of forage particle size on protein solubilisation during 6 h *in vitro* rumen fermentation

▨ : 3cm.; ▩ : 1cm.; ▪ : 0.25cm.; □ : CO₂; ▤ : N_{Liq}

Conclusion Our results show that particle size is a major source of variation when studying kinetics of protein breakdown. The mechanical damage affects the release of fermentable substrates as well as the accessibility of bacteria or their enzymes into the plant cells and the activity of plant endopeptidases. Larger particles may be more representative of actual animal behaviour, but they present practical problems for analytical purposes.

Acknowledgement This research was funded by Fondecyt Grant No. 1030918.

References

- Boudon, A., S. Mayne, J-L Peyraud & A.S. Laidlaw (2002). Effects of stage of maturity and chop length on the release of cell contents of fresh ryegrass (*Lolium perenne* L.) during ingestive mastication in steers fed indoors. *Animal Research*, 5, 349-365.
- Broderick, G.A. (1987). Determination of protein degradation rates using a rumen *in vitro* system containing inhibitors of microbial nitrogen metabolism. *British Journal of Nutrition*, 58, 463-475.
- Licitra, G., T.M. Hernández, & P.J. Van Soest (1996). Standardisation of procedures for nitrogen fractionation of ruminant feeds. *Animal Feed Science and Technology*, 57, 347-358.