




## The Performance of Cattle on Lowland Species-Rich Neutral Grassland at Three Contrasting Grazing Pressures

B. A. Griffith  
*IGER North Wyke, UK*

J. R.B. Tallwin  
*IGER North Wyke, UK*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium3/18>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

## The performance of cattle on lowland species-rich neutral grassland at three contrasting grazing pressures

B.A. Griffith and J.R.B. Tallowin

IGER North Wyke, Okehampton, Devon EX20 2SB UK, Email: jerry.tallowin@bbsrc.ac.uk

**Keywords:** species rich grassland, cattle, animal performance, botanical composition

**Introduction** Grazing is an essential management practice for maintaining the nature conservation value of lowland semi-natural neutral grassland to control succession and create different faunal habitats via structural heterogeneity within the pasture (Duffey *et al.*, 1974). However, there is a paucity of information on what would constitute a sustainable grazing intensity that will deliver the wildlife objectives and what the consequences of this management would be on growth rate of livestock and overall pasture output. An experiment was designed to quantify the ecological and agronomic consequences of imposing different grazing intensities on species-rich neutral grassland. The results will provide sward-based criteria for the integration of such species-rich grassland into commercial livestock systems.

**Methods** A randomised block experiment with three continuous grazing treatments, based on maintaining sward surface heights of 6-8cm (Severe), 8-10cm (Moderate) or 10-12cm (Lenient), using continental-cross beef heifers was imposed each year from 2000-2004 on species-rich grassland at a site in Somerset, England. The grassland comprised a mixture of *Lolium perenne*-*Cynosurus cristatus* (MG6) and *Centaurea nigra* – *Cynosurus cristatus* (MG5) grassland (Rodwell, 1992). A fourth treatment grazed to a surface sward height of 6-8 cm and receiving 130 kg N/ha per year was imposed on agriculturally improved land at the same site. This acted as a control and provided a means of comparison with a conventionally-managed beef system. Treatment targets were maintained by weekly adjustment of animals using a 'put and take' grazing system. Three 'Core' animals remained on each of the plots at all times. The following measurements were made: daily live-weight gain of the core animals, cumulative grazing day totals, weekly surface sward heights, herbage quality (by pluck sampling), sward canopy structure and botanical composition in 10 1m<sup>2</sup> fixed quadrats per paddock.

**Results** The species-rich grassland supported similar individual growth rates in the cattle to the fertilized control pasture (Table 1). Lower growth rates occurred on the severe compared with the lenient treatment on the species-rich grassland. The severe treatment carried significantly more live weight both per day and over the grazing season than the lenient and moderate treatments. The live weight carried was similar between the moderate and lenient treatments. There was no difference in live weight produced between the three grazing severities on the species-rich grassland. The fertilized pasture produced on average 91% more live weight over the grazing season than the species rich grassland.

**Table 3** Animal performance: values represent means of the four experimental years

	Severe	Moderate	Lenient	Improved	s.e.d. (df = 3)	significance
LWG (kg/day)	0.73	0.81	0.92	0.86	0.044	$P < 0.001$
LW carried per day (t/ha)	1.09	0.94	0.85	1.82	0.044	$P < 0.001$
LW carried per season (t/ha)	184.8	147	129.3	315.1	10.64	$P < 0.001$
Output (kg/ha per year)	350.6	319.8	305.5	620.2	47.76	$P < 0.001$

Although canopy structure was affected by grazing pressure on the species-rich grassland, there was no significant overall change in botanical diversity, which remained at 16-18 plant species m<sup>2</sup>, across treatments between 2000 and 2004.

**Conclusion** Under moderate to lenient grazing pressure predictable agronomic output and high individual animal growth rates can be achieved on lowland species-rich neutral grassland whilst maintaining the biodiversity interests – at least in the short term.

### References

- Duffey, E., M.G. Morris, J. Sheail, L.K. Ward, D.A. Wells & T.C.E. Wells (1974) Grassland Ecology and Wildlife Management. Chapman & Hall, London.
- Rodwell J.S. (1992) British Plant Communities. Volume 3. Grasslands and montane communities. Cambridge University Press, Cambridge.