

International Grassland Congress Proceedings

XX International Grassland Congress

An Ecological and Economic Risk Avoidance Drought Management Decision Support System

R. K. Heitschmidt USDA, Agricultural Research Service

L. T. Vermeire USDA, Agricultural Research Service

Follow this and additional works at: https://uknowledge.uky.edu/igc

Part of the Agricultural Science Commons, Agronomy and Crop Sciences Commons, Plant Biology Commons, Plant Pathology Commons, Soil Science Commons, and the Weed Science Commons This document is available at https://uknowledge.uky.edu/igc/20/satellitesymposium3/36 The XX International Grassland Congress took place in Ireland and the UK in June-July 2005. The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society. Proceedings Editor: D. A. McGilloway Publisher: Wageningen Academic Publishers, The Netherlands © Wageningen Academic Publishers, The Netherlands, 2005 The copyright holder has granted the permission for posting the proceedings here.

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

An ecological and economic risk avoidance drought management decision support system R.K. Heitschmidt and L.T. Vermeire

USDA Agricultural Research Service, Fort Keogh Livestock & Range Research Laboratory, 243 Fort Keogh Road, Miles City, MT 59301 USA, Email: rod@larrl.ars.usda.gov

Keywords: net primary production, precipitation, rangeland, temporal dynamics

Introduction Ecologists have long recognized the fundamental impacts of drought on rangeland structure and function. Simulation models have been developed to increase our understanding of these impacts as they relate to forage production, particularly for predictive purposes. Although the capacity of these models to accurately predict quantity and quality of forage produced under varying climatic conditions is often quite good, their ability to serve as an effective and proactive drought management decision support system is often limited. This is in large part because their complexity impedes their use by on-the-ground managers. The objective of this research was to develop a very simple and user-friendly drought management decision support system for rangeland managers in the Northern Great Plains of North America.

Materials and methods The proportion of herbage produced on a monthly basis was estimated using aboveground net primary production (ANPP) estimates from 14 native rangeland data sets (Heitschmidt *et al.* 1995; 1999; 2004) collected over 11 years at the Fort Keogh Livestock and Range Research Laboratory located near Miles City, MT. Correlations coefficients were calculated between monthly precipitation and annual ANPP estimates. Published (http://hprcc.unl.edu/index.html) site-specific, monthly, rainfall probabilities were also incorporated into the system to enhance predictive capacity.

Results Both total and perennial grass production were significantly (P<0.10) correlated with precipitation in January and April whereas cool-season perennial grass production, the over-whelming dominant functional group in this herbage complex, was significantly correlated with precipitation in January and May. In these analyses, the correlation in January is viewed with suspicion (i.e., a spurious correlation) as precipitation is snow and only averages 12 mm which is < 4% of the 338 mm of the annual total. The correlation with spring precipitation is biologically sound in that the sum total of precipitation in April and May averages 84 mm. In a similar study at this location, Kruse (2004) also found significant relationships between total herbage production and precipitation in April and May.

Analyses of the proportion of ANPP of perennial grasses produced on a monthly basis showed 35%, 69% and 91% was produced by 1 May, 1 June, and 1 July, respectively. The 95% confidence limit for the estimate on 1 July was 24% which indicates that at least 67% of annual production of perennial grasses would be completed by 1 July in 19 out of 20 years. Thus, in this region grazing managers can estimate appropriate end-of-growing season stocking rates by early July with a relatively high level of confidence. Moreover, confidence in these decisions can be further bolstered by examining long-term, post-1 July probabilities of rainfall. For example, the probability of receiving 25.4 mm of precipitation in July and August at Miles City, MT is 59% and 42%, respectively, whereas probability estimates for receiving 50.8 mm in July and August are 22% and 17%, respectively.

Conclusions The results provide insight into temporal dynamics of forage production in the Northern Great Plains of USA. Results demonstrate that forage production is largely a function of precipitation received prior to 1 July. Thus, proactive stocking rate adjustments can be made by early July, with considerable certainty, thereby reducing ecological and economic risks that arise from late-season forage demand/availability imbalances.

References

Heitschmidt, R. K., E. E. Grings, M. R., Haferkamp & M. G. Karl (1995). Herbage dynamics on two Northern Great Plains range sites. *Journal of Range Management*, 48, 211-217.

Heitschmidt, R. K., M. R. Haferkamp, M. G. Karl & A. L. Hild (1999). Drought and grazing: I. Effects on quantity of forage produced. *Journal of Range Management*, 52, 440-446.

Heitschmidt, R.K., K. D. Klement & M. R. Haferkamp (2004). Interaction effects of drought and grazing on Northern Great Plains rangelands. *Journal of Range Management* (in press)

Kruse, R. E. (2004). Beef cattle management decisions relating to drought in the Northern Great Plains. M.S. thesis, Department Animal & Range Sciences, Montana State University, Bozeman, MT, 138 pp.