

1 **Performance based on sEMG activity is related to psychosocial components: differences**
2 **between back and abdominal endurance tests.**

3

4 Van Damme Benedicte^{1,2}, Stevens Veerle^{1,2}, Van Tiggelen Damien^{1,2}, Perneel Christiaan³, Crombez
5 Geert⁴, Danneels Lieven²

6

7

8 ¹Center for Musculoskeletal Medicine & Rehabilitation, Military Hospital Queen Astrid, Brussels,
9 Belgium;

10 ²Department of Rehabilitation Sciences and Physiotherapy, Faculty of Medicine and Health Sciences,
11 Ghent University, Ghent, Belgium;

12 ³Department of Mathematics, Royal Military Academy, Brussels, Belgium;

13 ⁴Department of Experimental Clinical and Health Psychology, Faculty of Psychology and Educational
14 Sciences, Ghent University, Ghent, Belgium.

15

16

17 Corresponding author: Benedicte Van Damme (Benedicte.VanDamme@mil.be)

18 Military Hospital Queen Astrid, Center for Musculoskeletal Medicine & Rehabilitation, Bruynstraat 1,
19 1120 Brussels, Belgium

20 (32) 2-264-47-28

21

22

23

24

25

26

27

1 Introduction

2 Physical and psychosocial deconditioning is common in patients with chronic LBP [Brox et al, 2005;
3 Frost et al, 1998]. Research showed that the clinical evolution and prognosis of LBP is influenced by
4 psychosocial factors such as pain related fear [Crombez et al, 1999; Sullivan et al, 2009], pain
5 catastrophizing [Burton et al, 1995; Meyer et al, 2009], pain and functional self-perceived disability
6 [Vlaeyen & Linton, 2000; Thomas & France, 2008], self-reported health [Ahrens et al, 2010] and
7 depression [Ryan et al, 2010]. All these psychosocial influences make it hard to ensure that physical
8 measurements in patients with persistent non-specific LBP provide an accurate representation of the
9 real physical condition.

10 Biering Sorenson [Biering Sorenson, 1984] developed a test to measure back muscle fatigability and
11 demonstrated that a shorter position-holding time during this test predicted LBP within the next years
12 [Demoulin et al, 2006]. However, a clear influence of psychosocial factors on the endurance time of
13 the Biering-Sorenson test (B-S test) was demonstrated, in healthy subjects as well as in patients with
14 chronic LBP [Larivière et al, 2010; Coorevits et al, 2008]. In order to obtain a more objective
15 measurement of the intrinsic muscle fatigue, a measure for the decrease of the median frequency
16 during static submaximal testing, recorded by surface electromyography (sEMG), can be used
17 [Coorevits et al, 2008]. A regression coefficient of the median frequency slope towards lower
18 frequencies can be used as a fatigue index for the investigated muscle. This regression coefficient is
19 often normalized by the intercept, which is the crossing point of the slope and the Y-axis [Konrad,
20 2005]. A combination of the use of the regression coefficient of the median frequency, time to
21 exhausting and self-reporting questionnaires, could be useful to understand the different influencing
22 factors on physical performance.

23 In contrast to the almost systematic use of the B-S test and its modified versions to analyze back
24 muscle endurance, much more variation in position and procedure is observed in abdominal
25 endurance tests. Brox et al. [2005] utilized a dynamic abdominal endurance test to demonstrate that
26 deconditioning (deterioration of impairment and disability) was more related to psychological
27 measures and physical measures of abdominal and back muscle endurance than to general

1 cardiovascular fitness (estimated by a submaximal bicycle ergometer test). However, static endurance
2 tests are preferred since variation in muscle length throughout the test can alter the frequency content
3 of the EMG signal [Mannion et al, 1996; Sparto et al, 1999].

4 In conclusion, research on the relation between psychosocial deconditioning and physical
5 performance is scarce and often lacks standardized and specialized physical testing or sufficient
6 patients. Therefore, the aim of the present study was to investigate the influence of psychosocial
7 components on the duration of standardized back and abdominal static endurance tests in patients
8 with persistent non-specific LBP. Determining good performance may lead to a better understanding
9 of factors that make patients perform less than they physically can during these tests. The hypothesis
10 was that underperformance during back and abdominal endurance tests in patients with persistent
11 non-specific LBP is influenced and may be predicted by psychosocial components.

12 **Materials and Methods**

13 *Patients*

14 Three-hundred and thirty-two patients (291 men and 41 women; military and civilian employees) from
15 19 to 63 years old with persistent non-specific LBP, with or without leg pain, were included in this
16 study after screening by a medical doctor specialized in physical medicine at the Military Hospital
17 Queen Astrid in Brussels. When needed additional investigations were performed to assure that no
18 identifiable specific anatomical or neurophysiological causative factors were present. Pregnant
19 women, patients with previous spinal surgery, nerve root entrapment with neurological deficit, patients
20 with specific LBP diseases and patients with a BMI \geq 33 were excluded from this study. Study
21 population characteristics are shown in Table 1. The study was approved by the Ethical Committee of
22 the Ghent University.

23

24 *Psychosocial assessment questionnaires*

25 All subjects received self-report questionnaires before performing physical tests. General questions
26 related to the sociodemographic status and characteristics of their LBP were registered. Several

1 validated questionnaires were used as a measure of physical and/or psychological influence of the
2 LBP on daily life. A numerical scale was used to indicate the average pain over the last week. Pain
3 Catastrophizing was measured by the Pain Catastrophizing Scale (PCS) and its three subscales
4 (rumination, magnification and helplessness) [Sullivan et al, 1995]. The Hospital Anxiety or Depression
5 Scale (HADS) [Zigmond & Snaith, 1983] determined the levels of anxiety and depression that a patient
6 was experiencing. The Distress Risk Assessment Method (DRAM) consisting of the Modified Somatic
7 Perceptions Questionnaire (MSPQ) [Mannion et al, 2011; Main, 1983] and the Modified Zung
8 Depression Index (MZDI) [Mannion et al, 2011; Zung et al, 1965] were used as screening tools for
9 depression and somatic pain respectively. The Short Form Health Survey (SF-36) [Ware & Gandek,
10 1998] and its different subscales gave a general measure of health, both mental and physical
11 components. Depending on the mother tongue, the participants filled in the questionnaires in Dutch or
12 French (validated versions).

13

14 *Physical Assessment – endurance tests*

15 An isometric endurance test for the abdominal muscles (AE test) [Stevens et al, 2008] was performed
16 (Figure 1). The patient was seated on a bench in a straight-knee position, with the trunk unsupported
17 at a 45-degree angle. The hands were placed on the shoulders with the arms flexed alongside the
18 trunk. Neutral position of the head and lumbar spine was respected.

19 For the evaluation of the isometric endurance of the trunk extensor muscles patients performed a
20 modified version of the B-S test (Figure 2) [Demoulin et al, 2006; Coorevits et al, 2008; Demoulin et al,
21 2013; Stevens et al, 2006]. The patient was lying on a bench in a prone position with the anterior-
22 superior iliac spines at the rotation point of the bench. The lower body was fixed to the table by two
23 straps, one around the pelvis and one on the ankles. The patients had to hold their hands touching
24 their foreheads, with their elbows out to the side and leveled with the trunk. Patients were also
25 instructed to hold their head in a neutral position, and to look downward. The test was started with the
26 upper body in an about 70° downward position so that a concentric contraction of the trunk extensor
27 muscles was needed initially to reach the horizontal position.

1 The patient was asked to isometrically maintain these positions. This was checked by visual
2 evaluation. The time the patient held these positions was recorded. Verbal encouragement was given
3 by the tester during both endurance tests to ensure that the maximal effort was produced by the
4 patient.

5 *EMG recording*

6 sEMG was used to quantify the rate of development of muscle fatigue. After appropriate skin
7 preparation in order to get a good electrode-skin contact and to reduce skin impedance, 8 pairs of
8 circular Ag/AgCl sensor surface electrodes (Ambu® Blue Sensor M, Ambu A/S, Ballerup, DK) were
9 placed parallel to the muscle fibres [Ng et al, 1998], bilaterally, of 2 deep stabilizing and 2 superficial
10 torque producing abdominal and back muscle groups as follows: The inferior fibres of the internal
11 obliques (IO) (midway between the anterior iliac spine and the symphysis pubis, above the inguinal
12 ligament) [Stevens et al, 2008; Van Damme et al, 2012], the external obliques (EO) (15 cm lateral to
13 the umbilicus) [Cholewicki et al, 1997; Hubley & Vezina, 2002], the lumbar multifidus (LMF) (above
14 and below the L5 spinous process, parallel to the line between the posterosuperior iliac spine and the
15 L1–L2 interspinous space) [Danneels et al, 2001; Ng et al, 1997], the thoracic part of the iliocostalis
16 (ICLT) (above and below the L1 level, midway between the midline and the lateral aspect of the body)
17 [Stevens et al, 2008; Van Damme et al, 2012; Danneels et al, 2002]. A reference electrode was placed
18 on the thoracic cage.

19 Electromyographic signals were recorded using an 8-channel sEMG system (Myosystem 2000,
20 Noraxon U.S.A. Inc., Scottsdale, AZ) connected to a computer. The raw sEMG signals were recorded
21 at a sampling rate of 1000Hz, amplified (overall gain 1000, common mode rate rejection ratio 115 dB),
22 filtered to produce a bandwidth of 10 – 500 Hz and analog digital conversion (12-bit resolution) was at
23 1000 Hz. Each recorded sEMG signal that was stored on the computer was divided in intervals of 5 s.
24 The median frequency (MF) of the sEMG power spectrum was calculated at each 5-s interval with
25 Fast-Fourier Transform algorithms using the Noraxon MyoResearch software v2.11. During a
26 sustained isometric contraction of the muscles, the MF spectral shifts indicate local muscle fatigue and
27 decrease over time [Sung et al, 2009]. The MF was defined as the frequency that divided the

1 spectrum into two equal areas. Finally, linear regression analyses were performed on the calculated
2 MF's as a function of time. The initial MF (MF_{init}) was defined as the intercept of the regression line
3 and the MF slope (MF_{slope}) was determined as the slope of the regression line. MF_{slope} was normalized
4 to MF_{init} [Biering Sorenson, 1984], to deal with inter subject and inter location differences in
5 subcutaneous tissue layers. The MF_{slope} , normalized to MF_{init} , gave a measure of fatigability for each
6 recording site for each individual [Mannion et al, 2011; Coorevits et al, 2008].

7

8 *Data analysis*

9 The outcome of each questionnaire was calculated following the instructions of the original designers.
10 SPSS 21.0 was used for statistical analyses. Statistical significance was accepted at the 5% level.

11 Patients were dichotomized as underperformers and performers, by comparing their real endurance
12 time, to the expected time of endurance derived from the normalized MF_{slope} (NMF_{slope}). This was done
13 for the B-S test and the AE test separately. The NMF_{slope} [Coorevits et al, 2008] provided a measure of
14 fatigability for each muscle on both sides, for each subject. The endurance time was regressed on the
15 NMF_{slope} for the most fatigable region of the LMF during the B-S test and for the most fatigable region
16 of the IO during the AE test. The deep stabilizing muscles were selected because, in this group, they
17 showed generally a steeper slope than the superficial torque-producing muscles. Endurance appears
18 to be limited by the most fatigable region of a muscle group [Mannion & Dolan, 1994]. Stepwise
19 multiple regression analyses (Probability of F-to-enter ≤ 0.05 ; Probability of F-to-remove ≥ 0.10) were
20 used to determine a linear model for the endurance time of the two endurance tests. The influence of
21 gender on the slope and the intercept of the regression line were analyzed. For this reason, the
22 dummy variable gender and $gender * NMF_{slope}$ were introduced as two extra independent variables into
23 the model. Using the regression equation derived from the group data, the "expected" endurance time
24 for a given subject was determined for both exercises. Depending on whether their real endurance
25 time fell short (a) or was equal to or exceeded (b) the expected time, they were classified as
26 underperformers (a) or as performers (b) for that specific exercise. A similar method was used by
27 Mannion et al. (2011), but in the present model gender was also added in the regression model.

1
2
3
4
5
6
7
8
9

For the AE test and the B-S test separately, independent *t*-tests were performed to examine the differences between underperformers and performers on the different variables and self-reporting questionnaires. Stepwise multiple regression analysis (Probability of F-to-enter ≤ 0.05 ; Probability of F-to-remove ≥ 0.10) was performed using the variables that were found significantly different between the two groups in a bivariate analysis as predictor variables. For the regression analyses, the normality of the errors, independency of the errors and the homoscedasticity assumption were verified.

10 **Results**

11 The linear regression analysis for the B-S test (Figure 3) provided two different parallel regression lines, one for each gender. The regression line for the female patients was situated below the model of the male patients. In the AE test no significant differences were found between the regression line of male and female patients. This means that the prediction line (Figure 4) was the same for male and female subjects.

16 *Biering-Sorenson Test*

17 Independent *t*-tests between the underperformers and performers on the B-S test (Table 2) showed that the mean age ($p=0.007$) and mean BMI ($p<0.001$) of the underperformers were higher than these of the performers. No significant difference was found in hours of sport per week (sport hrs/week) between the two groups ($p=0.988$). The two groups had a similar NMF_{slope} of the LMF ($NMF_{slope}LMF$), but the performers group demonstrated a significantly longer endurance time on the B-S test ($p<0.001$). The scores on some of the subscales of the SF-36 were significantly higher for the performers: Physical Function ($p<0.001$), General Health ($p<0.001$) and total Physical Health ($p=0.046$).

25 Stepwise multiple regression analysis (Table 3) of all the variables that were found significant in 26 bivariate analysis revealed that a greater BMI ($\beta=-221.748$; $p<0.001$) and lower scores on the Physical

1 Function subscale of the SF-36 ($\beta=355.209$; $p=0.001$) were significant predictors of underperformance
2 on the B-S test. These components explained more than 10% ($R^2=0.113$) of the variance.

3 Abdominal Endurance Test

4 Independent *t*-tests between the underperformers and performers on the AE test (Table 4) showed
5 that the underperformers group of the AE test had a significantly higher BMI ($p=0.014$). No significant
6 difference was found in sport hrs/week between the two groups ($p=0.482$). The two groups had a
7 similar NMF_{slope} of the IO ($NMF_{\text{slope}}|IO$), but the performers group demonstrated a significantly longer
8 endurance time on the AE test ($p<0.001$). The underperformers group scored significantly higher on
9 the DRAM MZDI ($p=0.018$) and on the PCS scale ($p=0.020$), as well as on two of its subscales:
10 magnification ($p=0.025$) and helplessness ($p=0.012$). This group showed also significantly lower
11 scores on the SF-36 and some of its subscales: General Health ($p=0.009$), Social Functioning
12 ($p=0.048$), Role Emotional ($p=0.008$), total Physical Health ($p=0.25$) and total Mental Health ($p=0.006$).

13
14 Stepwise multiple regression analysis (Table 5) of all the variables that were found significant in
15 bivariate analysis revealed that a higher BMI ($\beta=-2748.282$; $p=0.005$) and a lower score on the total
16 mental health subscale of the SF-36 ($\beta=547.415$; $p=0.011$) were predictive for underperformance on
17 the AE test. These components explained more than 4% of the variance ($R^2=0.049$).

18 **Discussion**

19 The present study examined the relationship between psychosocial components and
20 underperformance -performance less than expected- on abdominal and back muscle endurance tests
21 in patients with persistent non-specific LBP. The main findings were that underperformance on back
22 muscle endurance tests was more likely to be influenced by physical components (self-reporting on
23 physical health), whereas abdominal muscle endurance tests seemed more affected by psychosocial
24 components (self-reporting of mental health). Performing less than expected as in underperformance,
25 does not necessarily involve low performance, but may even present high performance in well-trained
26 subjects.

1

2 *Back muscle endurance*

3 For the B-S test underperformers had statistical significant lower scores on different subscales of the
4 SF-36: Physical Function, General Health and total Physical Health. All these scales reflect a physical
5 component. So, we could suggest that more negative perceptions of physical health could induce
6 underperformance on the B-S test.

7 The results described above are in line with observations in elderly patients with chronic LBP [Ledoux
8 et al, 2012]. Functional capacity, measured by endurance and peak torque during prone and side
9 bridge positions, was very much dependent on physical components (physical activity and disability
10 levels), and not on psychosocial components as depression scores or pain catastrophizing. In the
11 current study baseline physical activity, reported in sport hrs / week was not different between the
12 group underperformers and good performers, but a more detailed questioning about physical activity
13 could have given more information. Disability levels in se were not reported in the current study, but
14 reporting on physical health was significantly different between the two groups in the B-S test.

15 Demoulin et al. [2013] did not find any association between endurance time on the B-S test and pain-
16 related fear measures, but no subdivision was made between performance groups. Mannion et al.
17 [2011] found that underperformers on the B-S test had more negative back beliefs, greater
18 psychological disturbance, greater catastrophizing and lower exercise self-efficacy compared with the
19 performers. In contrast to the results of the present study, the underperformers showed significant
20 greater catastrophizing and psychological disturbance. Larivière et al. [2010] suggested that pain
21 catastrophizing is related to outcome on the B-S test. These different results might be explained by the
22 population characteristics. In contrast to all active working patients in the present study, Larivière et al.
23 [2010] observed a population in which 10 of the 27 patients were not at work (7 due to back
24 problems). In the study of Mannion et al. [2011] the group symptoms were similar to the present
25 group, but there were a lot more female subjects (57%). In contrast to the present study, Mannion et
26 al. [2011] did not take gender into account while making the subdivisions in underperformers and
27 performers group. However, several studies demonstrated a clear gender difference in muscle

1 fatigability during the B-S test. Kankaanpää et al. [1998], for example, demonstrated sex differences in
2 paraspinal muscle fatigability during the B-S test and explained this by gender differences in muscle
3 anatomy and physiology. Therefore, gender cannot be thought away in determining the performance
4 group. In addition, gender differences have been found consistently in catastrophizing, with women
5 reporting significantly higher scores than men on the PCS [D'Eon et al, 2004; Thorn et al, 2004]. In
6 several studies, performance was evaluated by a Functional Capacity Evaluation (FCE) or other
7 physical tests and no strong relationship was found between the FCE-physical outcome and
8 psychological factors [Schiphorst et al, 2008; Reneman et al, 2007; Smeets et al, 2007]. This is in line
9 with the results found in the present study, although the physical tests were very different.

10 *Abdominal muscle endurance*

11 Underperformers on the AE test had significantly higher scores on the DRAM, the MZDI and on the
12 PCS and two of its subscales: magnification and helplessness. The SF-36 and some of its subscales
13 (general health, social functioning, role emotional, total physical health and mental health) of this
14 group showed lower scores, indicating more problems with these health components. Patients who
15 were more emotionally distressed and patients with high levels of catastrophizing tended to
16 underperform on the AE test. No studies were found which specifically investigated the relation
17 between underperformance on abdominal muscle tests and psychological distress. Brox et al. [2005]
18 investigated healthy controls and patients with sub-acute and chronic LBP and found that patients with
19 chronic LBP demonstrated significantly higher pain, self-reported functional disability and fear-
20 avoidance and lower abdominal and back muscle endurance times than sub-acute patients. A study of
21 Sullivan et al. [2002] revealed that pain catastrophizing was significantly predictive for low
22 performance on a repeated lifting task in patients with chronic LBP. However, performing less than
23 expected as in underperformance, does not necessarily involve low performance, but may even
24 present high performance in e.g. well-trained subjects.

25 Multiple regression analysis demonstrated that a greater BMI and lower scores on the Physical
26 Function subscale of the SF-36 were significant predictors of underperformance on the B-S test and
27 that a higher BMI and lower scores on the total mental health subscale of the SF-36 were predictive

1 for underperformance on the AE test. It is striking that performance on back muscle endurance tests
2 was more likely influenced by physical components, while the abdominal muscle endurance test
3 seemed more affected by psychological components. Different subjective observations were made,
4 which could explain this discrepancy. Some aspects, typically observed during the AE test, could
5 influence the patient's motivation to perform. In general, subjects reported more pain discomfort during
6 the AE test. The sitting position is for many patients a pain provocative position. Although the neutral
7 position of the back was respected during the AE test, patients did not like to sit in this position and
8 needed a lot more motivation to hold it. Previous research demonstrated that pain catastrophizing is in
9 strong relation with pain [Sullivan et al, 2001]. Catastrophizing, in the absence of pain, is not
10 associated with impaired physical function or with reduced motivation to perform physical maneuvers.
11 However, under conditions where movement is associated with pain, catastrophizing appears to
12 contribute to a reduction in physical outcome [Sullivan et al, 2002] and this could explain
13 underperformance. In contrast to the lumbar extension control position during the B-S test, the
14 abdominal endurance demanded more control to avoid lumbar flexion. Higher catastrophizing and
15 emotionally distressed patients may fear flexion positions more than extension postures. Flexion-
16 related pain disorders are the most common disorders observed in clinical practice [O'Sullivan, 2000].
17 In the sitting flexion endurance test, patients also experienced vibrations in the abdominal muscles;
18 these vibrations were generally not observed during the B-S test. Due to the uncomfortable sensation
19 of these vibrations, the patient's performance could have been disturbed. In addition, the visual focus
20 was also very different between the two endurance tests. During the B-S test, the patients' vision was
21 orientated to the ground, which created less visual disturbance in comparison to the view of the test
22 environment, equipment and testers during the AE test. Highly test-anxious persons often divide their
23 attention between task-irrelevant and task-relevant variables and become more distractible, whereas
24 low-test anxious persons focus their attention more fully on the task [Pijpers et al, 2005; Eubank et al,
25 2000]. This could explain why patients who were more psychologically distressed tend to
26 underperform on the AE-test compared to subjects who were not psychologically distressed, and
27 could also clarify why we did not find such a difference in the B-S test. However, the relative shift in
28 attention away from environmental cues, and towards internal monitoring of feelings, thoughts and

1 movements – which is more likely to happen in the B-S-test - has also been demonstrated to have a
2 detrimental effect on motor performance especially in anxious people [Maxwell et al, 2000; Janelle,
3 2002].

4 Patients who are more psychologically distressed could be more influenced by all these different
5 factors. Further exploration of the causes of this statistical difference between the AE test and the B-S
6 test is needed.

7 Although the results of the multiple regression analyses were in accordance to the clear statistical
8 significant differences between the group of the underperformers and the good performers, the effect
9 sizes of the regression analyses were small. Only about 10% in the B-S test and 4% in the AE test of
10 the variance could be explained by the defined variables. In sociological and biological measurements
11 high percentages are almost never achieved, however it is often possible to identify about 25% of the
12 variance of a relationship [Botz & Doering, 2002]. This was not achieved in the present study.

13 However, interpretation of linear regression analysis is not straightforward [Schneider et al, 2010] and
14 some statisticians believe that being able to improve an outcome by 4% can be clinically useful.

15 In previous research, BMI was observed as a significant predictor of endurance time in the B-S test
16 [Mbada et al, 2009]. This could be explained by the association between BMI and the rate of MF
17 decrease during this test. A higher BMI was shown to create a greater fatigability and a lower
18 endurance time of the paraspinal muscles [Kankaanpää et al, 1998]. The present study examined this
19 more thoroughly, and found in addition a direct association between BMI and underperformance, both
20 on the abdominal as well as on the back muscle endurance test. A high BMI was even a predictor of
21 underperformance on these two endurance tests. In the literature, BMI is often associated with lower
22 physical performance but also with a higher occurrence of psychosocial problems [Vaidya, 2006;
23 Fabricatore & Wadden, 2004].

24 The method used in this study to determine the group of underperformers and performers could be
25 questioned, because controversy exists concerning the use of MF_{slope} to determine the time to
26 exhaustion [Bouillard et al, 2012]. Because parameters derived from the EMG power spectrum are
27 less dependent on the force level of the muscle compared to amplitude parameters and appear to be

1 more sensitive to the myoelectric manifestations of muscle fatigue [Ng et al, 1996; Potvin & Bent,
2 1997], the use of spectral EMG variables as fatigue index was preferred over the EMG amplitude
3 parameters [Roy & Oddsson, 1998]. MF has been suggested to be the most suitable parameters for
4 describing localized muscle fatigue [Kankaanpää et al, 1998; Larivière et al, 2002] and may provide an
5 objective measure of muscle fatigue [Arnall et al, 2002; Mannion et al, 1998; Ng et al, 1996; Nicolaisen
6 & Jorgensen, 1985; Roy et al, 1989]. In contrast, Bouillard et al [2012] demonstrated that other
7 physiological (e.g. motor unit synchronization) and non-physiological (e.g., change in fiber pennation
8 angle, change in the muscular temperature) factors may affect the sEMG signal and thus its changes
9 over time. Other parameters have also been proposed, such as the Dimitrov spectral index, which was
10 shown to be higher correlated with endurance time than the initial slope of the MF in a small healthy
11 population performing isotonic biceps brachii contractions [Lee et al, 2011]. In addition, the MF was
12 demonstrated to be not a suitable indicator for dynamic contractions [Van Dieën et al, 1996] and
13 contractions at approximately 9-10% of maximal voluntary isometric contraction (MVIC) [Gonzalez-Izal
14 et al, 2010]. However, since most of the studies analyzing back muscle fatigue have used MF [Allison
15 & Henry, 2001; Biedermann et al, 1990; Champagne et al, 2008; Dederling et al, 2000; Elving et al,
16 1999; Kankaanpää et al, 1998; Larivière et al, 2002; Mannion et al, 2011; Müller et al, 2010; Ng et al,
17 2002; Peach & McGill, 1998; Süüden et al, 2008], and the endurance tests were not dynamically
18 performed and achieved higher levels than 9 to 10% of MVIC, the method applied in the present study
19 was chosen. However, the accuracy of the method using MF_{slope} to estimate the endurance time is
20 not known and thus some participants may not have been well classified. In addition, the low R^2
21 values could induce some doubt about the usefulness of the linear regression models. But R^2 is not a
22 way to decide if a model is good or bad and that prudence is warranted in interpreting R^2 values
23 (Achen, 1977; Kennedy, 2008; Goldberger, 1991; King, 1986).

24

25 Finally, the aim of this study was not to make a model to predict underperformance on endurance test,
26 but to demonstrate, in addition to previous research, that endurance time on the B-S test and the AE
27 test is not a clear measure of intrinsic muscle fatigue. Psychosocial components and reporting on
28 physical health are interacting with performance on these tests and should be taken into account while

1 making conclusions based on these tests. In addition, differences found between the AE test and B-S
2 test in this study indicate that these interactions could be test-specific. So these results cannot be
3 generalized to all performance measures. There is a clear need for the use of sEMG in measuring
4 intrinsic muscle fatigue in research and/or clinical settings. Using sEMG in the clinical practice is not
5 feasible, but clinical practitioners, who want to measure endurance, should be aware that psychosocial
6 components could influence performance.

7

8 **References**

- 9 Achen, Christopher H. Measuring Representation: Perils of the Correlation Coefficient. *American*
10 *Journal of Political Science* 1977;21:805-815
- 11 Ahrens C, Schiltewolf M, Wang H. Health-related quality of life (SF-36) in chronic low back pain and
12 comorbid depression. *Schmerz* 2010;24:251
- 13 Allison GT, Henry SM. Trunk muscle fatigue during a back extension task in standing. *Manual Therapy*
14 2001;6(4):221-228
- 15 Arnall FA, Koumantakis GA, Oldham JA, Cooper RG. Between-days reliability of electromyographic
16 measures of paraspinal muscle fatigue at 40, 50 and 60% levels of maximal voluntary contractile
17 force. *Clinical Rehabilitation* 2002;16:761-771
- 18 Biedermann HJ, Shanks GL, Forrest WJ, Inglis J. Median frequency estimates of paraspinal muscles:
19 reliability analysis. *Electromyography and Clinical Neurophysiology* 1990;30:83-88
- 20 Biering Sorenson. Physical Measurements as Risk Indicators for Low-Back Trouble Over a One-Year
21 Period. *Spine* 1984;9:105-223
- 22 Bortz J, Doering N. *Forschungsmethoden und evaluation*. 3rd ed., Berlin, Germany: Springer; 2002

1 Bouillard K, Frère J, Hug F, Guével A. Prediction of time-to-exhaustion in the first dorsal interosseous
2 muscle from early changes in surface electromyography parameters. *Muscle & Nerve* 2012;45(6):835-
3 840

4 Brox J, Storheim K, Holm I, Friis A, Reikeras O. Disability, Pain, Psychological factors and physical
5 performance in healthy controls, patients with sub-acute and chronic low back pain: A case-control
6 study. *J Rehabil Med* 2005;37:95-99

7 Burton, A. Kim PhD; Tillotson, K. Malcolm; Main, Chris J. PhD; Hollis, Sally MSc. Psychosocial
8 Predictors of Outcome in Acute and Subchronic Low Back Trouble. *Spine* 1995;20:635-748

9 Champagne A, Descarreaux M, Lafond D. Back and hip extensor muscles fatigue in healthy subjects :
10 task-dependency effect of two variants of the Sorensen test. *European Spine Journal*
11 2008;17(12):1721-1726

12 Coorevits P, Danneels L, Cambier D, Ramon H, Vanderstraeten G. Assessment of the validity of the
13 Biering-Sorensen test for measuring back muscle fatigue based on EMG median frequency
14 characteristics of back and hip muscles. *J Electromyogr Kines* 2008;18:997-1005

15 Cholewicki J, Panjabi M, Khachatryan A. Stabilizing Function of Trunk Flexor-Extensor Muscles
16 Around a Neutral Spine Posture. *Spine* 1997; 22:2207-2212

17 Crombez G, Vlaeyen J, Heuts P, Lysens R. Pain-related fear is more disabling than pain itself:
18 evidence on the role of pain-related fear in chronic back pain disability. *Pain* 1999;80:329-339

19 Danneels L, Coorevits P, Cools A, Vanderstraeten G, Cambier D, Witvrouw E, De Cuyper H.
20 Differences in electromyographic activity in the multifidus muscle and the iliocostalis lumborum
21 between healthy subjects and patients with sub-acute and chronic low back pain. *Eur Spine J*
22 2002;11:13-19

- 1 Danneels L, Vanderstraeten G, Cambier D, Witvrouw E, Stevens V, De Cuyper H. A functional
2 subdivision of hip, abdominal, and back muscles during asymmetric lifting. *Spine* 2001;26:114-121
- 3 Dederling A, Roos af Hjelmsäter M, Elfving B, Harms-Ringdahl K, Németh G. Between-days reliability
4 of subjective and objective assessments of back extensor muscle fatigue in subjects without lower-
5 back pain. *Journal of Electromyography and Kinesiology* 2000;10(3):151-158
- 6 Demoulin C, Huijnen, Somville PR, Grosdent S, Salamun I, Crielaard JM et al. Relationship between
7 different measures of pain-related fear and physical capacity of the spine in patients with chronic low
8 back pain. *Spine J* 2013;13:1039-1047
- 9 Demoulin C, Vanderthommen M, Duysens C, Crielaard J. Spinal muscle evaluation using the
10 Sorensen test: a critical appraisal of the literature. *Joint Bone Spine* 2006;73:43-50
- 11 Elfving B, Németh G, Arvidsson I, Lamontagne M. Reliability of EMG spectral parameters in repeated
12 measurements of back muscle fatigue. *Journal of Electromyography and Kinesiology* 1999;9:235-243
- 13 Eubank M, Collins D, Smith N. The influence of anxiety direction on processing bias. *Journal of Sport*
14 *and Exercise Psychology* 2000;22:292-306
- 15 D'Eon JL, Harris CA, Ellis JA. Testing factorial validity and gender invariance of the pain
16 catastrophizing scale. *J Behav Med* 2004;27:361-372
- 17 Fabricatore A, Wadden T. Psychological aspects of obesity. *Clin Dermatol* 2004;22:332-337
- 18 Frost H, Lamb S, Klaber Moffett J, Fairbank J, Moser, J. A fitness programme for patients with chronic
19 low back pain: 2-year follow-up of a randomised controlled trial. *Pain* 1998;75:273-279
- 20 Goldberger A. *A Course in Econometrics*. Harvard University Press, 1991, p. 177

1 Gonzalez-Izal M, Malanda A, Navarro-Amezqueta I, Gorostiaga EM, Mallor F. EMG spectral indices
2 and muscle power fatigue during dynamic contractions. *Journal of Electromyography and Kinesiology*
3 2010;20(2):233-240

4 Hubley-Kozey C, Vezina M. Muscle activation during exercises to improve trunk stability in men with
5 low back pain. *Arch Phys Med Rehabil* 2002;83:1100-1108

6 Janelle C. Anxiety, arousal and visual attention: A mechanisitic account of performance variability.
7 *Journal of Sports Sciences* 2002;20:237-51

8 Kankaanpää M, Laaksonen D, Taimela S, Kokko S, Airaksinen O, Hänninen O. Age, sex, and body
9 mass index as determinants of back and hip extensor fatigue in the isometric Sørensen back
10 endurance test. *Arch Phys Med Rehabil* 1998;79:1069-1075

11 Kennedy, Peter. *A Guide to Econometrics*. San Francisco, CA: Wiley-Blackwell, 2008.
12

13 King G. How Not to Lie with Statistics: Avoiding Common Mistakes in Quantitative Political Science.
14 *American Journal of Political Science* 1986;30(3):666–687

15 Konrad P. *ABC of EMG*. Powered by Noraxon Inc, USA. 2005, p. 50

16 Larivière C, Arsenault AB, Gravel D, Cagnon D, Loisel P. Evaluation of measurement strategies to
17 increase the reliability of EMG indices to assess back muscle fatigue and recovery. *Journal of*
18 *Electromyography and Kinesiology* 2002;12:91-102

19 Larivière C, Bilodeau M, Forget R, Vadeboncoeur R, Mecheri H. Poor back muscle endurance is
20 related to pain catastrophizing in patients with chronic low back pain. *Spine* 2010;35:1178-1186

21 Ledoux E, Dubois JD, Descarreaux M. Physical and psychosocial predictors of functional trunk
22 capacity in older adults with and without low back pain. *J Manip Physiol Ther* 2012;35:338-345

1 Lee K, Lee S, Choi A, Choi C, Mun J. Endurance time prediction of biceps brachii muscle using
2 Dimitrov spectral index of surface electromyogram during isotonic contractions. *International Journal of*
3 *Precision Engineering and Manufacturing* 2011;12(4):711-717

4 Main C. The modified somatic perception questionnaire (MSPQ). *J Psychosom Res* 1983;27:503-514

5 Mannion AF, Dolan P. Electromyographic Median Frequency changes during isometric contraction of
6 the back extensors to fatigue. *Spine* 1994;19:1223-1229

7 Mannion A, Dolan P, Adams M. Psychological questionnaires: do “abnormal” scores precede or follow
8 first-time low back pain? *Spine* 1996;21:2603-2611

9 Mannion AF, Dumas GA, Stevenson JM, Cooper RG. The influence of muscle fiber size and type
10 distribution on electromyographic measures of back muscle fatigability. *Spine* 1998;23(5):576-584

11 Mannion A, O’Riordan D, Dvorak J, Masharawi Y. The relationship between psychological factors and
12 performance on the Biering-Sorensen back muscle endurance test. *Spine J* 2011;11:849-857

13 Maxwell J, Masters R, Eves F. From novice to know-how: A longitudinal study of implicit motor
14 learning. *Journal of Sports Sciences* 2000;18:111-20

15 Mbada C, Ayanniyi O, Adedoyin R. Reference values of static back extensor muscle endurance in
16 healthy Nigerian adults. *Med Princ Pract* 2009;18:345-350

17 Meyer K, Tschopp A, Sprott H, Mannion A. Association between catastrophizing and self-rated pain
18 and disability in patients with chronic low back pain. *J Rehabil Med* 2009;41:620-625

19 Müller R, Strässle K, Wirth B. Isometric back muscle endurance: An EMG study on the criterion validity
20 of the Ito test. *Journal of Electromyography and Kinesiology* 2010;20(5):845-850

21 Ng J, Kippers V, Richardson C. Muscle fibre orientation of abdominal muscles and suggested surface
22 EMG electrode positions. *Electromyogr Clin Neurophysiol* 1998;38:51-58

1 Ng JK-F, Richardson CA. Reliability of electromyographic power spectral analysis of back muscle
2 endurance in healthy subjects. Archives of Physical Medicine and Rehabilitation 1996;77:259-264

3 Ng J, Richardson CA, Jull GA. Electromyographic amplitude and frequency changes in the iliocostalis
4 lumborum and multifidus muscles during a trunk holding test. Phys Ther 1997;77:954-961

5 Ng JK-F, Richardson CA, Parnianpour M, Kippers V. Fatigue-related changes in torque output and
6 electromyographic parameters of trunk muscles during isometric axial rotation exertion. An
7 investigation in patients with back pain and in healthy subjects. Spine 2002;27(6):637-646

8 Nicolaisen T, Jorgensen K. Trunk strength, back muscle endurance and low-back trouble.
9 Scandinavian Journal of Rehabilitation Medicine 1985;17:121-127

10 O'Sullivan P. Lumbar segmental 'instability': clinical presentation and specific stabilizing exercise
11 management. Manual Therapy 2000:2-12

12 Peach JP, McGill SM. Classification of low back pain with the use of spectral electromyogram
13 parameters. Spine 1998;23(10):1117-1123

14 Potvin JR, Bent LR. A validation of techniques using surface EMG signals from dynamic contractions
15 to quantify muscle fatigue during repetitive tasks. Journal of Electromyography and Kinesiology
16 1997;7(2):131-139

17 Pijpers J, Oudejans R, Bakker F. Anxiety-induced changes in movement behaviour during the
18 execution of a complex whole-body task. The Quarterly Journal of Experimental Psychology Section
19 A: Human Experimental Psychology 2005;58:421-45

20 Reneman MF, Schiphorts Preuper HR, Kleen M, Geertzen J, Dijkstra P. Are pain intensity and pain
21 related fear related to functional capacity evaluation performances of patients with chronic low back
22 pain? J Occup Rehabil 2007;17:247-258

- 1 Roy SH, De Luca CJ, Casavant DA. Lumbar muscle fatigue and chronic low back pain. *Spine*
2 1989;14(9):992-1001
- 3 Roy SH, Oddsson LIE. Classification of paraspinal muscle impairments by surface electromyography.
4 *Physical Therapy* 1998;78(8):838-851
- 5 Ryan C, Gray G, Newton M, Granat M. The relationship between psychological distress and free-living
6 physical activity in individuals with chronic low back pain. *Manual Ther* 2010;15:185-189.
- 7 Schiphorst Preuper HR, Reneman MF, Boonstra AM, Dijkstra P, Versteegen G, Geertzen J, Brouwer
8 S. Relationship between psychological factors and performance-based and self-reported disability in
9 chronic low back pain. *Eur Spine J* 2008;17:1448-1456
- 10 Schneider A, Hommel G, Blettner M. Linear regression analysis part 14 of a series on evaluation of
11 scientific publications. *Deutsches Arzteblatt International* 2010;107:776-782
- 12 Smeets R, van Geel A, Kester A, Knottnerus J. Physical capacity tasks in chronic low back pain: What
13 is the contributing role of cardiovascular capacity, pain and psychological factors? *Disabil Rehabil*
14 2007;29:577-586
- 15 Sparto P, Parnianpour M, Barria E, Jagadeesh J. Wavelet Analysis of Electromyography For Back
16 Muscle Fatigue Detection During Isokinetic Constant-Torque Exertions. *Spine* 1999;24: 1757-1858
- 17 Stevens V, Bouche K, Mahieu N, Coorevits P, Vanderstraeten G, Danneels L. Trunk muscle activity in
18 healthy subjects during bridging stabilization exercises. *BMC Musculoskelet Disord* 2006;7:75
- 19 Stevens V, Parlevliet T, Coorevits P, Mahieu N, Bouche K, Vanderstraeten G, Danneels L. The effect
20 of increasing resistance on trunk muscle activity during extension and flexion exercises on training
21 devices. *J Electromyogr Kines* 2008;18:434-445
- 22 Sullivan M, Bishop S, Pivik J. The Pain Catastrophizing Scale: Development and validation. *Psychol*
23 *Assessment* 1995;7:524-532

1 Sullivan M, Rodgers W, Wilson P, Bell G, Murray T, Fraser S. An experimental investigation of the
2 relation between catastrophizing and activity intolerance. *Pain* 2002;100:47-53

3 Sullivan M, Thibault P, Andrikonyte J, Butler H, Catchlove R, Larivière C. Psychological influences
4 on repetition-induced summation of activity-related pain in patients with chronic low back pain. *Pain*
5 2009;141:70-78

6 Sullivan M, Thorn B, Haythornthwaite J, Keefe F, Martin M, Bradley LA, Lefebvre J. Theoretical
7 Perspectives on the relation between catastrophizing and pain. *Clin J Pain* 2001;17:52-64

8 Sung PS, Lammers AR, Danial P. Different parts of erector spinae muscle fatigability in subjects with
9 and without low back pain. *Spine J* 2009;9:115–20

10 Süüden E, Erelina J, Gapeyeva H, Pääsuke M. Low back muscle fatigue during Sorensen endurance
11 test in patients with chronic low back pain: relationship between electromyographic spectral
12 compression and anthropometric characteristics. *Electromyography and Clinical Neurophysiology*
13 2008;48(3-4):185-192

14 Thomas J, France C. The relationship between pain-related fear and lumbar flexion during natural
15 recovery from low back pain. *Eur Spine J* 2008;17:97-103

16 Thorn BE, Clements KL, Wald LC, Dixon KE, Kersh BC, Boothby JL, Chaplin WF. Personality factors
17 in the explanation of sex differences in pain catastrophizing and response to experimental pain. *Clin J*
18 *Pain* 2004;20:275-282

19 Vaidya V. Psychosocial aspects of obesity. *Adv Psychosom Med* 2006;27:73-85

20 Van Damme B, Stevens V, Van Tiggelen D, Duvigneaud N, Neyens E, Danneels L. Velocity of
21 isokinetic trunk exercises influences back muscle recruitment patterns in healthy subjects. *J*
22 *Electromyogr Kines* 2012;23:378–386

- 1 Van Dieën JH, Boke B, Oosterhuis W, Toussaint HM. The influence of torque and velocity on erector
2 spinae muscle fatigue and its relationship to change of electromyogram spectrum density. *European*
3 *Journal of Applied Physiology* 1996;72(4):310-315

- 4 Vaidya V. Psychosocial aspects of obesity. *Adv Psychosom Med* 2006;27:73-85

- 5 Vlaeyen J, Linton S. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of
6 the art. *Pain* 2000;85:317-332

- 7 Ware J, Gandek B. Overview of the SF-36 health survey and the international quality of life
8 assessment (IQOLA) project. *J Clin Epidemiol* 1998;51:903-912

- 9 Zigmond S, Snaith, R. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand*
10 1983;67:361–370

- 11 Zung W, Richards C, Short M. Self-rating depression scale in an outpatient clinic: further validation of
12 the SDS. *Arch Gen Psychiatry* 1965;13:508

- 13

1 **Caption to illustrations**

2 Figure 1 The abdominal endurance test

3 Figure 2 A modified version of the Biering-Sorensen test

4 Figure 3 Plot of the real endurance time (ms) versus the normalized slope of the multifidus, for male
5 and female patients, during a modified version of the Biering-Sorensen Test. The predicted endurance
6 time is given by the regression equation for each gender separately. The regression line for the male
7 was: $Y = 122244 + (6638904 * x)$. For female subjects the regression line was $Y = (122244 - 18289) + ($
8 $6638904 * x)$.

9 Figure 4 Plot of the real endurance time (ms) versus the normalized slope of the internal obliques, for
10 male and female patients, during an abdominal endurance test. The predicted endurance time is given
11 by the regression equation ($Y = 135858,264 + (11104256,44 * x)$), which was found identical between the
12 two genders.

Fig 1



Fig 2





