

Short Communication

Bee pathogens found in *Bombus atratus* from Colombia: A case study



Viviana Gamboa^a, Jorgen Ravoet^b, Marleen Brunain^b, Guy Smagghe^c, Ivan Meeus^c, Judith Figueroa^a, Diego Riaño^d, Dirk C. de Graaf^{b,*}

^a Faculty of Veterinary Medicine and Zootechnology, National University of Colombia, Bogotá, Colombia

^b Laboratory of Molecular Entomology and Bee Pathology, Faculty of Sciences, Ghent University, Ghent, Belgium

^c Department of Crop Protection, Faculty of Bioscience Engineering, Ghent University, Ghent, Belgium

^d Faculty of Sciences, Military University Nueva Granada, Bogotá, Colombia

ARTICLE INFO

Article history:

Received 24 February 2015

Revised 21 May 2015

Accepted 22 May 2015

Available online 29 May 2015

Keywords:

Bombus atratus

Pathogen

Acute Bee Paralysis Virus

Black Queen Cell Virus

Sacbrood Virus

Lake Sinai Virus

Apicystis bombi

Critidia bombi

Nosema ceranae

ABSTRACT

Bombus atratus bumblebees from Colombia that were caught in the wild and from breeding programs were screened for a broad set of bee pathogens. We discovered for the first time Lake Sinai Virus and confirmed the infection by other common viruses. The prevalence of *Apicystis bombi*, *Critidia bombi* and *Nosema ceranae* was remarkably high. According to other studies the former two could have been co-introduced in South America with exotic bumble bees as *Bombus terrestris* or *Bombus ruderatus*. Given the fact that none of these species occur in Colombia, our data puts a new light on the spread of these pathogens over the South American continent.

© 2015 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The neotropical bumblebee *Bombus atratus* is one of the 12 *Bombus* species from Colombia. Besides, it is one of those species that allowed to develop technologies for breeding in captivity in order to be applied for the pollination of different cultivated crops and fruits (Liévano et al., 1991).

In the context of the global decline of pollinators, there is an increasing interest in determining the pathogen diversity of bumblebee species. Imported bee species are considered a putative threat for native plant-pollinator networks through the spillover of pathogens (Meeus et al., 2011; Arbetman et al., 2013). Especially in North and South America there is considerable data to suggest that pathogens are an important driver of bumble bee decline (Cameron et al., 2011; Arbetman et al., 2013) and that at least some of them could have been introduced with exotic, Palearctic bumble bee species (*Bombus terrestris*; *Bombus ruderatus*) commercially reared and shipped worldwide for pollination services (Plischuk et al., 2011; Arbetman et al., 2013). Given the

fact that none of these allochthonous species occur in Colombia makes this country particularly interesting for further research.

Here we present a comprehensive pathogen screen of a limited set of samples of *B. atratus* from Colombia that were caught in the wild and from breeding programs. We discovered for the first time Lake Sinai Virus (LSV) in this bumblebee species and found a strain of *Apicystis bombi* with most resemblance to the European clade.

2. Materials and methods

2.1. Sampling

In 2013, five foraging *B. atratus* specimens were collected with entomological nets in the municipality of Chía (Colombia). We also sampled seven nests reared and maintained in the facilities of the campus of the Military University Nueva Granada located in the municipality of Cajicá (Colombia). Two bumblebees were collected from each colony. All bumblebees were killed by freezing and preserved in RNAlater (Ambion) at 4 °C for transport to Belgium.

2.2. Nucleic acid extraction

The bumblebees were individually homogenized in PBS. RNA was isolated using the RNeasy Lipid Tissue kit (Qiagen) and DNA

* Corresponding author.

E-mail address: dirk.degraaf@ugent.be (D.C. de Graaf).

was extracted using the DNeasy Blood and Tissue kit (Qiagen), all in accordance to the manufacturer's instructions.

2.3. PCR analysis

Around 500 ng RNA was retro-transcribed using random hexamer primers with the RevertAid First Strand cDNA Synthesis Kit (Thermo Scientific). Using previously described PCR protocols, samples were screened for the presence of bacteria, microsporidia, protists and viruses (detailed description is given as Supplementary data).

Amplicons of the ABPV complex, *A. bombi* ITS region and *Spiroplasma* spp. were cloned using the TOPO TA Cloning Kit for sequencing (Invitrogen, USA) and isolated with the GeneJet Plasmid Miniprep kit (Thermo Scientific). DNA sequences obtained by direct sequencing of amplicons or by sequencing plasmids with M13 primers were BLAST-searched for confirmation.

2.4. Sequence analysis

Trypanosomatid amplicons were aligned with 18S sequences from *Critidilia bombi* and *Critidilia expoeki* (unpublished information) in order to resolve their identity.

Sequences of *A. bombi* ITS regions were aligned in Geneious R7 using clustalW and subsequently trimmed. The best fitting nucleotide substitution model with maximum likelihood was selected using the Bayesian Information Criterion as implemented in MEGA6. Phylogenetic trees were inferred via maximum likelihood (ML) with PhyML 3.0 (Guindon et al., 2010) using the HKY85 model with approximate non-parametric likelihood ratio test branch support based on a Shimodaira-Hasegawa-like (aLRT SH-like) procedure (Anisimova and Gascuel, 2006).

3. Results and discussion

An overview of the detected pathogens is given in Table 1. All tested bumblebees were found infected by at least four different species of pathogen, but up to eight species could be found in a single specimen. Deformed Wing Virus (DWV) and *Nosema ceranae* were omnipresent (both in captive-bred and wild samples), and *C. bombi* was discovered in all captive-bred samples. Overall, *A. bombi* was detected in 12 out of 19 samples (63%) and LSV in 13 out of 19 samples (68%). To the best of our knowledge this is the first report of the occurrence of LSV in *B. atratus*. After its first discovery in the USA (Runckel et al., 2011), the field of action of LSV was subsequently extended to honey bees and solitary bees of Europe (Ravoet et al., 2013, 2014; Granberg et al., 2013), and now to bumblebees of the northwest of South America.

The common honey bee viruses like Acute Bee Paralysis Virus (ABPV), Black Queen Cell Virus (BQCV), DWV and Sacbrood Virus (SBV) were already detected in South American honey bees (Antúnez et al., 2006; Teixeira et al., 2008; Mendoza et al., 2014). They are now considered multi-host pathogens, infecting many arthropod species even beyond the pollinator community (Li et al., 2011; Peng et al., 2011; Zhang et al., 2012; Levitt et al., 2013). All except ABPV were found in *B. atratus* from Brazil (Reynaldi et al., 2013). The scope of the microsporidium *N. ceranae* is so far only restricted to Hymenopteran insect species. Originating from Asian honey bees (*Apis cerana*) (Fries et al., 1996), it was subsequently found in European honey bees (*Apis mellifera*) (Higes et al., 2006; Huang et al., 2007), bumblebees (Graystock et al., 2013; Fürst et al., 2014) including *B. atratus* (Plischuk et al., 2009) and solitary bees (Ravoet et al., 2014). Its occurrence in South American honey bees is well-documented (Klee et al., 2007; Invernizzi et al., 2009; Martinez et al., 2012;




Fig. 1. Phylogenetic analysis of *Apicystis bombi* internal transcribed spacers (ITS). This region includes the partial 18S rRNA, complete ITS 1, 5.8S rRNA gene, ITS 2, and the partial 28S rRNA gene. Samples originating from Argentina are indicated in purple, from Colombia in blue, from Europe in black, from Mexico in red and the universal haplotype in green. Branch support is designated by aLRT statistics. Only values higher than 70% are shown.

Table 1

Overview of detected pathogens in *Bombus atratus* from Colombia, sampled along 2013. ABPV, Acute Bee Paralysis Virus; BQCV, Black Queen Cell Virus; DWV, Deformed Wing Virus; LSV, Lake Sinai Virus; SBV, Sacbrood Virus; *Nosema ceranae*, *N. ceranae*; *Apicystis bombi*; *Critidilia bombi*, *C. bombi*; *Spiroplasma apis*, *S. apis*.

Sample	Virus					<i>N. ceranae</i>	Microsporidia		Bacteria	Pathogen diversity
	ABPV	BQCV	DWV	LSV	SBV		<i>A. bombi</i>	<i>C. bombi</i>		
Nest 702,1	0	1	1	1	0	1	0	1	0	5
Nest 702,2	0	0	1	1	1	1	0	1	0	6
Nest 706,1	0	1	1	0	0	1	1	1	0	5
Nest 706,2	0	1	1	1	0	1	1	1	0	6
Nest 718,1	0	1	1	0	1	1	0	1	1	6
Nest 718,2	0	1	1	0	0	1	0	1	0	4
Nest 720,1	0	1	1	1	0	1	0	1	0	5
Nest 720,2	1	1	1	1	1	1	0	1	0	7
Nest 724,1	1	1	1	1	0	1	1	1	0	8
Nest 724,2	0	1	1	0	0	1	1	1	0	5
Nest FITXC,1	0	1	1	1	0	1	0	1	0	5
Nest FITXC,2	0	0	1	0	0	1	1	1	0	4
Nest R102,1	0	1	1	0	0	1	1	1	0	5
Nest R102,2	0	1	1	1	0	1	1	1	0	6
Subtotal: number (%)	2 (14)	12 (86)	14 (100)	8 (57)	3 (21)	14 (100)	7 (50)	14 (100)	1 (7)	
Wild 1,1	0	0	1	1	0	1	1	0	0	4
Wild 1,2	0	0	1	1	0	1	1	1	0	5
Wild 2,1	0	0	1	1	0	1	1	1	0	5
Wild 2,2	0	1	1	1	0	1	1	1	0	6
Wild 3	0	1	1	1	0	1	1	1	0	6
Subtotal: number (%)	0 (0)	2 (40)	5 (100)	5 (100)	0 (0)	5 (100)	5 (100)	4 (80)	0 (0)	
Total: number (%)	2 (11)	14 (74)	19 (100)	13 (68)	3 (16)	19 (100)	12 (63)	18 (95)	1 (5)	

[Teixeira et al., 2013](#); [Mendoza et al., 2014](#)). Recent reports demonstrated that the prevalence of DWV and *N. ceranae* in honey bees and bumblebees is linked, and that *Apis* is the likely source of DWV spillover to wild pollinators ([Fürst et al., 2014](#)).

Our discovery of the neogregarine *A. bombi* in *B. atratus* from Colombia was most interesting. Moreover, we could successfully amplify its ITS regions in some samples and found strong sequence similarity with several published *A. bombi* haplotypes (Genbank accession number of the found sequences: KP055606–KP055610). A phylogenetic analysis revealed that they are part of a clade with the universal haplotype (Genbank: KF322212) and several European haplotypes (Genbank: KF322207–KF322208, KF322210–KF322212; see Fig. 1). This suggests a European origin of the Colombian *A. bombi*. This is even more remarkable as this protist was reported to be possibly co-introduced with *B. terrestris* or *B. ruderatus* in Argentina ([Plischuk et al., 2011](#); [Arbetman et al., 2013](#)). Given the fact *B. terrestris* does not occur in Colombia, our data suggests that the route of invasion of *A. bombi* here should be different. This could indicate that the cosmopolitan prevalence of *A. bombi* was caused by another host than *B. terrestris*. Second the mutation rate of the ITS marker within neogregarine parasites is perhaps too low to resolve the differences between the native Palearctic and Neotropic parasites. Indeed, the high prevalence of *A. bombi* in introduced species in Argentina could have occurred by a spillback principle, in which native parasites find a new reservoir in exotic bees. This introduced reservoir can then disrupt natural host parasite associations. We conclude that there is a need for more molecular markers, with controls of regions without bee transports, to calculate the molecular mutation rate.

The prevalence of the trypanosome *C. bombi* was remarkably high (95%). In 2009 this protist was still lacking in an Argentinean survey covering five of the eight known native bumblebee species, including *B. atratus*, though it was found in 21.6% of the invasive *B. terrestris* ([Plischuk and Lange, 2009](#)). Recently, it was found in the native bumblebees in Chile ([Schmid-Hempel et al., 2014](#)). We believe that our data on the occurrence of *A. bombi* and *C. bombi* in an environment that lacks the invasive *Bombus* species, puts a new light on the spread of these pathogens over the South American continent.

Acknowledgments

The authors wish to thank the group of Beekeeping Science and Technology from the National University of Colombia and the group of research of Wild Bees Biodiversity and Ecology from the Military University Nueva Granada, particularly to the head of the group José Ricardo Cure. Picture of *Bombus atratus* was from Graciela Alcantara (<http://www.ecoregistros.org/Graciela Alcantara>).

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.jip.2015.05.013>.

References

- Anisimova, M., Gascuel, O., 2006. Approximate likelihood-ratio test for branches: a fast, accurate, and powerful alternative. *Syst. Biol.* 55, 539–552.
- Antúnez, K., D'Alessandro, B., Corbella, E., Ramallo, G., Zunino, P., 2006. Honeybee viruses in Uruguay. *J. Invert. Pathol.* 93 (1), 67–70.
- Arbetman, M.P., Meeus, I., Morales, C.L., Aizen, M.A., Smagghe, G., 2013. Alien parasite hitchhikes to Patagonia on invasive bumblebee. *Biol. Invasions* 15 (3), 489–494.
- Cameron, S.A., Lozicer, J.D., Strange, J.P., Koch, J.B., Cordes, N., Soltend, L.F., Griswold, T.L., 2011. Patterns of widespread decline in North American bumble bees. *Proc. Natl. Acad. Sci. USA* 108 (2), 662–667.
- Fries, I., Feng, F., Da Silva, A., Slemenda, S.B., Pieniazek, N., 1996. *Nosema ceranae* n. sp. (Microspora, Nosematidae), morphological and molecular characterization of a microsporidian parasite of the Asian honey bee *Apis cerana* (Hymenoptera, Apidae). *Eur. J. Protistol.* 32 (3), 356–365.
- Fürst, M.A., McMahon, D.P., Osborne, J.L., Paxton, R.J., Brown, M.J.F., 2014. Disease associations between honeybees and bumblebees as a threat to wild pollinators. *Nature* 506, 364–366.
- Granberg, F., Vicente-Rubiano, M., Rubio-Guerri, C., Karlsson, O.E., Kukielka, D., Belák, S., Sánchez-Vizcaíno, J.M., 2013. Metagenomic detection of viral pathogens in Spanish honeybees: co-infection by aphid lethal paralysis, Israel acute paralysis and lake sinai viruses. *PLoS One* 8 (2), e57459.
- Graystock, P., Yates, K., Darvill, B., Goulson, D., Hughes, W.O.H., 2013. Emerging dangers: deadly effects of an emergent parasite in a new pollinator host. *J. Invert. Pathol.* 114 (2), 114–119.
- Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59, 307–321.
- Higes, M., Martín-Hernández, R., Meana, A., 2006. *Nosema ceranae*, a new microsporidian parasite in honeybees in Europe. *J. Invert. Pathol.* 92 (2), 93–95.

- Huang, W.F., Jiang, J.H., Chen, Y.W., Wang, V.H., 2007. A *Nosema ceranae* isolate from the honeybee *Apis mellifera*. *Apidologie* 38, 30–37.
- Invernizzi, C., Abud, C., Tomasco, I.H., Harriet, J., Ramallo, G., Campá, J., Katz, H., Gardiol, G., Mendoza, Y., 2009. Presence of *Nosema ceranae* in honeybees (*Apis mellifera*) in Uruguay. *J. Invert. Pathol.* 101 (2), 150–153.
- Klee, J., Besana, A.M., Genersch, E., Gisder, S., Nanetti, A., Tam, D.Q., Chinh, T.X., Puerta, F., Ruz, J.M., Kryger, P., Message, D., Hatjina, F., Korpela, S., Fries, I., Paxton, R.J., 2007. Widespread dispersal of the microsporidian *Nosema ceranae*, an emergent pathogen of the western honey bee, *Apis mellifera*. *J. Invert. Pathol.* 96 (1), 1–10.
- Levitt, A.L., Singh, R., Cox-Foster, D.L., Rajotte, E., Hoover, K., Ostiguy, N., Holmes, E.C., 2013. Cross-species transmission of honey bee viruses in associated arthropods. *Virus Res.* 176, 232–240.
- Li, J., Peng, W., Wu, J., Strange, J.P., Boncristiani, H., Chen, Y., 2011. Cross-species infection of deformed wing virus poses a new threat to pollinator conservation. *J. Econ. Entomol.* 104 (3), 732–739.
- Liévano, A., Ospina, R., Nates, G., 1991. Distribución altitudinal del género *Bombus* en Colombia (Hymenoptera: Apidae). *Trianea* 4, 541–550.
- Martínez, J., Leal, G., Conget, P., 2012. *Nosema ceranae* an emergent pathogen of *Apis mellifera* in Chile. *Parasitol. Res.* 111 (2), 601–607.
- Meeus, I., Brown, M.J., de Graaf, D.C., Smagghe, G., 2011. Effects of invasive parasites on bumble bee declines. *Conserv. Biol.* 25 (4), 662–671.
- Mendoza, Y., Antúnez, K., Branchicella, B., Anido, M., Santos, E., Invernizzi, C., 2014. *Nosema ceranae* and rRNA viruses in European and Africanized honeybee colonies (*Apis mellifera*) in Uruguay. *Apidologie* 45, 224–234.
- Peng, W.J., Li, J., Boncristiani, H., Strange, J.P., Hamilton, M., Chen, Y., 2011. Host range expansion of honey bee black queen cell virus in the bumble bee, *Bombus huntii*. *Apidologie* 42, 650–658.
- Plischuk, S., Lange, C.E., 2009. Invasive *Bombus terrestris* (Hymenoptera: Apidae) parasitized by a flagellate (Euglenozoa: Kinetoplastea) and a neogregarine (Apicomplexa: Neogregarinorida). *J. Invert. Pathol.* 102 (3), 261–263.
- Plischuk, S., Martín-Hernández, R., Prieto, L., Lucía, M., Botías, C., Meana, A., Abrahamovich, A.H., Lange, C., Higes, M., 2009. South American native bumblebees (Hymenoptera: Apidae) infected by *Nosema ceranae* (Microsporidia), an emerging pathogen of honeybees (*Apis mellifera*). *Environ. Microbiol. Rep.* 1 (2), 131–135.
- Plischuk, S., Meeus, I., Smagghe, G., Lange, C.E., 2011. *Apicystis bombi* (Apicomplexa: Neogregarinorida) parasitizing *Apis mellifera* and *Bombus terrestris* (Hymenoptera: Apidae) in Argentina. *Environ. Microbiol. Rep.* 3 (5), 565–568.
- Ravoet, J., Maharramov, J., Meeus, I., De Smet, L., Wenseleers, T., Smagghe, G., de Graaf, D.C., 2013. Comprehensive bee pathogen screening in Belgium reveals *Critchidia mellifcae* as a new contributory factor to winter mortality. *PLoS One* 8 (8), e72443.
- Ravoet, J., De Smet, L., Meeus, I., Smagghe, G., Wenseleers, T., de Graaf, D.C., 2014. Widespread occurrence of honey bee pathogens in solitary bees. *J. Invert. Pathol.* 122, 55–58.
- Reynaldi, F., Sguazza, G.H., Albicoro, F.J., Pecoraro, M.R., Galosi, C.M., 2013. First molecular detection of co-infection of honey bee viruses in asymptomatic *Bombus atratus* in South America. *Braz. J. Biol.* 73 (4), 797–800.
- Runckel, C., Flenniken, M.L., Engel, J.C., Ruby, J.G., Ganem, D., Andino, R., DeRisi, J.L., 2011. Temporal analysis of the honey bee microbiome reveals four novel viruses and seasonal prevalence of known viruses, *Nosema*, and *Critchidia*. *PLoS One* 6, e20656.
- Schmid-Hempel, R., Eckhardt, M., Goulson, D., Heinemann, D., Lange, C., Plischuk, S., Escudero, L.R., Salathé, R., Scriven, J.J., Schmid-Hempel, P., 2014. The invasion of southern South America by imported bumblebees and associated parasites. *J. Anim. Ecol.* 83 (4), 823–837.
- Teixeira, E.W., Chen, Y., Message, D., Pettis, J., Evans, J.D., 2008. Virus infections in Brazilian honey bees. *J. Invert. Pathol.* 99 (1), 117–119.
- Teixeira, E.W., Dos Santos, L.G., Sattler, A., Message, D., Marques, M.L.F.A., Fonseca, M.M., Grassi-Sella, M.L., Franco, T.M., 2013. *Nosema ceranae* has been present in Brazil for more than three decades infecting Africanized honey bees. *J. Invert. Pathol.* 114 (3), 250–254.
- Zhang, X., He, S.Y., Evans, J.D., Pettis, J.S., Yin, G.F., Chen, Y.P., 2012. New evidence that deformed wing virus and black queen cell virus are multi-host pathogens. *J. Invert. Pathol.* 109 (1), 156–159.