

University of Dundee

Monocyclic Nitro-heteroaryl Nitrones with Dual Mechanism of Activation

Pacheco, Juliana Da Silva; Costa, Débora De Souza; Cunha-Júnior, Edézio Ferreira; Andrade-Neto, Valter Viana; Fairlamb, Alan H.; Wyllie, Susan

Published in:
ACS Medicinal Chemistry Letters

DOI:
[10.1021/acsmchemlett.1c00193](https://doi.org/10.1021/acsmchemlett.1c00193)

Publication date:
2021

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Pacheco, J. D. S., Costa, D. D. S., Cunha-Júnior, E. F., Andrade-Neto, V. V., Fairlamb, A. H., Wyllie, S., Goulart, M. O. F., Santos, D. C., Silva, T. L., Alves, M. A., Costa, P. R. R., Dias, A. G., & Torres-Santos, É. C. (2021). Monocyclic Nitro-heteroaryl Nitrones with Dual Mechanism of Activation: Synthesis and Antileishmanial Activity. *ACS Medicinal Chemistry Letters*, 12(9), 1405-1412. <https://doi.org/10.1021/acsmchemlett.1c00193>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Monocyclic Nitro-heteroaryl Nitrones with Dual Mechanism of Activation: Synthesis and Antileishmanial Activity

Juliana da Silva Pacheco^{1,4,†}, Débora de Souza S. Costa^{2,†}, Edézio Ferreira Cunha-Júnior^{1,5}, Valter Viana Andrade-Neto¹, Alan Fairlamb⁴, Susan Wyllie⁴, Marília O. F. Goulart⁶, Danyelle C. Santos⁶, Thaisa L. Silva⁶, Marina A. Alves⁷, Paulo R. R. Costa², Eduardo Caio Torres-Santos^{1*}, Ayres G. Dias^{3*}

¹Laboratório de Bioquímica de Tripanosomatídeos, Instituto Oswaldo Cruz, FIOCRUZ, RJ, Brazil; ²Laboratório de Química Bioorgânica, Instituto de Pesquisas de Produtos Naturais, Universidade Federal do Rio de Janeiro, RJ, Brazil; ³Universidade do Estado do Rio de Janeiro, Centro de Tecnologia e Ciências, Departamento de Química Orgânica, RJ, Brazil. ⁴University of Dundee, School of Life Sciences, Division of Biological Chemistry and Drug Discovery, Dow Street, Dundee, DD1 5EH, Scotland, United Kingdom; ⁵Laboratório de Imunoparasitologia, Unidade Integrada de Pesquisa em Produtos Bioativos e Biociências, Campus UFRJ-Macaé; ⁶Instituto de Química e Biotecnologia, Universidade Federal de Alagoas; ⁷Laboratório de Apoio ao Desenvolvimento Tecnológico, Universidade Federal do Rio de Janeiro, RJ, Brazil.

[†]Equal contribution

ABSTRACT: New 5-nitro-furannitrones (**1a-c**), 5-nitro-thiophennitrones (**2a-c**) and 4-nitroarylnitronone **4** were synthesized in one step, in high yields. Compounds **1a** and **1b** had the most potent leishmanicidal activity against intracellular amastigote forms of *L. amazonensis* and *L. infantum* (from 0.019 μ M to 2.76 μ M), with excellent selectivity (from 39 to 5673). The comparison of the leishmanicidal activity in promastigotes of wild type *L. Donovanii* with those obtained in species overexpressing nitroreductases NRT1 or NRT2 shows that **1a,b** are activated by both, a feature that could slow the development of resistance. The redox potential (E_{redox}) of these compounds obtained by cyclic voltammetry (0.57 V vs. Normal Hydrogen Electrode, for both compounds), compared to the other compounds shows that the reduction of the nitro group is modulated by the nitronone group. Formation of reactive oxygen species was not observed in promastigotes of *L. infantum* and murine macrophages. Compound **1b** was administered orally to mice infected by *Leishmania infantum* and reduced the parasite load on the spleen by 76.6% and 95.0% with doses of 50 and 100mg/kg, respectively, administered twice a day, for five days. In the liver, the parasite load suppression was above 75% with either treatment.

1. Introduction

Leishmaniasis is among the seven most important tropical diseases, primarily affecting poor people in developing countries. Environmental changes such as deforestation and population migration contribute to its broad geographic distribution and increasing incidence. Cutaneous, mucocutaneous and visceral manifestations are the main clinical forms of this neglected disease, caused by protozoan parasites belonging to the genus *Leishmania* (fam. *Trypanosomatidae*).^{1,2} According to the World Health Organization (WHO), cutaneous leishmaniasis is the most common form, with an estimated 0.6 million to 1 million new cases occurring annually. The visceral form is fatal if left untreated with an estimated 50,000 to 90,000 new cases occurring each year.³

The chemotherapeutic potential of nitroheteroaromatic compounds for treating diseases caused by trypanosomatids was first reported in the 1950s. Nitrofurazone was reported as having activity against *Trypanosoma bruceigambiense*⁴ followed by several other analogues in the following decades.⁵⁻⁹ However, severe toxicity issues limited their use in the clinic.¹⁰ Despite the fact that many nitro heteroaromatic compounds are still used today as therapeutics with acceptable safety profiles, the nitro group is regarded not only as a toxicological alert, but also as a no-go for many medicinal chemists. However, the knowledge of the role of specific and unique nitroreductases in their mode of action has brought nitro-compounds to the pipeline again.¹⁰⁻¹² The role of

nitro group in medicinal chemistry has been recently reviewed.^{13,14}




Figure 1. Nitroheteroaryl compounds with antiparasitic activity.

Currently, the arsenal of antitrypanosomal nitro-heteroaromatic drugs in clinical use is composed of benznidazole for treatment of the acute stage of Chagas Disease, nifurtimox for congenital Chagas' disease and nifurtimoxin combination with eflornithine

(NECT) for the late stage of Human African trypanosomiasis. These compounds are part of the list of essential medicines of WHO since 2009^{15,16} (Figure 1).

Regarding leishmaniasis, fexinidazole (Figure 1) was the first nitro-heteroaromatic compound to reach clinical trials^{17,18}, and its use was recently registered for sleeping sickness, as the first oral drug (DNDi, 2018¹⁹). The antitrypanosomal and leishmanicidal activities of 8-nitroquinolin-2-(1H)-one and derivatives was also reported (Pedron *et al.*, 2018²⁰). Delamanid, a bicyclic imidazole derivative approved for the treatment of drug resistant tuberculosis, showed potent activity against *Leishmania donovani in vitro* and *in vivo* and has been proposed for the oral therapy of visceral-leishmaniasis.¹²

All these compounds are pro-drugs and need to be activated by a nitroreductase. NTR1, a mitochondrial oxygen-insensitive nitroreductase, is known to transform monocyclic nitro-compounds such as nifurtimox into a hydroxylamine and an acyclic unsaturated nitrile, responsible for the biological effect. Wyllie, Fairlamb and co-workers identified a putative oxygen-sensitive NAD(P)H oxidase as another activating nitroreductase (NTR2)¹³ which preferentially reduces bicyclic nitro-compounds. Over expression of NTR2 in wild type parasites rendered cells hyper-sensitive to bicyclic nitro-heteroaromatic compounds, such as delamanid, but only marginally to the monocyclic nitro-heteroaromatic compounds, nifurtimox and fexinidazole sulfone, known to be activated by NTR1.²¹




Figure 2. Nitro-heteroaryl nitrones (1-3), 4 and aldehydes (5-7).

In this note we report the synthesis and leishmanicidal activity of the compounds shown in Figure 2. The importance of the new structural pattern bearing nitro and nitrone groups in the same molecule (**1a-c** and **2a-c**) is evidenced by comparing the potency and selectivity of these compounds with **3a** and **3b**. The bifunctional compound **4** was also studied to highlight the importance of the nitro-heteroaromatic moiety for the leishmanicidal activity. The nitro-aldehydes **5a,b** and **7**, used as starting materials to prepare the nitro-aryl nitrones, were also evaluated.


Promastigotes and amastigotes of *L. amazonensis* and *L. infantum* were used in our study. *Leishmania donovani* overexpressing NTR1 or NTR2 were also used and the results compared with those obtained with wild type.

It is worth mentioning that leaving systems have a good tolerance for the nitrone function. For example, the detoxification of natural amines and enamines is accomplished by the oxidation to the corresponding nitrone metabolites^{22,23}. Also, nitrones have been detected as metabolites of several drugs and two in widespread use, chloridiazepoxide²⁴ and minoxidil²⁵, have this function in their chemical structure. In addition, nitrones are also important spin-scavengers, allowing the detection and identification of highly unstable free radical intermediates in the laboratory and in biological medium.²⁶⁻²⁸ Due to its presumed antioxidant properties PBN (α -phenyl-*N*-*t*-butyl-nitron) and some derivatives were evaluated as neuroprotectors in neurodegenerative diseases and some of them reached pre-clinical trials for the treatment of stroke.²⁹⁻³¹ Arylnitrones were also reported to be protective against microvascular damage in ischemia/reperfusion in the ‘hamster cheek pouch’ assay.³²⁻³⁶ It has been suggested that the mechanism of biological action of these compounds is mainly related with their ability to modulate cytokine production.³⁷⁻⁴¹

2. Results and Discussion

2.1. Chemistry

Several methods are reported to prepare nitrones and we choose to use the protocol described by Dondoni and coworkers.⁴²



Scheme 1. Synthesis of heteroaryl nitrones **1-3**.

While the synthesis of *N*-*t*-butyl nitrones (**1a** and **2a**) required the use of *N*-*t*-butylhydroxylamine hydrochloride and reflux in the presence of Et₃N in CH₂Cl₂ to go to completion, *N*-methyl and *N*-benzyl nitrones (**1b,c**; **2b,c**; **3a,b** and **4**) were prepared in good yields by the reaction between the corresponding aromatic aldehydes (**5** and **7**) with less sterically hindered and less expensive *N*-methyl or *N*-benzylhydroxylamine hydrochloride in the presence of Et₃N, in CH₂Cl₂ at room temperature. In all cases *Z*-geometry was observed in the resulting nitrones, as confirmed by NOE experiments. In addition, *N*-methyl nitrones can be prepared in the absence of solvent in few minutes when the reagents are macerated (SI).^{43,44} Nitro-heteroaryl nitrones **1a-c** and **2a-c** are described for the first time in this note.


2.2. In vitro activity against *L. infantum*, *L. amazonensis* and *L. donovani*

First, the synthesized compounds were screened for activity against promastigotes (insect stage) of the causative species of cutaneous (*L. amazonensis*) and visceral (*L. infantum*) leishmaniasis (Table 1). Next, active compounds from the first screening were evaluated against intracellular amastigotes of the same species. To assess antiparasitic selectivity, cytotoxicity was measured on murine peritoneal macrophages (Table 1, entry 1). Product **1a** with 108 μM was the least toxic to macrophages with a similar CC_{50} to miltefosine (96.5 μM), used as standard. Products **1b**

($\text{CC}_{50} = 49.9 \mu\text{M}$) and **2b** ($\text{CC}_{50} = 24.2 \mu\text{M}$) also showed low toxicity.

The antileishmanial activity in promastigotes of *L. amazonensis* shown in Table 1, entry 2). Products **1a** ($\text{EC}_{50} = 1.99 \mu\text{M}$), **1b** ($\text{EC}_{50} = 9.37 \mu\text{M}$), **2a** ($\text{EC}_{50} = 11.32 \mu\text{M}$), **2b** ($\text{EC}_{50} = 9.72 \mu\text{M}$) and **5b** ($\text{EC}_{50} = 5.11 \mu\text{M}$) were the most potent. Using a cut off of 10 μM we then assessed whether these products could inhibit the proliferation of intracellular amastigotes within macrophages. Compounds **1a** ($\text{EC}_{50} = 2.76 \mu\text{M}$), **1b** ($\text{EC}_{50} = 1.07 \mu\text{M}$) and **2b** ($\text{EC}_{50} = 3.81 \mu\text{M}$) were the most potent derivatives (Table 1, entry 3).

Table 1. IC_{50} for leishmanicidal activity of nitro-heteroarylnitrones in *Leishmania sp.*



Entry	Cell	1a	1b	5a	2a	2b	5b	Miltefosine
1	Macrophages CC_{50} (μM)	107.8 ± 0.1	49.9 ± 0.8	27.8 ± 0.2	29.8 ± 1.9	24.2 ± 0.3	6.25 ± 0.43	96.5 \pm 3.4
	<i>L. amazonensis</i>							
2	Promastigotes EC_{50} (μM)	1.99 ± 0.06	9.37 ± 0.09	25.20 ± 2.65	11.32 ± 0.74	9.72 ± 3.90	5.11 ± 0.54	7.01 ± 0.98
3	Intracellular amastigotes EC_{50} (μM)	2.76 ± 0.84	1.07 ± 0.11	> 57.9	-	3.81 ± 1.84	> 11.54	3.4 ± 1.00
4	Selectivity Index (SI)**	39	47	<0.5	-	6	<0.5	28
	<i>L. infantum</i>							
5	Promastigotes EC_{50} (μM)	0.36 ± 0.10	0.88 ± 0.19	25.68 ± 5.44	13.26 ± 2.57	3.45 ± 0.52	1.39 ± 0.04	9.81 ± 0.43
6	Intracellular amastigotes EC_{50} (μM)	0.019 ± 0.01	0.169 ± 0.13	ND	ND	ND	ND	0.5 ± 0.40
7	Selectivity Index (SI)**	5670	295	-	-	-	-	193

* IC_{50} and CC_{50} values correspond to the average of three independent experiments conducted in triplicate \pm standard error. N.D.- Not determined.

From the selectivity index (SI) between CC_{50} in macrophages and EC_{50} in intracellular amastigotes of *L. amazonensis*, **1a** with an SI of 39 and **1b** with an SI of 47 (Table 1, entry 4) were identified as the best meeting DNDi's target product profile of > 100-fold selectivity (Katsuno *et al.*, 2015⁴⁵). At this point, product **2b** (SI of 6) was disregarded. Likewise, nitroaldehydes **5a** and **5b** showed no selectivity (Table 1, entry 1). Compound **4** (not shown) was inactive, unveiling the importance of the heterocyclic aromatic moiety for antileishmanial activity. Nitrones **3a** and **3b** were inactive against macrophages ($\text{CC}_{50} > 100 \mu\text{M}$) as well as *L. amazonensis* promastigotes due to the lack of the nitro group at C5 in the heteroaromatic ring. As an outcome of this first screening against *L. amazonensis*, it is clear that both groups

(nitro and nitronium) are required in the heterocyclic scaffold for a potent biological activity and low toxicity.

The second screening with *L. infantum* promastigotes (Table 1, entry 5) revealed that the products **1a** ($\text{EC}_{50} = 0.36 \mu\text{M}$), **1b** ($\text{EC}_{50} = 0.88 \mu\text{M}$), **2b** ($\text{EC}_{50} = 3.45 \mu\text{M}$) and **5a** ($\text{EC}_{50} = 1.39 \mu\text{M}$) were preferentially more potent on this visceral species in comparison with *L. amazonensis*. Further studies with products **1a** and **1b** against intracellular amastigotes revealed the potency in the nanomolar range for **1a** ($\text{EC}_{50} = 0.019 \mu\text{M}$) and **1b** ($\text{EC}_{50} = 0.169 \mu\text{M}$) which are, respectively, 14-fold and 6-fold more potent for intracellular amastigotes of *L. infantum* than of *L. amazonensis* (Table 1, entry 5). Together with their cytotoxicity profile, a selectivity index (SI) of 5,760 obtained for product **1a**

and 295 for product **1b** make both of them very promising hit compounds for antileishmanial drug discovery (Table 1, entry 6). The prodrug profiles of these compounds were determined comparing results obtained with promastigotes of wild type *L. donovani* with those obtained with genetically modified promastigotes of *L. donovani* over expressing the nitroreductase 1 (NTR1^{high}) or nitroreductase 2 (NTR2^{high}) genes (Table 2). Miltefosine, currently used in the treatment of leishmaniasis, nifurtimox, a commercial nitro-drug mainly activated by NTR1¹² and delamanid, a bicyclic nitro drug activated by NTR2²¹, were used as reference drugs standards. Table 2, entry 1 shows that **1a** is almost equally strongly activated by LdNTR1 (17 times) and LdNTR2 (21 times) while **1b** is even more activated (38 times) than nifurtimox (17 times) by LdNTR1 but less activated by LdNTR2 (6.1 times) than **1a** (21 times) and delamanid (12.5 times). In contrast, **1c** with a bulky benzyl substitution was less sensitive to both types of activation.

Thiophene derivative **2a** presented a similar profile of **1a** being equally activated by LdNTR1 and LdNTR2 (17 and 18 times, respectively). However, **2b** was less sensitive to activation by both nitroreductases (4.6 and 7 times) than **1b**. As observed with **1c**, **2c** was also less sensitive to the bioactivation by both nitroreductases.

We evaluated how the nitroderivatives would act on peritoneal macrophages and on *L. infantum*, regarding the production of ROS, initially labelling parasites with 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA), using antimycin A or nifurtimox as positive ROS-production standard.⁴⁶ Figure 2a displays the effect of our compounds on the production of ROS in *L. amazonensis* promastigotes. Interestingly, nitroheteroaryl nitrones **1a,b** and **2a,b** did not induce the production of important amounts of ROS when compared with untreated parasites.

Table 2. IC₅₀ for leishmanicidal activity of LQB's nitrones in *Leishmania donovani* overexpressing NTR1 or NTR2.



Entry	<i>L. donovani</i>	1a	1b	1c	2a	2b	2c	Reference Drugs
1	Wild Promastigotes	0.87 ± 0.026	1.16 ± 0.005	0.99 ± 0.036	4.55 ± 0.149	1.12 ± 0.042	0.41 ± 0.011	Nifurtimox 1.89 ± 0.052
	EC ₅₀ (μM)							Delamanid 0.0015 ± 0.00003
2	NTR1 ^{high} Promastigotes	0.05 ± 0.001	0.03 ± 0.001	1.22 ± 0.006	0.27 ± 0.009	0.24 ± 0.043	0.30 ± 0.019	Nifurtimox 0.11 ± 0.002
3	Ratio WT/ NTR1 ^{high}	17	38	0.81	17	4.6	1.3	17
4	NTR2 ^{high} Promastigotes	0.04 ± 0.005	0.19 ± 0.030	1.33 ± 0.037	0.31 ± 0.023	0.25 ± 0.021	0.76 ± 0.013	Delamanid 0.00012 ± 0.00001
5	Ratio WT/ NTR2 ^{high}	21	6.1	0.74	18	4.5	0.53	12.5

* IC₅₀ values correspond to the average of three independent experiments conducted in triplicate ± standard error.

However, nitroaldehyde **5b**, but not **5a**, was able to trigger ROS in a dose and time-dependent manner. The production of ROS in murine peritoneal macrophages follows the same trend (Figure 2b). For nitroaldehyde **5b** an increased ROS production was observed and can be correlated with its high toxicity. Next, we decided to evaluate the ROS production using a mixture of compounds **5b** and **1a** (Figures 2a). These compounds were combined in a 1:1 molar ratio and incubated with the parasite or murine macrophages at the EC₅₀ concentration, respec-

tively 0.88 μM and 16.25 μM. It is noteworthy that the generation of ROS was controlled during the 4h of incubation, confirming the antioxidant activity of **1a**. These results are in agreement with the known radical scavenging capacity of the nitronium functional group⁴⁷, present in the structure of **1a**. It is worth noting that the aldehyde group also stabilizes the nitronium anion radical formed (see E^o, *infra*); however, the aldehyde group is electrophilic and can combine with endogenous amino groups to exert toxicity.




Figure 2. Treatment with nitro-heteroaryl nitrones generate ROS in *L. amazonensis* and mammalian cells) *L. amazonensis* promastigotes treated with nitro-heterocyclic compounds for a 4 h kinetic; concentration of the nitro-heterocyclic compounds was based on the anti-promastigote EC₅₀. b) Murine macrophages incubated for 4h with 0.5 x EC₅₀, EC₅₀ and 2 x EC₅₀ nitro-heterocyclic compounds, concentrations based on anti-promastigote assay. ROS generation was quantified using H₂DCFDA (Molecular Probes). Results are presented as means ± SD; n=3; p<0.05, **p<0.01 and ***p<0.001, (Untreated) x (treated); #p<0.001, (5a x (1b+5a)).

3. Mechanistic considerations

The antiparasitic mechanism of action of nitroaromatic compounds at the molecular level is not yet fully understood, but certainly involves activation through the reduction of the nitro group. Taking in account the structural similarity between **1a,b** and nifurtimox, we considered different mechanisms for the antiparasitic activity of this compound, which have been proposed by several groups.^{11,48–50} In *T. cruzi* the one electron reduction to the corresponding nitro anion radical followed by recycling nifurtimox at the expense of superoxide formation was discarded (Figure 3a, left), based on the fact that ROS production was only observed in doses of nifurtimox much higher than the therapeutic dose.³⁹ More recently, it has been suggested that nifurtimox is preferentially reduced by NTR1^{11,21,48} through a sequential two-electron transfer pathway, leading to the corresponding nitroso and hydroxylamine derivatives (Figure 3a). An acyclic olefin conjugated with two electron withdrawing groups (strong 1,4-addition acceptor) is also formed from this hydroxylamine (highlighted in blue). These metabolites play a role in the antiparasitic activity and toxicity of nifurtimox.^{10,14,48} Reduction of **1a,b** by *LdNTR1* (Figure 3b) could lead to similar me-

tabolites observed for nifurtimox, although we have no experimental evidence to support this proposal. As shown in Figure 2, except for **5a**, we also did not observe oxidative stress in our experiments.

On the other hand, nifurtimox is only marginally activated by NTR2, the new nitroreductase discovered by Wyllie and coworkers. NTR2 reduce preferentially bicyclic nitro compounds, but the mechanism of this reduction via one or two electron reduction pathways is still not clear.^{13,21}

While nifurtimox is a good structural model for the activation of our nitro-nitrones by NTR1, in delamanid and other bicyclic nitro-heteroaryl compounds activated by NTR2 the heterocyclic moiety is quite different and so, we do not have a structural model for the reduction pathway involving our compounds and NTR2. However, a one electron route suggested in Figure 3b seems a good working hypothesis. This pathway can be estimated by cyclic voltammetry⁵¹, which allows to measure the redox potential (E⁰) of a studied system based on a one electron transfer pathway.




Figure 3. Activation of nifurtimox and nitro-heteroaryl nitrones **1a,b** by nitroreductases of leishmania.

Pedron and coworkers reported that in 8-nitroquinolin-2-(1H)-ones (Figure 4a) the antiparasitic activity in *L. infantum* could be correlated with the redox potential measured by cyclic voltammetry and only compounds with values above -0.6V vs NHE, in aprotic medium, presented antileishmanial activity.⁵² An intramolecular hydrogen bond involving the nitro group and the

N-H at the lactam moiety in 8-nitroquinolone and the formation of an aromatic oxyanion is responsible for the important increase in the redox potential from -0.84 V in 8-nitroquinoline to -0.54 V. Nevertheless, these 8-nitroquinolin-2-(1H)-ones are activated by NTR1 (2e transfer) and, despite that, the leishmanicidal activity could be correlated with the redox potential (E°), based on a mono-electronic transfer process.




Figure 4. Stabilization of anion radicals resulting from 1e nitro reduction and redox potential of some nitro aromatic compounds measured by cyclic voltammetry in DMSO.

In Figure 4b we show that the nitron group can stabilize the resulting nitro-anion radical formed from **4** by resonance through the aromatic ring and E° value increase from -0.85 V in nitrobenzene to -0.65 V. So, **4** is reduced easier than nitrobenzene. Interestingly, the aldehyde group in *p*-nitrobenzaldehyde **7** stabilizes still better the radical anion (E° -0.55 V). A similar trend was observed in nitro-nitrones **1a,b** and **2a,b**, having E° in the range of -0.56 V to -0.59 V (**4**, E° 0.65 V) showing a slight dependence on the nature of the heterocyclic (O x S) and the

substituent at the nitrogen atom. The aldehyde group in **5a** is also capable of stabilizing the anion radical better than nitrones (E° -0.38 V). However, while nitron group is in general not toxic, the aldehyde group is electrophilic and can react with important bioamines, explaining the higher toxicity of aldehydes over nitrones.

However, the data in Figure 4 also show that an appropriate E° value is a necessary condition to be a substrate of these enzymes, but other factors must be considered concerning the antiparasitic activity, such as the molecular recognition between substrate and enzyme. For example, based on E° , **4** could be reduced by NTRs, but this compound doesn't have leishmanicidal activity. Another point to be considered is the eventual formation of acyclic strong Michael acceptors, which have been correlated with the antileishmanial effect. In **4**, a ring opening would involve the cleavage of a strong C-C bond with the destruction of the high stabilized aromatic ring and this pathway seems unfavorable when compared with **1a,b**. We could also speculate that these nitro-furan nitrones are more prone to undergo this pathway than nitro-thiophenenitrones (**2a,b**), once thiophenes are more aromatic than furans.

4. *In vitro* pharmacokinetic and physicochemical properties of compound **1b**.

Drug candidates frequently fail at different stages of drug discovery and development. Wang and Urban reported that the most prominent cause of failure is poor pharmacokinetic properties.⁵⁶ Initially, *in silico* characteristics were evaluated by the pKCSM tool. Compound **1b** showed compliance with the rule of 5 and solubility in aqueous medium > 10 μ M (confirmed experimentally). It did not show any toxicity characteristic, except for the AMES test, which is expected to give positive because of the presence of the nitro group, but without significance to humans (SI).⁵⁶

Since **1b** has a potent leishmanicidal activity, excellent bioselectivity and, in addition, is the less expensive to synthesize among the nitro-heteroarylnitrones, we performed a preliminary evaluation of its pharmacokinetic parameters *in vitro* (SI); a summary of these results is shown in Figure 5. Compound **1b** showed chemical stability at pH 7, without degradation greater than 25%; however, at pH 2, almost 75% of the compound was degraded within 6 h. This compound showed good metabolic stability evaluated in human microsomal fraction S9 and reasonable mouse plasma stability (Figure 5). These data provide vital information since the desirable candidate for leishmaniasis must be administered orally.⁵⁴ The low stability in acidic pH could possibly be minimized through appropriate formulation. Our initial solution, however, was the co-administration with omeprazole.

5. Efficacy in experimental Visceral Leishmaniasis.

As a proof of concept, we evaluated the effectiveness of compound **1b** administered orally in mice infected with *L. infantum* (Figure 6). The animals were treated every 12 h for five consecutive days. As a strategy to minimize the possible degradation of compound **1b** in the stomach, a proton pump inhibitor was administered orally 1h before treatment (omeprazole). Sima *et al.*, demonstrated that the administration of 20mg/kg of omeprazole (ip) raises the stomach pH in rats from 3.5 to 6.7.⁵⁵ Im-

portantly, omeprazole does not present leishmanicidal activity, as shown in Figures 6a and 6b.




Figure 5. Chemical, microsomal and plasmatic stability of compound **1b**.




Figure 6. Efficacy of **1b** prototype in an experimental model of visceral leishmaniasis. BALB/c mice (n=5) were infected intraperitoneally with 1×10^8 *L. infantum* promastigotes. Seven days after infection, treatment with oral omeprazole 10mg was started 1 h before treatment with **1b** derivative at concentrations of 25, 50 and 100mg/kg twice a day, orally. Control groups were treated with 22mg/kg of miltefosine twice a day and vehicle. The parasitic load in the spleen (a) and liver (b) was estimated by a limiting dilution test. * $p < 0.05$ and *** $p < 0.001$ treated groups vs omeprazole group.

Treatment with **1b** showed a reduction in parasite load on the spleen of 76.6% and 95.0% with doses of 50 and 100mg/kg, respectively, being 2 to 3 times less potent than miltefosine. In the liver, parasite suppression was above 75% at the three doses used, without the difference in efficiency between them, being as active as miltefosine. The liver is one of the major sites for parasite burden and after 5 days of treatment, the results for compound **1b** meet the criteria for a lead that should be profiled for further development. It is worth to note that **1a** and **2a** have in common a bulky group (*t*-butyl) attached to the nitrogen atom while in **1b** and **2b** a small methyl group occupies this place.

6. Conclusion

Nitro-heteroaryl nitrones **1a** and **1b** are non-chiral compounds prepared in one step, in excellent yields, from commercially available 5-nitrofurfural, 5-nitro-2-formyl-thiophene and *N*-*t*-BuNHOH or *N*-MeNHOH. The prodrug profile of these compounds is unusual as they are activated by both *Ld*NRT1 and *Ld*NRT2 in *L. donovani*. Reliance on a single enzyme for prodrug activation may leave drugs such as nifurtimox and fexinidazole vulnerable to the emergence of drug resistance, thus the activation by both NRT1 and NRT2 at the same time is a promising option to overcome this problem. The examples shown in this work are, to the best of our knowledge, the first in which monocyclic nitro-compounds are markedly activated by NRT2. A criteria list proposed by Katsuno and co-workers⁴⁵ to define antileishmanial hit compounds includes novel structure, easy synthesis (up to 5 reaction steps), $IC_{50} < 10 \mu M$ (against intracellular *L. donovani*) and selectivity index > 100 (in comparison with mammalian cells). Of the tested nitro-heteroaryl nitrones, compounds **1a** and **1b** meet all these criteria. Efficacy in an experimental model was demonstrated with $> 70\%$ reduction in liver parasite burden after at most 5 doses at 50mg per kg delivered orally once or twice per day. Altogether, these results indicate that the strategy of joining nitro and nitrone in the same scaffold was successful and suggest that the study with compounds **1a** and **1b** should be extended to assays in other animal models. New patterns of substitution at the nitrogen atom can be used to modulate the chemical stability and selectivity for nitroreductases.

6. REFERENCES

- (1) Desjeux, P. Leishmaniasis: Current Situation and New Perspectives. *Comp. Immunol. Microbiol. Infect. Dis.* **2004**, *27* (5), 305–318. <https://doi.org/10.1016/j.cimid.2004.03.004>.
- (2) Torres-Guerrero, E.; Quintanilla-Cedillo, M. R.; Ruiz-Esmenjaud, J.; Arenas, R. Leishmaniasis: A Review. *F1000Res* **2017**, *6*, 750. <https://doi.org/10.12688/f1000research.11120.1>.
- (3) Leishmaniasis <https://www.afro.who.int/health-topics/Leishmaniasis> (accessed Aug 23, 2020).
- (4) Williamson, J. Chemotherapy and Chemoprophylaxis of African Trypanosomiasis. *Exp. Parasitol.* **1962**, *12*, 274–322. [https://doi.org/10.1016/0014-4894\(62\)90075-9](https://doi.org/10.1016/0014-4894(62)90075-9).
- (5) Schmidt, P.; Eichenberger, K.; Ilvespää, A. O.; Wilhelm, M. Nitroheterocycles with Antiparasitic Effects.

- Ann. N. Y. Acad. Sci.* **1969**, 160 (2), 530–535. <https://doi.org/10.1111/j.1749-6632.1969.tb15872.x>.
- (6) Winkelman, E.; Raether, W.; Sinharay, A. Chemotherapeutically Active Nitro Compounds. 4,5-Nitroimidazoles (Part II). *Arzneimittelforschung* **1978**, 28 (3), 351–366.
- (7) Papadopoulou, M. V.; Bloomer, W. D.; Rosenzweig, H. S.; Wilkinson, S. R.; Kaiser, M. Novel Nitro(Triazole/Imidazole)-Based Heteroarylamides/Sulfonamides as Potential Antitrypanosomal Agents. *Eur J Med Chem* **2014**, 87, 79–88. <https://doi.org/10.1016/j.ejmech.2014.09.045>.
- (8) Hamilton-Miller, J. M.; Brumfitt, W. The Versatility of Nitro Compounds. *J. Antimicrob. Chemother.* **1976**, 2 (1), 5–8. <https://doi.org/10.1093/jac/2.1.5>.
- (9) Raether, W.; Hänel, H. Nitroheterocyclic Drugs with Broad Spectrum Activity. *Parasitol. Res.* **2003**, 90 Supp 1, S19–39. <https://doi.org/10.1007/s00436-002-0754-9>.
- (10) Patterson, S.; Wyllie, S. Nitro Drugs for the Treatment of Trypanosomatid Diseases: Past, Present, and Future Prospects. *Trends Parasitol.* **2014**, 30 (6), 289–298. <https://doi.org/10.1016/j.pt.2014.04.003>.
- (11) Hall, B. S.; Bot, C.; Wilkinson, S. R. Nifurtimox Activation by Trypanosomal Type I Nitroreductases Generates Cytotoxic Nitrile Metabolites. *J. Biol. Chem.* **2011**, 286 (15), 13088–13095. <https://doi.org/10.1074/jbc.M111.230847>.
- (12) Wyllie, S.; Patterson, S.; Fairlamb, A. H. Assessing the Essentiality of Leishmania Donovanii Nitroreductase and Its Role in Nitro Drug Activation. *Antimicrob. Agents Chemother.* **2013**, 57 (2), 901–906. <https://doi.org/10.1128/AAC.01788-12>.
- (13) Patterson, S.; Fairlamb, A. H. Current and Future Prospects of Nitro-Compounds as Drugs for Trypanosomiasis and Leishmaniasis. *Curr. Med. Chem.* **2019**, 26 (23), 4454–4475. <https://doi.org/10.2174/0929867325666180426164352>.
- (14) Nepali, K.; Lee, H.-Y.; Liou, J.-P. Nitro-Group-Containing Drugs. *J. Med. Chem.* **2019**, 62 (6), 2851–2893. <https://doi.org/10.1021/acs.jmedchem.8b00147>.
- (15) Priotto, G.; Kasparian, S.; Mutombo, W.; Nguouma, D.; Ghorashian, S.; Arnold, U.; Ghabri, S.; Baudin, E.; Bular, V.; Kazadi-Kyanza, S.; Ilunga, M.; Mutangala, W.; Pohlig, G.; Schmid, C.; Karunakara, U.; Torreele, E.; Kande, V. Nifurtimox-Eflornithine Combination Therapy for Second-Stage African Trypanosoma Brucei Gambiense Trypanosomiasis: A Multicentre, Randomised, Phase III, Non-Inferiority Trial. *Lancet* **2009**, 374 (9683), 56–64. [https://doi.org/10.1016/S0140-6736\(09\)61117-X](https://doi.org/10.1016/S0140-6736(09)61117-X).
- (16) Carrilero, B.; Murcia, L.; Martínez-Lage, L.; Segovia, M. Side Effects of Benzimidazole Treatment in a Cohort of Patients with Chagas Disease in Non-Endemic Country. *Rev Esp Quimioter* **2011**, 24 (3), 123–126.
- (17) Wyllie, S.; Patterson, S.; Stojanovski, L.; Simeons, F. R. C.; Norval, S.; Kime, R.; Read, K. D.; Fairlamb, A. H. The Anti-Trypanosome Drug Fexinidazole Shows Potential for Treating Visceral Leishmaniasis. *Science Translational Medicine* **2012**, 4 (119), 119re1–119re1. <https://doi.org/10.1126/scitranslmed.3003326>.
- (18) Sundar, S.; Singh, A. Recent Developments and Future Prospects in the Treatment of Visceral Leishmaniasis. *Therapeutic Advances in Infection* **2016**, 3 (3–4), 98–109. <https://doi.org/10.1177/2049936116646063>.
- (19) Eperon, G.; Balasegaram, M.; Potet, J.; Mowbray, C.; Valverde, O.; Chappuis, F. Treatment Options for Second-Stage Gambiense Human African Trypanosomiasis. *Expert Review of Anti-infective Therapy* **2014**, 12 (11), 1407–1417. <https://doi.org/10.1586/14787210.2014.959496>.
- (20) Pedron, J.; Boudot, C.; Brossas, J.-Y.; Pinault, E.; Bourgeade-Delmas, S.; Sourmia-Saquet, A.; Boutet-Robinet, E.; Destere, A.; Tronnet, A.; Bergé, J.; Bonduelle, C.; Deraeve, C.; Prativiel, G.; Stigliani, J.-L.; Paris, L.; Mazier, D.; Corvaisier, S.; Since, M.; Malzert-Fréon, A.; Wyllie, S.; Milne, R.; Fairlamb, A. H.; Valentin, A.; Courtioux, B.; Verhaeghe, P. New 8-Nitroquinolinone Derivative Displaying Submicromolar *In Vitro* Activities against Both *Trypanosoma Brucei* and *Cruzi*. *ACS Med. Chem. Lett.* **2020**, 11 (4), 464–472. <https://doi.org/10.1021/acmedchemlett.9b00566>.
- (21) Wyllie, S.; Roberts, A. J.; Norval, S.; Patterson, S.; Foth, B. J.; Berriman, M.; Read, K. D.; Fairlamb, A. H. Activation of Bicyclic Nitro-Drugs by a Novel Nitroreductase (NTR2) in Leishmania. *PLoS Pathog* **2016**, 12 (11), e1005971. <https://doi.org/10.1371/journal.ppat.1005971>.
- (22) Barbara, J. E.; Kazmi, F.; Parkinson, A.; Buckley, D. B. Metabolism-Dependent Inhibition of CYP3A4 by Lapatinib: Evidence for Formation of a Metabolic Intermediate Complex with a Nitroso/Oxime Metabolite Formed via a Nitron Intermediate. *Drug Metab Dispos* **2013**, 41 (5), 1012. <https://doi.org/10.1124/dmd.113.051151>.
- (23) Rodriguez, R. J.; Miranda, C. L. Isoform Specificity of N -Deacetyl Ketoconazole by Human and Rabbit Flavin-Containing Monooxygenases. *Drug Metab Dispos* **2000**, 28 (9), 1083.
- (24) Gogas, K. R.; Lechner, S. M.; Markison, S.; Williams, J. P.; McCarthy, W.; Grigoriadis, D. E.; Foster, A. C. 6.04 - Anxiety. In *Comprehensive Medicinal Chemistry II*; Taylor, J. B., Triggle, D. J., Eds.; Elsevier: Oxford, 2007; pp 85–115. <https://doi.org/10.1016/B0-08-045044-X/00164-4>.
- (25) Atkins, J. M.; Mitchell, H. C.; Pettinger, W. A. Increased Pulmonary Vascular Resistance with Systemic Hypertension. Effect of Minoxidil and Other Antihypertensive Agents. *Am. J. Cardiol.* **1977**, 39 (6), 802–807. [https://doi.org/10.1016/s0002-9149\(77\)80030-1](https://doi.org/10.1016/s0002-9149(77)80030-1).
- (26) Floyd, R. A.; Soong, L. M. Spin Trapping in Biological Systems. Oxidation of the Spin Trap 5,5-Dimethyl-1-Pyrroline-1-Oxide by a Hydroperoxide-Hematin System. *Biochem. Biophys. Res. Commun.* **1977**, 74 (1), 79–84. [https://doi.org/10.1016/0006-291x\(77\)91377-8](https://doi.org/10.1016/0006-291x(77)91377-8).
- (27) Gomez-Mejiba, S. E.; Zhai, Z.; Della-Vedova, M. C.; Muñoz, M. D.; Chatterjee, S.; Towner, R. A.; Hensley, K.; Floyd, R. A.; Mason, R. P.; Ramirez, D. C. Immuno-Spin Trapping from Biochemistry to Medicine: Advances, Challenges, and Pitfalls. Focus on Protein-Centered Radicals. *Biochimica et Biophysica Acta (BBA) - General Subjects* **2014**, 1840 (2), 722–729. <https://doi.org/10.1016/j.bbagen.2013.04.039>.
- (28) Janzen, E. G., B., B. J. Detection and identification of short-lived free radicals by an electron spin resonance trapping technique. *Journal of the American Chemical Society*, 90(21), 5909–5910 | 10.1021/ja01023a051 <https://doi.org/10.1021/ja01023a051> (accessed Aug 23, 2020).

- (29) Green, A. R.; Ashwood, T.; Odergren, T.; Jackson, D. M. Nitrones as Neuroprotective Agents in Cerebral Ischemia, with Particular Reference to NXY-059. *Pharmacol. Ther.* **2003**, *100* (3), 195–214. <https://doi.org/10.1016/j.pharmthera.2003.07.003>.
- (30) Lees, K. R.; Zivin, J. A.; Ashwood, T.; Davalos, A.; Davis, S. M.; Diener, H.-C.; Grotta, J.; Lyden, P.; Shuaib, A.; Hårdemark, H.-G.; Wasiewski, W. W. NXY-059 for Acute Ischemic Stroke. *N Engl J Med* **2006**, *354* (6), 588–600. <https://doi.org/10.1056/NEJMoa052980>.
- (31) Floyd, R. A.; Castro Faria Neto, H. C.; Zimmerman, G. A.; Hensley, K.; Townner, R. A. Nitro-Based Therapeutics for Neurodegenerative Diseases: Their Use Alone or in Combination with Lanthionines. *Free Radic. Biol. Med.* **2013**, *62*, 145–156. <https://doi.org/10.1016/j.freeradbiomed.2013.01.033>.
- (32) Kim, S.; de A Vilela, G. V. M.; Bouajila, J.; Dias, A. G.; Cyrino, F. Z. G. A.; Bouskela, E.; Costa, P. R. R.; Nepveu, F. Alpha-Phenyl-N-Tert-Butyl Nitro (PBN) Derivatives: Synthesis and Protective Action against Microvascular Damages Induced by Ischemia/Reperfusion. *Bioorg. Med. Chem.* **2007**, *15* (10), 3572–3578. <https://doi.org/10.1016/j.bmc.2007.02.033>.
- (33) Dias, A. G.; Santos, C. E. V.; Cyrino, F. Z. G. A.; Bouskela, E.; Costa, P. R. R. N-Tert-Butyl and N-Methyl Nitrones Derived from Aromatic Aldehydes Inhibit Macromolecular Permeability Increase Induced by Ischemia/Reperfusion in Hamsters. *Bioorg. Med. Chem.* **2009**, *17* (11), 3995–3998. <https://doi.org/10.1016/j.bmc.2009.04.004>.
- (34) Marco-Contelles, J. Recent Advances on Nitrones Design for Stroke Treatment *J Med Chem* **2020**, *63* (22), 13413–13427. <https://doi.org/10.1021/acs.jmedchem.0c00976>
- (35) Rosselin, M.; Poeggeler, B.; Durand, G. Nitro Derivatives as Therapeutics: From Chemical Modification to Specific-targeting. *Curr Top Med Chem* **2017**, *17*, 1–17. <https://doi.org/10.2174/1568026617666170303115324>
- (36) Chamorro I, B.; Diez-Iriepa, D.; Merás-Sáiz, B.; Chioua, M.; García-Vieira, D.; Iriepa, I.; Hadjipavlou-Litina, D.; López-Muñoz, F.; Martínez-Murillo, R.; González-Nieto, D.; Fernández, I.; Marco-Contelles, J.; Oset-Gasque, M. J. Synthesis, antioxidant properties and neuroprotection of α -phenyl-*tert*-butylnitro derived *HomoBisNitrones* in *in vitro* and *in vivo* ischemia models *Sci. Rep* **2020**, *10* (1): 14150–14168. <https://doi.org/10.1038/s41598-020-70690-y>
- (37) Miyajima, T.; Kotake, Y. Spin Trapping Agent, Phenyl N-Tert-Butyl Nitro, Inhibits Induction of Nitric Oxide Synthase in Endotoxin-Induced Shock in Mice. *Biochem. Biophys. Res. Commun.* **1995**, *215* (1), 114–121. <https://doi.org/10.1006/bbrc.1995.2440>.
- (38) Kotake, Y.; Sang, H.; Miyajima, T.; Wallis, G. L. Inhibition of NF-KappaB, INOS MRNA, COX2 MRNA, and COX Catalytic Activity by Phenyl-N-Tert-Butylnitro (PBN). *Biochim. Biophys. Acta* **1998**, *1448* (1), 77–84. [https://doi.org/10.1016/s0167-4889\(98\)00126-8](https://doi.org/10.1016/s0167-4889(98)00126-8).
- (39) Sang, H.; Wallis, G. L.; Stewart, C. A.; Kotake, Y. Expression of Cytokines and Activation of Transcription Factors in Lipopolysaccharide-Administered Rats and Their Inhibition by Phenyl N-Tert-Butylnitro (PBN). *Arch. Biochem. Biophys.* **1999**, *363* (2), 341–348. <https://doi.org/10.1006/abbi.1998.1086>.
- (40) Mallick, S.; Halder, S.; Dutta, A.; Dey, S.; Paul, K.; Maiti, S.; Bandyopadhyay, C.; Saha, B.; Pal, C. Chromone Linked Nitro Derivative Induces the Expression of INOS2 and Th1 Cytokines but Reduces the Th2 Response in Experimental Visceral Leishmaniasis. *Int. Immunopharmacol.* **2013**, *15* (4), 772–779. <https://doi.org/10.1016/j.intimp.2013.02.013>.
- (41) Costa, D. S. S.; Martino, T.; Magalhães, F. C.; Justo, G.; Coelho, M. G. P.; Barcellos, J. C. F.; Moura, V. B.; Costa, P. R. R.; Sabino, K. C. C.; Dias, A. G. Synthesis of N-Methylarylnitrones Derived from Alkyloxybenzaldehydes and Antineoplastic Effect on Human Cancer Cell Lines. *Bioorg. Med. Chem.* **2015**, *23* (9), 2053–2061. <https://doi.org/10.1016/j.bmc.2015.03.014>.
- (42) Dondoni, A.; Franco, S.; Junquera, F.; Merchán, F. L.; Merino, P.; Tejero, T. Synthesis of N-Benzyl Nitrones. *Synthetic Communications* **1994**, *24* (18), 2537–2550. <https://doi.org/10.1080/00397919408010565>.
- (43) Yavuz, S.; Ozkan, H.; Colak, N.; Yildirim, Y. Fast Method for Synthesis of Alkyl and Aryl-N-Methylnitrones. *Molecules* **2011**, *16* (8), 6677–6683. <https://doi.org/10.3390/molecules16086677>.
- (44) Colacino, E.; Nun, P.; Colacino, F. M.; Martinez, J.; AndLamaty, F. Solvent-Free Synthesis of Nitrones in a Ball-Mill. *Tetrahedron* **2008**, *64* (23), 5569–5576. <https://doi.org/10.1016/j.tet.2008.03.091>.
- (45) Katsuno, K.; Burrows, J. N.; Duncan, K.; Hooft van Huijsduijnen, R.; Kaneko, T.; Kita, K.; Mowbray, C. E.; Schmatz, D.; Warner, P.; Slingsby, B. T. Hit and Lead Criteria in Drug Discovery for Infectious Diseases of the Developing World. *Nat Rev Drug Discov* **2015**, *14* (11), 751–758. <https://doi.org/10.1038/nrd4683>.
- (46) Quinlan, C. L.; Gerencser, A. A.; Treberg, J. R.; Brand, M. D. The Mechanism of Superoxide Production by the Antimycin-Inhibited Mitochondrial Q-Cycle. *J. Biol. Chem.* **2011**, *286* (36), 31361–31372. <https://doi.org/10.1074/jbc.M111.267898>.
- (47) Janzen, E. G.; Blackburn, B. J. Detection and Identification of Short-Lived Free Radicals by Electron Spin Resonance Trapping Techniques (Spin Trapping). Photolysis of Organolead, -Tin, and -Mercury Compounds. *J. Am. Chem. Soc.* **1969**, *91* (16), 4481–4490. <https://doi.org/10.1021/ja01044a028>.
- (48) Bot, C.; Hall, B. S.; Alvarez, G.; Di Maio, R.; González, M.; Cerecetto, H.; Wilkinson, S. R. Evaluating 5-Nitrofurans as Trypanocidal Agents. *Antimicrob. Agents Chemother.* **2013**, *57* (4), 1638–1647. <https://doi.org/10.1128/AAC.02046-12>.
- (49) Boiani, M.; Piacenza, L.; Hernández, P.; Boiani, L.; Cerecetto, H.; González, M.; Denicola, A. Mode of Action of Nifurtimox and N-Oxide-Containing Heterocycles against *Trypanosoma Cruzi*: Is Oxidative Stress Involved? *Biochem. Pharmacol.* **2010**, *79* (12), 1736–1745. <https://doi.org/10.1016/j.bcp.2010.02.009>.
- (50) Hall, B. S.; Wilkinson, S. R. Activation of Benznidazole by Trypanosomal Type I Nitroreductases Results in Glyoxal Formation. *Antimicrob. Agents Chemother.* **2012**, *56* (1), 115–123. <https://doi.org/10.1128/AAC.05135-11>.
- (51) Hillard, E. A.; de Abreu, F. C.; Ferreira, D. C. M.; Jaouen, G.; Goulart, M. O. F.; Amatore, C. Electrochemical Parameters and Techniques in Drug Devel-

- opment, with an Emphasis on Quinones and Related Compounds. *Chem. Commun. (Camb.)* **2008**, No. 23, 2612–2628. <https://doi.org/10.1039/b718116g>.
- (52) Pedron, J.; Boudot, C.; Hutter, S.; Bourgeade-Delmas, S.; Stigliani, J.-L.; Sourmia-Saquet, A.; Moreau, A.; Boutet-Robinet, E.; Paloque, L.; Mothes, E.; Laget, M.; Vendier, L.; Pratiel, G.; Wyllie, S.; Fairlamb, A.; Azas, N.; Courtioux, B.; Valentin, A.; Verhaeghe, P. Novel 8-Nitroquinolin-2(1H)-Ones as NTR-Bioactivated Antikinetoplastid Molecules: Synthesis, Electrochemical and SAR Study. *Eur J Med Chem* **2018**, *155*, 135–152. <https://doi.org/10.1016/j.ejmech.2018.06.001>.
- (53) Dixit, M., & Kumar, A. In Vitro Gene Genotoxicity Test Methods. *In Vitro Toxicology*, 67–89 | <https://doi.org/10.1016/B978-0-12-804667-8.00004-3> <https://sci-hub.tw/10.1016/B978-0-12-804667-8.00004-3> (accessed Sep 18, 2020).
- (54) Field, M. C.; Horn, D.; Fairlamb, A. H.; Ferguson, M. A. J.; Gray, D. W.; Read, K. D.; De Rycker, M.; Torrie, L. S.; Wyatt, P. G.; Wyllie, S.; Gilbert, I. H. Anti-Trypanosomatid Drug Discovery: An Ongoing Challenge and a Continuing Need. *Nat. Rev. Microbiol.* **2017**, *15* (4), 217–231. <https://doi.org/10.1038/nrmicro.2016.193>.
- (55) Šíma, M.; Kutinová-Canová, N.; Ryšánek, P.; Hořínková, J.; Moškořová, D.; Slanař, O. Gastric PH in Rats: Key Determinant for Preclinical Evaluation of PH-Dependent Oral Drug Absorption. *Prague Med Rep* **2019**, *120* (1), 5–9. <https://doi.org/10.14712/23362936.2019.5>.
- (56) Jianling Wang; Laszlo Urban. The Impact of Early ADME Profiling on Drug Discovery and Development Strategy. *Drug Discovery World Fall* **2004**.

(Word Style "TF_References_Section"). References are placed at the end of the manuscript. Authors are responsible for the accuracy and completeness of all references. Examples of the recommended formats for the various reference types can be found at <http://pubs.acs.org/page/4authors/index.html>. Detailed information on reference style can be found in The ACS Style Guide, available from Oxford Press.

ASSOCIATED CONTENT

Supporting Information

(Word Style "TE_Supporting_Information"). A listing of the contents of each file supplied as Supporting Information should be included. For instructions on what should be included in the Supporting Information as well as how to prepare this material for publication, refer to the journal's Instructions for Authors.

SYNOPSIS TOC (Word Style "SN_Synopsis_TOC"). If you are submitting your paper to a journal that requires a synopsis graphic and/or synopsis paragraph, see the In To format double-column figures, schemes, charts, and tables, use the following instructions:

The Supporting Information is available free of charge on the ACS Publications website.

brief description (file type, i.e., PDF)

brief description (file type, i.e., PDF)

AUTHOR INFORMATION

Corresponding Author

* (Word Style "FA_Corresponding_Author_Footer"). Give contact information for the author(s) to whom correspondence should be addressed.

PresentAddresses

†If an author's address is different than the one given in the affiliation line, this information may be included here.

Author Contributions

The manuscript was written through contributions of all authors. / All authors have given approval to the final version of the manuscript. / ‡These authors contributed equally. (match statement to author names with a symbol)

Funding Sources

Any funds used to support the research of the manuscript should be placed here (per journal style).

Notes

Any additional relevant notes should be placed here.

ACKNOWLEDGMENT

(Word Style "TD_Acknowledgments"). Generally the last paragraph of the paper is the place to acknowledge people (dedications), places, and financing (you may state grant numbers and sponsors here). Follow the journal's guidelines on what to include in the Acknowledgement section.

ABBREVIATIONS

CCR2, CC chemokine receptor 2; CCL2, CC chemokine ligand 2; CCR5, CC chemokine receptor 5; TLC, thin layer chromatography..

REFERENCES

(Word Style "TF_References_Section"). References are placed at the end of the manuscript. Authors are responsible for the accuracy and completeness of all references. Examples of the recommended formats for the various reference types can be found at <http://pubs.acs.org/page/4authors/index.html>. Detailed information on reference style can be found in The ACS Style Guide, available from Oxford Press.