

Effect of Nintedanib on Lung Function in Patients With Systemic Sclerosis–Associated Interstitial Lung Disease: Further Analyses of a Randomized, Double-Blind, Placebo-Controlled Trial

Toby M. Maher,¹ Maureen D. Mayes,² Michael Kreuter,³ Elizabeth R. Volkman,⁴ Martin Aringer,⁵ Ivan Castellvi,⁶ Maurizio Cutolo,⁷ Christian Stock,⁸ Nils Schoof,⁹ Margarida Alves,⁹ and Ganesh Raghu,¹⁰ on behalf of the SENSIS Trial Investigators

Objective. In the SENSIS trial in subjects with systemic sclerosis–associated interstitial lung disease (SSc-ILD), nintedanib reduced the rate of decline in forced vital capacity (FVC) over 52 weeks by 44% versus placebo. This study was undertaken to investigate the effects of nintedanib on categorical changes in FVC and other measures of ILD progression.

Methods. In post hoc analyses, we assessed the proportions of subjects with categorical changes in FVC % predicted at week 52 and the time to absolute decline in FVC of $\geq 5\%$ predicted or death and absolute decline in FVC of $\geq 10\%$ predicted or death.

Results. A total of 288 subjects received nintedanib and 288 subjects received placebo. At week 52, in subjects treated with nintedanib and placebo, respectively, 55.7% and 66.3% had any decline in FVC % predicted, 13.6% and 20.1% had a decline in FVC of $>5\%$ to $\leq 10\%$ predicted, and 3.5% and 5.2% had a decline in FVC of $>10\%$ to $\leq 15\%$ predicted; 34.5% and 43.8% had a decrease in FVC of $\geq 3.3\%$ predicted (proposed minimal clinically important difference [MCID] for worsening of FVC), while 23.0% and 14.9% had an increase in FVC of $\geq 3.0\%$ predicted (proposed MCID for improvement in FVC). Over 52 weeks, the hazard ratio (HR) for an absolute decline in FVC of $\geq 5\%$ predicted or death with nintedanib versus placebo was 0.83 (95% confidence interval [95% CI] 0.66–1.06) ($P = 0.14$), and the HR for an absolute decline in FVC of $\geq 10\%$ predicted was 0.64 (95% CI 0.43–0.95) ($P = 0.029$).

Conclusion. These results suggest that nintedanib has a clinically relevant benefit on the progression of SSc-ILD.

INTRODUCTION

Systemic sclerosis (SSc) is a rare and heterogeneous autoimmune disease characterized by microvascular damage and

progressive fibrosis of the skin and internal organs (1). Interstitial lung disease (ILD) is a common manifestation of SSc and the leading cause of death in subjects with SSc (2). The course of SSc-associated ILD (SSc-ILD) is unpredictable, but a decline in forced

ClinicalTrials.gov identifier: NCT02597933.

Supported by Boehringer Ingelheim.

¹Toby M. Maher, MD: National Heart and Lung Institute, Imperial College London and NIHR Clinical Research Facility, Royal Brompton Hospital, London, UK, and Keck School of Medicine, University of Southern California, Los Angeles; ²Maureen D. Mayes, MD: University of Texas McGovern Medical School, Houston; ³Michael Kreuter, MD: Thoraxklinik University of Heidelberg, and the German Center for Lung Research, Heidelberg, Germany; ⁴Elizabeth R. Volkman, MD: University of California, David Geffen School of Medicine, Los Angeles; ⁵Martin Aringer, MD: University Medical Center and Faculty of Medicine Carl Gustav Carus, TU Dresden, Dresden, Germany; ⁶Ivan Castellvi, MD: Hospital de la Santa Creu i Sant Pau, Barcelona, Spain; ⁷Maurizio Cutolo, MD: University of Genova, IRCCS San Martino Polyclinic Hospital, Genova, Italy; ⁸Christian Stock, PhD: Boehringer Ingelheim Pharma GmbH & Co. KG, Ingelheim am Rhein, Germany; ⁹Nils Schoof, PhD, Margarida Alves, MD: Boehringer Ingelheim International GmbH, Ingelheim am Rhein, Germany; ¹⁰Ganesh Raghu, MD: University of Washington, Seattle.

Dr. Maher has received consulting fees, speaking fees, and/or honoraria from AstraZeneca, Blade Therapeutics, Boehringer Ingelheim, Bristol Myers Squibb, Galápagos, GlaxoSmithKline, Indalo, Pliant, Respivant, Roche, and Theravance (less than \$10,000 each). Dr. Mayes has received consulting fees,

speaking fees, and/or honoraria from Boehringer Ingelheim, Corbus, Eicos Sciences, Galápagos, and Mitsubishi Tanabe (less than \$10,000 each). Dr. Kreuter has received consulting fees, speaking fees, and/or honoraria from Boehringer Ingelheim and Roche (less than \$10,000 each). Dr. Volkman has received consulting fees, speaking fees, and/or honoraria from Boehringer Ingelheim (more than \$10,000). Dr. Aringer has received consulting fees, speaking fees, and/or honoraria from Boehringer Ingelheim (less than \$10,000). Dr. Castellvi has received consulting fees, speaking fees, and/or honoraria from Boehringer Ingelheim, Bristol Myers Squibb, Janssen-Cilag, Roche, and Sanofi (less than \$10,000 each). Dr. Cutolo has received consulting fees, speaking fees, and/or honoraria from Bristol Myers Squibb (less than \$10,000). Drs. Stock, Schoof, and Alves are employees of Boehringer Ingelheim. Dr. Raghu has received consulting fees, speaking fees, and/or honoraria from Roche-Genentech (less than \$10,000) and from Boehringer Ingelheim (more than \$10,000).

Address correspondence to Toby M. Maher, Keck School of Medicine, University of Southern California, Los Angeles, CA 90033. Email: toby.maher@med.usc.edu.

Submitted for publication March 17, 2020; accepted in revised form October 29, 2020.

vital capacity (FVC) in subjects with SSc-ILD is an indicator of ILD progression and is associated with mortality (3–7). While there is no established definition for progression of ILD, in 2015, the Outcome Measures in Rheumatology (OMERACT) connective tissue disease–associated ILD (CTD-ILD) Working Group agreed that a relative decline in FVC % predicted of $\geq 10\%$, or a relative decline in FVC % predicted of $\geq 5\%$ to $< 10\%$ with a relative decline in diffusing capacity for carbon monoxide (DL_{CO}) % predicted $\geq 15\%$, represents clinically meaningful progression of SSc-ILD (8).

Nintedanib, an intracellular inhibitor of tyrosine kinases (9), has been approved in many countries for the treatment of idiopathic pulmonary fibrosis (IPF) and SSc-ILD. Clinical trials in subjects with IPF (INPULSIS) (10), with SSc-ILD (SENSCIS) (11), and with various forms of progressive fibrosing ILDs (INBUILD) (12), including progressive autoimmune disease-related ILDs (13), have shown that nintedanib reduces the rate of decline in FVC (milliliters/year). The objective of the current analyses was to investigate the effects of nintedanib on categorical changes in FVC and other measures of ILD progression in subjects with SSc-ILD in the SENSCIS trial.

PATIENTS AND METHODS

Trial design. The design of the SENSCIS trial (ClinicalTrials.gov identifier: NCT02597933) has been published previously, together with the trial protocol (11). Briefly, eligible subjects had SSc with onset of first non-Raynaud's phenomenon symptom < 7 years before screening, extent of fibrotic ILD $\geq 10\%$ on a high-resolution computed tomography (HRCT) scan (based on assessment of the whole lung), FVC $\geq 40\%$ predicted, and DL_{CO} of the lung (corrected for hemoglobin) 30–89% predicted. Subjects receiving prednisone ≤ 10 mg/day or equivalent and/or stable therapy with mycophenolate or methotrexate for ≥ 6 months prior to randomization were allowed to participate. Spirometers were provided to the sites and the results were confirmed centrally. Spirometry was performed in accordance with guidelines issued by the American Thoracic Society and European Respiratory Society (14), including daily calibration of the spirometer. Percent predicted values for FVC were calculated using the Global Lung Initiative equations based on the subject's age, sex, race, and height (15). The method used to measure DL_{CO} and the equation used to calculate percent predicted values for DL_{CO} were chosen by the site.

Subjects were randomized 1:1 to receive nintedanib 150 mg twice a day or placebo, stratified by the presence of anti-topoisomerase I (anti-topo I) antibodies. Subjects could continue to receive treatment in a blinded manner until the last subject had reached week 52, but for ≤ 100 weeks. Subjects who discontinued treatment prematurely were asked to remain in the trial and attend visits as originally planned. The trial was conducted in accordance with the trial protocol, the principles of the Declaration of Helsinki, and the International Council for Harmonisation Guidelines for Good Clinical Practice, and was approved by local authorities. All subjects provided written informed consent.

End points. The following were assessed in the nintedanib and placebo groups based on data at week 52: the proportion of subjects with categorical absolute declines or increases in FVC % predicted or categorical relative declines or increases in FVC (milliliters) (as listed in the Results section); the proportion of subjects who met proposed thresholds for minimal clinically important differences (MCID) for improvement in FVC (absolute increase of $\geq 3.0\%$ predicted), stable FVC (absolute increase of $< 3.0\%$ predicted or decrease of $< 3.3\%$ predicted), and worsening of FVC (absolute decrease of $\geq 3.3\%$ predicted) based on data from Scleroderma Lung Studies I and II, anchored to the health transition question from the Medical Outcomes Study Short Form-36 (16); and time to 1) an absolute decline in FVC of $\geq 5\%$ predicted or death; 2) an absolute decline in FVC of $\geq 10\%$ predicted or death; and 3) an absolute decline in FVC of $\geq 10\%$ predicted or absolute decline in FVC of $\geq 5\%$ to $< 10\%$ predicted with an absolute decline in DL_{CO} of $\geq 15\%$ predicted, or death. The proportions of subjects who met proposed thresholds for improvement in FVC, stable FVC, and worsening of FVC at week 52 were also assessed in subgroups defined by the following baseline characteristics: FVC of $< 80\%$ versus $\geq 80\%$ predicted, extent of fibrotic ILD on HRCT of $< 20\%$ versus $\geq 20\%$, time since onset of first non-Raynaud's phenomenon symptom ≤ 3 versus > 3 years, and glucocorticoid use.

Statistical analysis. Analyses were conducted post hoc in subjects who received ≥ 1 dose of trial medication (intention-to-treat). The proportions of subjects with categorical declines or increases in FVC % predicted and FVC measured in milliliters at week 52, and the proportions of subjects who met proposed thresholds for improvement in FVC, stable FVC, and worsening of FVC at week 52 were compared between treatment groups using a Cochran-Mantel-Haenszel test, stratified by anti-topo I antibody status. In the subgroup analyses, the proportions of subjects who met proposed thresholds for improvement in FVC, stable FVC, and worsening of FVC at week 52 were compared between treatment groups using a logistic regression model, including trial medication (nintedanib/placebo), anti-topo I antibody status, subgroup, and treatment-by-subgroup interaction as terms. Missing values were imputed using a worst value carried forward approach. Exploratory interaction *P* values were calculated to assess potential heterogeneity in the treatment effect of nintedanib versus placebo across the subgroups. Time-to-event end points were analyzed based on data from 52 weeks (± 7 days) using a Cox regression model with a term for treatment and stratified by anti-topo I antibody status. Analyses were not adjusted for multiplicity. Hazard ratios (HRs) or odds ratios and 95% confidence intervals (95% CIs) were calculated.

RESULTS

Characteristics of the study subjects. A total of 576 subjects received ≥ 1 dose of trial medication (288 received nintedanib and 288 received placebo). The baseline




Figure 1. Proportions of subjects with systemic sclerosis-associated interstitial lung disease (SSc-ILD) treated with nintedanib or placebo in the SENSICIS trial who had the indicated absolute increases and declines in forced vital capacity (FVC) % predicted at week 52. A post-baseline FVC measurement was not available for 1 patient.

characteristics of subjects in the SENSICIS trial have been described previously (11). The mean \pm SD age was 54.0 ± 12.2 years, 75.2% of subjects were female, and 67.2% were White. The mean \pm SD FVC was $2,500 \pm 777$ milliliters and 72.5 ± 16.7 % predicted, and the mean \pm SD DL_{CO} was 53.0 ± 15.1 % predicted. Baseline characteristics were similar between the treatment groups (11).

Categorical changes in FVC % predicted. In total, 46 subjects (16.0%) in the nintedanib group and 31 subjects (10.8%) in the placebo group had missing FVC data at week 52. At week 52, 55.7% of subjects in the nintedanib group and 66.3% of subjects in the placebo group had a decline in FVC % predicted. At week 52 the proportions of subjects in the nintedanib group with an absolute decline in FVC of $> 5\%$ to $\leq 10\%$ predicted and an absolute decline in FVC of $> 10\%$ to $\leq 15\%$ predicted were 13.6% and 3.5%, respectively, while the proportions of subjects in the

placebo group with an absolute decline in FVC of $> 5\%$ to $\leq 10\%$ predicted and an absolute decline in FVC of $> 10\%$ to $\leq 15\%$ predicted were 20.1% and 5.2%, respectively (Figure 1). The proportions of subjects who met thresholds for relative declines or increases in FVC (in milliliters) are shown in Supplementary Figure 1 (available on the *Arthritis & Rheumatology* website at <http://onlinelibrary.wiley.com/doi/10.1002/art.41576/abstract>). In the nintedanib and placebo groups, respectively, 34.5% versus 43.8% of subjects had an absolute decrease in FVC of $\geq 3.3\%$ predicted at week 52 (the proposed MCID for worsening of FVC), while 23.0% versus 14.9% had an absolute increase in FVC of $\geq 3.0\%$ predicted at week 52 (the proposed MCID for improvement in FVC) (Figure 2). Exploratory interaction *P* values did not indicate heterogeneity in the effect of nintedanib versus placebo between subgroups classified by FVC % predicted, extent of fibrotic ILD on HRCT, time since onset of first non-Raynaud's phenomenon symptom at baseline, or glucocorticoid




Figure 2. Proportions of subjects with SSc-ILD treated with nintedanib or placebo in the SENSICIS trial who met the proposed threshold for worsening of FVC (decrease of $\geq 3.3\%$ predicted), stable FVC (increase of $< 3.0\%$ predicted or decrease of $< 3.3\%$ predicted), or improvement in FVC (increase of $\geq 3.0\%$ predicted) at week 52. A post-baseline FVC measurement was not available for 1 patient. OR = odds ratio; 95% CI = 95% confidence interval (see Figure 1 for other definitions).

Table 1. Proportions of subjects with absolute declines in FVC % predicted or death over 52 weeks in the SENSICIS trial*

	Nintedanib (n = 288)	Placebo (n = 288)
Absolute decline in FVC $\geq 5\%$ predicted or death		
Subjects with event, no. (%)	124 (43.1)	145 (50.3)
Hazard ratio (95% confidence interval)	0.83 (0.66–1.06)	
P	0.14	
Absolute decline in FVC $\geq 10\%$ predicted or death		
Subjects with event, no. (%)	40 (13.9)	62 (21.5)
Hazard ratio (95% confidence interval)	0.64 (0.43–0.95)	
P	0.03	
Absolute decline in FVC $\geq 10\%$ predicted or absolute decline in FVC $\geq 5\%$ to $<10\%$ predicted with absolute decline in DLco $\geq 15\%$ predicted, or death		
Subjects with event, no. (%)	39 (13.5)	69 (22.9)
Hazard ratio (95% confidence interval)	0.58 (0.39–0.87)	
P	0.008	

* FVC = forced vital capacity; DLco = diffusing capacity for carbon monoxide.

use ($P > 0.05$ for treatment-by-subgroup interactions) (Supplementary Figure 2, available on the *Arthritis & Rheumatology* website at <http://onlinelibrary.wiley.com/doi/10.1002/art.41576/abstract>).

Time to lung function decline or death. Over 52 weeks, an absolute decline in FVC of $\geq 5\%$ predicted or death occurred in 43.1% of the subjects in the nintedanib group and 50.3% of the subjects in the placebo group (HR 0.83 [95% CI 0.66–1.06]; $P = 0.14$), and an absolute decline in FVC of $\geq 10\%$ predicted or death occurred in 13.9% of the subjects in the nintedanib group and 21.5% of the subjects in the placebo group (HR 0.64 [95% CI 0.43–0.95]; $P = 0.029$) (Table 1 and Figure 3). Over 52 weeks, an absolute decline in FVC of $\geq 10\%$ predicted, absolute decline in FVC of $\geq 5\%$ to $<10\%$ predicted with an absolute decline in DLco of $\geq 15\%$ predicted, or death occurred in 13.5% and 22.9% of subjects in the nintedanib and placebo groups, respectively (HR 0.58 [95% CI 0.39–0.87]; $P = 0.008$) (Table 1).

DISCUSSION

These findings from the SENSICIS trial provide further evidence that nintedanib has a clinically relevant effect on the

progression of SSc-ILD. Although there is no established definition for the progression of ILD, declines in FVC of $>10\%$ predicted have been used to assess the proportion of subjects with clinically relevant ILD progression in previous studies of SSc-ILD (6,7) and other ILDs (17–20), based on the association between decline in FVC and mortality. In addition, thresholds for improvement and worsening of FVC, derived based on anchoring to the health transition question from the Medical Outcomes Study Short Form-36 in the Scleroderma Lung Studies I and II, have been proposed as MCIDs at a population level (16). In a recent European Delphi consensus study, physicians experienced in the management of SSc-ILD agreed that the progression of SSc-ILD can be assessed using changes in % predicted values for FVC and DLco and that measurement of lung function is an effective tool in long-term follow-up of ILD progression in patients with SSc-ILD (21).

Several studies have shown that a decline in FVC is associated with mortality in patients with SSc-ILD. A study of 171 patients at a single center showed that patients who died within 4 years of SSc-ILD diagnosis had a higher annual rate of decline in FVC than those who died between 4 and 8 years after their SSc-ILD diagnosis or who survived for >8 years (22). Data from Scleroderma Lung Study I ($n = 158$) showed that absolute declines in FVC of $\geq 10\%$




Figure 3. Time to **A**, absolute decline in FVC of $\geq 5\%$ predicted or death and **B**, absolute decline in FVC of $\geq 10\%$ predicted or death, over 52 weeks in patients with SSc-ILD treated with nintedanib or placebo in the SENSICIS trial. See Figure 1 for definitions.

predicted or DL_{CO} of $\geq 15\%$ predicted over 2 years were associated with mortality over a median follow-up period of 8 years, while data from Scleroderma Lung Study II (n = 142) showed that an absolute decline in FVC of $\geq 10\%$ predicted at 1 year was associated with mortality over a median follow-up period of 4 years (7). In a large cohort of subjects in the European Scleroderma Trials and Research (EUSTAR) database (n = 857), an absolute decline in FVC of $\geq 10\%$ predicted, or an absolute decline in FVC of $\geq 5\%$ predicted with a decline in DL_{CO} of $\geq 15\%$ predicted, over 12 months was predictive of mortality over a maximum follow-up period of 5 years (5). Recent data from a well-characterized Norwegian cohort (n = 391) showed that ILD progression defined as severe (absolute decline in FVC of $> 10\%$ predicted or decline in FVC of 5–10% predicted with decline in DL_{CO} of $\geq 15\%$ predicted) or moderate (absolute decline in FVC of 5–10% predicted with decline in DL_{CO} of $< 15\%$ predicted) over a mean follow-up period of almost 4 years was associated with lower survival compared with stable FVC (change $< 5\%$ predicted), with 10-year survival rates of 59% versus 78% (6). In a long-term UK study of 162 patients, a relative decline in FVC (in milliliters) of $\geq 10\%$, or a relative decline in FVC (in milliliters) of 5–9% with a relative decline in DL_{CO} of $> 15\%$ at 1 year was strongly associated with mortality over 15 years (4).

Subgroup analyses of the data from the SENSICIS trial suggest that the proportions of subjects who met proposed thresholds for worsening and improvement in FVC were consistent across subgroups based on FVC % predicted, time since onset of first non-Raynaud's phenomenon symptom, extent of fibrotic ILD on HRCT, and glucocorticoid use at baseline. Previous analyses have shown that the proportions of subjects who met these thresholds were similar across subgroups by anti-topo I antibody status (23), SSc subtype (limited versus diffuse cutaneous SSc) (24), and mycophenolate use at baseline (25). Taken together, these data support a benefit of nintedanib in reducing the proportion of patients with clinically relevant progression of ILD, and increasing the proportion of patients with stable or increased FVC, across a broad population of subjects with SSc-ILD, consistent with the effects of nintedanib previously demonstrated in patients with IPF (26–28).

Strengths of our study include the participation of a large number of well-characterized subjects with SSc-ILD and the highly standardized procedure used for measurement of FVC. Limitations of our analyses include that they were conducted post hoc and, as such, should be considered exploratory. The present study did not assess whether the categorical changes in FVC translated into meaningful improvements/declines in patient-reported outcomes. Our analyses were not adjusted for multiple testing or for confounding factors such as use of mycophenolate. The number of deaths was too small to enable associations between FVC decline and mortality to be studied.

In conclusion, these further analyses of FVC decline in the SENSICIS trial support a clinically meaningful effect of nintedanib on slowing the progression of SSc-ILD.

ACKNOWLEDGMENTS

We would like to thank the subjects who participated in this trial.

AUTHOR CONTRIBUTIONS

All authors were involved in drafting the article or revising it critically for important intellectual content, and all authors approved the final version to be published. Dr. Maher had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study conception and design. Maher, Mayes, Stock, Schoof, Alves, Raghu.

Acquisition of data. Maher, Mayes, Kreuter, Aringer, Castellvi, Cutolo.

Analysis and interpretation of data. Maher, Mayes, Kreuter, Volkman, Aringer, Castellvi, Cutolo, Stock, Schoof, Alves, Raghu.

ROLE OF THE STUDY SPONSOR

Boehringer Ingelheim was involved in the design of the study, the interpretation of the data, and the writing of the manuscript. The authors had the final decision to submit the manuscript for publication. Writing assistance was provided by Julie Fleming and Wendy Morris (Fleishman-Hillard Fishburn, London, UK; supported financially by Boehringer Ingelheim).

REFERENCES

1. Van den Hoogen F, Khanna D, Fransen J, Johnson SR, Baron M, Tyndall A, et al. 2013 classification criteria for systemic sclerosis: an American College of Rheumatology/European League against Rheumatism collaborative initiative. *Arthritis Rheum* 2013;65:2737–47.
2. Elhai M, Meune C, Boubaya M, Avouac J, Hachulla E, Balbir-Gurman A, et al. Mapping and predicting mortality from systemic sclerosis. *Ann Rheum Dis* 2017;76:1897–905.
3. Goh NS, Desai SR, Veeraraghavan S, Hansell DM, Copley SJ, Maher TM, et al. Interstitial lung disease in systemic sclerosis: a simple staging system. *Am J Respir Crit Care Med* 2008;177:1248–54.
4. Goh NS, Hoyle RK, Denton CP, Hansell DM, Renzoni EA, Maher TM, et al. Short-term pulmonary function trends are predictive of mortality in interstitial lung disease associated with systemic sclerosis. *Arthritis Rheumatol* 2017;69:1670–8.
5. Sobanski V, de Vries-Bouwstra J, Hoffmann-Vold A, Huscher D, Alves M, Matucci-Cerinic M, et al. Association of short-term longitudinal changes in clinical and physiologic variables with overall survival in patients with systemic sclerosis-associated interstitial lung disease [abstract]. *Arthritis Rheumatol* 2018;70 Suppl 10. URL: <https://acrabstracts.org/abstract/association-of-short-term-longitudinal-changes-in-clinical-and-physiologic-variables-with-overall-survival-in-patients-with-systemic-sclerosis-associated-interstitial-lung-disease-ssc-ild/>.
6. Hoffmann-Vold AM, Fretheim H, Halse AK, Seip M, Bitter H, Wallenius M, et al. Tracking impact of interstitial lung disease in systemic sclerosis in a complete nationwide cohort. *Am J Respir Crit Care Med* 2019;200:1258–66.
7. Volkman ER, Tashkin DP, Sim M, Li N, Goldmuntz E, Keyes-Elstein L, et al. Short-term progression of interstitial lung disease in systemic sclerosis predicts long-term survival in two independent clinical trial cohorts. *Ann Rheum Dis* 2019;78:122–30.
8. Khanna D, Mittoo S, Aggarwal R, Proudman SM, Dalbeth N, Matteson EL, et al. Connective Tissue Disease-associated Interstitial

- Lung Diseases (CTD-ILD): report from OMERACT CTD-ILD Working Group. *J Rheumatol* 2015;42:2168–71.
9. Wollin L, Distler JH, Redente EF, Riches DW, Stowasser S, Schlenker-Herceg R, et al. Potential of nintedanib in treatment of progressive fibrosing interstitial lung diseases. *Eur Respir J* 2019; 54:1900161.
 10. Richeldi L, du Bois RM, Raghu G, Azuma A, Brown KK, Costabel U, et al. Efficacy and safety of nintedanib in idiopathic pulmonary fibrosis. *N Engl J Med* 2014;370:2071–82.
 11. Distler O, Highland KB, Gahlemann M, Azuma A, Fischer A, Mayes MD, et al. Nintedanib for systemic sclerosis-associated interstitial lung disease. *N Engl J Med* 2019;380:2518–28.
 12. Flaherty KR, Wells AU, Cottin V, Devaraj A, Walsh SL, Inoue Y, et al. Nintedanib in progressive fibrosing interstitial lung diseases. *N Engl J Med* 2019;381:1718–27.
 13. Wells AU, Flaherty KR, Brown KK, Inoue Y, Devaraj A, Richeldi L, et al. Nintedanib in patients with progressive fibrosing interstitial lung diseases: subgroup analyses by interstitial lung disease diagnosis in the randomised, placebo-controlled INBUILD trial. *Lancet Respir Med* 2020;8:453–60.
 14. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J* 2005;26:319–38.
 15. Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH, et al. Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *Eur Respir J* 2012;40:1324–43.
 16. Kafaja S, Clements PJ, Wilhalme H, Tseng CH, Furst DE, Kim GH, et al. Reliability and minimal clinically important differences of forced vital capacity: results from the Scleroderma Lung Studies (SLS-I and SLS-II). *Am J Respir Crit Care Med* 2018;197:644–52.
 17. Zappala CJ, Latsi PI, Nicholson AG, Colby TV, Cramer D, Renzoni EA, et al. Marginal decline in forced vital capacity is associated with a poor outcome in idiopathic pulmonary fibrosis. *Eur Respir J* 2010;35:830–6.
 18. Solomon JJ, Chung JH, Cosgrove GP, Demoruelle MK, Fernandez-Perez ER, Fischer A, et al. Predictors of mortality in rheumatoid arthritis-associated interstitial lung disease. *Eur Respir J* 2016;47:588–96.
 19. Gimenez A, Storrer K, Kuranishi L, Soares MR, Ferreira RG, Pereira CA. Change in FVC and survival in chronic fibrotic hypersensitivity pneumonitis [letter]. *Thorax* 2017;73:391–2.
 20. Paterniti MO, Bi Y, Rekić D, Wang Y, Karimi-Shah BA, Chowdhury BA. Acute exacerbation and decline in forced vital capacity are associated with increased mortality in idiopathic pulmonary fibrosis. *Ann Am Thorac Soc* 2017;14:1395–402.
 21. Hoffmann-Vold A, Maher TM, Philpot EE, Ashrafzadeh A, Barake R, Barsotti S, et al. The identification and management of interstitial lung disease in systemic sclerosis: evidence-based European consensus statements. *Lancet Rheumatol* 2020;2:E71–83.
 22. Guler SA, Winstone TA, Murphy D, Hague C, Soon J, Sulaiman N, et al. Does systemic sclerosis-associated interstitial lung disease burn out? Specific phenotypes of disease progression. *Ann Am Thorac Soc* 2018;15:1427–33.
 23. Mayes MD, Highland KB, Gahlemann M, Fischer A, Raghu G, Girard M, et al. Effect of anti-topoisomerase I antibody status on decline in lung function in patients with systemic sclerosis-associated interstitial lung disease (SSc-ILD): data from the SENSICIS trial [poster]. American College of Rheumatology: Poster presented at the 2019 Annual Meeting, Atlanta, Georgia. November 8–13, 2019. URL: http://ILDPosters2019.com/pdf/ACR_SENSICISATASubgroups_Mayes.pdf.
 24. Kuwana M, Highland KB, Gahlemann M, Denton CP, Fischer A, Mayes MD, et al. Effects of nintedanib in patients with diffuse and limited cutaneous systemic sclerosis and interstitial lung disease: subgroup analysis of the SENSICIS trial. American College of Rheumatology: Poster presented at the 2019 Annual Meeting, Atlanta, Georgia. November 8–13, 2019. URL: http://ILDPosters2019.com/pdf/ACR_SENSICIScSScvsdcSSc_Kuwana.pdf.
 25. Highland KB, Distler O, Kuwana M, Allanore Y, Assassi S, Azuma A, et al. Efficacy and safety of nintedanib in patients with systemic sclerosis-associated interstitial lung disease treated with mycophenolate: subgroup analysis of the SENSICIS trial. *Lancet Respir Med* 2021;9:96–106.
 26. Costabel U, Inoue Y, Richeldi L, Collard HR, Tschöepe I, Stowasser S, et al. Efficacy of nintedanib in idiopathic pulmonary fibrosis across prespecified subgroups in INPULSIS. *Am J Respir Crit Care Med* 2016;193:178–85.
 27. Brown KK, Flaherty KR, Cottin V, Raghu G, Inoue Y, Azuma A, et al. Lung function outcomes in the INPULSIS trials of nintedanib in idiopathic pulmonary fibrosis. *Respir Med* 2019;146:42–8.
 28. Richeldi L, Crestani B, Azuma A, Kolb M, Selman M, Stansen W, et al. Outcomes following decline in forced vital capacity in patients with idiopathic pulmonary fibrosis: results from the INPULSIS and INPULSIS-ON trials of nintedanib. *Respir Med* 2019;156:20–5.