

<https://helda.helsinki.fi>

Disease control and psychiatric comorbidity among adolescents with chronic medical conditions : a single-centre retrospective study

Kallio, Mira

2023-02

Kallio , M , Tornivuori , A , Miettinen , P , Kolho , K-L , Culnane , E , Sawyer , S & Kosola , S
2023 , ' Disease control and psychiatric comorbidity among adolescents with chronic medical
conditions : a single-centre retrospective study ' , BMJ Paediatrics Open , vol. 7 , no. 1 ,
e001605 . <https://doi.org/10.1136/bmjpo-2022-001605>

<http://hdl.handle.net/10138/356287>

<https://doi.org/10.1136/bmjpo-2022-001605>

cc_by_nc

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Disease control and psychiatric comorbidity among adolescents with chronic medical conditions: a single-centre retrospective study

Mira Kallio ^{1,2}, Anna Tornivuori,^{1,3} Päivi Miettinen,^{1,2} Kaija-Leena Kolho,^{1,2,4} Evelyn Culnane,⁵ Susan Sawyer,^{6,7,8,9} Silja Kosola¹

To cite: Kallio M, Tornivuori A, Miettinen P, *et al*. Disease control and psychiatric comorbidity among adolescents with chronic medical conditions: a single-centre retrospective study. *BMJ Paediatrics Open* 2023;**7**:e001605. doi:10.1136/bmjpo-2022-001605

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjpo-2022-001605>).

Received 8 July 2022

Accepted 16 January 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Mira Kallio; mira.kallio@hus.fi

ABSTRACT

Background To investigate disease control, psychiatric comorbidity, substance use and their possible associations in adolescents with chronic medical conditions before transfer to adult healthcare.

Methods We collected clinical data from the year preceding transfer of care and psychiatric data from the records of the paediatric hospital in Helsinki, Finland (population base 1.7 million). Participants were grouped into three disease and/or adherence control categories (good, some evidence of concern, poor) based on clinical data from the medical records of the year preceding the transfer of care. Participants completed the Adolescent's Substance Use Measurement Questionnaire before transfer of care and were divided into four risk subgroups accordingly.

Results In total, 253 adolescents (mean age 17.3 years, SD 1.2) from six paediatric subspecialties participated in this study. Disease control and/or adherence were rated as good in 28% (n=70), moderate in 42% (n=105) and poor in 30% (n=76) in the year before participants transferred to adult health services. A quarter of participants had at least one psychiatric diagnosis during adolescence. Adolescents with concomitant psychiatric diagnoses more often had poor disease control of their chronic medical condition than adolescents with only a medical condition (44% vs 26%; n=25 of 59 vs 51 of 194, respectively). More than half of adolescents (56%) were abstinent or used substances infrequently; 10% (n=26) reported hazardous substance use.

Conclusions Psychiatric comorbidity in adolescents with chronic medical conditions is common. Its negative association with disease control and possible substance use should be considered in the transition process to adult health services.

INTRODUCTION

The landscape of chronic illness in children and adolescents with chronic medical conditions has changed. Over the past few decades, treatment success has resulted in improved survival of many of these young people, which together with an increased incidence of particular chronic medical disorders in childhood,¹⁻³ results in increasing numbers

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Adolescents with chronic medical disorders have a higher prevalence of mental health disorders than their peers, but the impact of mental health disorders on disease control prior to transfer to adult healthcare is poorly understood.

WHAT THIS STUDY ADDS

⇒ Adolescents with chronic medical disorders and comorbid psychiatric diagnoses more frequently have poor control of their medical condition than those without psychiatric comorbidities, which highlights the necessity of a holistic approach to care.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Ensuring that mental health disorders are identified and treated is an important aspect of caring for adolescents and young adults with chronic medical conditions, including at the time they transfer to adult services. Future research to measure the potential impact of adolescent mental health disorders on transfer to adult health service is indicated.

of adolescents requiring regular follow-up in adult health services. Living with a chronic medical condition can pose many challenges during adolescence, not least due to the requirement for regular medication and other treatments, frequent medical appointments and the uncertainty of future treatment such as surgical procedures. These demands for health management and monitoring, together with the impact of the medical condition itself, such as pain and fatigue,⁴ can reduce participation in normative activities with effects on peer relationships, extracurricular participation and school attendance.⁵

Chronic medical conditions are associated with increased rates of mental health problems.^{6,7} In the USA, 25% of children with epilepsy have anxiety and/or depression compared with the general population aged

12–17 years old, of whom 6% have a current diagnosis of depression and 11% anxiety.^{8 9} During a decade of follow-up, 17% of Swedish children with inflammatory bowel disease (IBD) received a diagnosis of any psychiatric disorder compared with 12% of the general population.¹⁰ Population surveys that assess self-reported symptoms of depression or anxiety yield even higher proportions of affected young people.¹¹ So the proportion of adolescents with chronic medical conditions who struggle with symptoms of mental health disorders is expected to be even higher.

Mental health problems, together with pain and fatigue,⁴ can affect disease control, adherence behaviours, and health-related quality of life in children and adolescents with chronic medical conditions.^{12 13} For example, depression is associated with higher healthcare costs and emergency department visits in children and adolescents with IBD,¹⁴ but which young people are most at risk is still incompletely understood.

Adolescents living with chronic medical conditions often engage in risk behaviours, such as substance use.¹⁵ Substance use in adolescence is frequently associated with other mental health disorders, and the incidence of both substance use and mental health disorders increases across adolescence and into adulthood.^{16–18} This accumulation of risk behaviours and mental health disorders among adolescents and young adults suggests it is important to identify which adolescents with chronic medical disorders are at greatest risk so that they can be appropriately managed, including at the time they transfer to adult health services.

In this study of adolescents with different chronic medical conditions, we aimed to investigate disease control, rates of psychiatric comorbidity and risky substance use and their associations before adolescent care is transferred to adult health services. Our hypothesis was that psychiatric comorbidity is common and associated with poorer disease control and higher rates of substance use.

METHODS

Study design

This study is part of an international prospective cohort study named the Bridge, for which the study protocol is available online.¹⁹ In this cross-sectional analysis, we combined retrospective data with questionnaire responses from one study site. The Bridge trial has been registered with ClinicalTrials.gov (ID number NCT04631965).

Patient and public involvement

Fifteen adolescents with chronic conditions participated in the design of the study questionnaires.

Participants

Participants were recruited from the New Children's Hospital, Helsinki, Finland (hospital catchment area 1.7 million inhabitants) between September 2017 and

August 2019. In Finland, practically all adolescents with chronic medical conditions are managed by public health services, with few cared for by private doctors. Clinicians from six different paediatric subspecialty clinics (endocrinology, gastroenterology, rheumatology, nephrology and organ transplants, neurology and cardiology) identified adolescents who were expected to transfer to adult clinics within the coming 6–12 months. Inclusion criteria were that participants were aged 15–24 years and had a chronic medical condition that had been diagnosed at least 6 months earlier and was expected to require regular follow-up in adult health services. Eligible participants also needed to have sufficient cognitive and linguistic ability to communicate in Finnish, English or Swedish. Adolescents received a movie ticket (value €10) after completing the survey in acknowledgement of their time.

A research nurse, uninvolved in patient care, personally met 306 consecutive adolescents during their routine hospital visits and provided oral and written information about the Bridge Study.

Demographics

Background information was obtained from the baseline survey, including gender, residential area, home language and family composition (living with two parents or any other family type).

Disease control

To determine disease control, we gathered data and laboratory test results from the electronic medical records for the year preceding the final visit at the New Children's Hospital. The final visit was specified as the outpatient appointment where the clinician made a formal referral to the respective adult health service. We divided participants into one of three categories according to the condition for which they were recruited: (1) good disease control and/or adherence; (2) some evidence of concern; and (3) poor disease control and/or adherence or more severe condition (see study protocol for further details).¹⁹ Where appropriate, we used published cut-offs for disease control. For adolescents with diabetes, a mean glycosylated haemoglobin (the mean of all measurements across the year preceding transfer of care) ≤ 53 mmol/mol was categorised as good control/adherence, 54–69 mmol/mol was considered evidence of some concern, while ≥ 70 mmol/mol was classified as having poor control/adherence. For adolescents with rheumatic disease, the 10-joint Juvenile Arthritis Disease Activity Score was completed. For those with oligoarthritis, cut-off points of ≤ 0.5 (good), 0.6–2.8 (some concern) and > 2.8 (poor) were used, while for those with polyarthritis, the cut-off points were ≤ 0.7 , 0.8–4 and > 4 , as previously applied.^{19–21} For adolescents with IBD, good control/adherence required minimal pain (Visual Analogue Scale, VAS 1–2), at least 80% of faecal calprotectin results < 100 $\mu\text{g/g}$ and always < 300 $\mu\text{g/g}$, medication unchanged or reduced, and no inpatient care; indicators of some evidence of concern were VAS 3–5, $< 80\%$ of faecal

calprotectin results within target range or exceeded 300 µg/g even once, but no significant medication changes nor inpatient care; and for poor control/adherence any of the following: VAS ≥6, significant changes in medication, need for corticosteroids and/or commencement of biological medication or an episode of inpatient care. For rare conditions without standard criteria of disease control and/or adherence, experienced clinicians subjectively categorised participants according to their symptoms, clinical and laboratory findings, need for inpatient care and changes in medication. MK first reviewed medical records of every participant, of whom 28 (11%) were difficult to categorise. For these 28 adolescents, SK made an independent assessment of disease control. Six adolescents were categorised differently, and the final grading was based on a consensus between the two reviewers. For some analyses, we combined the three smallest subspecialties (neurology, cardiology and nephrology/organ transplant) into 'others'.

Psychiatric diagnosis and treatment

Psychiatric diagnoses and treatment details were collected from the electronic medical records of the Helsinki University Hospital, which is the only public provider of adolescent psychiatric care in the study catchment area. Data were collected from ages 13 to 18 years, the age range of adolescent psychiatry in the study area. These data were collected separately from the disease control data, and in a different order of patients. Psychiatric diagnoses were divided into four categories: (1) mood disorders; (2) anxiety disorders; (3) others and (4) multiple diagnoses (ie, diagnosis from at least two different diagnostic categories). The New Children's Hospital has no general guideline regarding psychiatric assessment, and adolescents may have received referrals to adolescent psychiatry from community healthcare centres and student health services. Outpatient appointments with mental health professionals over these 6 years were divided into three categories: (1) no appointments; (2) one to three appointments and (3) more than three appointments. We also recorded the age at first psychiatric diagnosis, any admissions for psychiatric inpatient care and, if so, the number of days spent in psychiatric inpatient care.

Substance use

Participants completed the Adolescent Substance Use Measurement (ADSUME), a brief measure that was developed and validated in Finland to evaluate adolescent substance use and its possible consequences during the previous year. ADSUME is reliable when compared with validated adult questionnaire The Alcohol Use Disorders Identification Test (AUDIT) and may even be superior in detecting hazardous substance use among adolescents.²² We calculated a total score (0–31 points) of the first nine questions (online supplemental appendix 1) to yield a numerical score, which we used in analyses. We also divided participants into subgroups according

to guidelines from the Finnish Institute for Health and Welfare: (1) abstinence or experimental use, 0–3 points; (2) recurring use, 4–6 points; (3) risky use, 7–9 points; and (4) hazardous use, 10 points or more.²³

Data analysis

Categorical data are presented as frequencies (with percentages). For continuous variables, means and SDs or medians (with IQR) are used. Associations between disease control, psychiatric diagnoses and family composition were compared using Fisher's exact test (two-sided). To evaluate the associations between categorical data and continuous variables, we used non-parametric tests (Mann-Whitney U and Kruskal-Wallis). Finally, ordinal logistic regression with disease control as the dependent variable was conducted. Data analyses were performed using IBM SPSS Statistics V.25 and a p value of <0.05 was considered significant.

RESULTS

In total, 279 (91.2%) adolescents consented to participate, and 253 (82.7%) adolescents completed the baseline survey after a maximum of two reminders. Demographic and clinical data of the 253 study participants are summarised in [tables 1 and 2](#). The mean age at diagnosis of the medical condition was 9.0 years (SD 5.3) and the mean age at first psychiatric diagnosis was 13.7 years (SD 2.7). Age at transfer differed greatly between subspecialties: rheumatology patients were the youngest to transfer (mean age 16.2 years) and gastroenterology were the oldest (mean age 19.0 years; $p<0.001$ for difference between groups).

Among all study participants, 27.7% (n=70) were categorised as having good disease control and/or adherence, 41.5% (n=105) were rated as having some evidence of concern, and 30.0% (n=76) were classified as having poor disease control and/or adherence (or more severe condition) during the year preceding the transfer of care. Only 16% of adolescents with gastroenterological diseases and 14% of adolescents with diabetes had good disease control (mean of 4.6 measurements). Among adolescents with rheumatological disease, 39% had good disease control, while it was 48% in the group of 'others'. Gender and age at diagnosis showed no association with different disease control groups. Older age at transfer of care was associated with poorer disease control ($p=0.045$).

In this cohort, 23.3% of adolescents (n=59) had a psychiatric diagnosis, of whom 37.2% (n=22) had at least two different diagnoses. Anxiety disorders were most common (n=38), comprising 15.0% of all participants and 64.4% of adolescents with a psychiatric diagnosis. Females were over-represented in anxiety disorders and males in the 'other psychiatric diagnosis' group. Nine adolescents (3.6%) received inpatient psychiatric care, with a median of 22 days (range 9–956 days) per person. Age at medical diagnosis and age at psychiatric diagnosis showed no association ($p=0.32$).

Table 1 Demographic and clinical data of 253 adolescents

	N	%
Gender		
Male	118	46.6
Female	132	52.2
Other	3	1.2
Paediatric subspecialty		
Diabetes	92	36.4
Rheumatology	66	26.1
Gastroenterology	45	17.8
Cardiology	19	7.5
Neurology	18	7.1
Kidney or liver disease with/without transplant	13	5.1
Lives with		
Mother and father	158	62.5
Other than with two parents	95	37.5
Disease control*		
Good control and/or adherence	70	27.7
Some evidence of concern	105	41.5
Poor control and/or adherence or more severe condition	76	30.0
Psychiatric diagnoses	59	23.3
Anxiety disorders	38	15.0
Mood disorders	18	7.1
Others†	26	10.3
Many diagnoses	22	8.7
Outpatient visits in adolescent psychiatry		
None	159	62.8
1–3	36	14.2
More than 3	58	22.9
Median ADSUME total scores (IQR)‡	3	(0–6)
Females	2	(0–6)
Males	3	(0–6)
ADSUME risk category‡	N	%
Abstinence or experimental use (0–3 points)	142	56.1
Recurring use (4–6 points)	57	22.5
Risky use (7–9 points)	27	10.7
Hazardous use (10 points or more)	26	10.3

*Data available for 251 adolescents as transfer of two adolescents was delayed.
†Others: for example, conduct disorder, attention deficit-hyperactivity disorder, psychotic disorder, eating disorders, developmental disorders.
‡One response missing.
ADSUME, Adolescent Substance Use Measurement.

More than half of participants (n=142, 56.1%) were abstinent or reported only experimental substance use (table 1). Approximately half of the adolescents reported using alcohol in the last year. The median ADSUME total score was 3.0 and the maximum score was 27 points.

Table 2 Demographic and clinical data (n=253)

	Mean, years (median)	SD
Age at first medical diagnosis	9.0 (10.1)	5.3
Age at first psychiatric diagnosis	13.7 (14.3)	2.7
Age when completing the ADSUME	17.2 (16.8)	1.2
Age at transfer of care	17.3 (17.1)	1.2
Diabetes	17.2	0.7
Rheumatology	16.2	0.5
Gastroenterology	19.0	0.9
Others	17.8	1.2
Time between ADSUME and transfer of care	0.2 (0.0)	0.4
Time between medical diagnosis and transfer of care	8.3 (7.4)	8.3
Time between psychiatric diagnosis and transfer of care	4.0 (3.2)	4.0

ADSUME, Adolescent Substance Use Measurement.

Table 3 and figure 1 summarise the associations between disease control, psychiatric diagnoses, substance use and family composition. Adolescents with psychiatric diagnoses more often had poor disease control and/or adherence or more severe disease/condition

Table 3 Family composition, disease control and substance use, by psychiatric diagnosis (n=253). P < 0.05 is considered as significant.

	No psychiatric diagnosis (n=194) N (%)	Psychiatric diagnosis (n=59) N (%)	P value
Living with two parents	128 (66)	30 (51)	0.05
Disease control*			
Good control and/or adherence	59 (30)	11 (19)	0.13
Some evidence of concern	84 (43)	21 (37)	0.45
Poor control and/or adherence or more severe condition	51 (26)	25 (44)	0.01
ADSUME score, median (IQR)†	3 (0–6)	0 (0–6)	0.56

Other factors compared using Fisher's exact test; ADSUME median scores compared using the Mann-Whitney U test. Bold values signifies P < 0.05.
*Transfer of two adolescents has been delayed.
†One response missing.
ADSUME, Adolescent Substance Use Measurement.




Figure 1 Associations between disease control, psychiatric diagnoses, substance use (ADSUME) and family composition. $P < 0.05$ is considered significant. *Good control and/or adherence, some evidence of concern or poor control and/or adherence or more severe condition. **Abstinence or experimental use, repetitive use, risky use or hazardous use. ***Living with mother and father or with someone else (for example, with only one parent, mixed family or any other third party). ADSUME, Adolescent Substance Use Measurement.

than adolescents without psychiatric diagnoses ($p=0.01$). Hazardous substance use was more frequent among adolescents with psychiatric diagnoses (8% vs 19%), but total ADSUME points showed no direct association with psychiatric diagnoses nor disease control. Adolescents who lived with two parents ($n=158$, 62.5%) had fewer psychiatric diagnoses than adolescents living in other kinds of families ($p=0.05$). In ordinal regression, only psychiatric diagnoses were significant for disease control (table 4).

DISCUSSION

In this cross-sectional study of adolescents with chronic medical conditions, nearly one-third were assessed to have poor disease control in the year preceding transfer

to adult health services and almost a quarter had received a psychiatric diagnosis during adolescence. Adolescents with a comorbid psychiatric diagnosis more frequently had poor control of their medical condition than those without a psychiatric diagnosis.

In our study, 23% of the adolescents received a mental health diagnosis between the ages of 13 and 18 years. Adolescents living with two parents had fewer psychiatric diagnoses than their peers living in other kinds of families. The protective effect of living in a nuclear family when considering psychiatric morbidity has also been reported among the general population of children and adolescents.^{24 25} Overall, the prevalence of psychiatric diagnoses was higher in our study than in other diagnostic studies.^{8 10} Regular assessment by medical professionals might have contributed to unnecessary referrals aided by the relative ease of referral to public mental health professionals. However, as the portion of adolescents with three or more outpatient visits in the department of adolescent psychiatry is comparable with the proportion of adolescents with a psychiatric diagnosis, overdiagnosis seems unlikely.

In this study, 22.5% of adolescents reported using substances regularly, while a further 21.0% were categorised as risky or hazardous users. This is consistent with national Finnish data that found 15%–24% of students aged 17–18 years old reported monthly binge drinking.²⁶ Conversely, only 1% of adolescents in our cohort had used cannabis during the past year, in comparison with 8%–12% of students aged 17–18 years old in Finland.²⁶ Although alcohol consumption has decreased among adolescents in many developed countries, adolescent binge drinking and the accompanying risky behaviour remain significant public health problems.²⁷

The association between poor disease control and psychiatric comorbidity found in our study may be bidirectional: mental health problems may affect self-management resources and thus worsen disease control, while living with a severe disease may increase the emotional burden experienced. The burden of a

Table 4 Results of ordinal regression with disease control (good, moderate or poor) as the dependent variable. $P < 0.05$ is considered as significant.

	Estimate	SE	Wald	df	P value	95% CI	
						Lower	Upper
Threshold: poor to moderate dc	-2.963	1.718	2.975	1	0.085	-6.330	0.404
Threshold: moderate to good dc	-1.111	1.708	0.423	1	0.516	-4.459	2.237
Male vs female	-0.147	0.240	0.377	1	0.539	-0.617	0.323
Two parents vs other family	0.359	0.246	2.116	1	0.146	-0.125	0.842
Psychiatric dg vs no dg	0.636	0.291	4.794	1	0.029	0.067	1.206
Age (continuous)	-0.158	0.096	2.675	1	0.102	-0.347	0.031

Pseudo- r^2 : Cox and Snell 0.046, Nagelkerke 0.052.
 Bold values signifies $P < 0.05$.
 dc, disease control; dg, diagnosis.

chronic condition also includes a sense of interrupted social maturation for which substance use can sometimes present as a means of catching up although adolescents are aware of the harmful effects of substance use.²⁸

Considering the modest proportion of adolescents with good disease control before transfer of care, additional strategies may be required to optimise health outcomes in adolescents whose disease is challenging to treat, who lack motivation or skills for self-management, or have mental health or substance use problems.^{29 30} The treatment of anxiety and depression in adolescents is expected to improve symptoms and positively influence disease control and healthcare utilisation,^{31–33} while some adolescents will need concurrent transition of psychiatric and medical care. A child's illness is a significant stressor for parents,³⁴ and family psychosocial support and interventions to minimise financial hardship also warrant consideration. Our data affirm the importance of regularly discussing substance use and its effects with adolescents with chronic medical conditions especially given its potential effect on specific medical conditions, such as a reduced seizure threshold in adolescents with epilepsy or hypoglycaemia in adolescents with diabetes.^{35 36}

A strength of this study is the high response rate of adolescents recruited from six different hospital departments to intentionally span a range of conditions, increasing the generalisability of our results. However, this also provided some challenges, especially regarding how disease control was defined as by definition, this differs by medical condition. Although the same researcher collected data on disease control and psychiatric diagnoses, we attempted to reduce bias by collecting these data on separate occasions, using separate files, and in different patient order. While we needed to categorise disease control subjectively in adolescents with rare diseases, the likelihood of misclassification was reduced by having two independent reviewers. As we had access to all electronic medical records in the study catchment area, including psychiatry, we were able to use specific psychiatric diagnoses and estimate disease control of medical conditions based on clinical and laboratory findings. While psychiatric diagnoses were made by the psychiatry department, standardised assessments were not routinely used. Further, some psychiatric diagnoses may have been missed if the adolescent had moved to or from another hospital district. The public health system in Finland results in adolescents receiving the same treatment regardless of socioeconomic status, with provision of psychiatric care in the private sector uncommon. In countries without public healthcare, associations between mental health problems and poor disease control may be even more significant than in our study.

CONCLUSIONS

In conclusion, psychiatric morbidity and poor disease control of chronic medical conditions are prevalent and closely intertwined among adolescents, reinforcing the

importance of collaboration between medical and psychiatric care.³⁷ Maximising mental health before adolescents transfer to adult health services appears especially important given the associations of both poor mental health and the transfer of care itself with unsatisfactory adherence to treatment and increased rates of emergency visits and hospitalisation.^{12 38 39}

Author affiliations

¹Pediatric Research Center, New Children's Hospital, Helsinki, Finland

²Department of Pediatrics, Helsinki University Hospital and University of Helsinki, Helsinki, Finland

³Nursing Science, University of Turku, Turku, Finland

⁴Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

⁵Transition Support Service, Royal Children's Hospital, Parkville, Victoria, Australia

⁶Department of Adolescent Medicine, Royal Children's Hospital, Parkville, Victoria, Australia

⁷Department of Paediatrics, The University of Melbourne, Parkville, Victoria, Australia

⁸Centre for Adolescent Health, Murdoch Children's Research Institute and the Royal Children's Hospital, Parkville, Victoria, Australia

⁹Health Services, Murdoch Children's Research Institute, Melbourne, Victoria, Australia

Acknowledgements We wish to thank all the participants of the study as well as research nurse Satu Lindström. We also acknowledge Kristiina Aalto, Eero Jokinen, Eeva-Liisa Metsähonkala, Tiina Remes and Elisa Ylinen for clinical advice regarding disease control.

Contributors MK collected data, carried out initial analyses, drafted the manuscript and reviewed and revised the manuscript. AT collected data and reviewed and revised the manuscript. PM and K-LK participated in designing the study and reviewed and revised the manuscript. EC and SS participated in study design and reviewed and revised the manuscript. SK designed the study, coordinated and supervised data collection and critically reviewed the manuscript for important intellectual content. SK is responsible for the overall content as the guarantor. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Funding This study is supported by grants from the Foundation for Medical Research, the Helsinki University Hospital (TYH2019312), the Foundation for Paediatric Research and the Paulo Foundation.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not required.

Ethics approval This study involves human participants and was approved by the Ethics Committee for Women's and Children's Health and Psychiatry at the Helsinki University Hospital (HUS/1547/2017). Participants gave informed consent to link their routine clinical data with individual questionnaire responses.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is

properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Mira Kallio <http://orcid.org/0000-0003-0188-5341>

REFERENCES

- 1 Raissadati A, Nieminen H, Jokinen E, *et al*. Progress in late results among pediatric cardiac surgery patients: a population-based 6-decade study with 98 % follow-up. *Circulation* 2015;131:347–53;
- 2 Benchimol EI, Fortinsky KJ, Gozdyra P, *et al*. Epidemiology of pediatric inflammatory bowel disease: a systematic review of international trends. *Inflamm Bowel Dis* 2011;17:423–39.
- 3 Patterson CC, Karuranga S, Salpea P, *et al*. Worldwide estimates of incidence, prevalence and mortality of type 1 diabetes in children and adolescents: results from the international diabetes federation diabetes atlas, 9th edition. *Diabetes Res Clin Pract* 2019;157:107842.
- 4 Nap-van der Vlist MM, Dalmeijer GW, Grootenhuys MA, *et al*. Fatigue in childhood chronic disease. *Arch Dis Child* 2019;104:1090–5.
- 5 Maes M, Van den Noortgate W, Fustolo-Gunnink SF, *et al*. Loneliness in children and adolescents with chronic physical conditions: a meta-analysis. *J Pediatr Psychol* 2017;42:622–35.
- 6 Brady AM, Deighton J, Stansfeld S. Psychiatric outcomes associated with chronic illness in adolescence: a systematic review. *J Adolesc* 2017;59:112–23.
- 7 Robinson ME, Simard M, Larocque I, *et al*. Risk of psychiatric disorders and suicide attempts in emerging adults with diabetes. *Diabetes Care* 2020;43:484–6.
- 8 LaGrant B, Marquis BO, Berg AT, *et al*. Depression and anxiety in children with epilepsy and other chronic health conditions: national estimates of prevalence and risk factors. *Epilepsy Behav* 2020;103(Pt A):106828.
- 9 Ghandour RM, Sherman LJ, Vladutiu CJ, *et al*. Prevalence and treatment of depression, anxiety, and conduct problems in US children. *J Pediatr* 2019;206:256–67.
- 10 Butwicka A, Olén O, Larsson H, *et al*. Association of childhood-onset inflammatory bowel disease with risk of psychiatric disorders and suicide attempt. *JAMA Pediatr* 2019;173:969–78.
- 11 Finnish Institute for Health and Welfare. Well-being of children and young people – school health promotion study 2021. 2021. Available: <https://thl.fi/en/web/thlfi-en/statistics-and-data/statistics-by-topic/social-services-children-adolescents-and-families/well-being-of-children-and-young-people-school-health-promotion-study>
- 12 Bernstein CM, Stockwell MS, Gallagher MP, *et al*. Mental health issues in adolescents and young adults with type 1 diabetes: prevalence and impact on glycemic control. *Clin Pediatr (Phila)* 2013;52:10–5.
- 13 Reed-Knight B, Lee JL, Greenley RN, *et al*. Disease activity does not explain it all: how internalizing symptoms and caregiver depressive symptoms relate to health-related quality of life among youth with inflammatory bowel disease. *Inflamm Bowel Dis* 2016;22:963–7.
- 14 Wong JJ, Sceats L, Dehghan M, *et al*. Depression and health care use in patients with inflammatory bowel disease. *J Crohns Colitis* 2019;13:19–26.
- 15 Sawyer SM, Drew S, Yeo MS, *et al*. Adolescents with a chronic condition: challenges living, challenges treating. *Lancet* 2007;369:1481–9.
- 16 Shrier LA, Harris SK, Kurland M, *et al*. Substance use problems and associated psychiatric symptoms among adolescents in primary care. *Pediatrics* 2003;111(6 Pt 1):e699–705.
- 17 Deas D, Brown ES. Adolescent substance abuse and psychiatric comorbidities. *J Clin Psychiatry* 2006;67:e02.
- 18 Kessler RC, Berglund P, Demler O, *et al*. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National comorbidity survey replication. *Arch Gen Psychiatry* 2005;62:593–602.
- 19 Kosola S, Culfane E, Loftus H, *et al*. Bridge study protocol: an international, observational cohort study on the transition of healthcare for adolescents with chronic conditions. *BMJ Open* 2021;11:e048340.
- 20 Backström M, Tynjälä P, Ylijoki H, *et al*. Finding specific 10-joint juvenile arthritis disease activity score (JADAS10) and clinical JADAS10 cut-off values for disease activity levels in non-systemic juvenile idiopathic arthritis: a Finnish multicentre study. *Rheumatology (Oxford)* 2016;55:615–23.
- 21 DiMeglio LA, Acerini CL, Codner E, *et al*. ISPAD clinical practice consensus guidelines 2018: glycemic control targets and glucose monitoring for children, adolescents, and young adults with diabetes. *Pediatr Diabetes* 2018;19 Suppl 27:105–14.
- 22 Piskanen M. Promoting adolescents' abstinence from substance use. an early intervention model for school and student health care. 2007. Available: <http://urn.fi/URN:ISBN:978-951-27-0510-8>
- 23 Pirskanen M, Laukkanen E, Pietilä AM. A formative evaluation to develop a school health nursing early intervention model for adolescent substance use. *Public Health Nurs* 2007;24:256–64.
- 24 Angarne-Lindberg T, Wadsby M. Psychiatric and somatic health in relation to experience of parental divorce in childhood. *Int J Soc Psychiatry* 2012;58:16–25.
- 25 Perales F, Johnson SE, Baxter J, *et al*. Family structure and childhood mental disorders: new findings from Australia. *Soc Psychiatry Psychiatr Epidemiol* 2017;52:423–33.
- 26 Helakorpi S, Kivimäki H. Well-being of children and young people – school health promotion study 2021: large share of children and young people satisfied with their lives – a sense of loneliness has become more common. Statistical report 42/2021. Finland National Institute for Health and Welfare; 2022. Available: <https://www.julkari.fi/handle/10024/143418>
- 27 Siqueira L, Smith VC, ABUSE COS. Binge drinking. *Pediatrics* 2015;136:e718–26.
- 28 Weitzman ER, Salimian PK, Rabinow L, *et al*. Perspectives on substance use among youth with chronic medical conditions and implications for clinical guidance and prevention: a qualitative study. *PLoS One* 2019;14.
- 29 Andrade DM, Bassett AS, Bercovici E, *et al*. Epilepsy: transition from pediatric to adult care. recommendations of the Ontario epilepsy implementation Task force. *Epilepsia* 2017;58:1502–17.
- 30 Healy SA, Fantaneanu TA, Whiting S. The importance of mental health in improving quality of life in transition-aged patients with epilepsy. *Epilepsy Behav* 2020;112:107324.
- 31 Bennett S, Shafran R, Coughtrey A, *et al*. Psychological interventions for mental health disorders in children with chronic physical illness: a systematic review. *Arch Dis Child* 2015;100:308–16.
- 32 Thabrew H, Stasiak K, Hetrick SE, *et al*. E-health interventions for anxiety and depression in children and adolescents with long-term physical conditions. *Cochrane Database Syst Rev* 2018;8.
- 33 Lehtimäki S, Martic J, Wahl B, *et al*. Evidence on digital mental health interventions for adolescents and young people: systematic overview. *JMIR Ment Health* 2021;8.
- 34 Mutanen A, Kosola S, Merras-Salmio L, *et al*. Long-term health-related quality of life of patients with pediatric onset intestinal failure. *J Pediatr Surg* 2015;50:1854–8.
- 35 Samokhvalov AV, Irving H, Mohapatra S, *et al*. Alcohol consumption, unprovoked seizures, and epilepsy: a systematic review and meta-analysis. *Epilepsia* 2010;51:1177–84.
- 36 Tetzschner R, Nørsgaard K, Ranjan A. Effects of alcohol on plasma glucose and prevention of alcohol-induced hypoglycemia in type 1 diabetes-A systematic review with GRADE. *Diabetes Metab Res Rev* 2018;34.
- 37 Fazel M, Townsend A, Stewart H, *et al*. Integrated care to address child and adolescent health in the 21st century: a clinical review. *JCPP Advances* 2021;1. 10.1002/jcv2.12045 Available: <https://onlinelibrary.wiley.com/doi/10.1002/jcv2.12045>
- 38 Kosola S, Yliinen E, Finne P, *et al*. Implementation of a transition model to adult care may not be enough to improve results: national study of kidney transplant recipients. *Clin Transplant* 2019;33.
- 39 Rollo A, Salardi S, Ciavarella A, *et al*. Transition from pediatric to adult care. eight years after the transition from pediatric to adult diabetes care: metabolic control, complications and associated diseases. *J Endocrinol Invest* 2014;37:653–9.