

Risk perception and risk attitudes in Tokyo: A report of the first administration of DOSPERT+M in Japan

Alan Schwartz* Kimihiko Yamagishi[†] Norimichi Hirahara[‡] Hirotaka Onishi[§]
James Barnes IV[¶] Adam Rosman* Maggie Garcia* Sam Lee* Shoshana Butler*

Abstract

Background: The Domain-Specific Risk Taking scale (DOSPERT) has been used to measure risk perceptions and attitudes in several nations and cultures. Takahashi translated DOSPERT to Japanese but DOSPERT responses from Japan have never been reported. Butler et al. (2012) developed an additional medical risk domain subscale to be added to DOSPERT to form DOSPERT+M.

Objective: To describe the translation of the medical risk domain subscale to Japanese and to characterize domain-specific risk attitudes in Tokyo.

Methods: Members of a probability-weighted online panel representative of the Tokyo metro area were randomized to complete pairs of DOSPERT+M tasks (risk attitude, risk perception, benefit perception). We explored relationships among domains through correlational and factor analysis; we tested the hypothesis that the medical risk domain and DOSPERT's health/safety domains were uncorrelated.

Participants: One hundred eighty panelists.

Results: Six of the original DOSPERT items (two each in the ethics, health/safety, and financial domains) are not useable in Japan according to the Japanese Marketing Research Association code because they ask about participation in illegal activities; we thus used abbreviated versions of those domains leaving out these items. The DOSPERT+M items generally did not cluster cleanly into the expected domains, although items within the same domain usually were intercorrelated. Participants demonstrated domain-specific conventional risk attitudes, although nearly half of those assessed were perceived-risk neutral in all domains. Unlike our recently reported findings in the U.S. population, DOSPERT+M medical domain scores were associated with health/safety domain scores, although they were often more strongly associated with scores in other domains, such as recreational activities.

Conclusion: The DOSPERT (and DOSPERT+M) instruments are problematic in Japan but Japanese citizens may also differ from those of other nations in their risk attitudes and perceptions.

Keywords: DOSPERT, DOSPERT+M, risk attitude, risk perception, medical decision making, Japan

1 Introduction

The Domain-Specific Risk Taking scale (DOSPERT) (Blais & Weber, 2006; Blais & Weber, 2009; Weber, Blais, & Betz, 2002) measures risk attitudes (willingness

to engage), risk perceptions, and expected benefits for a set of activities organized into five domains: ethical, financial, health/safety, recreational, and social risk taking. Conceptually, DOSPERT assumes that risk perceptions and attitudes may differ among domains, and has often been used to examine whether willingness to engage in an activity is negatively associated with perceptions of the riskiness of the activities and positively associated with expected benefits (the so-called risk-return framework) (Sarin & Weber, 1993).

Studies that have compared conventional risk attitudes (willingness to engage in activities defined objectively as risky) with perceived risk attitudes (willingness to engage in activities that the respondent themselves perceives as risky) across domains suggest that people may be more consistent in their perceived risk attitudes than in their conventional risk attitudes (Weber et al., 2002; Weber & Milliman, 1997).

The work reported here was funded in part by U.S. National Science Foundation grant SES-1023831, a Fulbright Foundation Senior Specialist grant, and a Tokyo Institute of Technology Visiting Scholar appointment to Alan Schwartz.

Copyright: © 2013. The authors license this article under the terms of the Creative Commons Attribution 3.0 License.

*Department of Medical Education (mc 591), University of Illinois at Chicago, 808 S. Wood St., 986 CME, Chicago, IL 60612. E-mail: alansz@uic.edu.

[†]Graduate School of Decision Science & Technology, Tokyo Institute of Technology

[‡]Department of Healthcare Quality Assessment, Graduate School of Medicine, University of Tokyo

[§]International Research Center for Medical Education, Graduate School of Medicine, University of Tokyo

[¶]Department of Internal Medicine, Stanford University

Table 1: DOSPERT+M medical risk items in English and Japanese

English	Japanese
Donating one kidney to a patient you do not know	移植のため、知らない患者に腎臓を片方提供する
Giving blood	献血をする
Participating in a clinical trial to determine whether a new drug is effective	新開発の薬の効果を確かめる治験（医療実験）に参加する
Taking daily medication to relieve allergy symptoms	アレルギー症状を抑えるため毎日薬を飲む
Knee replacement surgery to treat arthritis	膝の関節炎を治すために人工関節の手術をする
Receiving general rather than local anesthesia when having a wisdom tooth removed	親知らずを抜くときに局所麻酔ではなく全身麻酔をする

A particular strength of DOSPERT is that it is available in several translations and risk attitudes and perceptions have been studied in participants in Quebec (Blais & Weber, 2006), Germany (Johnson, Wilke, & Weber, 2004), China (Cheung, Wu, & Tao, 2012; Hu & Xie, 2012), and South Africa (Szrek, Chao, Ramlagan, & Peltzer, 2012). Most of these studies have found within-nation domain-specificity in risk attitudes and perceptions (often with less interdomain variation in perceived-risk attitudes) and differences in levels of risk attitude and risk perception across nations or cultures (although not all made direct comparisons among cultures within the same study). A Japanese translation of DOSPERT was performed between 2006–2009 by Yusuke Takahashi at the University of Illinois (Takahashi, n.d.) but use of a portion of this translation only been reported in two Japanese publications (Hirahara & Yamagishi, 2009, 2011).

DOSPERT is also one of the few instruments Harrison et al. (2005) recommends for assessing health-related or clinical risks. However, DOSPERT does not include a domain that focuses on health care activities; the health/safety scale is limited to preventive behaviors, and two Japanese studies suggest that health/safety risk perceptions are not related to perceptions of treatment risks by breast or blood cancer patients (Hirahara & Yamagishi, 2009, 2011). Butler et al. (2012) developed and provided initial psychometric evidence for an add-on medical risk domain consisting of six activities, modeled after the other DOSPERT domains. Adding the new domain's items to the standard DOSPERT items produces a 36-item instrument that we refer to as DOSPERT+M. Rosman et al. (2013) administered DOSPERT+M to a nationally-representative U.S. sample and found that the medical risk domain appeared to measure a different construct than the health/safety domain. We sought to determine whether the same distinction would be observed in Japan, as a question of generalizability (rather than a

cross-cultural comparison).

The goals of this study were to (1) translate the medical risk domain items to Japanese to create a Japanese DOSPERT+M, (2) administer the Japanese DOSPERT+M instrument to a representative sample of the Tokyo metro population, and (3) characterize risk attitudes, including medical risk attitudes, among the sample and examine the associations among domains. We take the DOSPERT+M instrument as pre-existing; our focus is application rather than item selection or the development of validity evidence per se. We specifically sought to examine whether the DOSPERT factor structure could be recovered in the data, whether perceived-risk attitudes were more stable across domains than willingness to take risks, and whether the DOSPERT health/safety scale was or was not associated with the DOSPERT medical scale.

2 Translation

2.1 Initial Translation

Takahashi's Japanese DOSPERT, including the 30 DOSPERT items (based on the Blais and Weber, 2006 "revised" DOSPERT) and the 3 DOSPERT response tasks/scales was used as the basis for the Japanese DOSPERT+M. The DOSPERT+M uses 7 anchors in the expected benefits task, but the original DOSPERT uses only the two extremes and midpoint for anchors in this task (and thus only those three anchors appear in Takahashi's translation). The 6 medical risk items and the 4 missing expected benefits anchors were translated to Japanese using the approach advocated by Weidmer et al. (Weidmer, Hurtado, Weech-Maldonado, Ngo-Metzger, & Bogen, 2006) for the translation of the Consumer Assessment of Healthcare Providers and Systems (CAHPS) surveys administered by the U.S. Agency for Healthcare Research and Quality. Specifically, two independent for-

Table 2: DOSPERT activities illegal in Japan (Research and Development, personal communication, 2012)

Activity	DOSPERT domain
6. Taking some questionable deductions on your tax return	Ethics
8. Betting a day's income at a high-stake poker game	Financial
14. Betting a day's income on the outcome of a sporting event	Financial
17. Driving a car without wearing a seat belt	Health/safety
20. Riding a motorcycle without a helmet	Health/safety
30. Not returning a wallet you found that contains \$200	Ethics

ward translations were produced by co-authors who are native Japanese speakers (NH and HO). A third co-author (KY) served as the translation reviewer and compared the original English item with each of the two Japanese translations. The translation reviewer selected which translation was preferable or reconciled the two translations into a third translation. Finally, all three translators reviewed the final versions of items and anchors and approved their use. Table 1 presents the medical risk domain items in English and Japanese translation.

2.2 Application Issues

Following the DOSPERT+M translation, during survey preparation, Research and Development, the Japanese survey company managing the online panel indicated that six of the original DOSPERT items were not useable in Japan according to the Japanese Marketing Research Association code because survey researchers are not permitted to ask about participation in illegal activities. These items, and their domains, are listed in Table 2.

For the ethics and health/safety items, we presented reversed items (e.g. "Not taking questionable deductions on your tax return") to respondents but omitted them from the data analysis as they clearly were not going to measure the same construct. Instead, we constructed abbreviated subscales for these domains without the illegal activity items. Gambling privately is illegal in Japan, but regulated gambling is allowed. Accordingly, for the financial items, we replaced these items with acceptable forms of gambling: "Betting a day's income at a Pachinko game" (一日分の収入を、パチンコに費やす) and "Spending a day's income for gambling such as a horse race, bicycle

race, or boat race" (一日分の収入を、競馬、競輪、競艇といった賭け事に使う).

The two earlier Japanese studies employed only the health/safety items of the original Japanese DOSPERT translation (Hirahara & Yamagishi, 2009, 2011) and only the risk perception task. These studies were conducted as part of interviews with patient support groups and not as large-scale survey research, did not ask about willingness to engage in activities, and were not examined in the context of the Japanese Marketing Research Association code. We thus faced this problem for the first time.

3 Study Methods

3.1 Instrument

The 36 DOSPERT+M items, each describing a risky activity, can be presented in any of the three standard DOSPERT tasks (risk-taking, risk perception, or expected benefit). In the risk-taking task, participants indicate their likelihood of participating in the activity on a 1–7 scale from extremely unlikely to extremely likely. In the risk perception task, participants indicate how risky they believe the activity is on a 1–7 scale from not at all risky to extremely risky. In the expected benefit task, participants indicate the benefits they believe they would receive from each activity on a 1–7 scale from no benefits at all to great benefits. To reduce respondent fatigue, each participant was block-randomized to receive two of the three tasks presented in either order (30 participants in each of the 6 task-pair/ordering combinations).

3.2 Participants

DOSPERT+M was administered online by Research and Development, a Japanese survey research company. The sample was probability-weighted to be representative of the greater Tokyo metropolitan area and stratified by gender. Each participant was also asked a series of demographic questions related to age, gender, location (Tokyo, Saitama, Chiba, or Kanagawa), marital status, household income, whether participant had children, and occupation. On Japan's prefecture list by population rank, these places are 1st, 2nd, 5th, and 6th, respectively, and the stratified sample from these areas is reasonably representative of Japan's manufacturing and service industry region (Japan Ministry of Land, Infrastructure, Transport and Tourism, 2011). Data collection took place between August 23, 2012 and August 29, 2012.

3.3 Data Analysis

We examined the relationship among risk-taking, risk perception, and expected benefits in each domain by

computing correlations among domain subscale scores in each task. Because participants were randomized to complete two of three tasks, each intra-domain (inter-task) correlation is based on a different group of participants.

Following Weber et al. (2002), we computed conventional risk attitudes for each domain on the 120 subjects who completed the risk-taking task based on whether the domain scale score was less than 1 SD below the mean (risk averse), from 1 SD below to 1 SD above the mean (risk neutral) or more than 1 SD above the mean (risk seeking). These attitudes are “conventional” in that they assume that every DOSPERT item is a risky activity and base risk attitude on stated willingness to participate in the activity. We compute perceived risk attitudes by considering willingness to engage in activities that respondents themselves perceive to be risky. Because of our design, we were not able to compute perceived risk attitudes controlling for expected benefit by regressing risk taking on both risk perception and expected benefit as in Weber et al. 2002), so we computed perceived risk attitudes for each domain on the 60 subjects who completed both risk-taking and risk perception tasks based on whether the coefficient for the risk perception scale score was significantly positive (risk-seeking), significantly negative (risk averse), or nonsignificant (risk neutral) when risk taking was regressed on risk perception (only) in each domain.

We conducted confirmatory factor analyses for each task excluding the medical domain items to determine whether we could recover the 6-factor solution found in Weber et al. (2002), in which each domain except the financial is a factor constituted of that domain’s items, and the financial domain items are split between two factors, one composed of the three gambling items, and the other composed of the three investment items. We then conducted a similar analysis including the medical domain items and hypothesizing a 7th factor composed of those items. We considered the fit acceptable if RMSEA was less than 0.06. If the fit was not acceptable, we investigated the factor structure through exploratory factor analyses with and without medical items, extracting six factors by maximum likelihood methods and using an oblimin rotation, and examined the factor pattern matrix and inter- factor correlations. (Fabrigar, Wegener, MacCallum, & Strahan, 1999). Our planned sample size of 120 per task was deemed to be sufficient for factor analysis based on the rule of thumb of 3–5 participants per item.

We also conducted multiple linear regressions to determine whether variance in domain task scores (risk-taking, risk perception, and expected benefits) was associated with demographic factors. Age, gender, cohabitation (married/living with partner or not), and occupational status (employed full or part time outside the home vs. not) were included as predictors.

Table 3: Demographic characteristics.

Characteristic	N	%
Gender		
Male	90	50
Female	90	50
Age (years)		
18-29	28	16
30-39	34	19
40-49	34	19
50-59	28	16
60+	56	31
Marital Status		
Single	64	36
Married	106	59
Widowed	5	3
Divorced	5	3
Has children		
No	95	53
Yes	85	47
Occupational Status		
Employee	58	32
Self-employed	24	13
Part-time job	21	12
Student	15	8
Homemaker	40	22
Retired	7	4
Unemployed	15	8
Household Income (in JPY)		
< 2M	13	8
2–3.999M	29	19
4–5.999M	27	18
6–7.999M	35	23
8–9.999M	17	11
10–11.999M	13	8
12–14.999M	12	8
15–19.999M	5	3
>= 20M	2	1

Analyses were conducted using R 3.0 and the packages reshape, GPArotation, psych, nFactors, Hmisc, gmodels, sem, and semPlot (Bernaards & IJennrich, 2005; Epskemp, 2013; Fox, Nie, & Byrnes, 2013; Harrell, 2013; R Core Team, 2013; Raiche, 2010; Revelle, 2013; Warnes, 2013; Wickham & Hadley, 2007).

4 Results

4.1 Participants

Table 3 presents the demographic characteristics of the 180 study participants.

4.2 Relationships among Tasks

Within the medical subscale, risk-taking scores were not significantly correlated with risk perception scores ($r=.08$) in the subsample that completed those two tasks. Risk taking was significantly positively correlated with benefit perception scores ($r=.42$, $p=.0008$) in the subsample that completed those two tasks. Benefit and risk perception scores were negatively correlated ($r=-.44$, $p=.0005$) in the subsample that completed those two tasks, a pattern labeled the “halo effect” by other researchers (Finucan, Alhakami, Slovic, & Johnson, 2000). A similar pattern of associations was found for the social subscale; for recreation, financial, abbreviated ethical, and abbreviated health/safety subscales, risk-taking and risk perception were also significantly negatively correlated ($r=-.38$, $r=-.38$, $r=-.32$ and $r=-.46$, respectively).

4.3 Relationships among Domains

Table 4 presents the correlations among domain subscale scores in each task, which exhibit a positive manifold. Medical domain scores were least strongly correlated with the financial domain scores, but were correlated with health/safety domain scores.

4.4 Risk attitudes

Table 5 presents conventional and perceived risk attitudes by domain. For perceived risk attitudes, missing subjects are those where regression failed due to lack of variance in predictor and outcome variable. Similar to the findings of Weber et al. (2002), conventional risk attitudes across participants reflected primarily risk neutrality in all domains (although with a dearth of risk averse individuals in the financial and recreational domain). Perceived-risk attitudes were largely neutral or perceived-risk averse.

Among the 52 individuals for whom both conventional and perceived risk attitudes could be obtained for every domain, 23 (44%) had the same perceived risk attitude (perceived risk neutrality) in all domains; only 7 (14%) had the same conventional risk attitude (risk neutrality) in all domains (McNemar test chi-squared(1 df) = 10.2, $p=0.001$).

4.5 Confirmatory Factor Analysis

Confirmatory factor analyses of the risk-taking and expected benefits tasks with (7 factor) and without (6 factor) the medical domain items provided acceptable fit (RMSEA = .055 for risk-taking, RMSEA = .035 for expected benefits). In each case path coefficients between items and the expected factors were significantly positive, except that the path coefficients for items R2 (“Going down a ski run that is beyond your ability”) and H2 (“Engaging

in unprotected sex”) were not significant in the expected benefits model.

In the risk perception task, the confirmatory analysis did not fit as well (RMSEA = .072 for 6 factors without medical items, RMSEA = .074 for 7 factors with medical items), although all of the path coefficients were significant and positive except for item M2 (“Giving blood”). Accordingly, we proceeded to examine responses in this task with exploratory factor analysis.

4.6 Exploratory Factor Analysis

Table 7 presents the pattern matrix from exploratory factor analysis of the responses in the risk perception task including the medical domain items with loadings with absolute value smaller than 0.3 suppressed. Parallel analysis suggested a 5-factor solution was most appropriate for the data, but the fit was more interpretable with a 6-factor solution, which we report here.

The social items, financial (gambling) items, and financial (investment) items loaded onto separate factors. However, the (abbreviated) ethics subscale shared common variance with the health/safety scale in this task that was unexpected (this was also observed in exploratory analyses omitting the medical items). Several recreational domain items loaded together with the medical domain items. Item M6 (general anesthesia) did not load strongly on any factor.

4.7 Demographic Predictors of Medical Domain Scores

The regression model for medical risk-taking scores did not account for significant variance in the scores ($F(4,115)=1.22$, $p=0.31$). For risk perception of medical activities, the model reached significance ($F(4,115)=3.34$, $p=0.01$) and two predictors were associated with risk perceptions. Married respondents had lower scores for risk perception ($B=-0.34$, $SE=.15$, $p=.022$) and older respondents had higher scores ($B=0.014$ per year of age, $SE=0.005$, $p=.005$). For expected benefit of medical activities, the regression model reached significance ($F(4,115)=3.7$, $p=0.007$). The only significant demographic predictor was age: older respondents had lower scores for expected benefit ($B=-0.028$ per year of age, $SE=0.008$, $p=.0004$).

5 Discussion

5.1 Summary of results

In this study, we report the first attempted application of DOSPERT (and DOSPERT+M) in Japan. In the process,

Table 4: Correlations among domains and Cronbach's α for each domain.

	Risk-taking task					
	Medical	Social	Recreation	Ethics (abbr.)	Health/safety (abbr.)	Financial
Medical	.62	.42**	.47**	.40**	.27**	.15
Social		.71	.30**	.16	.32**	.11
Recreation			.79	.45**	.15	.41**
Ethics (abbreviated)				.51	.40**	.13
Health/safety (abbreviated)					.65	.03
Financial						.77

	Risk perception task					
	Medical	Social	Recreation	Ethics (abbr.)	Health/safety (abbr.)	Financial
Medical	.59	.11	.47**	.32**	.41**	.18
Social		.69	.14	.00	.09	.35**
Recreation			.71	.45**	.52**	.19*
Ethics (abbreviated)				.75	.65**	.06
Health/safety (abbreviated)					.68	.17
Financial						.83

	Expected benefit task					
	Medical	Social	Recreation	Ethics (abbr.)	Health/safety (abbr.)	Financial
Medical	.80	.62**	.48**	.45**	.47**	.41**
Social		.81	.58**	.34**	.34**	.48**
Recreation			.81	.55**	.53**	.30**
Ethics (abbreviated)				.80	.76**	.28**
Health/safety (abbreviated)					.74	.25**
Financial						.84

* $p < .05$, ** $p < .01$

Note: Cronbach's α on main diagonals, correlations on off-diagonals.

Table 5: Conventional and perceived risk attitudes by domain.

Domain	Conventional risk attitudes (n=120)			Perceived risk attitudes (n=60)			Missing perceived risk attitude
	Risk averse	Risk neutral	Risk seeking	Perceived-risk averse	Perceived-risk neutral	Perceived-risk seeking	
Medical	23 (19%)	78 (65%)	19 (16%)	8 (13%)	49 (82%)	1 (2%)	2 (3%)
Ethical	28 (23%)	65 (54%)	27 (23%)	6 (10%)	45 (75%)	2 (3%)	7 (12%)
Financial	0 (0%)	96 (80%)	24 (20%)	7 (12%)	44 (73%)	0 (0%)	9 (15%)
Health/safety	25 (21%)	76 (63%)	19 (16%)	4 (7%)	52 (87%)	0 (0%)	4 (7%)
Recreational	0 (0%)	99 (83%)	21 (18%)	6 (10%)	52 (87%)	0 (0%)	2 (3%)
Social	23 (19%)	78 (65%)	19 (16%)	9 (15%)	46 (77%)	0 (0%)	5 (8%)

Table 6: Exploratory factor analysis of responses to the DOSPERT+M risk perception task (6-factor solution).

	Factor					
	1	2	3	4	5	6
E2	0.40					
E3	0.50					
E4	0.59					
E5	0.34					
F1				0.56		
F2		0.75				
F3				0.47		
F4		0.79				
F5				0.97		
F6		0.77				
H1	0.86					
H2	0.46					
H5	0.49					
H6					0.36	
M1			0.54			
M2			0.30		-0.32	
M3		0.37	0.42			
M4			0.53			
M5	0.39		0.31			
M6						
R1	0.31					
R2					0.45	
R3					0.71	
R4			0.66			
R5			0.56			
R6			0.38		0.40	
S1						0.73
S2						0.40
S3						0.52
S4						0.56
S5						0.55
S6					0.35	

Note: Loadings with absolute value ≥ 0.3 shown.

we experienced practical difficulty due to regulations related to acceptable survey practices. Using our modified instrument and a Tokyo metropolitan area-representative panel sample, the DOSPERT+M items generally did not cluster cleanly into the expected domains, although items within the same domain usually were intercorrelated.

Participants demonstrated domain-specific conventional risk attitudes, although nearly half of those assessed were perceived risk neutral in all domains. In part, this is likely a consequence of the restricted numbers of items in each domain; researchers interested in this particular question may be better served by using the original (8 items per domain) DOSPERT of Weber et al. (2002) rather than the revised (6 items per domain) DOSPERT of Blais and Weber (2006).

Unlike our recently reported findings in the U.S. population, DOSPERT+M medical domain scores were associated with (abbreviated) health/safety domain scores, although they were often more strongly associated with scores in other domains, such as recreational activities. DOSPERT+M risk-taking scores were not associated with demographic factors; DOSPERT+M risk perception scores were lower for married respondents and increased with age. Expected benefit scores decreased with age.

5.2 Limitations

A major limitation of the study is the finding that several of the original DOSPERT scales cannot be used as written in large-sample survey research in Japan. Accordingly, conclusions drawn about the ethics and health/safety domains are based on abbreviated subscales and should be considered tentative; conclusions drawn about the financial domain, where two items required modification to make them suitable, also require caution. We obtained the responses of breast and blood cancer patients to the risk perception task using the 8-item health/safety scale administered by Hirahara and Yamagishi (2009, 2011) and found that in each of those studies, the full 8-item scale was highly correlated with the abbreviated scale ($r=0.94$ and $r=0.93$, respectively); however, it should be noted that the patient sample is highly unrepresentative and this was reflected in lower perceptions of the riskiness of these everyday items than in our population sample.

We did not measure actual decision behavior, and do not know whether medical domain subscale scores might be associated with actual medical decisions. This is an area for future investigation.

Out of concern for survey fatigue, we present only two of the three DOSPERT tasks to each participant. Although this is an improvement on past studies using DOSPERT that have simply omitted the expected benefits scale, it remains a limitation on our ability to conduct within-subject comparisons of responses to each task, and to measure the risk return attitude in the manner of Weber et al. (2002).

5.3 Implications

Although there is good evidence for domain-specificity of risk perceptions and risk attitudes overall, patterns of responses to particular DOSPERT (and DOSPERT+M) items do not universally reproduce the expected domains, and the relationships among domains may differ for different populations. These valuable instruments for studying risk require pilot-testing when introduced to new populations and new languages.

6 References

- Bernaards, C. A. & Jennrich, R. I. (2005). Gradient projection algorithms and software for arbitrary rotation criteria in factor analysis. *Educational and Psychological Measurement, 65*, 676–696.
- Blais, A., & Weber, E. (2006). A Domain-Specific Risk-Taking (DOSPERT) scale for adult populations. *Judgment and Decision Making, 1*, 33–47.
- Blais, A., & Weber, E. U. (2009). *The Domain-Specific Risk Taking (DOSPERT) scale for adult populations: Item selection and preliminary psychometric properties*. Toronto, Ontario: Defence R & D.
- Butler, S., Rosman, A., Seleski, S., Garcia, M., Lee, S., Barnes, J., & Schwartz, A. (2012). A medical risk attitude subscale for DOSPERT. *Judgment and Decision Making, 7*, 189–195.
- Cheung, H. Y., Wu, J., & Tao, J. (2012). Risk Perception and Risk-Taking Attitude: A Comparison Between Hong Kong and Mainland Chinese Undergraduate Students. *The Asia-Pacific Education Researcher, 1*–10.
- Epskamp, S. (2013). semPlot: Path diagrams and visual analysis of various SEM packages' output. R package version 0.3.2. <http://CRAN.R-project.org/package=semPlot>.
- Fabrigar, L., Wegener, D., MacCallum, R., & Strahan, E. (1999). Evaluating the use of exploratory factor analysis in psychological research. *Psychological Methods, 4*, 272–299.
- Finucane, M. L., Alhakami, A., Slovic, P., & Johnson, S. M. (2000). The affect heuristic in judgments of risks and benefits. *Journal of Behavioral Decision Making, 13*(1), 1–17.
- Fox, J., Nie, Z., & Byrnes, J. (2013). sem: Structural equation models. R package version 3.1-3. <http://CRAN.R-project.org/package=sem>.
- Harrell, F. E., Jr. (2013). Hmisc: Harrell Miscellaneous. R package. <http://CRAN.R-project.org/package=Hmisc>.
- Harrison, J. D., Young, J. M., Butow, P., Salkeld, G., & Solomon, M. J. (2005). Is it worth the risk? A systematic review of instruments that measure risk propensity for use in the health setting. *Social science & medicine, 60*, 1385–1396.
- Hirahara, N., & Yamagishi, K. (2009). 血液がん患者の示す治療リスク認知の楽観性に関するステージ差. 医療の質・安全学会誌 [A change in medical risk perception over treatment stages of blood cancer patients]. *The Japanese Journal of Quality and Safety in Healthcare, 4*, 34–40.
- Hirahara, N., & Yamagishi, K. (2011). 乳がん患者の示す治療リスク認知の楽観性 闘病ステージによる変化. 認知科学 [A change in medical risk perception over treatment stages of breast cancer patients]. *Cognitive Studies, 18*, 534–545.
- Hu, X., & Xie, X. (2012). Validation of the Domain-Specific Risk-Taking Scale in Chinese college students. *Judgment and Decision Making, 7*(2), 181–188.
- Japan Ministry of Land, Infrastructure, Transport and Tourism (2011). White Paper on Land, Infrastructure, Transport and Tourism in Japan, 2011. <http://www.mlit.go.jp/english/white-paper/2011.pdf>.
- Johnson, J., Wilke, A., & Weber, E. (2004). Beyond a trait view of risk taking: A domain-specific scale measuring risk perceptions, expected benefits, and perceived-risk attitudes in German-speaking populations. *Polish Psychological Bulletin, 35*(3), 153–163.
- R Core Team (2013). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.
- Raiche, G. (2010). nFactors: an R package for parallel analysis and non graphical solutions to the Cattell scree test. <http://CRAN.R-project.org/package=nFactors>.
- Revelle, W. (2013) psych: Procedures for personality and psychological research, Northwestern University, Evanston, Illinois, USA, <http://CRAN.R-project.org/package=psych>, Version = 1.3.2.
- Rosman, A., Garcia, M., Lee, S., Butler, S., & Schwartz, A. (2013). DOSPERT+M: A survey of medical risk attitudes in the United States. *Judgment and Decision Making, 8*(4), 470–481.
- Sarin, R. K., & Weber, M. (1993). Risk-value models. *European Journal of Operational Research, 70*(2), 135–149.
- Szrek, H., Chao, L.-W., Ramlagan, S., & Peltzer, K. (2012). Predicting (un)healthy behavior: a comparison of risk-taking propensity measures. *Judgment and Decision Making, 7*(6), 716–727.
- Takahashi, Y. (n.d.). Domain-Specific Risk Taking (DOSPERT) Scale - Japanese translation. Columbia University. https://sites.google.com/a/decisionsciences.columbia.edu/cds-wiki/home/dospert/DOSPERT_30_Japanese.doc.
- Warnes, G. R. (2013). gmodels: Various R programming tools for model fitting. <http://CRAN.R-project.org/>.

- org/package=gmodels.
- Weber, E. U., Blais, A. R., & Betz, N. E. (2002). A Domain-specific Risk-attitude Scale: Measuring risk perceptions and risk behaviors. *Journal of Behavioral Decision Making*, *15*, 263–290.
- Weber, E. U., & Milliman, R. A. (1997). Perceived risk attitudes: Relating risk perception to risky choice. *Management Science*, *43*(2), 123–144.
- Weidmer, B., Hurtado, M., Weech-Maldonado, R., Ngo-Metzger, Q., & Bogen, K. (2006). Guidelines for Translating CAHPS Surveys. Retrieved 10/26/2007, from https://www.cahps.ahrq.gov/content/resources/pdf/Guidelines_Translation.htm.
- Wickham, H. (2007). Reshaping data with the reshape package. *Journal of Statistical Software*, *21*(12).