



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Selective vulnerability of tripartite synapses in amyotrophic lateral sclerosis

**Citation for published version:**

Broadhead, MJ, Bonthron, C, Waddington, J, Smith, WV, Lopez, MF, Burley, S, Valli, J, Zhu, F, Komiyama, NH, Smith, C, Grant, SGN & Miles, GB 2022, 'Selective vulnerability of tripartite synapses in amyotrophic lateral sclerosis', *Acta Neuropathologica*. <https://doi.org/10.1007/s00401-022-02412-9>

**Digital Object Identifier (DOI):**

[10.1007/s00401-022-02412-9](https://doi.org/10.1007/s00401-022-02412-9)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Acta Neuropathologica

**Publisher Rights Statement:**

This is the author's peer-reviewed manuscript as accepted for publication.

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## **Selective Vulnerability of Tripartite Synapses in Amyotrophic Lateral Sclerosis.**

Broadhead, Matthew J.\*<sup>1,2</sup>

Bonthron Calum\*<sup>1</sup>

Waddington Julia<sup>1</sup>

Smith, William V.<sup>1</sup>

Lopez, Maite F.<sup>1</sup>

Burley, Sarah<sup>1</sup>

Valli, Jessica<sup>2</sup>

10 Zhu, Fei<sup>3</sup>

Komiyama, Noboru H.<sup>3,4</sup>

Smith, Colin<sup>5,6</sup>

Grant, Seth G.N.<sup>3,4</sup>

Miles, Gareth B.<sup>†1</sup>.

\* co-first author, † corresponding author

1. School of Psychology and Neuroscience, University of St Andrews, St Andrews, UK

2. Edinburgh Super-Resolution Imaging Consortium, Heriot Watt University, Edinburgh, UK.

20 3. Genes to Cognition Program, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh EH16 4SB, UK.

4. Simons Initiative for the Developing Brain (SIDB), Centre for Discovery Brain Sciences, University of Edinburgh, Edinburgh EH8 9XD, UK.

5. Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK

6. Euan MacDonald Centre for MND Research, The University of Edinburgh, Edinburgh, UK.

Contact Prof. Gareth B Miles: [gbm4@st-andrews.ac.uk](mailto:gbm4@st-andrews.ac.uk)

Revised Version 2022/02/22

## 30 **Abstract**

Amyotrophic Lateral Sclerosis (ALS) is a fatal neurodegenerative disorder. Separate lines of evidence suggest that synapses and astrocytes play a role in the pathological mechanisms underlying ALS. Given that astrocytes make specialised contacts with some synapses, called tripartite synapses, we hypothesise that tripartite synapses could act as the fulcrum of disease in ALS. To test this hypothesis, we have performed an extensive microscopy-based investigation of synapses and tripartite synapses in the spinal cord of ALS model mice and post-mortem human tissue from ALS cases. We reveal widescale synaptic changes at the early symptomatic stages of the SOD1<sup>G93a</sup> mouse model. Super-resolution microscopy reveals that large complex postsynaptic structures are lost in ALS mice. Most surprisingly, tripartite synapses are selectively lost while non-tripartite synapses remain in equal number to healthy controls. Finally, we also observe a similar selective loss of tripartite synapses in human post-mortem ALS spinal cords. From these data we conclude that tripartite synaptopathy is a key hallmark of ALS.

### **Key Words**

ALS/MND; Synapse; Astrocyte; Neurodegeneration

## Introduction

50 Amyotrophic Lateral Sclerosis (ALS) is a devastating form of motor neuron disease, characterised by loss of motor neurons (MNs) in the brain, brainstem and spinal cord, leading to progressive decline of motor control and fatal paralysis, typically within a few years of diagnosis [84]. The pathological mechanisms that precede MN cell death remain unclear. One line of evidence suggests that ALS, like many neurodegenerative disorders, could be caused by a loss or a change in the synapses between neurons within the central nervous system (CNS) prior to cell loss [23, 31, 77, 87]. Pre-symptomatic and early-symptomatic changes in cortical dendritic spines, neuromuscular junctions and the synaptic inputs to spinal cord MNs have all been reported [24, 25, 60, 63, 68, 72, 87]. Separate evidence indicates that astrocytes, the  
60 supportive glial cells of the CNS, may be responsible for non-cell autonomous degeneration of MNs [17, 26, 74, 80, 81, 85]. Astrocytes make specialised connections with chemical synapses, forming tripartite synapses, through which they help regulate synaptic structure and transmission [2, 59]. Despite evidence that glial cells may contribute to synaptic degeneration in neurodegenerative diseases [5, 39, 71], it is not known whether tripartite synapses are affected in ALS. We hypothesise that both synaptic and astrocytic dysfunction could be mechanistically linked, and that the tripartite synapse could act as a vulnerable fulcrum of disease in ALS.

To investigate how synapses and tripartite synapses are affected in ALS, we have performed a thorough, quantitative microscopy-based study of millions of excitatory  
70 synapses and markers of perisynaptic astrocytic processes (PAPs) in the spinal cord. This was conducted in mice with ALS-causing mutations and in post-mortem spinal cord tissue from human ALS patients. Using high-resolution and super-resolution microscopy, in combination with genetic and immunohistochemical labelling strategies

for visualising synapses and astrocytes, our findings reveal that tripartite synapses are selectively vulnerable to degeneration in ALS. This finding suggests that synaptic loss may be mechanistically linked to astrocytic pathology in ALS, and the identification of this vulnerable synapse subtype could pave the way for targeted therapeutic strategies.

## Methods

### 80 Animals and Ethics

All procedures performed on animals were conducted in accordance with the UK Animals (Scientific Procedures) Act 1986 and were approved by the University of St Andrews Animal Welfare and Ethics Committee. The following mouse lines were kindly provided by Dr Richard Mead, University of Sheffield: B6SJL-TgN(SOD1-G93A)1Gur/J (SOD1<sup>G93a</sup>), B6SJL-Tg(SOD1)2Gur/J (WTSOD1) and FVB/NJ-Tg(C9ORF72)500Lpwr/J (C9orf72). SOD1<sup>G93a</sup> and WTSOD1 were maintained on a congenic C57bl/6 background strain, while C9orf72 mice were maintained on an FVB background mouse strain. The ALS mouse lines were crossed with PSD95-eGFP<sup>+/+</sup> mice, originally obtained from Prof. Seth Grant (University of Edinburgh), to produce transgenic (Tg) ALS and control offspring expressing PSD95-eGFP<sup>+/+</sup>. Unlike the other mouse lines, C9orf72 mice were produced on an FVB background. The progeny of C9orf72 x PSD95-eGFP mice were a first generation FVB/C57bl/6 cross, displaying a light brown fur coat and black eyes (indicating they did not possess the recessive retinal degeneration 1 allele of Pde6brd1 which causes blindness in FVB mice).

Disease phenotype was monitored by routine weighing and behavioural scoring from the age of 30 days old (P30), with more frequent checks performed once the mice were over P60 or when mice were showing disease phenotypes. Mice were scored 0-4 in severity; 4 showing no phenotype, 3 showing hind-limb splay and tremors upon

tail raising, 2 showing gait abnormalities, 1 showing hind limb dragging, 0 showing  
100 inability to self-right after 30 seconds when placed on its back. Mice were euthanised  
through either cervical dislocation or perfusion if displaying a score of 2 or lower.  
Overall, SOD1<sup>G93a</sup> mice presented hind limb tremor and reduced hind-limb splay by  
76 days We observed no difference in the age of symptom onset between SOD1<sup>G93a</sup>  
mice and the cross-bred PSD95-eGFP mice, nor have we observed any difference in  
the age of symptom onset within our colonies over time. Our data is remarkably  
consistent with those of the source colonies [53] and suggests that little-to-no  
reduction in transgene copy number had occurred within our breeding colonies or  
experimental cohorts. No such overt phenotypes were observed in either the WTSOD1  
and C9orf72 mouse colonies.

#### 110 **Mouse Tissue Collection**

Mice were anaesthetised with pentobarbitol (30mg/kg dose; Dolethal), and the chest  
cavity opened to reveal the heart. The right atrium was severed and 10 ml ice cold 1x  
phosphate buffered saline (PBS) was perfused through the left ventricle, followed by  
10ml 4% paraformaldehyde (PFA; Alfa Aesar). The lumbar spinal cord was then  
dissected and incubated for a further 3-4 hrs in 4% PFA before being incubated in  
sucrose 30% w/v for 24 to 72 hours at 4°C until sunk. Tissue was then cryo-embedded  
in OCT compound and stored at -80°C. Cryosections were obtained using a Leica  
CM1860 cryostat at 20 µm thickness and adhered to Superfrost Gold Plus glass slides  
(VWR).

#### 120 **Mouse Tissue Processing and Immunohistochemistry**

Mouse tissue processing for visualising synapses and tripartite synapses is detailed  
in Broadhead et al., (2020) [6]. For visualisation of PSD95-eGFP only, slides were  
thawed at room temperature, briefly washed in PBS then deionised (DI) water before

being dried and mounted in lab-made Mowiol or Prolong Glass anti-fade for super-resolution microscopy with a 1.5 thickness coverslip. For immunohistochemistry, sections were first heated at 37°C for 30 mins to aid the adherence of the tissue to the glass slides and reduce tissue loss during subsequent wash steps. Slides were washed 3 times in PBS before being blocked and permeabilised in PBS containing 3% Bovine Serum Albumin (BSA) and 0.2% Triton X100 for 2 hours at room temperature.

130 Primary antibody incubation was performed in PBS containing 1.5% BSA, 0.1% Triton X100 for 2 nights at 4°C. Once incubated, slides are washed 5 times in PBS over the course of 1-2 hours. Secondary antibody incubation was performed in 0.1% Triton X100 for 2 hours at room temperature followed by a further 5 washes in PBS over the course of 1-2 hours. Dye labelling of astrocyte-specific primary antibodies (GFAP, EAAT2 and p-Ezrin) was performed with 10µg of each antibody using the Lightning-Link conjugation kits (Abcam) to address whether some proteins colocalised in astrocytes when the primary antibodies were raised in the same species (rabbit). See Table 1 for details on antibodies for mouse tissue experiments.

#### **Human Post-Mortem Tissue and Immunohistochemistry.**

140 Human cervical spinal cord samples were obtained from the Edinburgh Brain and Tissue Bank, by Prof. Colin Smith. Ethical approval for this work was provided by the East of Scotland Research Ethics Service REC 1 (16/ES/0084). Samples of the cervical spinal cord were obtained from patients who died from ALS displaying mutations in the C9orf72 gene or the SOD1 gene. Age matched patients who died from natural causes were used as controls. Mean average patient ages were 57.2 years (controls), 57.6 years (C9orf72) and 54 years (SOD1). Immunohistochemistry protocol was adapted from Curran et al., (2021) [15]. Sections were fixed, paraffin embedded and sectioned at 4 µm thickness and mounted to glass slides. For

immunohistochemistry, samples were first de-paraffinised with sequential 3 minute  
150 washes in Xylene, 100% ethanol, 95% ethanol, 70% ethanol and 50% ethanol. Antigen  
retrieval was performed by incubating samples in Citric Acid (pH 6.0) for 20 minutes  
in a steamer, providing a consistent heat of 95°C. Following antigen retrieval, sections  
were washed 3 times in PBS and then incubated in 1x Tris buffered Saline (TBS) with  
5% BSA, 0.2% Triton X100 for 2 hours at room temperature. Samples were then  
incubated in primary antibody solution consisting of TBS with 3% BSA, 0.2% Triton  
X100 for 2 nights at 4°C before being washed 5 times in TBS containing 0.2% Triton  
X100 over the course of 1 hour. Secondary antibody incubation was performed in TBS  
with 0.2% Triton X100 solution for 3 hours at room temperature followed by 5-6 washes  
in PBS with 0.2% Triton X100 over the course of 2 hours. Finally, sections were  
160 washed in DI water, dried and mounted in Mowiol. See Table 2 for antibodies used in  
human tissue experiments.

### **RNA FISH**

Cy5-labelled DNA probes were used against the G4C2 RNA repeat expansion. All  
solutions were made up in DEPC-treated water or PBS. Fixed spinal cord tissue was  
permeabilised with 0.1% Triton X100 for 1 hour. Tissue was treated with 2x saline  
sodium citrate (SSC) and 10% formamide in PBS for 5 minutes. Slides were then  
incubated with hybridization solution containing probes in 10 % formamide overnight  
at 37°C. After hybridization slides were washed once in 2x SSC containing 10%  
formamide, for 30 mins at 37°C, then washed in 2x SSC for 5 mins at room  
170 temperature. DAPI was used at 1:15,000 in PBS. Images were captured on a Zeiss  
Airyscan LSM800 at 63x.

### **High-Resolution Microscopy**



High-resolution confocal-like microscopy was performed using a Zeiss Axio Imager M2 Microscope equipped with an Apotome 2.0, which enables an XY resolution of 320 nm [6]. Illumination was provided by a HXP120 lamp, and images acquired using a digital MRm camera. Exposure times and illumination intensity were kept consistent for analysis within batches of data sets. To ‘map’ entire hemi-sections of spinal cords to study anatomical diversity between laminae, single optical sections were captured and tiled across half a transverse section of the spinal cord. Maps were captured using  
180 a single Z-stack plane approximately 3-5  $\mu\text{m}$  depth into the tissue that was kept at the same depth throughout the entire map. Images were stitched together to create whole montage images that were subsequently analysed.

### **Super-Resolution Microscopy**

Confocal and gated-stimulated emission depletion (g-STED) microscopy was performed using a Leica SP8 SMD g-STED microscope available at the Edinburgh Super-Resolution Imaging Consortium hosted by Heriot Watt University. Excitation was provided by a CW super-continuum white light laser source at 488nm to excite eGFP with depletion provided by a 594 nm and 775 nm laser. Images were acquired with a 100x 1.4NA STED objective lens with an optical zoom set to provide optimal xy-  
190 resolution.

### **Image Analysis**

Image analysis was performed in FIJI (Fiji is just ImageJ) [66]. High-resolution image analysis was performed as described previously [6]. A standard mouse spinal cord anatomical atlas was used to guide the delineation of Rexed’s laminae [76]. For clarity, our delineation of lamina VIII represents a heterogenous population of ventral neurons including medial MN pools, whilst laminae IX represents lateral MN pools. For mapping data sets, whole images were processed using a background subtraction and

gaussian smoothing. For the PSD95-eGFP-only data sets of excitatory synapse mapping, automated Moments-based thresholding was applied. For data sets  
200 involving detection of immunolabelled markers of synaptic or astrocytic structures, manual thresholding was typically required, and the researcher was blinded to the genotype of the mouse to avoid unconscious bias. Minimum and maximum size thresholds were set to minimise false positive detections of small dim structures of a few pixels in size and large clusters that were likely to be lipofuscin aggregates. Colocalisation between structures in high-resolution data sets was determined from binarized images with the criteria of showing overlap by at least 1 pixel.

Nanostructural analysis of PSD95 organisation from g-STED imaging was analysed as described previously [6, 7]. Briefly, confocal and g-STED images are processed with background subtraction and gaussian smoothing. Structures resolved in g-STED  
210 were deemed to be PSD95 nanoclusters (NCs), while larger diffraction-limited structures resolved with confocal microscopy were deemed to be whole PSDs. Structural parameters such as size, shape and intensity were obtained for PSDs and NCs, as well as how many NCs resided within a given PSD.

### **Statistical Analysis.**

Data analysis and graph preparation was performed in Microsoft Excel. Statistical analyses were performed using either SPSS (IBM) or Prism (Graphpad). Data were assessed for normality using the Shapiro Wilks test. Multi-factorial (Two-Way) ANOVA were used for mouse spinal cord synaptic and tripartite synaptic mapping (data presented in Figure 1 and 3, SI Fig. 3, 6 and 7), where the factors were genotype and  
220 anatomical subregion. Sulak's post-hoc comparison test was performed following Two-Way ANOVA's where more than two genotypes (e.g. controls, SOD1<sup>G93a</sup> and WTSOD1 mice) were compared, while Tukey's post-hoc comparison was performed

following Two-Way ANOVA's where only two groups were compared (e.g. controls and SOD1<sup>G93a</sup> mice). Two Sample T-tests were performed for PSD95 nanoscale analyses (presented in Fig. 2c-e), zDHHC2 expression (Fig. 2g,h), ChAT MN staining quantification (SI. Fig. 2b-c) and human ALS synapse analysis, when both SOD1 and C9orf72 patient cases were merged into one 'ALS' group and compared with controls (Fig. 4c,d). Pearson's Chi-squared test was performed to test for differences in the frequency of synapse subtypes in controls and SOD1<sup>G93a</sup> mice, in combination with  
230 Two-Sample T-Tests to compare the frequency of each specific synapse subtype between the two genotypes (Fig. 2e). Pearson's Correlation Analysis was performed to assess regional differences in tripartite synapse loss and the percentage of synapses contacted by astrocytes (SI. Fig. 7). One-Way ANOVAs with post-hoc Tukey's comparison tests were used to assess differences in synapses and tripartite synapses between Control, SOD1 and C9orf72 human patient spinal cords (Fig. 4c,d).

## Results

### Widescale Mapping of the Mouse Spinal Cord Reveals Synaptic Changes in ALS

To visualise and quantify synaptic changes in ALS mice using a standardised and  
240 robust approach, we cross-bred PSD95-eGFP<sup>+/+</sup> mice [7, 86] with SOD1<sup>G93a</sup> mice [35], WTSOD1 mice [35] and C9orf72 mice [48]. The use of PSD95-eGFP mice ensures that all mice expressed fluorescently tagged postsynaptic density (PSD) scaffolding molecule, PSD95, at virtually all excitatory synapses throughout the nervous system [86]. As sex-dependent differences in phenotypes have been observed in both the SOD1<sup>G93a</sup> and C9orf72 mice [10, 36, 48], males and females were analysed separately.

From here onwards, the terms SOD1<sup>G93a</sup>, WTSOD1 and C9orf72 refer to mice produced from breeding with homozygous PSD95-eGFP knock-in mice to facilitate synapse visualisation in heterozygous PSD95-eGFP progeny. The term 'Control' is used to refer to heterozygous PSD95-eGFP progeny not expressing the SOD1<sup>G93a</sup> or C9orf72 transgene.

Consistent with previous reports [53], SOD1<sup>G93a</sup> mice presented with hind-limb tremor and hind-limb splay by approximately 76 days of age, with males displaying a slightly earlier onset compared to females, though this was not statistically significant (73.5 days old for n=4 male SOD1<sup>G93a</sup> mice, 77.1 days for n=9 female SOD1<sup>G93a</sup> mice;  $t(11)=1.3$ ,  $p=0.21$ ).

At the point of termination, none of the 8 week SOD1<sup>G93a</sup> mice displayed hind limb tremor (5 males, 5 females); all the 12 week male (5/5) and some of the 12 week female (3/5) SOD1<sup>G93a</sup> mice displayed hind limb tremor, and all the 16 week mice SOD1<sup>G93a</sup> (6 males, 5 females) displayed hind limb tremor. 16 week old SOD1<sup>G93a</sup> mice were also lower in weight compared to controls (males:  $t(9)=5.2$ ,  $p=0.0006$ ; females:  $t(8)=5.3$ ,  $p=0.0007$ ). No signs of motor dysfunction or weight loss were observed in WTSOD1 mice up to 16 weeks. Similarly, no signs of acute motor dysfunction or weight loss were seen in C9orf72 mice up to 22 weeks. C9orf72 mice did display G4C2 RNA foci, a key molecular hallmark of disease, in the nuclei of spinal cord cells (Sl. Fig. 1a-b).

Tiled, high-resolution maps of PSD95-eGFP synapses were obtained across hemisections of spinal cords which we delineated into spinal laminae I-II, III-IV, V, VI, VII, VIII, IX and X. Each map of a mouse hemi-spinal cord yielded data on approximately 40,000 to 50,000 PSDs. In total, spinal cord synapse maps were obtained from 122 mice, from which over 5 million PSDs were quantified. In this study, our analysis

focuses on PSD density (the number per unit area) and PSD size (area). Overall, inter-regional mapping of PSD95 revealed anatomical diversity in synapse number and size comparable to our previous observations [6], with little difference associated with age or sex.

In both male and female SOD1<sup>G93a</sup> and control mice, synaptic maps were generated at 8 weeks (pre-symptomatic), 12 weeks (early-symptomatic) and 16 weeks of age (early-symptomatic with significant weight loss) (Fig. 1 a-b). At 8 and 12 weeks of age there was no difference between SOD1<sup>G93a</sup> and controls in PSD density (Fig. 1c) or PSD size (Fig. 1d). At 16 weeks of age, however, SOD1<sup>G93a</sup> males showed significantly reduced PSD density ( $F(1,72)=9.60$ ,  $P=0.0028$ ) and PSD size ( $F(1,72)=15.77$ ,  $P=0.0002$ ) compared to controls. These effects were also significant when SOD1<sup>G93a</sup> males were compared with both controls and WTSOD1 controls (Fig. 1e, PSD density:  $F(2,104)=7.49$ ,  $P=0.0009$ ; Fig. 1f, PSD size:  $F(2,104)=11.02$ ,  $P<0.0001$ ). These synaptic changes were widespread across the spinal cord, with the exception of the dorsal laminae I-II; while post-hoc Tukey's testing revealed that PSD density and size were most significantly affected in lamina VI and VII respectively.

The SOD1<sup>G93a</sup> 16-week old females displayed reduced PSD density when compared to control females ( $F(1, 64) = 10.26$ ,  $P=0.0021$ ), but no difference in PSD size ( $F(1, 64) = 3.23$ ,  $P=0.077$ ). Furthermore, these effects were not significant in comparison to the female 16-week-old WTSOD1 mice (Fig. 1g-h). Interestingly, it was noted that unlike the SOD1<sup>G93a</sup> males, the SOD1<sup>G93a</sup> females displayed more complex laminae-genotype-specific synaptic changes in both PSD density ( $F(14, 96) = 2.34$ ,  $P=0.008$ ) and size ( $F(14, 96) = 3.68$ ,  $P<0.0001$ ). The dorsal laminae I-II of SOD1<sup>G93a</sup> females displaying increased PSD density and size compared to WTSOD1, while ventral

laminae were more likely to show reduced PSD density and size compared to WTSOD1 and controls.

To address whether synaptic loss was associated with MN cell death, we performed immunolabelling using choline acetyltransferase (ChAT) specific antibody. We found  
300 no significant loss of MNs in 16-week old SOD1<sup>G93a</sup> male spinal cords (SI. Fig. 2a-d;  $t(6)=0.99$ ,  $p=0.36$ ). While other studies indicate MN cell loss can occur from approximately 16 weeks onwards [19], our data suggest that synaptic changes occur prior to overt MN cell death.

Next, we examined the spinal cords of mice carrying a mutation in the C9orf72 gene, one of the most common genes associated with familial and sporadic ALS (SI. Fig 3a-b). Synaptic maps from 12 week and 22 week old C9orf72 mutant mice of both sexes were obtained to capture reported pre-symptomatic and early-symptomatic stages of the disease phenotype in this model [48, 55, 58]. Mapping PSD95-eGFP synapses revealed no difference in PSD density or size in C9orf72 mice compared to controls at  
310 either 12 or 22 weeks of age in both males and females (SI. Fig. 3c-j). The lack of synaptic changes in C9orf72 mice may reflect the fact that no motor phenotypes were observed in any of our C9orf72 mice even by 22 weeks. Our observations are consistent with reports from other groups that the penetrance of the phenotype in this mouse model can be very low in some colonies [55, 58].

In summary, our extensive synaptic mapping reveals a loss of PSDs at excitatory synapses in both male and female SOD1<sup>G93a</sup> mice during early symptomatic stages. Our data also suggest that male SOD1<sup>G93a</sup> mice display structural changes in synapses, an effect not as clearly observed in female SOD1<sup>G93a</sup> mice, perhaps due to the delayed onset of disease in females. While we observed no synaptic changes in  
320 C9orf72 mutant mice, this mirrors the fact that no ALS-related phenotypes, such as

acute motor decline or weight loss, were observed in these mice. Given that our C9orf72 mice fail to robustly recapitulate an ALS disease phenotype, they were not included in any further analyses.

### **Super-Resolution Microscopy Reveals Loss of Multi-Nanocluster Synapses in the ALS Mouse Spinal Cord**

Our high-resolution microscopy data revealed both a reduced number and reduced average size of PSDs in 16-week SOD1<sup>G93a</sup> males. While confocal-level resolution is sufficient to measure synapse number and overall PSD size, the nanoscale organisation of synapses is best resolved using super-resolution microscopy. PSD95 forms nanoclusters (NCs) of approximately 100-150 nm diameter within the PSD [6, 7, 49, 56]. PSD size, which correlates with synaptic strength, is determined by the number of NCs per PSD (Fig. 2a). We performed g-STED microscopy of PSD95-eGFP expression in 16-week-old male SOD1<sup>G93a</sup> and control mice to quantify synaptic nanostructure (Fig. 2b; n=4 mice per group). PSD95 NC size did not differ between genotypes (diameter ~140nm; Fig. 2c;  $t(6)=0.31$ ,  $p=0.77$ ), and was consistent with previous measurements of PSD95 NC size using g-STED microscopy in both the brain and spinal cord [6, 7, 30]. Instead, SOD1<sup>G93a</sup> mice showed a reduced average number of NCs per PSD (~1.75 NCs per PSD) compared with controls (~2 NCs per PSD) (Fig. 2d;  $t(6)=2.8$ ,  $p=0.03$ ). This appears to be the result of a significant loss of large multi-NC-PSDs in SOD1<sup>G93a</sup> mice, while smaller 1 or 2 NC-PSDs were observed with similar frequency (Fig. 2e;  $\chi(2) = 14.7$ ,  $p=0.0006$ ). In controls, 23% of synapses were classed as containing 3 or more NCs, while 17% of SOD1<sup>G93a</sup> synapses contained 3 or more NCs.

The formation of PSD95 NCs at the synapse is dependent on palmitoylation of PSD95 at its N-terminal cysteine residue by the palmitoylating enzyme, zDHHC2. Having revealed a reduced number of PSD95 clusters per synapse in ALS mice, we assessed whether a change in zDHHC2 expression could be responsible. We analysed excitatory synapses (PSDs using PSD95-eGFP and presynaptic terminals using anti-VGLUT2 immunolabelling) for their expression of zDHHC2 by acquiring high-resolution images from the ventral horn of spinal cords from 16 week male control and SOD1<sup>G93a</sup> mice (Fig 2f). zDHHC2 displayed a punctate expression showing close association with both PSD95-eGFP and VGLUT2 puncta. While the overall density of zDHHC2 puncta was not significantly different between genotypes (Fig 2g;  $t(9)=2.1$ ,  $p=0.067$ ), there was a reduced number of PSDs colocalising with z-DHHC2 in SOD1<sup>G93a</sup> mice (Fig. 2h;  $t(9)=2.6$ ,  $p=0.031$ ). This finding suggests that reduced postsynaptic presence of the palmitoylation enzyme, zDHHC2, may be related to a reduced number of PSD95 NCs at the synapse and lead to incremental reductions in postsynaptic strength.

360

### **Widescale Mapping Reveals Tripartite Synapse Loss in the ALS Mouse Spinal Cord.**

Tripartite synapses are chemical synapses contacted by perisynaptic astrocytic processes (PAPs). Previously we used antibody labelling against PAP proteins EAAT2 and p-Ezrin to identify tripartite synapses in the mouse lumbar spinal cord based on colocalization or close association of PAP proteins with postsynaptic PSD95 and presynaptic VGLUT2 [6]. Immunolabelling for VGLUT2, p-Ezrin and EAAT2 are further validated here (SI. Fig. 4a-f; SI. Fig. 5a-b). p-Ezrin and EAAT2 puncta strongly colocalised with one another in the mouse spinal cord (SI. Fig 5a). Both PAP markers are closely associated with astrocytic cell bodies, branches and fine processes,

370



labelled with glutamine synthetase, but display little direct colocalization with astrocytic GFAP-positive primary arborescences (SI. Fig 5b). These observations further confirm that both astrocytic EAAT2 and p-Ezrin label PAPs, making them ideal markers, in combination with synaptic markers, for visualising tripartite synapses. We, therefore, performed extensive high-resolution mapping of tripartite synapses in SOD1<sup>G93a</sup> and control spinal cords, labelling PSDs (PSD95-eGFP), presynaptic terminals (VGLUT2) and PAPs (EAAT2 or p-Ezrin).

At 8 weeks, male SOD1<sup>G93a</sup> mice displayed an increased density of EAAT2 puncta ( $F(1, 56) = 11.19, p=0.0015$ ; SI. Fig 6a) and p-Ezrin puncta ( $F(1, 56) = 4.672, p=0.035$ ; SI. Fig. 6b) compared to controls. No changes were observed in male 12-week-old mice (SI. Fig 6c-d), but by 16 weeks, male SOD1<sup>G93a</sup> mice displayed significantly reduced EAAT2 ( $F(1, 64) = 11.31, p=0.0013$ ; SI. Fig 6e) and p-Ezrin ( $F(1, 48) = 8.828, p=0.0046$ ; SI. Fig. 6f) puncta density compared to controls. These data demonstrate age-dependent changes in PAP proteins, suggesting an early stage increase in p-Ezrin-positive PAPs and increased EAAT2 expression at PAPs, followed by a significant loss of both proteins by 16 weeks, following the onset of early motor deficits.

We next quantified the density of non-tripartite synapses and tripartite synapses in male control and SOD1<sup>G93a</sup> spinal cords at 8, 12 and 16 weeks. Using p-Ezrin, we observed no difference in the density of non-tripartite synapses between controls and SOD1<sup>G93a</sup> males at either 8 weeks (Fig. 3c,  $F(1, 56) = 2.24, P=0.1404$ ) 12 weeks (Figure 3e;  $F(1,56)=0.56, P=0.456$ ) or 16 weeks of age (Fig 3g;  $F(1,48)=2.4, p=0.126$ ). The density of tripartite synapses was also no different between control and SOD1<sup>G93a</sup> males at 8 weeks (Fig 3c;  $F(1,56)=0.2998, P=0.586$ ) and 12 weeks (Fig. 3e;  $F(1,56)=0.83, p=0.365$ ). However, there was a significant reduction in tripartite

synapse density at 16 weeks in SOD1<sup>G93a</sup> males (Fig 3g;  $F(1,48)=14.59$ ,  $P=0.0004$ ). It was also confirmed that female 16 week old SOD1<sup>G93a</sup> mice displayed reduced numbers of p-Ezrin associated tripartite synapses compared to controls (SI. Fig 7a;  $F(1,80)=5.65$ ,  $P=0.020$ ), while non-tripartite synapses were seemingly unaffected (SI. 400 Fig. 7b;  $F(1,80)=0.19$ ,  $P=0.66$ ).

Similarly, using EAAT2 as the PAP marker, the density of non-tripartite synapses was no different between controls and SOD1<sup>G93a</sup> males at either 8 weeks (Fig. 3d;  $F(1, 56)<0.0001$ ,  $p=0.99$ ), 12 weeks (Fig. 3f;  $F(1, 56)=1.37$ ,  $p=0.247$ ) or 16 weeks (Fig. 3h;  $F(1,64)=2.40$ ,  $p=0.126$ ). The density of EAAT2-associated tripartite synapses, however, was significantly increased in 8 week SOD1<sup>G93a</sup> males (Fig. 3d,  $F(1, 56)=7.21$ ,  $p=0.0095$ ), then subtly but significantly reduced in 12 week SOD1<sup>G93a</sup> males (Fig. 3f;  $F(1, 56) = 4.77$ ,  $p=0.033$ ) and more significantly reduced in 16 week SOD1<sup>G93a</sup> males (Fig. 3h;  $F(1, 64)=23.58$ ,  $p<0.0001$ ). Female 16 week old SOD1<sup>G93a</sup> mice also displayed reduced numbers of EAAT2-associated tripartite synapses compared to 410 controls (SI. Fig 7c;  $F(1,88)=8.42$ ,  $P=0.0047$ ), while non-tripartite synapses were unaffected (SI. Fig. 7d;  $F(1,88)=3.48$ ,  $P=0.066$ ).

Next, we investigated whether there is a relationship between synaptic vulnerability and astrocytic coverage of synapses in different spinal cord laminae (SI. Fig. 8a-d). In the 16 week old male SOD1<sup>G93a</sup> mice, there was a significant correlation between the percentage of synapses contacted by p-Ezrin and the loss of tripartite synapses in each region ( $r = 0.935$ ,  $p = 0.0006$ ; SI. Fig. 8a-b). For example, dorsal laminae I-II displayed the highest proportion of synapses contacted by p-Ezrin (28%) and the smallest degree of tripartite synapse loss compared to controls (-29%). This compares to ventral laminae VII with the smallest proportion of synapses contact by p-Ezrin 420 (17%) and the greatest degree of tripartite synapse loss compared to controls (-48%).

There was a similar correlation between the percentage of EAAT2-associated tripartite synapses and the loss of tripartite synapses, though this was not statistically significant ( $r = 0.345$ ,  $p = 0.401$ ; SI. Fig. 8c-d).

From these experiments, using two different PAP markers, our data consistently demonstrate that tripartite synapses are a selectively vulnerable synapse subtype in early-symptomatic stage ALS mice.

### **Selective loss of Tripartite Synapses in the Human ALS Spinal Cord**

430 Finally, we addressed whether human cases of ALS also demonstrated synaptic and tripartite synaptic degeneration. Post-mortem cervical spinal cord sections were obtained from human cases, including 6 healthy controls (5 male, 1 female), and 9 ALS patients (all male) comprising 4 SOD1 cases and 5 C9orf72 cases (SI. Table 1). Synapses and tripartite synapses were visualised in the ventral horn using immunolabelling for PSD95 and p-Ezrin to identify PSDs and PAPs respectively (Fig. 4a-b; SI. Fig. 9a-f).

When comparing control cases with all ALS cases (SOD1 and C9orf72), PSD density was significantly reduced in the ALS cervical spinal cord (Fig. 4c;  $t(13)=2.6$ ,  $p=0.021$ ). Interestingly, when control, SOD1 and C9orf72 cases were analysed separately, only C9orf72 cases displayed significantly reduced PSD density compared to controls 440 ( $F(2,12)=5.4$ ,  $P=0.0214$ ; Control versus C9orf72:  $p=0.0167$ ). We observed no difference in PSD size between the three groups ( $F(2,12)=0.16$ ,  $P=0.85$ ).

Next, we asked how tripartite synapses and non-tripartite synapses are affected in ALS patients (Fig. 4d). Similar to results from the SOD1<sup>G93a</sup> mouse model, the density of non-tripartite synapses was no different between control, SOD1 and C9orf72 patients ( $F(2,12)=1.44$ ,  $P=0.275$ ), while the density of tripartite synapses was

significantly reduced in C9orf72 cases compared to controls ( $F(2,12)=8.16$ ,  $P=0.0058$ , Control versus C9orf72:  $p=0.0044$ ).

Taken together these findings provide evidence that tripartite synapses are selectively vulnerable to degeneration in human ALS cases, as observed in the SOD1<sup>G93a</sup> mouse  
450 model.

## Discussion

This study set out to investigate how synapses and tripartite synapses are affected in the ALS spinal cord. To this end, we performed large-scale quantitative mapping of synapses and tripartite synapses in ALS mouse models and human post-mortem tissue. Our data revealed that, of the millions of synapses in the spinal cord, tripartite synapses appear to be selectively vulnerable to degeneration in ALS. We therefore describe ALS as a tripartite synaptopathy.

460 The functional, structural and molecular diversity of synapses throughout the nervous system, i.e. synaptome diversity, endows the nervous system with the capability to support a wide repertoire of behaviours from motor control to learning and memory [32, 33]. Over 130 different nervous system disorders, including ALS, can arise from mutations in postsynaptic proteins [4], and lead to wide-scale reprogramming of the synaptome [86]. Different diseases of the nervous system may display a unique synaptome signature, as certain subtypes of synapses may be more vulnerable to changes in one condition compared to another – thus leading to the specific behavioural consequences and symptoms of a given condition. By first identifying the vulnerable synapse subtypes of a given disease, we can establish therapeutic  
470 roadmaps to selectively target these synapse subtypes to treat the symptoms of disease.

The PSD95-eGFP mouse model provides a powerful tool with which to investigate the effects of diseases on synapses as PSD95 is abundantly expressed at the PSDs of excitatory synapses and forms a core molecular framework for synaptic machinery [13, 20, 28, 42, 86]. The same mouse model can be used for nanoscopic analyses of synapse subtypes based on postsynaptic substructure using super-resolution microscopy [6, 7].

This study also sought to identify whether astrocytes play a role in the vulnerability of synapses in ALS. Labelling the astrocytic component of the tripartite synapse was performed here using immunolabelling for either EAAT2 or p-Ezrin. Various studies support the use of either EAAT2 or Ezrin/p-Ezrin as robust markers for the PAPs which contact synapses [16, 27, 37, 46]. By performing tripartite synapse mapping with both PAP markers in parallel, we have observed strikingly similar findings which give us considerable confidence in our conclusion that tripartite synapses are vulnerable in ALS. Reduced levels of EAAT2 in the ALS spinal cord have been well documented [41, 47, 65]. The consistent reduction in both EAAT2 and the actin filament binding protein, p-Ezrin, suggest not just a molecular change in astrocytic protein expression but that the PAPs themselves are being lost.

Using the SOD1<sup>G93a</sup> mouse model, we observed synapse loss, structural changes in PSDs, changes in PAPs and a loss of tripartite synapses at early, symptomatic stages. While we observed significant synaptopathy in SOD1<sup>G93a</sup> mouse spinal cords, we observed no such alterations in C9orf72 mutant mice. The C9orf72 (C9BAC-500) mice were originally reported to display molecular hallmarks of ALS pathology, and acute motor deficits in 35% of the females carrying the mutation [48]. Further investigation by other groups using this model have shown different degrees of phenotypic penetrance [55, 58]. In our hands, the C9orf72 mice (the pure-bred colony and the

progeny of PSD95-eGFP cross-breeding) display no such acute motor phenotypes in either males or females, though we did observe RNA foci accumulating in cell nuclei – a key molecular hallmark of ALS pathology. Other key molecular hallmarks of ALS such as TAR-DNA Binding protein-43 (TDP-43) pathology have been observed in the brains of symptomatic end-stage C9orf72 mice, but were not observed in asymptomatic mice [48], and this remains untested in the spinal cord. We consider that either genetic or environmental differences in our mouse colonies may have led to significantly reduced penetrance of the disease phenotype in the C9orf72 mouse line [58]. Given that phenotypic penetrance can be low, and that acute ALS phenotypes can occur between 20 and 40 weeks of age, further data collection from larger numbers of older animals may have captured the potential synaptic phenotype of this model. On the other hand, another mouse model with a more reliable phenotype, such as TDP-43<sup>A315T</sup> which has been shown previously to display both synaptic and astrocytic changes [44, 45], may be a more suitable alternative for further investigation of tripartite synaptopathy.

We employed large-scale anatomical mapping of synapses and tripartite synapses in the lumbar mouse spinal cord. It was anticipated that some anatomical laminae, for example laminae VII-IX where MNs reside, may display more significant changes in synapses in ALS. Despite significant inter-regional diversity in PSD size and number [6], our data suggest that the loss of synapses occurs widely throughout the spinal cord and not simply localised to regions where MNs reside. This observation suggests that ALS may impact a broad range of sensory, integratory and motor circuitry. Interestingly, female SOD1<sup>G93a</sup> mice displayed significantly increased numbers of synapses in dorsal laminae I-II compared to controls, which is supported by previous reports of sensory-level alterations in ALS and other related disorders [22, 75]. p-Ezrin

associated tripartite synapse loss also appeared to be most prominent in ventral lamina VII, where the smallest percentage of synapses are contacted by p-Ezrin. It is plausible that greater glial coverage of synapses in some regions of the nervous system may in fact help preserve synapses from degeneration. Regional diversity in astrocyte properties could also contribute to variation in tripartite synapse vulnerability [54, 73]. It would be of interest to investigate the nature of tripartite synapses in regions of the nervous system more resistant to ALS neurodegeneration, such as the oculomotor nucleus [14], to further elucidate why some synapses are more vulnerable than others.

Our mapping analyses also suggested that while synapse and tripartite synapse loss occurred in both male and female SOD1<sup>G93a</sup> mice, structural changes in PSDs may be more prominent across the whole spinal cord in males. In female SOD1<sup>G93a</sup> mice however, PSD size was reduced in many ventral laminae but increased in some dorsal laminae. Other studies indicate a more discernible gender bias in SOD1<sup>G93a</sup> mice, with males showing an earlier onset and increased rate of disease progression, and some changes in specific subtypes of spinal synapses only occurring in males [10, 36, 40]. In addition, clinical ALS cases present with a higher incidence in males [52].

Our extensive analysis of SOD1<sup>G93a</sup> mouse spinal cords revealed significant postsynaptic changes at the early symptomatic stage of 16 weeks (112 days), with no overt changes at 12 weeks (84 days) or 8 weeks (56 days) of age, although early signs of tripartite synapse loss were observed at 12 weeks. Others have reported reduced sizes of clusters of other synaptic proteins such as scaffolding proteins Shank1 and Homer, and glutamatergic AMPA receptors on spinal cord MNs in pre-symptomatic 8 week old SOD1<sup>G93a</sup> mice [3]. Similarly, pre-symptomatic changes in cortical synapses and neuromuscular junctions have been observed as early as 4-6 weeks in mouse

models [18, 21, 24, 25, 29]. Our current data does not suggest that significant synaptic alterations occur pre-symptomatically in spinal cord circuits, but instead suggests that they occur at the early-symptomatic stages of the disease. Dying-forward (cortex to spinal cord) and dying-backward (periphery to spinal cord) mechanisms exist pre-symptomatically [18, 21], and may not lead to significant synaptic changes in the spinal cord until later in the disease progression. Furthermore, different techniques to visualise and quantify synaptic structure and function may confound direct comparisons across studies. For example, it is conceivable that changes in dendritic spines may occur without any apparent changes in the expression or organisation of PSD95 (Fogarty et al., 2015). Comparative analyses of synaptic changes in both the brain and spinal cord in the early stages of ALS may help decipher whether synaptic changes occur simultaneously in various parts of the CNS, or whether, for example, motor cortex dysfunction drives synaptic dysfunction in the spinal cord in a feedforward manner.

In accordance with other reports of changes in the structure of synapses in ALS [3, 63, 64], our high-resolution imaging revealed reduced PSD size in early-symptomatic SOD1<sup>G93a</sup> males. By employing super-resolution microscopy, we have performed nanoscopic analysis of the molecular organisation of synapses in ALS for the first time. This powerful microscopy technique has characterised the nature of ALS synaptopathy at the sub-diffraction scale, revealing a reduced number of large multi-NC synapses. PSD95 forms nanoscale postsynaptic domains that tether neurotransmitter receptor subdomains at the postsynaptic membrane, which align with presynaptic release sites to create trans-synaptic signalling nanocolumns [70]. Synaptic strength appears to correlate with the number of nanodomains within the synapse [7, 49, 56]. Our previous work has shown that multi-NC synapses are more



likely to be tripartite synapses, based on their association with PAP proteins EAAT2 and p-Ezrin [6]. This supports the concept that large multi-NC tripartite synapses are those which are selectively vulnerable to degeneration in ALS. Given that the number of PSD95 subdomains per synapse correlates with synaptic strength, it is possible that the loss of these large, high-fidelity synaptic connections significantly contributes to early-stage, progressive motor deficits. zDHHC2 is a palmitoylating enzyme that facilitates the clustering of PSD95 at the synapse. From our data, we also speculate that the dissociation of zDHHC2 from the PSD may lead to reduced clustering of PSD95 at the synapse [30], though the precise role of protein palmitoylation in neurodegenerative disease remains to be determined [12, 83].

Importantly, we were able to replicate our key observations of synapse and tripartite synapse loss in ALS mouse models when studying human cervical spinal cord tissue from ALS patients. Our analysis suggested that C9orf72 patients displayed significant synaptic loss and tripartite synapse vulnerability, while SOD1 patients did not show significant synapse or tripartite synapse loss compared to controls. In comparison, a previous investigation into synaptic changes in the prefrontal cortex of ALS patients revealed no differences in synaptic loss between SOD1 and C9orf72 associated cases, but did suggest that synaptic loss was most significant in cases displaying TDP-43 pathology – of which SOD1 cases showed none [38]. Together these data highlight the heterogenous nature of ALS. This heterogeneity is further underlined by the range of familial and sporadic cases of ALS. Whilst less data is currently available regarding the fate of synapses in sporadic ALS, given that C9orf72 mutations represent the most common genetic basis of both sporadic and familial ALS and since 3 out of 5 C9orf72 patients used in this study had no family history of ALS, our findings may be applicable to both familial and sporadic cases.

On the face of it, our data presents a dichotomy between the SOD1 and C9orf72 mouse models and human patients. However, the lack of a robust phenotype in the C9orf72 mouse model (in our hands) likely renders it inappropriate to directly compare the two genotypes across mouse models and human patients. In addition to this, it is already known that there are differences between mouse lines expressing ALS-linked mutations and the human genetic cases that these models are attempting to recapitulate. For example, the SOD1<sup>G93a</sup> model shows a rapid rate of disease onset compared to the slow progression observed in human SOD1<sup>G93a</sup> cases [61]. Furthermore, the reported presence of TDP-43 pathology in SOD1<sup>G93a</sup> mice at end stage conflicts with the lack of such pathology in human SOD1 cases [50, 67]. Together with our own data, this demonstrates that caution should be taken when attempting to directly correlate mouse models with human cases involving similar genetic origins. It should be noted that while the SOD1<sup>G93a</sup> mouse may not fully recapitulate human ALS, it nevertheless represents a useful entry point for pursuing mechanistic understanding of the disease. Indeed the use of such mouse models has been essential in many advances in the understanding of ALS pathology [34].

The mechanisms leading to tripartite synapse degeneration remain to be determined. Mislocalisation and aggregation of TDP-43 in the cytoplasm of both neurons and glia is correlated with synapse loss in human ALS [38], and could play a role in designating tripartite synapses for degeneration. However, there is mixed evidence of TDP-43 pathology in the SOD1<sup>G93a</sup> mouse model [43, 67, 69]. Alternatively, the loss of astrocytic EAAT2 in ALS, due to aberrant RNA processing, is known to induce glutamatergic excitotoxicity that could lead to synapse degeneration and ultimately cell death [41, 47, 51, 62]. Our own data supports this hypothesis given that synaptic loss

coincides with EAAT2 loss. Astrocytes also contribute to bi-directional signalling mechanisms that help modulate spinal cord mediated motor behaviour through purinergic signalling [1, 8, 9, 11, 78, 79]. Neuronal-derived glutamate release during locomotor network activity activates astrocytes through mGluR5 receptors, leading to ATP/adenosine release from astrocytes that inhibits neuronal activity through A<sub>1</sub> adenosine receptors [8]. Increased adenosine levels in ALS patients [82], combined with increased expression of the excitatory A<sub>2A</sub> adenosine receptors [57], could imbalance this modulatory mechanism, biasing the system toward excitotoxic actions at the tripartite synapse. Our current study provides evidence that tripartite synapses, and the specific signalling pathways they utilise to regulate neuronal activity, may act as a fulcrum of ALS. These novel findings regarding selective synaptic vulnerability highlight numerous future avenues to explore and provide the framework to investigate the mechanisms and therapeutic potential of ALS tripartite synaptopathy.

### **Acknowledgements**

We would like to acknowledge the following funders: Motor Neurone Disease (MND) Association UK (Miles/Apr18/863-791), the Euan MacDonald Centre and Chief Scientist Office, The European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Programme (695568 SYNNOVATE), Simons Foundation Autism Research Initiative (529085), and the Wellcome Trust (Technology Development grant 202932). We would also like to acknowledge the MRC Edinburgh Brain Bank for the provision of human post-mortem tissue, and kindly thank the patient donors and their families for their valued contribution to this work.

### **Data Availability**

The data used in this manuscript is freely available upon request.

## References

- 650 1. Acton D, Miles GB (2015) Stimulation of Glia Reveals Modulation of Mammalian Spinal Motor Networks by Adenosine. *PLoS One* 10:e0134488. doi: 10.1371/journal.pone.0134488
2. Araque A, Parpura V, Sanzgiri RP, Haydon PG (1999) Tripartite synapses: Glia, the unacknowledged partner. *Trends Neurosci* 22:208–215. doi: 10.1016/S0166-2236(98)01349-6
3. Bączyk M, Alami NO, Delestrée N, Martinot C, Tang L, Commisso B, Bayer D, Doisne N, Frankel W, Manuel M, Roselli F, Zytnicki D (2020) Synaptic restoration by cAMP/PKA drives activity-dependent neuroprotection to motoneurons in ALS. *J Exp Med* 217. doi: 10.1084/JEM.20191734
- 660 4. Bayés A, van de Lagemaat LN, Collins MO, Croning MDR, Whittle IR, Choudhary JS, Grant SGN (2011) Characterization of the proteome, diseases and evolution of the human postsynaptic density. *Nat Neurosci* 14:19–21. doi: 10.1038/nn.2719
5. Boillée S, Yamanaka K, Lobsiger CS, Copeland NG, Jenkins NA, Kassiotis G, Kollias G, Cleveland DW (2006) Onset and progression in inherited ALS determined by motor neurons and microglia. *Science* (80- ) 312:1389–1392. doi: 10.1126/SCIENCE.1123511
6. Broadhead MJ, Bonthron C, Arcinas L, Bez S, Zhu F, Goff F, Nylk J, Dholakia K, Gunn-Moore F, Grant SGN, Miles GB (2020) Nanostructural Diversity of Synapses in the Mammalian Spinal Cord. *Sci Rep* 10:8189. doi: 670 10.1038/s41598-020-64874-9
7. Broadhead MJ, Horrocks MH, Zhu F, Muresan L, Benavides-Piccione R, DeFelipe J, Fricker D, Kopanitsa M V., Duncan RR, Klenerman D, Komiyama NH, Lee SF, Grant SGN (2016) PSD95 nanoclusters are postsynaptic building blocks in hippocampus circuits. *Sci Rep* 6:24626. doi: 10.1038/srep24626
8. Broadhead MJ, Miles GB (2020) Bi-Directional Communication Between Neurons and Astrocytes Modulates Spinal Motor Circuits. *Front Cell Neurosci* 14. doi: 10.3389/fncel.2020.00030
9. Broadhead MJ, Miles GB (2021) A common role for astrocytes in rhythmic behaviours? *Prog Neurobiol* 102052. doi: 10.1016/j.pneurobio.2021.102052
- 680 10. Cacabelos D, Ramírez-Núñez O, Granado-Serrano AB, Torres P, Ayala V, Moiseeva V, Povedano M, Ferrer I, Pamplona R, Portero-Otin M, Boada J (2016) Early and gender-specific differences in spinal cord mitochondrial function and oxidative stress markers in a mouse model of ALS. *Acta Neuropathol Commun* 4:3. doi: 10.1186/s40478-015-0271-6
11. Carlsen EM, Perrier J-F (2014) Purines released from astrocytes inhibit excitatory synaptic transmission in the ventral horn of the spinal cord. *Front Neural Circuits* 8:60. doi: 10.3389/fncir.2014.00060
12. Cho E, Park M (2016) Palmitoylation in Alzheimer's disease and other neurodegenerative diseases. *Pharmacol Res* 111:133–151. doi: 690 10.1016/j.phrs.2016.06.008
13. Cizeron M, Qiu Z, Koniaris B, Gokhale R, Komiyama NH, Fransén E, Grant SGN (2020) A brainwide atlas of synapses across the mouse life span. *Science* (80- ) 369. doi: 10.1126/science.aba3163
14. Comley L, Allodi I, Nichterwitz S, Nizzardo M, Simone C, Corti S, Hedlund E (2015) Motor neurons with differential vulnerability to degeneration show distinct protein signatures in health and ALS. *Neuroscience* 291:216–229. doi:

10.1016/J.NEUROSCIENCE.2015.02.013

- 700 15. Curran OE, Qiu Z, Smith C, Grant SGN (2021) A single-synapse resolution survey of PSD95-positive synapses in twenty human brain regions. *Eur J Neurosci* 54:6864–6881. doi: 10.1111/ejn.14846
16. Derouiche A, Anlauf E, Aumann G, Mühlstädt B, Lavielle M (2002) Anatomical aspects of glia–synapse interaction: the perisynaptic glial sheath consists of a specialized astrocyte compartment. *J Physiol* 96:177–182. doi: 10.1016/S0928-4257(02)00004-9
17. Devlin A-C, Burr K, Borooah S, Foster JD, Cleary EM, Geti I, Vallier L, Shaw CE, Chandran S, Miles GB (2015) Human iPSC-derived motoneurons harbouring TARDBP or C9ORF72 ALS mutations are dysfunctional despite maintaining viability. *Nat Commun* 6:5999. doi: 10.1038/ncomms6999
- 710 18. Eisen A (2021) The Dying Forward Hypothesis of ALS: Tracing Its History. *Brain Sci* 11:1–9. doi: 10.3390/BRAINSCI11030300
19. Feeney SJ, McKelvie PA, Austin L, Bernadette Jean-Francois MJ, Kapsa R, Tombs SM, Byrne E (2001) Presymptomatic motor neuron loss and reactive astrocytosis in the *sod1* mouse model of amyotrophic lateral sclerosis. *Muscle and Nerve* 24:1510–1519. doi: 10.1002/mus.1176
20. Fernández E, Collins MO, Uren RT, Kopanitsa M V, Komiyama NH, Croning MDR, Zografos L, Armstrong JD, Choudhary JS, Grant SGN (2009) Targeted tandem affinity purification of PSD-95 recovers core postsynaptic complexes and schizophrenia susceptibility proteins. *Mol Syst Biol* 5. doi: 10.1038/msb.2009.27
- 720 21. Fischer LR, Culver DG, Tennant P, Davis AA, Wang M, Castellano-Sanchez A, Khan J, Polak MA, Glass JD (2004) Amyotrophic lateral sclerosis is a distal axonopathy: evidence in mice and man. *Exp Neurol* 185:232–40
22. Fletcher E V., Simon CM, Pagiazitis JG, Chalif JI, Vukojicic A, Drobac E, Wang X, Mentis GZ (2017) Reduced sensory synaptic excitation impairs motor neuron function via Kv2.1 in spinal muscular atrophy. *Nat Neurosci* 20:905–916. doi: 10.1038/nn.4561
23. Fogarty MJ (2019) Amyotrophic lateral sclerosis as a synaptopathy. *Neural Regen. Res.* 14:189–192
- 730 24. Fogarty MJ, Klenowski PM, Lee JD, Driberg-Thompson JR, Bartlett SE, Ngo ST, Hilliard MA, Bellingham MC, Noakes PG (2016) Cortical synaptic and dendritic spine abnormalities in a presymptomatic TDP-43 model of amyotrophic lateral sclerosis. *Sci Rep* 6. doi: 10.1038/srep37968
25. Fogarty MJ, Noakes PG, Bellingham MC (2015) Motor cortex layer V pyramidal neurons exhibit dendritic regression, spine loss, and increased synaptic excitation in the presymptomatic hSOD1G93A mouse model of amyotrophic lateral sclerosis. *J Neurosci* 35:643–647. doi: 10.1523/JNEUROSCI.3483-14.2015
- 740 26. Forsberg K, Andersen PM, Marklund SL, Brännström T (2011) Glial nuclear aggregates of superoxide dismutase-1 are regularly present in patients with amyotrophic lateral sclerosis. *Acta Neuropathol* 121:623–634. doi: 10.1007/s00401-011-0805-3
27. Foster JB, Zhao F, Wang X, Xu Z, Lin K, Askwith CC, Hodgetts KJ, Lin C liang G (2018) Pyridazine-derivatives Enhance Structural and Functional Plasticity of Tripartite Synapse Via Activation of Local Translation in Astrocytic Processes. *Neuroscience* 388:224–238. doi: 10.1016/j.neuroscience.2018.07.028

28. Frank RA, Grant SG (2017) Supramolecular organization of NMDA receptors and the postsynaptic density. *Curr Opin Neurobiol* 45:139–147. doi: 10.1016/j.conb.2017.05.019
- 750 29. Frey D, Schneider C, Xu L, Borg J, Spooren W, Caroni P (2000) Early and Selective Loss of Neuromuscular Synapse Subtypes with Low Sprouting Competence in Motoneuron Diseases. *J Neurosci* 20
30. Fukata Y, Dimitrov A, Boncompain G, Vielemeyer O, Perez F, Fukata M (2013) Local palmitoylation cycles define activity-regulated postsynaptic subdomains. *J Cell Biol* 202:145–161. doi: 10.1083/jcb.201302071
31. Gillingwater TH, Wishart TM (2013) Mechanisms underlying synaptic vulnerability and degeneration in neurodegenerative disease. *Neuropathol Appl Neurobiol* 39:320–334. doi: 10.1111/nan.12014
- 760 32. Grant SGN (2019) Synapse diversity and synaptome architecture in human genetic disorders. *Hum Mol Genet* 28:R219–R225. doi: 10.1093/hmg/ddz178
33. Grant SGN, Fransén E (2020) The Synapse Diversity Dilemma: Molecular Heterogeneity Confounds Studies of Synapse Function. *Front Synaptic Neurosci* 12. doi: 10.3389/fnsyn.2020.590403
34. Gurney ME, Fleck TJ, Himes CS, Hall ED (1998) Riluzole preserves motor function in a transgenic model of familial amyotrophic lateral sclerosis. *Neurology* 50:62–66. doi: 10.1212/WNL.50.1.62
35. Gurney ME, Pu H, Chiu AY, Dal Canto MC, Polchow CY, Alexander DD, Caliendo J, Hentati A, Kwon YW, Deng HX, Chen W, Zhai P, Sufit RL, Siddique T (1994) Motor neuron degeneration in mice that express a human Cu,Zn superoxide dismutase mutation. *Science* (80- ) 264:1772–1775. doi: 10.1126/science.8209258
- 770 36. Heiman-Patterson TD, Deitch JS, Blankenhorn EP, Erwin KL, Perreault MJ, Alexander BK, Byers N, Toman I, Alexander GM (2005) Background and gender effects on survival in the TgN(SOD1-G93A)<sup>1Gur</sup> mouse model of ALS. *J Neurol Sci* 236:1–7. doi: 10.1016/j.jns.2005.02.006
37. Heller JP, Rusakov DA (2017) The Nanoworld of the Tripartite Synapse: Insights from Super-Resolution Microscopy. *Front Cell Neurosci* 11:374. doi: 10.3389/fncel.2017.00374
- 780 38. Henstridge CM, Sideris DI, Carroll E, Rotariu S, Salomonsson S, Tzioras M, McKenzie CA, Smith C, von Arnim CAF, Ludolph AC, Lulé D, Leighton D, Warner J, Cleary E, Newton J, Swingler R, Chandran S, Gillingwater TH, Abrahams S, Spires-Jones TL (2018) Synapse loss in the prefrontal cortex is associated with cognitive decline in amyotrophic lateral sclerosis. *Acta Neuropathol* 135:213–226. doi: 10.1007/s00401-017-1797-4
39. Henstridge CM, Tzioras M, Paolicelli RC (2019) Glial Contribution to Excitatory and Inhibitory Synapse Loss in Neurodegeneration. *Front Cell Neurosci* 0:63. doi: 10.3389/FNCEL.2019.00063
- 790 40. Herron LR, Miles GB (2012) Gender-specific perturbations in modulatory inputs to motoneurons in a mouse model of amyotrophic lateral sclerosis. *Neuroscience* 226:313–323. doi: 10.1016/j.neuroscience.2012.09.031
41. Howland DS, Liu J, She Y, Goad B, Maragakis NJ, Kim B, Erickson J, Kulik J, DeVito L, Psaltis G, DeGennaro LJ, Cleveland DW, Rothstein JD (2002) Focal loss of the glutamate transporter EAAT2 in a transgenic rat model of SOD1 mutant-mediated amyotrophic lateral sclerosis (ALS). *Proc Natl Acad Sci U S A* 99:1604–1609. doi: 10.1073/pnas.032539299
42. Husi H, Grant SGN (2001) Isolation of 2000-kDa complexes of N-methyl-D-

- aspartate receptor and postsynaptic density 95 from mouse brain. *J Neurochem* 2:2006.0023. doi: 10.1038/msb4100041
- 800 43. Jeon GS, Shim YM, Lee DY, Kim JS, Kang MJ, Ahn SH, Shin JY, Geum D, Hong YH, Sung JJ (2019) Pathological Modification of TDP-43 in Amyotrophic Lateral Sclerosis with SOD1 Mutations. *Mol Neurobiol* 56:2007–2021. doi: 10.1007/s12035-018-1218-2
44. Jiang T, Handley E, Brizuela M, Dawkins E, Lewis KEA, Clark RM, Dickson TC, Blizzard CA (2019) Amyotrophic lateral sclerosis mutant TDP-43 may cause synaptic dysfunction through altered dendritic spine function. *DMM Dis Model Mech* 12. doi: 10.1242/dmm.038109
45. Ke YD, van Hummel A, Stevens CH, Gladbach A, Ippati S, Bi M, Lee WS, Krüger S, van der Hoven J, Volkerling A, Bongers A, Halliday G, Haass NK, Kiernan M, Delerue F, Ittner LM (2015) Short-term suppression of A315T mutant human TDP-43 expression improves functional deficits in a novel inducible transgenic mouse model of FTLTDP and ALS. *Acta Neuropathol* 130:661–678. doi: 10.1007/s00401-015-1486-0
- 810 46. Lavalie M, Aumann G, Anlauf E, Pröls F, Arpin M, Derouiche A (2011) Structural plasticity of perisynaptic astrocyte processes involves ezrin and metabotropic glutamate receptors. *Proc Natl Acad Sci U S A* 108:12915–9. doi: 10.1073/pnas.1100957108
47. Lin CLG, Bristol LA, Jin L, Dykes-Hoberg M, Crawford T, Clawson L, Rothstein JD (1998) Aberrant RNA processing in a neurodegenerative disease: The cause for absent EAAT2, a glutamate transporter, in amyotrophic lateral sclerosis. *Neuron* 20:589–602. doi: 10.1016/S0896-6273(00)80997-6
- 820 48. Liu Y, Pattamatta A, Zu T, Reid T, Bardhi O, Borchelt DRR, Yachnis ATT, Ranum LPWPW (2016) C9orf72 BAC Mouse Model with Motor Deficits and Neurodegenerative Features of ALS/FTD. *Neuron* 90:521–534. doi: 10.1016/j.neuron.2016.04.005
49. MacGillavry HD, Song Y, Raghavachari S, Blanpied TA (2013) Nanoscale Scaffolding Domains within the Postsynaptic Density Concentrate Synaptic AMPA Receptors. *Neuron* 78:615–622. doi: 10.1016/j.neuron.2013.03.009
- 830 50. Mackenzie IRA, Bigio EH, Ince PG, Geser F, Neumann M, Cairns NJ, Kwong LK, Forman MS, Ravits J, Stewart H, Eisen A, McClusky L, Kretzschmar HA, Monoranu CM, Highley JR, Kirby J, Siddique T, Shaw PJ, Lee VMY, Trojanowski JQ (2007) Pathological TDP-43 distinguishes sporadic amyotrophic lateral sclerosis from amyotrophic lateral sclerosis with SOD1 mutations. *Ann Neurol* 61:427–434. doi: 10.1002/ana.21147
51. Maragakis NJ, Rothstein JD (2006) Mechanisms of Disease: astrocytes in neurodegenerative disease. *Nat Clin Pract Neurol* 2:679–689. doi: 10.1038/ncpneuro0355
52. McCombe PA, Henderson RD (2010) Effects of gender in amyotrophic lateral sclerosis. *Gend Med* 7:557–570. doi: 10.1016/j.genm.2010.11.010
- 840 53. Mead RJ, Bennett EJ, Kennerley AJ, Sharp P, Sunyach C, Kasher P, Berwick J, Pettmann B, Battaglia G, Azzouz M, Grierson A, Shaw PJ (2011) Optimised and rapid pre-clinical screening in the SOD1 G93A transgenic mouse model of amyotrophic lateral sclerosis (ALS). *PLoS One* 6. doi: 10.1371/journal.pone.0023244
54. Molofsky A V., Kelley KW, Tsai HH, Redmond SA, Chang SM, Madireddy L, Chan JR, Baranzini SE, Ullian EM, Rowitch DH (2014) Astrocyte-encoded positional cues maintain sensorimotor circuit integrity. *Nature* 509:189. doi:

10.1038/NATURE13161

- 850 55. Mordes DA, Morrison BM, Ament XH, Cantrell C, Mok J, Eggan P, Xue C, Wang JY, Eggan K, Rothstein JD (2020) Absence of Survival and Motor Deficits in 500 Repeat C9ORF72 BAC Mice. *Neuron* 108:775-783.e4. doi: 10.1016/J.NEURON.2020.08.009
56. Nair D, Hosy E, Petersen JD, Constals A, Giannone G, Choquet D, Sibarita J-BJ-B (2013) Super-Resolution Imaging Reveals That AMPA Receptors Inside Synapses Are Dynamically Organized in Nanodomains Regulated by PSD95. *J Neurosci* 33:13204–13224. doi: 10.1523/JNEUROSCI.2381-12.2013
57. Ng S kah, Higashimori H, Tolman M, Yang Y (2015) Suppression of adenosine 2a receptor (A2aR)-mediated adenosine signaling improves disease phenotypes in a mouse model of amyotrophic lateral sclerosis. *Exp Neurol* 267:115–122. doi: 10.1016/j.expneurol.2015.03.004
- 860 58. Nguyen L, Laboissonniere LA, Guo S, Pilotto F, Scheidegger O, Oestmann A, Hammond JW, Li H, Hyysalo A, Peltola R, Pattamatta A, Zu T, Voutilainen MH, Gelbard HA, Saxena S, Ranum LPW (2020) Survival and Motor Phenotypes in FVB C9-500 ALS/FTD BAC Transgenic Mice Reproduced by Multiple Labs. *Neuron* 108:784-796.e3. doi: 10.1016/j.neuron.2020.09.009
59. Perea G, Navarrete M, Araque A (2009) Tripartite synapses: astrocytes process and control synaptic information. *Trends Neurosci* 32:421–431. doi: 10.1016/j.tins.2009.05.001
- 870 60. Perkins EM, Burr K, Banerjee P, Mehta AR, Dando O, Selvaraj BT, Suminaite D, Nanda J, Henstridge CM, Gillingwater TH, Hardingham GE, Wyllie DJA, Chandran S, Livesey MR (2021) Altered network properties in C9ORF72 repeat expansion cortical neurons are due to synaptic dysfunction. *Mol Neurodegener* 16:1–16. doi: 10.1186/s13024-021-00433-8
61. Picher-Martel V, Valdmanis PN, Gould P V., Julien JP, Dupré N (2016) From animal models to human disease: a genetic approach for personalized medicine in ALS. *Acta Neuropathol Commun* 4. doi: 10.1186/S40478-016-0340-5
62. Rothstein JD, Dykes-Hoberg M, Pardo CA, Bristol LA, Jin L, Kuncl RW, Kanai Y, Hediger MA, Wang Y, Schielke JP, Welty DF (1996) Knockout of glutamate transporters reveals a major role for astroglial transport in excitotoxicity and clearance of glutamate. *Neuron* 16:675–686. doi: 10.1016/S0896-6273(00)80086-0
- 880 63. Sasaki S, Iwata M (1995) Synaptic loss in the proximal axon of anterior horn neurons in motor neuron disease. *Acta Neuropathol* 90:170–5
64. Sasaki S, Iwata M (1996) Synaptic loss in anterior horn neurons in lower motor neuron disease. *Acta Neuropathol* 91:416–21
65. Sasaki S, Komori T, Iwata M (2000) Excitatory amino acid transporter 1 and 2 immunoreactivity in the spinal cord in amyotrophic lateral sclerosis. *Acta Neuropathol* 100:138–44
- 890 66. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A (2012) Fiji: an open-source platform for biological-image analysis. *Nat Methods* 9:676–682. doi: 10.1038/NMETH.2019
67. Shan X, Vocadlo D, Krieger C (2009) Mislocalization of TDP-43 in the G93A mutant SOD1 transgenic mouse model of ALS. *Neurosci Lett* 458:70–74. doi: 10.1016/j.neulet.2009.04.031



68. Starr A, Sattler R (2018) Synaptic dysfunction and altered excitability in C9ORF72 ALS/FTD. *Brain Res.* 1693:98–108
- 900 69. Tan CF, Eguchi H, Tagawa A, Onodera O, Iwasaki T, Tsujino A, Nishizawa M, Kakita A, Takahashi H (2007) TDP-43 immunoreactivity in neuronal inclusions in familial amyotrophic lateral sclerosis with or without SOD1 gene mutation. *Acta Neuropathol* 113:535–542. doi: 10.1007/s00401-007-0206-9
70. Tang A-H, Chen H, Li TP, Metzbower SR, MacGillavry HD, Blanpied TA (2016) A trans-synaptic nanocolumn aligns neurotransmitter release to receptors. *Nature* 536:210–214. doi: 10.1038/nature19058
71. Tong J, Huang C, Bi F, Wu Q, Huang B, Liu X, Li F, Zhou H, Xia X-G (2013) Expression of ALS-linked TDP-43 mutant in astrocytes causes non-cell-autonomous motor neuron death in rats. *EMBO J* 32:1917–1926. doi: 10.1038/emboj.2013.122
- 910 72. Tremblay E, Martineau É, Robitaille R (2017) Opposite Synaptic Alterations at the Neuromuscular Junction in an ALS Mouse Model: When Motor Units Matter. *J Neurosci* 37:8901. doi: 10.1523/JNEUROSCI.3090-16.2017
73. Tsai HH, Li H, Fuentealba LC, Molofsky A V., Taveira-Marques R, Zhuang H, Tenney A, Murnen AT, Fancy SPJ, Merkle F, Kessler N, Alvarez-Buylla A, Richardson WD, Rowitch DH (2012) Regional Astrocyte Allocation Regulates CNS Synaptogenesis and Repair. *Science* 337:358. doi: 10.1126/SCIENCE.1222381
- 920 74. Valori CF, Brambilla L, Martorana F, Rossi D (2014) The multifaceted role of glial cells in amyotrophic lateral sclerosis. *Cell Mol Life Sci* 71:287–297. doi: 10.1007/s00018-013-1429-7
75. Vaughan SK, Sutherland NM, Zhang S, Hatzipetros T, Vieira F, Valdez G (2018) The ALS-inducing factors, TDP43A315T and SOD1G93A, directly affect and sensitize sensory neurons to stress. *Sci Rep* 8. doi: 10.1038/s41598-018-34510-8
76. Watson C, Paxinos G, Kayalioglu G, Christopher & Dana Reeve Foundation. (2009) *The spinal cord : a Christopher and Dana Reeve Foundation text and atlas.* Elsevier/Academic Press
- 930 77. Wishart TM, Parson SH, Gillingwater TH (2006) Synaptic Vulnerability in Neurodegenerative Disease. *J Neuropathol Exp Neurol* 65:733–739. doi: 10.1097/01.jnen.0000228202.35163.c4
78. Witts EC, Nascimento F, Miles GB (2015) Adenosine-mediated modulation of ventral horn interneurons and spinal motoneurons in neonatal mice. *J Neurophysiol* 114:2305–2315. doi: 10.1152/jn.00574.2014
79. Witts EC, Panetta KM, Miles GB (2012) Glial-derived adenosine modulates spinal motor networks in mice. *J Neurophysiol* 107:1925–34. doi: 10.1152/jn.00513.2011
- 940 80. Yamanaka K, Chun SJ, Boillee S, Fujimori-Tonou N, Yamashita H, Gutmann DH, Takahashi R, Misawa H, Cleveland DW (2008) Astrocytes as determinants of disease progression in inherited amyotrophic lateral sclerosis. *Nat Neurosci* 11:251–3. doi: 10.1038/nn2047
81. Yamanaka K, Komine O (2018) The multi-dimensional roles of astrocytes in ALS. *Neurosci. Res.* 126:31–38
82. Yoshida Y, Une F, Utatsu Y, Nomoto M, Furukawa Y, Maruyama Y, Machigashira N, Matsuzaki T, Osame M (1999) Adenosine and neopterin levels in cerebrospinal fluid of patients with neurological disorders. *Intern Med* 38:133–139. doi: 10.2169/internalmedicine.38.133

- 950 83. Zaręba-Koziół M, Figiel I, Bartkowiak-Kaczmarek A, Włodarczyk J (2018) Insights into protein S-palmitoylation in synaptic plasticity and neurological disorders: Potential and limitations of methods for detection and analysis. *Front Mol Neurosci* 11:175. doi: 10.3389/fnmol.2018.00175
84. Zarei S, Carr K, Reiley L, Diaz K, Guerra O, Altamirano PF, Pagani W, Lodin D, Orozco G, China A (2015) A comprehensive review of amyotrophic lateral sclerosis. *Surg. Neurol. Int.* 6
85. Zhao C, Devlin AC, Chouhan AK, Selvaraj BT, Stavrou M, Burr K, Brivio V, He X, Mehta AR, Story D, Shaw CE, Dando O, Hardingham GE, Miles GB, Chandran S (2020) Mutant C9orf72 human iPSC-derived astrocytes cause non-cell autonomous motor neuron pathophysiology. *Glia* 68:1046–1064. doi: 10.1002/glia.23761
- 960 86. Zhu F, Cizeron M, Qiu Z, Benavides-Piccione R, Kopanitsa M V, Skene NG, Koniaris B, DeFelipe J, Fransén E, Komiyama NH, Grant SGN (2018) Architecture of the Mouse Brain Synaptome. *Neuron*. doi: 10.1016/j.neuron.2018.07.007
87. Van Zundert B, Peuscher MH, Hynynen M, Chen A, Neve RL, Brown RH, Constantine-Paton M, Bellingham MC (2008) Neonatal neuronal circuitry shows hyperexcitable disturbance in a mouse model of the adult-onset neurodegenerative disease amyotrophic lateral sclerosis. *J Neurosci* 28:10864–10874. doi: 10.1523/JNEUROSCI.1340-08.2008

970

## Figure Legends And Tables

**Figure 1. Mapping excitatory synapses in the spinal cord revealed early symptomatic stage synaptic changes in SOD1<sup>G93a</sup> mice.** **a.** High-resolution mapping of PSD95-eGFP was performed to analyse synaptic changes in SOD1<sup>G93a</sup> mice at different time points of the disease. **b.** Example images of postsynaptic puncta in control and SOD1<sup>G93a</sup> mice at different time points of disease. **c.** Heat maps of PSD density in male and female SOD1<sup>G93a</sup> mice compared to their respective controls at different ages. Sample sizes denoted as n = number of controls / number of SOD1<sup>G93a</sup>'s. **d.** Heat maps of PSD size in male and female SOD1<sup>G93a</sup> mice compared to their respective controls at different ages. **e.** Chart plotting PSD density in all 8 laminae of 16-week-old controls, WTSOD1 controls and SOD1<sup>G93a</sup> males. Sample sizes denoted as n = number of WTSOD1's / number of controls / number of SOD1<sup>G93a</sup>'s. **f.** Chart plotting PSD size in all 8 laminae of 16-week-old controls, WTSOD1 controls and SOD1<sup>G93a</sup> males. **g.** Chart plotting PSD density in all 8 laminae of 16-week-old controls, WTSOD1 controls and SOD1<sup>G93a</sup> females. **h.** Chart plotting PSD size in all 8 laminae of 16-week-old controls, WTSOD1 controls and SOD1<sup>G93a</sup> females.

**Figure 2. Nanoscopic Changes in Substructural Organisation of PSD95 in ALS Synapses.** **a.** Diagram of synapse subtypes, defined by the number of PSD95 nanoclusters (NCs) per PSD. PSDs can be visualised using confocal microscopy while NCs are resolved from correlative g-STED images. **b.** Spinal cord tissue from control and SOD1<sup>G93a</sup> mice were imaged using confocal and g-STED microscopy to visualise PSDs and NCs respectively. **c.** Graph showing the size of the PSD95 NCs was not different between controls and SOD1<sup>G93a</sup> mice. **d.** Graph showing the average number of NCs per PSD was significantly reduced in SOD1<sup>G93a</sup> mice. **e.** Graph showing the number of large 3+ NC-PSDs are specifically reduced in SOD1<sup>G93a</sup> mice compared to controls. **f.** High-resolution imaging of synapses (PSD95 and VGLUT2) and the palmitoylation enzyme, zDHHC2, was performed to investigate whether changes in zDHHC2 expression might explain the reduced number of PSD95 NCs at synapses. **g.** Graph showing no difference in zDHHC2 puncta density between control and SOD1<sup>G93a</sup> mice. **h.** Graph showing reduced number of PSDs expressing zDHHC2 in SOD1<sup>G93a</sup> mice.

**Figure 3. Mapping Tripartite Synapses in the SOD1<sup>G93a</sup> Mouse Spinal Cord. a.**

Example of high-resolution spinal cord mapping of tripartite synapses from 16-week-old male control and SOD1<sup>G93a</sup> mice using PSD95-eGFP, VGLUT2 and p-Ezrin to label the PSDs, presynaptic terminals and PAPs respectively. Cropped images display individual synaptic and tripartite synaptic structures. **b.** Example of high-resolution spinal cord mapping of tripartite synapses from 16-week-old male control and SOD1<sup>G93a</sup> mice using PSD95-eGFP, VGLUT2 and EAAT2 to label the PSDs, presynaptic terminals and PAPs respectively. Cropped images display individual synaptic and tripartite synaptic structures. **c.** Colour coded heat maps denote the difference in synapse or tripartite synapse density observed in SOD1<sup>G93a</sup> mice as a percentage of the respective controls. Bar charts display the mean synapse density from mice, averaged across all anatomical subregions for simplicity. From p-Ezrin labelling, no changes in non-tripartite or tripartite synapses were observed in 8 week SOD1<sup>G93a</sup> mice. **d.** From EAAT2 labelling, no changes in non-tripartite synapses were observed, but a significant increase in tripartite synapse numbers were observed in 8 week SOD1<sup>G93a</sup> mice compared to control. **e.** From p-Ezrin labelling, no changes in non-tripartite or tripartite synapses were observed in 12 week SOD1<sup>G93a</sup> mice. **f.** From EAAT2 labelling, no changes in non-tripartite synapses were observed, but a significant decrease in tripartite synapse numbers were observed in 12 week SOD1<sup>G93a</sup> mice compared to control. **g.** From p-Ezrin labelling, no changes in non-tripartite synapses were observed, but a significant decrease in tripartite synapse numbers were observed in 16 week SOD1<sup>G93a</sup> mice compared to control. **h.** From EAAT2 labelling, no changes in non-tripartite synapses were observed, but a significant decrease in tripartite synapse numbers were observed in 16 week SOD1<sup>G93a</sup> mice compared to control.

**Figure 4. Tripartite Synapses are selectively lost in the cervical spinal cord of human ALS cases. a.**

Diagram of human cervical spinal cord and location of image sampling. **b.** Example images of synapses (PSD95 – green) and astrocytic processes (p-Ezrin – magenta) in human spinal cord tissue. **c.** Graph showing the total PSD density (PSD95 puncta) in controls, all ALS cases, and then SOD1 and C9orf72 cases separated. **d.** Graph showing synapse density in patient groups when synapses are

separated based on whether they are non-tripartite synapses or tripartite synapses, as determined by the association of PSD95 with p-Ezrin.

1040

Table 1. Antibodies For Mouse Spinal Cord Tissue Labelling					
Primary Antibody Target	Species	Company	Code	Dilution	Secondary Antibody/Conjugation
VGLUT2	Mouse	Abcam	Ab79157	1:500	Anti-Ms Alexa Fluor 555
p-Ezrin	Rabbit	ProteinTech	26056-1-AP	1:500	Anti-Rb Alexa Fluor 647 / Dye-Conjugated Alexa Fluor 647
EAAT2	Rabbit	Bioss	BS1751R	1:500	Anti-Rb Alexa Fluor 647 / Dye-Conjugated Alexa Fluor 594
zDHHC2	Rabbit	Biorbyt	Orb385443	1:100	Anti-Rb Alexa Fluor 647
Glutamine Synthetase	Mouse	ProteinTech	66323-1-Ig	1:500	Anti-Ms FITC
ChAT	Chicken	Abcam	Ab34419	1:250	Anti-Ch Alexa Fluor 647
GFAP	Rabbit	DAKO	Z0334	1:1000	Dye-conjugated Alexa Fluor 594

Table 2. Antibodies For Human Spinal Cord Tissue Labelling					
Primary Antibody Target	Species	Company	Code	Dilution	Secondary Antibody/Conjugation
PSD95	Guinea Pig	Synaptic Systems	124-014	1:500	Gt Anti-Gp Alexa Fluor 647
p-Ezrin	Rabbit	ProteinTech	26056-1-AP	1:250	Gt Anti-Rb Cy3
VGLUT2	Mouse	Abcam	Ab79157	1:500	Dk Anti-Ms Alexa Fluor 555