

2021

Intraventricular Septation in the Context of Dilated Cardiomyopathy Associated With TTN Mutation

Karl R. Khandalavala
University of Nebraska Medical Center

Jean A. Ballweg
University of Nebraska Medical Center

Stanley J. Radio
University of Nebraska Medical Center, sradio@unmc.edu

Jeffrey A. Robinson
University of Nebraska Medical Center, jeffrey.robinson@unmc.edu

Tell us how you used this information in this [short survey](#).

Follow this and additional works at: https://digitalcommons.unmc.edu/com_peds_articles



Part of the [Pediatrics Commons](#)

Recommended Citation

Khandalavala, Karl R.; Ballweg, Jean A.; Radio, Stanley J.; and Robinson, Jeffrey A., "Intraventricular Septation in the Context of Dilated Cardiomyopathy Associated With TTN Mutation" (2021). *Journal Articles: Pediatrics*. 35.

https://digitalcommons.unmc.edu/com_peds_articles/35

This Article is brought to you for free and open access by the Pediatrics at DigitalCommons@UNMC. It has been accepted for inclusion in Journal Articles: Pediatrics by an authorized administrator of DigitalCommons@UNMC. For more information, please contact digitalcommons@unmc.edu.

IMAGING VIGNETTE

ADVANCED

CLINICAL VIGNETTE

Intraventricular Septation in the Context of Dilated Cardiomyopathy Associated With *TTN* Mutation



Karl R. Khandalavala, BS,^a Jean A. Ballweg, MD,^{b,c} Stanley J. Radio, MD,^d Jeffrey A. Robinson, MD^{b,c}

ABSTRACT

A 6-month-old infant boy presented with symptomatic heart failure. Dilated cardiomyopathy was found in association with a mutation in *TTN*. Structural heart disease included novel septation of the left ventricle with a fenestrated membrane resulting from aberrant congenital mitral valve apparatus formation. (**Level of Difficulty: Advanced.**) (J Am Coll Cardiol Case Rep 2021;3:1674-1676) © 2021 Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

CASE PRESENTATION

A previously healthy 6-month-old full-term infant boy was brought to an emergency department with a 2-day history of vomiting, pallor, and poor feeding. Upon transfer to a tertiary pediatric referral center, his blood pressure (BP) was 85/39 mmHg, pulse 145 beats/min, respirations 36/min, weight 6.4 kg (1st percentile), and length 69 cm (50th percentile). Significant physical examination findings included a gallop rhythm, hepatomegaly, and low-set ears with fatty and thickened helices. A chest x-ray demonstrated massive cardiomegaly with increased pulmonary vascular markings (Supplemental Figure 1). The presenting electrocardiogram demonstrated normal sinus rhythm, nonspecific interventricular block, right ventricular hypertrophy, and T-wave inversion in the inferior leads (Supplemental Figure 2). An echocardiogram demonstrated dilated cardiomyopathy with severe left ventricular dilation, systolic dysfunction (Supplemental Tables 1 to 3), and the unique finding of a septating band within the left ventricle (LV) (Figures 1A and 1B). This was further characterized as a multiply fenestrated membrane. Multiple left ventricular thrombi were noted, both within the true left ventricular cavity and within the false chamber. Genetic studies indicated a titin (*TTN*) mutation-associated dilated cardiomyopathy (1G) (Supplemental Table 4).

Despite maximal pharmacologic therapy for systolic failure, left ventricular ejection fraction remained depressed, and orthotopic cardiac transplantation was performed. Gross pathologic examination of the native explanted heart revealed severe biventricular dilation and hypertrophy, as well as interstitial fibrosis with myocytolysis. Within the LV, there was endocardial fibroelastosis with a “sheetlike” partition consisting of partially fenestrated fibromyxomatous tissue that was intertwined with and extended from the chordae

From the ^aCollege of Medicine, University of Nebraska Medical Center, Omaha, Nebraska, USA; ^bDivision of Cardiology, Department of Pediatrics, College of Medicine, University of Nebraska Medical Center, Omaha, Nebraska, USA; ^cThe Dr. C. C. and Mabel L. Criss Heart Center, Children’s Hospital and Medical Center, Omaha, Nebraska, USA; and the ^dDepartment of Pathology, College of Medicine, University of Nebraska Medical Center, Omaha, Nebraska, USA.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors’ institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

Manuscript received June 1, 2021; revised manuscript received August 17, 2021, accepted September 2, 2021.

tendineae of the mitral valve to an area of noncompaction-like change of the left ventricular endomyocardium (Figure 1C). Histologic examination confirmed endocardial fibroelastosis (EFE) (Figures 1D and 1E). In addition, features of a dilated cardiomyopathy, including subendocardial myocytolysis, interstitial fibrosis, and T cell (CD3+) lymphocyte infiltration were observed. The septating band itself consisted of valve tissue components, including fibromyxomatous stroma and minimal collagen with decreased amounts of elastin relative to normal chordae tendineae or valve leaflets.

ABBREVIATIONS AND ACRONYMS

BP = blood pressure

EFE = endocardial
fibroelastosis


LV = left ventricle

TTN = titin gene

DISCUSSION

Within current published reports, this case is not consistent with typical classifications of intraventricular findings such as intramyocardial delamination, congenital diverticula, congenital aneurysms, or EFE. Failure

FIGURE 1 Novel Intraventricular Septation in Multiple Imaging Modalities



(A) 2-dimensional (2D) image by transthoracic echocardiogram in the 4-chamber view, demonstrating severe dilation of the left ventricle with the mitral valve apparatus and intraventricular septation seen. (B) 2D image by transthoracic echocardiogram in the subcostal coronal view, demonstrating the fenestrated left ventricular septation within the severely dilated left ventricle. (C) Cross-section on gross examination of the explanted heart. Corresponding plane of dissection demonstrating left ventricle (LV), right ventricle (RV), and ventricular septum (VS) with endocardial fibroelastosis (EFE) (black arrows) and "apron" (blue star). Apical mural thrombus is also present (yellow arrow). (D) Microscopic section of the ventricular myocardium on Verhoeff-Van Gieson stain at 10× magnification, demonstrating EFE of VS highlighted as black region (yellow arrow). (E) Microscopic cross-section of the delaminated ventricular apron on Movat stain at 20× magnification, demonstrating thin strands of elastin (black) with prominent sea-green to blue myxomatous stroma.

of myocardial delamination has been implicated in congenital valvular conditions such as Ebstein's anomaly but does not result in defects containing multiple myocardial layers (1). Congenital LV diverticula and aneurysms involve dilations of the left ventricle wall that contain all three layers of the ventricular wall contracting synchronously within the heart, which was not observed in this case (2). The pathologic specimen contained features of EFE, which is generally limited to the endocardial layer but was observed here as a separate intracavity sheet or septation (3).

The case presentation, echocardiographic imaging, and gross pathologic changes fail to meet the diagnostic criteria for any known condition. The presence of EFE raises an interesting question about the functional consequences of this finding within the dilated cardiomyopathy diagnosis. Early identification of significant anatomic abnormalities, as observed in this case, may have an impact on clinical management. These patients may be at greater risk for medical treatment failure and require early cardiac transplantation.

FUNDING SUPPORT AND AUTHOR DISCLOSURES

The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

ADDRESS FOR CORRESPONDENCE: Dr Jeffrey A. Robinson, Pediatric Cardiology, Dr. C.C. and Mabel L. Criss Heart Center, Children's Hospital & Medical Center, 8200 Dodge Street, Omaha, Nebraska 68114-4113, USA. E-mail: jrobinson@ChildrensOmaha.org.

REFERENCES

1. Osiro S, Tiwari KJ, Mathenge N, Rodriguez JR, Tubbs RS, Loukas M. When lithium hurts: a look at Ebstein anomaly. *Cardiol Rev*. 2013;21:257-263.
2. Hamaoka K, Onaka M, Tanaka T, Onouchi Z. Congenital ventricular aneurysm and diverticulum in children. *Pediatr Cardiol*. 1987;8:169-175.
3. Weixler V, Hammer PE, Marx GR, Emani SM, Del Nido PJ, Friehs I. Flow disturbances and progression of endocardial fibroelastosis: a case report. *Cardiovasc Pathol*. 2019;42:1-3.

KEY WORDS congenital heart disease, dilated cardiomyopathy, endocardial

fibroelastosis, intraventricular septation, titin gene

APPENDIX For supplemental figures and tables, please see the online version of this paper.