

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



ISSN: 0015-5659

e-ISSN: 1644-3284

Mesenchymal stem cell-derived exosomes are beneficial to suppressing inflammation and promoting autophagy in intervertebral disc degeneration

Authors: Baicheng Yang, Xinming Yang

DOI: 10.5603/FM.a2023.0021

Article type: Original article

Submitted: 2023-01-04

Accepted: 2023-02-14

Published online: 2023-03-22

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited. Articles in "Folia Morphologica" are listed in PubMed.

Mesenchymal stem cell-derived exosomes are beneficial to suppressing inflammation and promoting autophagy in intervertebral disc degeneration

Baicheng Yang et al., MSC-exosomes in intervertebral disc degeneration

Baicheng Yang^{1,#}, Xinming Yang^{2,#}

¹Hebei North University, Zhangjiakou, Hebei, China

²Department of Orthopedics, the First Affiliated Hospital of Hebei North University, Zhangjiakou, Hebei, China

Address for correspondence: Xinming Yang, Department of Orthopedics, the First Affiliated Hospital of Hebei North University, Zhangjiakou 075000, Hebei, China, e-mail: yxm11200@126.com

#Contributed equally to this work

ABSTRACT

Intervertebral disc degenerative diseases (IDDD) is one the main causes of lumbago, and its main pathological mechanism is intervertebral disc degeneration (IDD). In previous reports, mesenchymal stem cell (MSC)-exosomes can slow down or even reverse degenerated nucleus pulposus (NP) cells in IDD. Thus, we attempted to clarify specific role of MSC-exosomes underlying IDD progression. In the present study, the harvested particles were identified as MSC-exosomes. MSC-exosomes facilitated activation of autophagy pathway in AGE-treated NP cells. MSC-exosomes repressed inflammatory response in AGE-treated NP cells. Autophagy pathway activation enhanced inflammatory response in AGE-stimulated NP cells. MSC-exosomes facilitated autophagy pathway activation and repressed inflammation in IDD rats. Autophagy inhibition exerted a protective role against inflammatory response in IDD rats. In conclusion, MSC-exosomes represses inflammation via activating autophagy pathway, which provides a potential novel insight for seeking therapeutic plans of IDD.

Key words: intervertebral disc degeneration, mesenchymal stem cell, exosomes, autophagy, inflammation

INTRODUCTION

Lumbago is the most common spinal surgery symptom with high morbidity and disability rate, seriously affecting human health and quality of life and also bringing a heavy burden to individuals, families and the whole society [1]. One of the main causes of lumbago is intervertebral disc degenerative diseases (IDDD), of which main pathological mechanism is intervertebral disc degeneration (IDD). Since most patients with early IDD have no obvious symptoms, they also miss the opportunity for early intervention and therapy [2, 3]. Currently, treatment methods for diseases caused by IDD include conservative therapy and surgery. Both of them can relieve clinical symptoms to a certain extent, but the long-term effect is often unfavorable [4]. Thus, it is of great clinical significance to figure out molecular biology to prevent or even reverse IDD with less side damage to patients.

The intervertebral disc (IVD) is the largest avascular structure in the human body [5]. The changes in the structure of IVD are caused by a combination of factors such as cellular aging, nutritional disorders, stress factors, inflammatory mediators, reduction of nucleus pulposus cells, abnormal metabolism of extracellular matrix (ECM), etc. [6]. To address IDD and related degenerative diseases, multiple studies have focused on maintaining amount of nucleus pulposus (NP) cells [7, 8]. Thus, bone marrow mesenchymal stem cells (BMSCs), which have the advantages of easy access, multi-directional differentiation, low immunogenicity, and mature acquisition and identification, have been widely researched [9]. The inflammatory response mediated by pro-inflammatory cytokines can directly cause the loss of water in IVD and exacerbate IDD process [10]; MSCs are capable of migrating to damaged tissues and releasing cytokines, inflammatory regulators, extracellular matrix components, and antimicrobial proteins in this area to create a suitable microenvironment for tissue repair [11]. With the deepening of research, it has been demonstrated that biological information can be transmitted between non-contact cells through paracrine or in the form of exosomes [12]. MSC-exosomes can upregulate autophagy marker proteins

LC3 and Beclin-1 and facilitate autophagosome formation [13]. Rat MSCs can produce positive trophic effects on degenerated NP cells by secreting exosomes, and NP degeneration degree can still be improved after non-contact co-culture of BMSCs and NP cells [14]. Moreover, MSC-exosomes are capable of attenuating NP cell apoptosis induced by proinflammatory cytokines [15]. These reports indicate that MSC-exosomes can slow down or even reverse degenerated NP cells in IDD. Thus, we hypothesized that MSC-exosomes delayed or even reversed IDD via autophagy activation and inflammation inhibition. Herein, we clarified potential role of MSC-exosomes in IDD by establishing animal models and AGE-treated cell models, which may provide a novel insight for targeted therapy of IDD.

MATERIALS AND METHODS

Isolation and identification of MSCs

Human bone marrow specimens were collected from healthy donors. The institutional review board of our hospital approved this research with informed consent signed by patients. MSCs from bone marrow got separation using density gradient centrifugation and adherence to tissue culture plastic. MSCs received amplification in Dulbecco's modified Eagle medium (DMEM) supplemented with 15% fetal bovine serum (FBS; Gibco) and 1% penicillin-streptomycin (Invitrogen). MSCs were photographed through microscopy (Olympus, USA). For detecting cell surface markers, MSCs were characterized by CD73, CD90 and CD105 positive expression as well as CD34 and HLA-DR negative expression through flow cytometry (BD Biosciences, USA) under manufacturer's guidance. The surface antigens were provided by BD Biosciences.

Isolation and identification of exosomes

MSCs received culture in DMEM deprived of FBS for 2 days. Then the collected culture media received centrifugation at 500 g for 10 min, then 2000 g for 30 min to eliminate dead cells and debris, and finally 10000 g for 1 h to eliminate large vesicles.

The supernatant containing cell-free culture media were delivered to a new tube without disturbing the pellet and added the Total Exosome Isolation reagent (Invitrogen) under manufacturer's guidance. After collection, morphology of isolated exosomes got observation through transmission electron microscopy (TEM) (Philips, NLD). The particles were characterized by exosomal markers including CD63, TSG101 and Alix via western blotting.

A cellular model of IDD

Human NP cells (cat no. CP-H097) provided by Procell (Wuhan, China) received culture in DMEM containing 15% FBS and 1% penicillin-streptomycin. The culture medium was replaced twice every week and NP cells in the second or third passage were harvested for further use. NP cells under AGEs (200 µg/mL) stimulation received co-culture with MSC-exosomes (cat no. CP-H166) at 10, 50, or 100 µg/mL for 24 h. The untreated NP cells were taken as negative controls. For rescue assays, AGE-stimulated NP cells co-cultured with MSC-exosomes received treatment with 5 mM of 3-MA (Sigma-Aldrich), a specific autophagy inhibitor, for 24 h.

An animal model of IDD

The Animal Experimentation Committee of our hospital approved the animal experiments. A total of 24 three-month-old Sprague-Dawley rats were utilized in our research. After all rats received anesthetization with 2% (w/v) pentobarbital (40 mg/kg), three IVDs (Co7/8, Co8/9 and Co9/10) in each rat were located by palpation on coccygeal vertebrae and validated using trial radiography. Moreover, three IVDs (Co7/8, Co8/9 and Co9/10) respectively received intradiscal injection of 2 µL of PBS, AGEs (200 µg/mL) or a mixture of AGEs (200 µg/mL) and exosomes (100 µg/mL) through a 33-gauge needle (Hamilton), termed Sham, Model or Model+exosomes group, with 6 rats in each group. The injections were performed every two weeks for two months. For rescue assays, IVDs in Model+exosomes group received intraperitoneal injection of 3-MA at 1.5 mg/100 g for 24 h.

MRI examination

MRI was performed on all rats using a 7.0 T animal specific MRI system (Bruker Pharmascan, Germany). Pfirrmann classification was used to assess the degree of IVD degeneration [16]. The average score of punctured IVDs received calculation as degeneration grading of rats.

Histological analysis

After MRI examination, all rats were sacrificed through intraperitoneal administration with overdose pentobarbital sodium. The specimens were decalcified and received fixation in formaldehyde followed by dehydration and were finally by embedded in paraffin. The slides of each disc received staining with hematoxylin-eosin (HE) and observed through microscopy.

Immunofluorescence

NP cells or tissues attached to slides received fixation with 4% paraformaldehyde and permeabilization with 0.2% Triton X-100 in PBS. After washed using PBS, slides were blocked with 2% BSA in PBS at 37°C for 2 h, followed by incubation with primary antibody against LC3B (ab48394, Abcam, Shanghai, China). After washed twice, slides received subsequent treatment with secondary goat anti-rabbit antibody at 37°C for 2 h. Nuclei received co-staining by DAPI (Beyotime) at 0.1 g/mL for 5 min. Images were obtained through microscopy.

Western blot

Total proteins got isolation from NP cells or mouse tissues through a protein extraction kit (Beyotime) under manufacturer' guidance. The primary antibodies including CD63, TSG101, Alix, LC3B, Beclin-1, ATG5, IL-6, IL-1 β , TNF- α and β -actin as well as horseradish peroxidase (HRP)-conjugated secondary antibodies were obtained from Abcam. Protein bands got visualization and detection through enhanced chemiluminescence.

Enzyme linked immunosorbent assay (ELISA)

The concentrations of proinflammatory cytokines were tested with corresponding ELISA kits (IL-6, IL-1 β and TNF- α ; ThermoFisher) under manufacturer's guidance.

Statistical analysis

Data were expressed as the mean \pm standard deviation of three independent assays. Statistical analysis was conducted using GraphPad Prism 7 software. Comparisons between two groups were assessed with Student's t-test and comparisons among multiple groups were assessed with one-way ANOVA followed by Tukey's post hoc test. A statistical significance was presented upon $P < 0.05$.

RESULTS

Identification of MSC and its derived exosomes

MSCs, which have the advantages of easy access, multi-directional differentiation, low immunogenicity, and mature acquisition and identification, have been widely used in IDD-related reports [9]. Herein, human MSCs were harvested from bone marrow aspirates of donors. We observed characteristics of MSCs through microscopy. As a result, it appeared like a spindle with an approximately 60-70% confluence (Figure 1A). Flow cytometry demonstrated CD90, CD73 and CD105 positive expression as well as HLA-DR and CD34 negative expression in MSCs (Figure 1B). Then, MSC-exosomes got isolation and purification from MSC culture medium and particle morphology got confirmation under TEM (Figure 1C). Additionally, western blotting depicted that level of exosomal marker proteins (CD63, TSG101 and Alix) presented a remarkable elevation in exosomes relative to MSCs (Figure 1D). Collectively, these harvested particles are identified as MSC-exosomes.

MSC-exosomes facilitate autophagy in AGE-stimulated NP cells

Previously, AGEs accumulation exerted a pathogenic role in IDD [17]. Thus, we mimicked IDD characteristics in vitro through establishing a cellular model in NP

cells under AGE treatment, termed AGEs group, and non-treated NP cells were taken as Control group. To clarify MSC-exosomes role in AGE-stimulated NP cells, AGEs group received additional exosomes treatment at different doses, termed Model+exos-10/50/100 group. Previously, MSC-exosomes upregulated autophagy marker proteins LC3 and Beclin-1 and facilitated autophagosome formation [13]. We attempted to clarify impact of MSC-exosomes on autophagy pathway in NP cells. Detecting autophagosomal marker LC3 through immunofluorescence is a reliable means of monitoring autophagy [18]. Thus, we measured fluorescent intensity of LC3 using immunofluorescence. As a result, LC3-labeled NP cells presented depletion in AGEs group and showed a dose-dependent elevation under co-treatment with MSC-exosomes (Figure 2A). Additionally, western blotting demonstrated the downregulation of LC3B-II, Beclin-1 and ATG5 in AGEs group and this effect was reversed by MSC-exosomes along with increase of concentration of MSC-exosomes (Figure 2B). Collectively, MSC-exosomes facilitate activation of autophagy pathway in AGE-treated NP cells.

MSC-exosomes suppress inflammation in AGE-stimulated NP cells

MSC-exosomes are capable of attenuating NP cell apoptosis induced by proinflammatory cytokines [15]. Thus, we attempted to clarify influence of MSC-exosomes on inflammatory response in NP cells. ELISA demonstrated that contents of proinflammatory cytokines (IL-6, IL-1 β and TNF- α) presented elevation in AGEs group and showed a dose-dependent depletion under co-treatment with MSC-exosomes (Figure 3A). Similarly, western blotting demonstrated the upregulation of IL-6, IL-1 β and TNF- α in AGEs group and this effect was reversed by MSC-exosomes along with increase of concentration of MSC-exosomes (Figure 3B). Collectively, MSC-exosomes repress inflammatory response in AGE-treated NP cells.

Inhibiting autophagy enhances inflammation in AGE-stimulated NP cells

We aimed to clarify whether autophagy pathway exerted a role in inflammatory

response in NP cells. Thus, we performed rescue assays using 3-MA, a specific autophagy inhibitor [19], in cellular models under AGEs treatment.

Immunofluorescence illustrated that LC3 fluorescence in AG-stimulated NP cells presented elevation under treatment with MSC-exosomes and showed depletion under co-treatment with 3-MA (Figure 4A). Moreover, ELISA depicted that contents of IL-6, IL-1 β and TNF- α in AG-stimulated NP cells presented depletion under treatment with MSC-exosomes and showed elevation under co-treatment with 3-MA (Figure 4B). Western blotting showed a similar trend to ELISA results (Figure 4C).

Collectively, autophagy pathway activation enhances inflammatory response in AGE-stimulated NP cells.

MSC-exosomes induce autophagy and repress inflammation in a rat model of IDD

We attempted to further elucidate role of MSC-exosomes underlying IDD progression. Thus, we mimicked IDD characteristics in vivo through establishing IDD rat models. All rats were divided into Sham, Model or Model+exosomes group in a random manner. Three independent discs of each rat received intradiscal injection with PBS, AGEs, or AGEs+MSC-exosomes, respectively. We evaluated IDD degree through Pfirrmann MRI grading system [20]. As a result, Pfirrmann scores presented elevation in Model group and showed depletion under co-treatment with MSC-exosomes (Figure 5A). Histology by H&E staining indicated that MSC-exosomes reversed AGE-triggered IVD degradation during IDD (Figure 5B). Moreover, immunofluorescence demonstrated that AGE-triggered depletion in LC3 fluorescence was rescued by MSC-exosomes (Figure 5C). Western blotting also depicted that AGE-triggered depletion in LC3B-II, Beclin-1 and ATG5 protein abundances was rescued by MSC-exosomes (Figure 5D), suggesting that autophagy pathway is suppressed in IDD rats while activated by MSC-exosomes. Furthermore, AGE-triggered increase in IL-6, IL-1 β and TNF- α concentrations was reversed by MSC-exosomes (Figure 5E). Western blotting showed a similar trend to ELISA results (Figure 5F). Collectively, MSC-exosomes facilitates autophagy pathway activation

and repress inflammation in IDD rats.

Autophagy inhibition exerts a protective role against inflammation in a rat model of IDD

We aimed to clarify whether autophagy pathway exerted a role in inflammatory response in IDD progression. Thus, we performed rescue assays using 3-MA in IDD rat models. MSC-exosomes-triggered depletion in Pfirrmann scores in IDD rats was rescued by 3-MA treatment (Figure 6A). Histology by H&E staining indicated that 3-MA countervailed the inhibitory impact of MSC-exosomes on IVD degradation during IDD (Figure 6B). Moreover, immunofluorescence demonstrated that MSC-exosomes-triggered elevation in LC3 fluorescence was rescued by 3-MA (Figure 6C). Furthermore, MSC-exosomes-triggered decrease in IL-6, IL-1 β and TNF- α concentrations was reversed by MSC-exosomes (Figure 6D). Western blotting showed a similar trend to ELISA results (Figure 6E). Collectively, autophagy inhibition exerts a protective role against inflammatory response in IDD rats.

Ethics approval and consent to participate

Animal procedures were approved by The First Affiliated Hospital of Hebei North University and this study was approved by the Ethics Committee of The First Affiliated Hospital of Hebei North University.

DISCUSSION

At the cellular level, IDD presents pathological characterization by decrease in NP cell amount and a metabolic imbalance of major products of the ECM (proteoglycan and collagen type II) [21-23]. Thus, finding a source of cells that can replenish degenerated NP cells can effectively improve IDD and simultaneously can achieve the purpose of early prevention and avoid further aggravation of IDD and serious consequences.

Autophagy is a crucial mechanism for cells to self-digest and recycle damaged components, especially under stressful conditions, and is a vital survival mechanism

for cells under nutrient deprivation [24, 25]. Autophagy exerts a protective role in IVD cells [26]. Oxidative stress, pH, stress load, inflammatory cytokine stimulation, and hyperglycemic hypertonic environment in IVD microenvironment can activate the autophagy activity of NP cells, while various extracellular environment stimuli, such as oxidative stress, can reduce ability of cells to proliferate [27-29]. Our research also validated that IDD degree can be remarkably improved through modulating autophagy level in NP cells. Biological information can be transmitted between cells that are not in contact with each other in the form of paracrine or exosomes, and MSCs can also treat IDD through secreting exosomes [30, 31]. MSC exosomes can alleviate ER stress-induced apoptosis by activating AKT and ERK signaling [32]. Herein, exosomes secreted by MSCs received successful detection through TEM and immunoblot. By establishing a co-culture model, we discovered that in NP cellular model under AGEs stimulation, autophagy-related proteins in AGE-stimulated NP cells presented downregulation, while autophagy-related proteins in AGE-stimulated NP cells with MSC-exosome co-culture presented upregulation along with elevation of MSC-exosomes concentration, fully suggesting that MSCs enhances autophagy level in AGE-stimulated NP cells through MSC-secreted exosomes. Additionally, we established an IDD rat model and obtained similar results to a cellular model.

Inflammatory response mediated by pro-inflammatory cytokines can directly cause loss of water in IVD and exacerbate IDD process [33]. The enhanced "pro-inflammatory" role and insufficient "anti-inflammatory" role, that is, the imbalance of "inflammatory arm", may be a vital reason for aggravated inflammatory response in IDD [34]. Previously, pro-inflammatory cytokine IL-1 presented upregulation and its inhibitor IL-1a presented depletion in IDD [35]. Kritschil et al illustrated that IGF-1, which exerted an anti-inflammatory role, presented downregulation in IDD [36]. Herein, in AGE-treated NP cellular model, pro-inflammatory cytokines in AGEs-treated NP cells presented upregulation, while pro-inflammatory cytokines presented downregulation along with elevation of MSC-exosomes concentration, fully suggesting that MSCs repress inflammatory response of NP cells through MSC-

secreted exosomes. Additionally, we established an IDD rat model and obtained similar results to a cellular model.

Autophagy is a widespread self-protection mechanism in eukaryotic cells [37]. Recent research has revealed that there is a close connection between autophagy and inflammation [38, 39]. Though inflammatory responses innately protect the body from infection or tissue damage, persistent or excessive inflammatory responses can result in irreversible damage to the body [40]. It has been demonstrated that autophagy suppresses the inflammatory response possibly through directly repressing inflammatory complexes and indirectly through removing inflammatory stimuli such as damaged organelles or pathogenic microorganisms, thereby protecting cells from excessively persistent inflammation [41, 42]. Herein, in AGE-stimulated NP cellular model co-cultured with MSC-exosomes, 3-MA, an autophagy inhibitor, counteracted promoting influence of MSC exosomes on autophagy activation and reversed inhibitory impact of MSC-exosomes on inflammation, fully suggesting that MSC-exosomes suppresses AGE-induced NP cell inflammation via activating autophagy pathway. These results were also validated in animal models.

In conclusion, MSC-exosomes can improve degeneration of IDD via activating autophagy pathway to suppress inflammation, providing a potential novel insight for seeking therapeutic plans of IDD. However, due to complex and diverse components in exosomes, it remains elusive which component or cytokine in exosomes and which pathway exerts the main role, which will be addressed one by one in our future research.

Funding

This work was supported by Scientific and Technological Winter Olympic Games Special Project of Hebei Province Technological Innovation-Guided Plan in 2020 (No. 20477707D). Key Laboratory of Medical Science Research in Hebei Province (20110176).

Conflict of interest: None declared

REFERENCES

1. Buchbinder, R., et al., Low back pain: a call for action. *Lancet*, 2018. 391(10137): p. 2384-2388.
2. Wang, Y., et al., The role of IL-1 β and TNF- α in intervertebral disc degeneration. *Biomed Pharmacother*, 2020. 131: p. 110660.
3. Loibl, M., et al., Controversies in regenerative medicine: Should intervertebral disc degeneration be treated with mesenchymal stem cells? *JOR Spine*, 2019. 2(1): p. e1043.
4. van den Berg, R., et al., Association Between Self-Reported Spinal Morning Stiffness and Radiographic Evidence of Lumbar Disk Degeneration in Participants of the Cohort Hip and Cohort Knee (CHECK) Study. *Phys Ther*, 2020. 100(2): p. 255-267.
5. Fournier, D.E., et al., Vascularization of the human intervertebral disc: A scoping review. *JOR Spine*, 2020. 3(4): p. e1123.
6. González Martínez, E., et al., [Biology and mechanobiology of the intervertebral disc]. *Neurocirugia (Astur)*, 2017. 28(3): p. 135-140.
7. He, R., et al., HIF1A Alleviates compression-induced apoptosis of nucleus pulposus derived stem cells via upregulating autophagy. *Autophagy*, 2021. 17(11): p. 3338-3360.
8. Ou, X., et al., Activation of SIRT1 promotes cartilage differentiation and reduces apoptosis of nucleus pulposus mesenchymal stem cells via the MCP1/CCR2 axis in subjects with intervertebral disc degeneration. *Int J Mol Med*, 2020. 46(3): p. 1074-1084.
9. Kos, N., L. Gradisnik, and T. Velnar, A Brief Review of the Degenerative Intervertebral Disc Disease. *Med Arch*, 2019. 73(6): p. 421-424.
10. Song, X.X., et al., Substance P Mediates Estrogen Modulation Proinflammatory Cytokines Release in Intervertebral Disc. *Inflammation*, 2021. 44(2): p. 506-517.
11. Luo, Z., et al., Bone Marrow Stromal Cell-Derived Exosomes Promote Muscle Healing Following Contusion Through Macrophage Polarization. *Stem Cells Dev*, 2021. 30(3): p. 135-148.
12. Webber, J.P., et al., Differentiation of tumour-promoting stromal myofibroblasts by cancer exosomes. *Oncogene*, 2015. 34(3): p. 290-302.
13. Zhao, S., Y. Liu, and Z. Pu, Bone marrow mesenchymal stem cell-derived exosomes attenuate D-GaIN/LPS-induced hepatocyte apoptosis by activating autophagy in vitro. *Drug Des Devel Ther*, 2019. 13: p. 2887-2897.
14. Shi, M., et al., Therapeutic effect of co-culture of rat bone marrow mesenchymal stem cells and degenerated nucleus pulposus cells on intervertebral disc degeneration.

- Spine J, 2021. 21(9): p. 1567-1579.
15. Li, M., et al., Exosomes Derived from Bone Marrow Mesenchymal Stem Cells Prevent Acidic pH-Induced Damage in Human Nucleus Pulposus Cells. *Med Sci Monit*, 2020. 26: p. e922928.
 16. Pfirrmann, C.W., et al., Magnetic resonance classification of lumbar intervertebral disc degeneration. *Spine (Phila Pa 1976)*, 2001. 26(17): p. 1873-8.
 17. Luo, R., et al., FAM134B-Mediated ER-phagy Upregulation Attenuates AGEs-Induced Apoptosis and Senescence in Human Nucleus Pulposus Cells. *Oxid Med Cell Longev*, 2021. 2021: p. 3843145.
 18. Tanida, I., T. Ueno, and E. Kominami, LC3 and Autophagy. *Methods Mol Biol*, 2008. 445: p. 77-88.
 19. Shi, Y., et al., Delayed treatment with an autophagy inhibitor 3-MA alleviates the progression of hyperuricemic nephropathy. *Cell Death Dis*, 2020. 11(6): p. 467.
 20. Hollenberg, A.M., et al., Bone morphogenic protein-2 signaling in human disc degeneration and correlation to the Pfirrmann MRI grading system. *Spine J*, 2021. 21(7): p. 1205-1216.
 21. Chen, J., et al., Tension induces intervertebral disc degeneration via endoplasmic reticulum stress-mediated autophagy. *Biosci Rep*, 2019. 39(8).
 22. Bai, Z., et al., Protective effects of autophagy and NFE2L2 on reactive oxygen species-induced pyroptosis of human nucleus pulposus cells. *Aging (Albany NY)*, 2020. 12(8): p. 7534-7548.
 23. Wu, X., et al., Targeting the IL-1 β /IL-1Ra pathways for the aggregation of human islet amyloid polypeptide in an ex vivo organ culture system of the intervertebral disc. *Exp Mol Med*, 2019. 51(9): p. 1-16.
 24. Kim, J., et al., Autophagic Organelles in DNA Damage Response. *Front Cell Dev Biol*, 2021. 9: p. 668735.
 25. Feng, Y., et al., Downregulation of autophagy by Met30-mediated Atg9 ubiquitination. *Proc Natl Acad Sci U S A*, 2021. 118(1).
 26. Madhu, V., A.R. Guntur, and M.V. Risbud, Role of autophagy in intervertebral disc and cartilage function: implications in health and disease. *Matrix Biol*, 2021. 100-101: p. 207-220.
 27. Shi, P.Z., et al., Urolithin a alleviates oxidative stress-induced senescence in nucleus pulposus-derived mesenchymal stem cells through SIRT1/PGC-1 α pathway. *World J Stem Cells*, 2021. 13(12): p. 1928-1946.
 28. Yin, H., et al., The REDD1/TXNIP Complex Accelerates Oxidative Stress-Induced Apoptosis of Nucleus Pulposus Cells through the Mitochondrial Pathway. *Oxid Med Cell Longev*, 2021. 2021: p. 7397516.
 29. Xiang, Q., et al., Allicin Attenuated Advanced Oxidation Protein Product-Induced

- Oxidative Stress and Mitochondrial Apoptosis in Human Nucleus Pulposus Cells. *Oxid Med Cell Longev*, 2020. 2020: p. 6685043.
30. Zhu, L., et al., Mesenchymal stem cells-derived exosomes ameliorate nucleus pulposus cells apoptosis via delivering miR-142-3p: therapeutic potential for intervertebral disc degenerative diseases. *Cell Cycle*, 2020. 19(14): p. 1727-1739.
 31. Zhu, G., et al., Exosomal miR-532-5p from bone marrow mesenchymal stem cells reduce intervertebral disc degeneration by targeting RASSF5. *Exp Cell Res*, 2020. 393(2): p. 112109.
 32. Liao, Z., et al., Exosomes from mesenchymal stem cells modulate endoplasmic reticulum stress to protect against nucleus pulposus cell death and ameliorate intervertebral disc degeneration in vivo. *Theranostics*, 2019. 9(14): p. 4084-4100.
 33. Liu, Z.M., et al., Suramin attenuates intervertebral disc degeneration by inhibiting NF- κ B signalling pathway. *Bone Joint Res*, 2021. 10(8): p. 498-513.
 34. Zhao, F., et al., Magnoflorine Alleviates "M1" Polarized Macrophage-Induced Intervertebral Disc Degeneration Through Repressing the HMGB1/Myd88/NF- κ B Pathway and NLRP3 Inflammasome. *Front Pharmacol*, 2021. 12: p. 701087.
 35. Chen, Y., et al., Association of interleukin 1 gene polymorphism with intervertebral disc degeneration risk in the Chinese Han population. *Biosci Rep*, 2018. 38(4).
 36. Kritschil, R., et al., Effects of suppressing bioavailability of insulin-like growth factor on age-associated intervertebral disc degeneration. *JOR Spine*, 2020. 3(4): p. e1112.
 37. Li, M. and D.W. Liu, [Advances in the research of effects of regulation of cell autophagy on wound healing]. *Zhonghua Shao Shang Za Zhi*, 2017. 33(10): p. 625-628.
 38. Deretic, V., Autophagy in inflammation, infection, and immunometabolism. *Immunity*, 2021. 54(3): p. 437-453.
 39. Levine, B. and G. Kroemer, Biological Functions of Autophagy Genes: A Disease Perspective. *Cell*, 2019. 176(1-2): p. 11-42.
 40. Choudhary, V., et al., Pathogen-Associated Molecular Pattern-Induced TLR2 and TLR4 Activation Increases Keratinocyte Production of Inflammatory Mediators and is Inhibited by Phosphatidylglycerol. *Mol Pharmacol*, 2020. 97(5): p. 324-335.
 41. Chung, C., et al., Crosstalks between inflammasome and autophagy in cancer. *J Hematol Oncol*, 2020. 13(1): p. 100.
 42. Ravanan, P., I.F. Srikumar, and P. Talwar, Autophagy: The spotlight for cellular stress responses. *Life Sci*, 2017. 188: p. 53-67.

Figure 1




Figure 1. Characterization of MSC and its derived exosomes were identified. A, Characteristics of MSCs through microscopy. **B,** Characteristics of MSCs through flow cytometry. **C,** Characteristics of MSC-exosomes through TEM. **D,** Characteristics of MSC-exosomes through western blot.

Figure 2




Figure 2. MSC-exosomes facilitated autophagy in AGE-stimulated NP cells. A, Immunofluorescence detected LC3 fluorescence in NP cells in Control/AGEs/Model+exos-10/Model+exos-50/Model+exos-100 group. B, Western blot detected levels of autophagy-related proteins in NP cells under different conditions.

Figure 3




Figure 3. MSC-exosomes suppressed inflammation in AGE-stimulated NP cells. A, ELISA detected levels of proinflammatory cytokines in NP cells in Control/AGEs/Model+exos-10/Model+exos-50/Model+exos-100 group. B, Western blot detected levels of inflammation-related proteins in NP cells under different conditions.

Figure 4




Figure 4. Inhibiting autophagy enhanced inflammation in AGE-stimulated NP cells.

A, Immunofluorescence detected LC3 fluorescence in NP cells in

AGEs/AGEs+exos/AGEs+exos+3-MA group. B, ELISA detected levels of

proinflammatory cytokines in NP cells in NP cells under different conditions. C,

Western blot detected levels of inflammation-related proteins in NP cells under different conditions.

Figure 5




Figure 5. MSC-exosomes induced autophagy and repressed inflammation in a rat model of IDD. A, Pfirrmann MRI grading system evaluated pathological changes in discs in Sham/Model/Model+exosomes group. B, H&E staining evaluated pathological changes in discs in each group. C, Immunofluorescence detected LC3 fluorescence in discs in each group. D, Western blot detected levels of autophagy-related proteins in discs in each group. E, ELISA detected levels of proinflammatory cytokines in discs in each group. F, Western blot detected levels of inflammation-related proteins in discs in each group.

Figure 6




Figure 6. Autophagy inhibition exerted a protective role against inflammation in a rat model of IDD. A, Pfirrmann MRI grading system evaluated pathological changes in

discs in Exosomes/Exosomes+3-MA group. B, H&E staining evaluated pathological changes in discs in each group. C, Immunofluorescence detected LC3 fluorescence in discs in each group. D, ELISA detected levels of proinflammatory cytokines in discs in each group. E, Western blot detected levels of inflammation-related proteins in discs in each group.