



## **How multislice computed tomography (MSCT) of the coronary arteries can change the chronic total occlusion (CTO) recanalization procedure?**

**Authors:** Anna Kańtoch, Bernadeta Chyrchel, Sławomir Surowiec, Michał Chyrchel,  
Łukasz Rzeszutko, Andrzej Surdacki, Stanisław Bartuś, Leszek Bryniarski

**Article type:** Clinical vignette

**Received:** February 10, 2023

**Accepted:** March 19, 2023

**Early publication date:** March 26, 2023

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them

commercially.

# **How multislice computed tomography (MSCT) of the coronary arteries can change the chronic total occlusion (CTO) recanalization procedure?**

**Short title:** MSCT-coronary angiography can change the CTO recanalization procedure

Anna Kańtoch<sup>1</sup>, Bernadeta Chyrchel<sup>1,2</sup>, Sławomir Surowiec<sup>1</sup>, Michał Chyrchel<sup>1,2</sup>,  
Łukasz Rzeszutko<sup>1,2</sup>, Andrzej Surdacki<sup>1,2</sup>, Stanisław Bartuś<sup>1,2</sup>, Leszek Bryniarski<sup>1,2</sup>

<sup>1</sup>Department of Cardiology and Cardiovascular Interventions, University Hospital, Kraków, Poland

<sup>2</sup>2<sup>nd</sup> Department of Cardiology, Institute of Cardiology, Jagiellonian University Medical College, Kraków, Poland

## **Correspondence to:**

Prof. Leszek Bryniarski, MD, PhD, FESC,  
Department of Cardiology and Cardiovascular Interventions,  
Institute of Cardiology, Jagiellonian University Medical College,  
Jakubowskiego 2, 30–688 Kraków, Poland,  
phone: +48 12 400 22 50,  
e-mail: l\_bryniarski@poczta.fm

The Euro CTO Club guidelines advise against performing ad hoc coronary intervention (PCI) of chronic total occlusion (CTO) — it should be done after a careful analysis of the coronary angiography [1]. However, performing multislice computed tomography (MSCT) of the coronary arteries before coronary angiography may change this strategy.

MSCT of the coronary arteries was performed in a 58-year-old man with arterial hypertension, hypercholesterolemia, type 2 diabetes mellitus and CCS (Canadian Cardiovascular Society) class II angina for approximately 12 months. MSCT showed a 10 mm occlusion (**Figure 1A, 1C**) in the proximal segment of the dominant right coronary artery (RCA). In addition, there was critical stenosis distal to the occlusion.

Angiography (**Figure 1B, 1D**), in contrast to the MSCT results, revealed a 40 mm long RCA CTO with bifurcation at distal cap. The CTO had an ambiguous proximal cap and the distal part of the vessel could be visualized with ipsilateral and contralateral collaterals (J-CTO score 2 — intermediate category of difficulty).

Based on the information from the MSCT (visible entry and length — CT-RECTOR score — 0 — easy difficulty category), in contrast to the angiography result, the dedicated CTO operator decided to perform ad hoc PCI CTO from the right femoral arterial access.

Using Gaia Second and Confianza wires and a microcatheter, the lesion was crossed in 25 seconds. After pre-dilation, a Xience Pro drug-eluting stent (3.5 × 48 mm) was implanted, followed by post-dilation with an NC balloon to 18 atmospheres. The CTO procedure lasted 20 minutes (45 minutes with angiography), radiation dose 0.229 Gy, fluoroscopy time — 16.1 minutes, contrast — 200 ml. The periprocedural period was uncomplicated.

In conclusion, distal RCA stenosis blocked retrograde flow of contrast, mimicking a much longer CTO lesion in angiography. Based on the MSCT reconstruction, ad hoc PCI CTO could be performed.

Coronary CT angiography has become a significant step forward in evaluating the benefit-risk balance of the CTO PCI procedure [2]. The main purpose of using MSCT prior to CTO PCI is to quantify the structure of atherosclerotic plaque and to provide detailed anatomical information about coronary vascularity [2].

According to the literature, pre-procedural coronary CT-guidance for CTO was associated with fewer direct periprocedural complications, including periprocedural myocardial infarction and coronary perforation [3]. The limitation of the intra-procedural use of CT may be limited by the need for additional doses of radiation and contrast in patients undergoing PCI [4]. However, CT-guided CTO procedures have been found to have significantly higher success rates than procedures performed without CT [4, 5]. What is more, CT-guided PCI is associated with a shorter procedure duration, so it can be hypothesized that the dose of radiation and contrast during these procedures may be lower than in patients not undergoing CT scans [4].

Coronary CT angiography is becoming the basic tool in the treatment of CTO from pre-procedural evaluation, intra-procedural control to follow-up [4]. A new horizon in interventional cardiology could be the use of CT scans directly in the catheterization lab for real-time PCI [4].

## **Article information**

**Conflict of interest:** None declared.


**Funding:** None.

**Open access:** This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, which

allows downloading and sharing articles with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at [kardiologiapolska@ptkardio.pl](mailto:kardiologiapolska@ptkardio.pl).

## REFERENCES

1. Galassi AR, Werner GS, Boukhris M, et al. Percutaneous recanalisation of chronic total occlusions: 2019 consensus document from the EuroCTO Club. *EuroIntervention*. 2019; 15(2): 198–208, doi: [10.4244/EIJ-D-18-00826](https://doi.org/10.4244/EIJ-D-18-00826), indexed in Pubmed: [30636678](https://pubmed.ncbi.nlm.nih.gov/30636678/).
2. Tzimas G, Gulsin GS, Takagi H, et al. Coronary CT Angiography to Guide Percutaneous Coronary Intervention. *Radiol Cardiothorac Imaging*. 2022; 4(1): e210171, doi: [10.1148/ryct.210171](https://doi.org/10.1148/ryct.210171), indexed in Pubmed: [35782760](https://pubmed.ncbi.nlm.nih.gov/35782760/).
3. Hong SJ, Kim BK, Cho I, et al. CT-CTO Investigators. Effect of Coronary CTA on Chronic Total Occlusion Percutaneous Coronary Intervention: A Randomized Trial. *JACC Cardiovasc Imaging*. 2021; 14(10): 1993–2004, doi: [10.1016/j.jcmg.2021.04.013](https://doi.org/10.1016/j.jcmg.2021.04.013), indexed in Pubmed: [34147439](https://pubmed.ncbi.nlm.nih.gov/34147439/).
4. Melotti E, Belmonte M, Gigante C, et al. The Role of Multimodality Imaging for Percutaneous Coronary Intervention in Patients With Chronic Total Occlusions. *Front Cardiovasc Med*. 2022; 9: 823091, doi: [10.3389/fcvm.2022.823091](https://doi.org/10.3389/fcvm.2022.823091), indexed in Pubmed: [35586657](https://pubmed.ncbi.nlm.nih.gov/35586657/).
5. Rolf A, Werner GS, Schuhbäck A, et al. Preprocedural coronary CT angiography significantly improves success rates of PCI for chronic total occlusion. *Int J Cardiovasc Imaging*. 2013; 29(8): 1819–1827, doi: [10.1007/s10554-013-0258-y](https://doi.org/10.1007/s10554-013-0258-y), indexed in Pubmed: [23793727](https://pubmed.ncbi.nlm.nih.gov/23793727/).



**F**

**figure 1. A.** MSCT — a 10 mm occlusion (blue arrow) in the proximal segment of the dominant right coronary artery (RCA). **B.** Right coronary artery (RAO 30° projection) — proximal occlusion (yellow arrow), blunt stump visible. From the ipsilateral collateral circulation, the middle and distal sections fill. End of occlusion (red arrow). The length of the occlusion assessed angiographically — approx. 40–45 mm. **C.** MSCT (RAO 18) — a 10 mm occlusion in the proximal segment of the dominant right coronary artery (RCA). **D.** Right coronary artery (LAO 30° projection) — proximal occlusion (yellow arrow), blunt stump visible. From the ipsilateral collateral circulation, the middle and distal sections fill. End of occlusion (red arrow). Angiographically assessed occlusion length — approx. 40–45 mm