



A randomized comparison of HBP versus RVP: effect on left ventricular function and biomarkers of collagen metabolism

Authors: Jan Mizner, Petr Waldauf, Domenico Grieco, Hana Linkova, Oana Ionita, Pugazhendhi Vijayaraman, Robert Petr, Radka Raková, Jana Vesela, Petr Stros, Dalibor Herman, Pavel Osmancik, Karol Curila

Article type: Original article

Received: November 1, 2022

Accepted: January 2, 2023

Early publication date: March 9, 2023

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

A randomized comparison of HBP versus RVP: effect on left ventricular function and biomarkers of collagen metabolism

Short title: A comparison of His bundle pacing versus right ventricular pacing

Jan Mizner¹, Petr Waldauf², Domenico Grieco³, Hana Linkova¹, Oana Ionita¹, Pugazhendhi Vijayaraman⁴, Robert Petr¹, Radka Raková¹, Jana Vesela¹, Petr Stros¹, Dalibor Herman¹, Pavel Osmancik¹, Karol Curila¹

¹Cardiocenter, 3rd Faculty of Medicine, Charles University and University Hospital Kralovske Vinohrady, Prague, Czech Republic

²Department of Anesthesia and Intensive Care, Charles University and University Hospital Kralovske Vinohrady, Prague, Czech Republic

³Department of Cardiology, Policlinico Casilino of Rome, Rome, Italy

⁴Geisinger Heart Institute, Wilkes Barre, Pennsylvania, USA

Correspondence to:

Karol Curila, MD, PhD, MSc,

Cardiocenter, 3rd Faculty of Medicine,

Charles University,

Srobarova 50, 100 34 Prague, Czech Republic,

phone: +420 2 6716 2621,

e-mail: karol.curila@fnkv.cz, ORCID identifier: 0000-0003-3523-6358

WHAT'S NEW?

In patients at high risk of pacing-induced cardiomyopathy, His bundle pacing does not deteriorate left ventricular function in contrast to right ventricular pacing, which leads to its decline. Higher Galectin 3 and ST2-IL levels in patients treated with right ventricular pacing are associated with decline of left ventricular ejection fraction after six months of pacing. Galectin 3 and ST2-IL may improve the identification of patients in whom right ventricular pacing will not be associated with decline in left ventricular function.

ABSTRACT

Background: Right ventricular pacing (RVP) can result in pacing-induced cardiomyopathy (PICM). It is unknown whether specific biomarkers reflect differences between His bundle pacing (HBP) and RVP and predict a decrease in left ventricular function during RVP.

Aims: To compare the effect of HBP and RVP on the LV ejection fraction (LVEF) and to study how they affect serum markers of collagen metabolism.

Methods: Ninety-two high-risk PICM patients were randomized to HBP or RVP. Their clinical characteristics, echocardiography, and serum levels of TGF- β 1, MMP-9, ST2-IL, TIMP-1, and Gal-3 were studied before and six months after pacemaker implantation.

Results: Fifty-three patients were randomized to HBP and 39 patients to RVP. HBP failed in 10 patients, which crossed over to the RVP group. Patients with RVP had significantly lower LVEF compared to HBP after six months of pacing (-5% and -4% in as-treated and intention-to-treat analysis, respectively). Levels of TGF- β 1 after 6 months were lower in HBP than RVP (mean difference -6 ng/ml; $P = 0.009$) and preimplant Gal-3 and ST2-IL levels were higher in RVP patients with a decline in the LVEF $\geq 5\%$ compared to those with a decline of $< 5\%$ (mean difference 3 ng/ml and 8 ng/ml; $P = 0.02$ for both).

Conclusion: In high-risk PICM patients, HBP was superior to RVP in providing more physiological ventricular function, as reflected by higher LVEF and lower levels of TGF- β 1. Among RVP patients, LVEF declined more in those with higher baseline Gal-3 and ST2-IL levels than those with lower levels.

Key words: His bundle pacing, markers of collagen metabolism, right ventricular pacing

INTRODUCTION

Myocardial pacing of the right ventricle (RVP) is responsible for declining left ventricular (LV) function and heart failure in some patients. The highest risk of these adverse consequences is seen in older patients with a high burden of RV pacing, decreased left ventricular function, coronary artery disease (CAD), and wider spontaneous or paced QRS complexes [1]. His bundle pacing (HBP) preserves synchronous ventricular activation and represents the most physiological method of ventricular pacing[2, 3]. The pacing method is more complex, with longer procedure times and higher radiation doses, and requires more sophisticated equipment [4]. For these reasons, HBP is best suited for patients who would gain the most from physiological ventricular activation. However, the benefit of HBP in high-risk populations has never been described.

Although the RV pacing is unphysiological, most patients tolerate it even for extended periods [5]. Currently, we cannot precisely identify (before pacemaker implantation) which patients will experience deterioration in ventricular function after RV pacing. The period after which pacing-induced cardiomyopathy (PICM) starts to develop is estimated to be 2–3 years. However, subtle changes in LV function (i.e., decline $\geq 5\%$) can present sooner, and these patients are at the highest risk of further heart failure [6]. Remodeling and altered LV function are present together with changes in the ventricular microstructure. These changes are reflected by perfusion changes in particular ventricular segments, abnormal myocardial metabolism, increased fibrosis, and myocardial disarray [7]. It was already shown that subtle myocardial microstructure changes in patients after myocardial infarction or heart failure could be evaluated using collagen metabolism biomarkers [7]. However, their significance in patients with a permanent pacemaker has never been established. Demonstrating their relevance to LV performance in these patients could be an important marker of increased risk of further heart failure.

Our study aimed to assess the effect of RVP and HBP on LV function in patients at high risk of heart failure after cardiac pacing. Another goal was to identify laboratory markers that can predict or detect the adverse effects of RV pacing on LV performance.

METHODS

Patients

This was a prospective open-labeled randomized study with the anticipated recruitment of 120 patients. The project was approved by the Ethics Committee of the Faculty Hospital Kralovske Vinohrady, Prague, CZ; all subjects signed informed consent before enrollment. Only patients with conduction disease and an indication for permanent cardiac pacing per 2013 ESC Guidelines were enrolled. Patients had to have a permanent conduction disease with an anticipated high burden of the RV pacing and a life expectancy greater than two years. Also, at least one of the following criteria had to be fulfilled: (1) left ventricular ejection fraction $\leq 60\%$; (2) QRS duration > 115 ms; (3) presence of ischemic heart disease (defined as previous myocardial infarction or coronary intervention due to significant occlusion of coronary arteries or angina pectoris requiring pharmacologic treatment).

Exclusion criteria were as follows: a severe valvular disease with a planned intervention, cardiac surgery due to valvular disease or CAD in the last three months, permanent or persistent atrial fibrillation, dilated or hypertrophic cardiomyopathy, an indication for ICD or CRT implantation, and active myocarditis. Patients were randomized into the HBP or RVP

arm with a 4:3 ratio; the anticipated His bundle pacing success rate was 80–90%. After randomization, patients were informed which arm of the study they were enrolled in. After pacemaker implantation, outpatient clinic follow-ups were at six weeks and six months. During these visits, the pacemaker was checked (with data collection), clinical status was assessed, and a physical examination was performed. Blood sampling and echocardiography were performed before pacemaker implantation and at the six-month follow-up visit.

Pacemaker implantation

His bundle pacing was performed using Select Secure leads (model 3830, 69 cm, Medtronic Inc., Minneapolis, MN, US) delivered through a fixed-curve sheath (C315 HIS, Medtronic, Minneapolis, MN, US) preferentially from the left subclavian approach. The end of the sheath was delivered to the tricuspid annulus over the guidewire, and then the pacing lead was advanced through the sheath 1–2 mm beyond the tip of the catheter. The His bundle area was mapped in unipolar settings using an electrophysiology system (Lab system Pro, Boston Scientific, Marlborough, MA, US) at a sweep speed of 200 mm/s. After the His bundle signal was identified, the lead was fixed by 3–5 clockwise rotations, and pacing from the lead tip was initiated. For the implant procedure to be considered successful, selective, or nonselective, His bundle capture had to be present during the pacing with a pacing output below 2.5 V at 1 ms.

RV septal pacing was performed using Tendril^R (Abbott, Little Canada, MN, US) or Ingevity^R (Boston Scientific, Marlborough, MA, US) pacing leads, preferably from the left subclavian approach. Once the lead was placed in the RV outflow tract/pulmonary artery, the stylet was pre-shaped, and the lead was fixed in the RV septum using the RAO projection and counter-clockwise torque on the leads' stylet. The lead tip septal position was verified in the RAO 30° and LAO 30° projections.

Echocardiography

Echocardiography assessments were performed one day before pacemaker implantation and six months after by three cardiac sonographers using a GE Vivid E95 Cardiovascular Ultrasound (Boston, MA, US). Two evaluators blinded to the studied groups measured and calculated end-diastolic and end-systolic volumes from the apical 4- and 2- chamber views, and LVEF was calculated using the formula: $EF = [(LVEDV - LVESV) \div LVEDV]$ (modified Simpson's method). [Definitions: EF, ejection fraction; LVEDV, left ventricular end-diastolic volume; LVESV, left ventricular end-systolic volume]. The mean value of

LVEF calculated by each evaluator was used for statistical analyses.

Blood sample collections and quantification of cytokines

Approximately four mL of peripheral venous blood were collected from each patient. Blood samples were centrifugated at 950 g for 20 minutes. Serum samples were aliquoted and stored at -80°C . Samples were thawed prior to quantifying Transforming Growth Factor β 1 (TGF- β 1), Matrix Metalloproteinase 9 (MMP-9), Suppression of Tumorigenicity 2 Interleukin (ST2-IL), Tissue Inhibitor of Metalloproteinase 1 (TIMP-1), and Galectin 3 (Gal-3) levels. Per the manufacturer's instructions, the measurements of the selected biomarkers were performed using specific Quantikine ELISA kits (R&D Systems, Minneapolis, MN, US).

Statistical analysis

Statistical analysis was performed using Software: R version 4.0.5 (March 31, 2021). Exploratory data analysis was performed for all variables. Categorical data are presented as count with frequency and continuous data as mean with standard deviation (SD) or alternatively median with interquartile ranges (IQR) for nonparametric data. Kolmogorov and Smirnov tests were used for normality testing, and further statistical analysis included a linear mixed effect model with random intercept, Student's t-test, Fisher's exact test, and χ^2 test.

For the linear mixed effect model, the fixed part of the model is represented by the interaction between two binary parameters: stimulation site (His vs. septum) and visitation (Day 0 vs. Day 180). The random part of the model is represented by the random intercept, which is the patient ID. A maximum likelihood estimator was used to fit models (function lmer of package lme4).[8] Post hoc analysis was performed using the emmeans package. Intention-to-treat and as-treated analysis were performed. For nonparametric data, the Wilcoxon test and Mann-Whitney U test were used. A $P < 0.05$ was considered statistically significant. The area under the curve (AUC) of the receiver operating characteristic (ROC) curve was calculated for ST2-IL and Gal3 to assess their predictive value for LVEF deterioration. The optimal cutting points of both markers were calculated using maximization of the Youden index (sensitivity + [specificity - 1]). This was a pilot feasibility trial, and no power calculation was performed prior to the initiation of the study.

RESULTS

Ninety-two patients were randomized into the study. The mean age was 78 years, and all had

AV conduction disease as the pacing indication. Planned patient recruitment was not reached, and randomization was stopped due to challenges during the COVID pandemic. Fifty-three patients were randomized in the His bundle pacing (HBP group), and 39 were randomized to right ventricular pacing (RVP group). Lead placement in the HB region failed in 10 of 53 patients (19%) randomized to the HBP group. The lead was then successfully placed in the RV with myocardial capture in all patients. However, two of these patients (20%) required ventricular lead revision due to pacing threshold rise. The reasons for lead implant failure in the HB region were as follows: (1) in two patients, the HB signal was not found; (2) in four patients, the distal HV block could not be corrected by HB pacing; and (3) in four patients, pacing the HB region did not lead to conductive tissue capture with QRS narrowing. As a result, 49 patients had RVP (47 septal and two apical lead positions), and 43 had HBP. No difference in clinical characteristics was observed between groups relative to intention-to-treat and as-treated analyses (Table 1).

HBP required a longer fluoroscopy time (in intention-to-treat analysis), higher acute and chronic pacing thresholds and presented with lower acute and chronic ventricular sensing than RVP. However, there was no difference in rates of lead repositions due to higher pacing thresholds between HBP and RVP group (Table 2).

There was no difference between HBP and RVP groups in the preimplant LVEF in both intention-to-treat and as-treated comparisons. However, the LVEF significantly decreased after six months of RVP but remained the same in the HBP group. Also, the LVEF was significantly lower in RVP than in the HBP group after six months of follow-up in both as-treated ($P < 0.001$) and intention-to-treat analysis ($P = 0.008$) (Figure 1).

The decline in the LVEF of $\geq 5\%$ after six months of pacing was observed in 13 of 46 patients (28%) in the RVP group but in none in the HBP group. Among patients with RVP, a decline in LVEF $\geq 10\%$ was observed in nine patients (20%); and in eight patients (17%), the resultant LVEF was $\leq 45\%$ after six months of pacing.

There was no difference in baseline serum levels of TGF- β 1, MMP-9, ST2-IL, TIMP-1, and Gal-3 between patients with HBP vs. patients with RVP (both as-treated and intention-to-treat comparison). In the RVP group, in an *as-treated* comparison, a significant decline in the levels of ST2-IL and TIMP-1 was observed after six months of pacing, but no difference in the serum levels of TGF- β 1, MMP-9, and Gal-3 was detected. In the HBP group, a significant decline in the serum level of ST2-IL, MMP-9, and TGF- β 1 was seen after six months of pacing; the levels of Gal-3 and TIMP-1 remained statistically the same. When comparing differences in serum levels of studied biomarkers between HBP and RVP six months after the

pacemaker implantation, the only difference was observed in the levels of TGF- β 1, which were significantly lower in the HBP group than in the RVP group (Figure 2).

To determine whether cytokine levels before pacemaker implantation could predict an LVEF decline of $\geq 5\%$, we compared cytokine levels in patients with RVP and an LVEF decline of $\geq 5\%$ (13 patients) vs. cytokine levels in patients with RVP and LVEF decline $< 5\%$ (36 patients). Patients in both groups did not differ with respect to age, gender, preimplant LVEF, QRS duration during spontaneous rhythm, the prevalence of CAD, myocardial infarction, hypertension, or DM.

Patients with an LVEF decline $\geq 5\%$ after six months of RVP had higher baseline levels of Gal-3 and ST2-IL. After six months, the elevations of both markers persisted and were higher than in patients with an LVEF decline $< 5\%$ in the primary analysis and also after adjustment to the baseline levels of both molecules (Figure 3 and Supplementary material, Figure S1). During RVP, a decline in TIMP-1 was observed in patients without deterioration of LVEF ($P = 0.04$). No difference in serum levels of the other studied biomarkers was found before and after six months of RVP (Figure 3). The ROC analysis showed an AUC of 0.79 for Gal-3 and 0.71 for ST2-IL relative to the prediction of a decline in LVEF $\geq 5\%$ (Figure 4). Gal-3 serum concentrations ≥ 8.88 ng/ml was 100% sensitive and 61% specific, with a positive predictive value of 45%, a negative predictive value of 100%, and an accuracy of 72%; ST2-IL concentrations ≥ 19 ng/ml showed 90% specificity and 52% specificity, with a positive predictive value of 38%, a negative predictive value of 94%, and an accuracy of 71% for detection of patients with a decline in LVEF $\geq 5\%$ after six months of RV pacing.

In the HBP group, patients with higher baseline Gal 3 (>8.88 ng/ml) and ST2-IL (>19 ng/ml) levels did not differ in LVEF change after 6 months of follow-up in comparison to patients with lower baseline Gal 3 and ST2-IL levels (LVEF change 1 vs. 1 % and 1 vs. 1 %; $P = 0.66$ and $P = 0.72$, respectively).

DISCUSSION

This study compared the effect of His bundle pacing and RV myocardial pacing on the LVEF in patients at high risk of pacing-induced cardiomyopathy. Also, this is the first trial studying fibrosis biomarkers in patients with pacemakers. We showed that adverse effect on LV function with a decline in LVEF $\geq 5\%$ after pacing was not uncommon and affected almost 1/3 of patients with RV pacing, with the LVEF falling below 45% in 17% of the group. Contrary to this, HBP preserved LV function in all patients. We also showed that initiation of permanent cardiac pacing resulted in changes in the serum levels of some of the studied

biomarkers, with serum TGF- β 1 levels reflecting different ventricular activation during HBP and RVP. Lastly, patients with a decline in the LVEF $\geq 5\%$ due to unphysiological RV pacing had significantly higher serum levels of Gal-3 and ST2-IL than patients with a $< 5\%$ decline in LVEF, both at the baseline and after six months of RV pacing.

HBP vs. RVP

His bundle pacing is well established, and guidelines support treatment options in selected patients with bradycardia [9]. However, data from randomized trials supporting its use in a wider spectrum of patients are missing. So far, only one randomized trial comparing His bundle pacing to right ventricular septal pacing in patients with conduction disease has been published [10]. It used a crossover design, with HBP and RV pacing being utilized in the same patient for 12 months, and the number of randomized patients was small. Moreover, the studied population differed from our group, e.g., only patients with narrow QRS complexes (the average was 93 ms), and most were without coronary artery disease. The study showed that HBP preserved LVEF and ventricular synchrony better than right ventricular septal pacing, which resulted in a significant decline in the LVEF (mean decline of $4 \pm 1\%$). A similar level of LVEF deterioration during RVP occurred in a shorter period in our study; possibly reflecting the higher risk profile of our patients. Coronary artery disease was present in 1/3 of our patients, and the average QRS duration was 126 ms; both have been associated with a higher risk of adverse LV remodeling during pacing.[1] Considering the relationship between the severity of the LVEF decline and the duration of unphysiological RV pacing, it is possible that the difference in LVEF between HBP and RVP would be even greater with a longer follow-up. In our study, a decrease of LVEF $\geq 5\%$ was seen only in patients with RV pacing. Although a 5% decline in LVEF could be considered clinically negligible, it was previously shown that patients who demonstrate a slight decrease in LVEF soon after the pacemaker implantation were at the highest risk of further PICM [11]. It is often defined as a decline in the LVEF of more than 10% and/or an LVEF $< 50\%$ [1]. Using this definition, 20% of patients in our high-risk population developed PICM after six months of pacing. This agrees with the numbers reported by other investigators; however, it occurred longer after pacemaker implantation than in our study [1].

The difference in serum levels of studied cytokines between HBP vs. RVP

In patients with bradycardia and pacemaker implantation, we studied serum levels of collagen metabolism and fibrosis biomarkers, which were already shown to play a role in adverse

ventricular remodeling in different clinical scenarios [12–15]. Right ventricular myocardial pacing leads to unphysiological ventricular activation with adverse remodeling and LVEF deterioration in some patients [7]. These changes should be reflected in serum levels of biomarkers of fibrosis, although they have yet to be studied in patients with pacemakers. We showed that cardiac pacing (HBP or RVP) led to a decline in the serum levels of some of the studied cytokines; however, after six months of pacing, the groups differed only in the levels of TGF- β 1. TGF- β 1 is a pleiotropic cytokine critically involved in cardiac injury, repair, remodeling, and fibrogenesis. It also exerts potent matrix-preserving actions by suppressing the activity of MMPs and by inducing the synthesis of protease inhibitors, such as TIMP-1. Elevated TGF- β 1 levels in experimental in vivo models of heart failure were associated with increased myocardial stiffness, fibrosis, and LV diastolic dysfunction [16]. We found that TGF- β 1 declined after the institution of HBP but remained the same in RVP patients. This may reflect the normalization of atrioventricular synchrony with truly physiological ventricular activation in HBP patients [17]. In RVP patients, AV synchrony was also normalized, but at the cost of unphysiological ventricular activation due to RV pacing, which is associated with worsening LV diastolic function. [18].

New pacing strategies, such as His bundle pacing and left bundle branch area pacing, reduce the risk of adverse LV remodeling and heart failure in bradycardia patients [3, 19]. However, because they are more complex, the techniques may be best suited for those with the highest risk of LVEF deterioration after RVP. This remains a challenge because we still cannot accurately predict which patients will have a decline in LVEF due to RVP. Our theory was that the detrimental effect of RVP would be seen mostly in patients susceptible to the harmful effect of RV pacing, i.e., with a pre-existing condition, like increased myocardial fibrosis, which could be reflected in serum levels of studied biomarkers. Therefore, we compared these biomarkers in patients with an LVEF decline of $\geq 5\%$ vs. those with preserved LVEF during RVP (i.e., $< 5\%$). The only cytokines that showed different preimplant levels were Gal-3 and ST2-IL, both known as prognostic biomarkers in heart failure patients and involved in collagen metabolism and ventricular remodeling. [14, 15]. Data on their significance in patients with pacemakers are scarce. However, it was already shown that higher preimplant Gal-3 levels were negatively associated with response to cardiac resynchronization therapy and higher levels of myocardial fibrosis in ventricular myocardium, as seen on preimplant cardiac magnetic resonance [20]. It is possible that increased levels of Gal-3 and ST2-IL in our patients with a more significant decline in LVEF during RVP reflected a higher degree of pre-implant myocardial fibrosis, which led to a more

deleterious effect of RV pacing on LV performance. On the other hand, patients without significant myocardial fibrosis have a greater ability to compensate for dyssynchronous ventricular activation during RVP while maintaining the LVEF.

Limitations

This was a single-center study with echocardiographic follow-up restricted to six months, which prohibited tracking LVEF changes and clinical outcomes over a more extended period. Potential bias could have been present during the evaluation of echocardiographic measurements. Although the evaluator was blinded to the randomization of patients in the studied groups, the position of the pacing lead in the His bundle or RV septal region could be seen during the evaluation. An LVEF decline of 5%, which was used to compare groups, is relatively small and difficult to measure precisely, especially in patients with LV dyssynchrony due to pacing. The burden of ventricular pacing was taken from the programmer's printouts, and we did not study the incidence of fused pacing beats during Holter-ECG monitoring, which could lead to a higher burden of ventricular pacing as was, in fact, present. Finally, the number of patients in the RVP group and, more specifically, those with a decline in the LVEF after pacing was small, preventing more robust conclusions about the PICM prediction based on specific levels of studied molecules.

CONCLUSIONS

In patients at high risk of PICM, right ventricular pacing led to a decline in left ventricular ejection fraction compared to His bundle pacing, which preserved LV function after six months of pacing. Gal-3 and ST2-IL have the potential to better identify patients in which right ventricular pacing does not pose a significant risk. Further studies with more patients, longer follow-up, and clinical endpoints are needed to verify their predictive powers relative to pacing-induced cardiomyopathy.

Supplementary material

Supplementary material is available at https://journals.viamedica.pl/kardiologia_polska

Article information

Conflict of interest: None declared.

Funding: This paper was supported by the Charles University Research Center program

Cooperatio — Cardiovascular science, National Institute for Research of Metabolic and Cardiovascular Diseases (Programme EXCELES, ID Project No. LX22NPO5104) - Funded by the European Union – Next Generation EU and Ministry of Health of the Czech Republic, grant number NU21-02-00584.


Open access: This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, which allows downloading and sharing articles with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at kardiologiapolska@ptkardio.pl.

REFERENCES

1. Khurshid S, Epstein AE, Verdino RJ, et al. Incidence and predictors of right ventricular pacing-induced cardiomyopathy. *Heart Rhythm*. 2014; 11(9): 1619–1625, doi: [10.1016/j.hrthm.2014.05.040](https://doi.org/10.1016/j.hrthm.2014.05.040), indexed in Pubmed: [24893122](https://pubmed.ncbi.nlm.nih.gov/24893122/).
2. Curila K, Jurak P, Halamek J, et al. Ventricular activation pattern assessment during right ventricular pacing: Ultra-high-frequency ECG study. *J Cardiovasc Electrophysiol*. 2021; 32(5): 1385–1394, doi: [10.1111/jce.14985](https://doi.org/10.1111/jce.14985), indexed in Pubmed: [33682277](https://pubmed.ncbi.nlm.nih.gov/33682277/).
3. Abdelrahman M, Subzposh FA, Beer D, et al. Clinical outcomes of his bundle pacing compared to right ventricular pacing. *J Am Coll Cardiol*. 2018; 71(20): 2319–2330, doi: [10.1016/j.jacc.2018.02.048](https://doi.org/10.1016/j.jacc.2018.02.048), indexed in Pubmed: [29535066](https://pubmed.ncbi.nlm.nih.gov/29535066/).
4. Vijayaraman P, Herweg B, Dandamudi G, et al. Outcomes of His-bundle pacing upgrade after long-term right ventricular pacing and/or pacing-induced cardiomyopathy: Insights into disease progression. *Heart Rhythm*. 2019; 16(10): 1554–1561, doi: [10.1016/j.hrthm.2019.03.026](https://doi.org/10.1016/j.hrthm.2019.03.026), indexed in Pubmed: [30930330](https://pubmed.ncbi.nlm.nih.gov/30930330/).
5. Dreger H, Maethner K, Bondke H, et al. Pacing-induced cardiomyopathy in patients with right ventricular stimulation for >15 years. *Europace*. 2012; 14(2): 238–242, doi: [10.1093/europace/eur258](https://doi.org/10.1093/europace/eur258), indexed in Pubmed: [21846642](https://pubmed.ncbi.nlm.nih.gov/21846642/).
6. Fang F, Zhang Q, Chan JYS, et al. Early pacing-induced systolic dyssynchrony is a strong predictor of left ventricular adverse remodeling: analysis from the Pacing to Avoid Cardiac Enlargement (PACE) trial. *Int J Cardiol*. 2013; 168(2): 723–728, doi: [10.1016/j.ijcard.2012.08.005](https://doi.org/10.1016/j.ijcard.2012.08.005), indexed in Pubmed: [22944596](https://pubmed.ncbi.nlm.nih.gov/22944596/).

7. Ahmed FZ, Khattar RS, Zaidi AM, et al. Pacing-induced cardiomyopathy: pathophysiological insights through matrix metalloproteinases. *Heart Fail Rev.* 2014; 19(5): 669–680, doi: [10.1007/s10741-013-9390-y](https://doi.org/10.1007/s10741-013-9390-y), indexed in Pubmed: [23856884](https://pubmed.ncbi.nlm.nih.gov/23856884/).
8. Bates D, Mächler M, Bolker B, et al. Fitting linear mixed-effects models using **lme4**. *Journal of Statistical Software.* 2015; 67(1), doi: [10.18637/jss.v067.i01](https://doi.org/10.18637/jss.v067.i01).
9. Kusumoto FM, Schoenfeld MH, Barrett C, et al. 2018 ACC/AHA/HRS Guideline on the Evaluation and Management of Patients With Bradycardia and Cardiac Conduction Delay: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Circulation.* 2019; 140(8): e382–e482, doi: [10.1161/CIR.0000000000000628](https://doi.org/10.1161/CIR.0000000000000628), indexed in Pubmed: [30586772](https://pubmed.ncbi.nlm.nih.gov/30586772/).
10. Kronborg MB, Mortensen PT, Poulsen SH, et al. His and para-His pacing in AV block: feasibility and electrocardiographic findings. *J Interv Card Electrophysiol.* 2011; 31(3): 255–262, doi: [10.1007/s10840-011-9565-1](https://doi.org/10.1007/s10840-011-9565-1), indexed in Pubmed: [21465234](https://pubmed.ncbi.nlm.nih.gov/21465234/).
11. Chan JYS, Fang F, Zhang Q, et al. Biventricular pacing is superior to right ventricular pacing in bradycardia patients with preserved systolic function: 2-year results of the PACE trial. *Eur Heart J.* 2011; 32(20): 2533–2540, doi: [10.1093/eurheartj/ehr336](https://doi.org/10.1093/eurheartj/ehr336), indexed in Pubmed: [21875860](https://pubmed.ncbi.nlm.nih.gov/21875860/).
12. Morishita T, Uzui H, Mitsuke Y, et al. Association between matrix metalloproteinase-9 and worsening heart failure events in patients with chronic heart failure. *ESC Heart Fail.* 2017; 4(3): 321–330, doi: [10.1002/ehf2.12137](https://doi.org/10.1002/ehf2.12137), indexed in Pubmed: [28772055](https://pubmed.ncbi.nlm.nih.gov/28772055/).
13. Dobaczewski M, Chen W, Frangogiannis NG. Transforming growth factor (TGF)- β signaling in cardiac remodeling. *J Mol Cell Cardiol.* 2011; 51(4): 600–606, doi: [10.1016/j.yjmcc.2010.10.033](https://doi.org/10.1016/j.yjmcc.2010.10.033), indexed in Pubmed: [21059352](https://pubmed.ncbi.nlm.nih.gov/21059352/).
14. Imran TF, Shin HJ, Mathenge N, et al. Meta-Analysis of the Usefulness of Plasma Galectin-3 to Predict the Risk of Mortality in Patients With Heart Failure and in the General Population. *Am J Cardiol.* 2017; 119(1): 57–64, doi: [10.1016/j.amjcard.2016.09.019](https://doi.org/10.1016/j.amjcard.2016.09.019), indexed in Pubmed: [28247849](https://pubmed.ncbi.nlm.nih.gov/28247849/).
15. Aimo A, Vergaro G, Passino C, et al. Prognostic Value of Soluble Suppression of Tumorigenicity-2 in Chronic Heart Failure: A Meta-Analysis. *JACC Heart Fail.* 2017; 5(4): 280–286, doi: [10.1016/j.jchf.2016.09.010](https://doi.org/10.1016/j.jchf.2016.09.010), indexed in Pubmed: [27816512](https://pubmed.ncbi.nlm.nih.gov/27816512/).
16. Brooks WW, Conrad CH. Myocardial fibrosis in transforming growth factor beta(1)heterozygous mice. *J Mol Cell Cardiol.* 2000; 32(2): 187–195, doi: [10.1006/jmcc.1999.1065](https://doi.org/10.1006/jmcc.1999.1065), indexed in Pubmed: [10722796](https://pubmed.ncbi.nlm.nih.gov/10722796/).

17. Pastore G, Aggio S, Baracca E, et al. Hisian area and right ventricular apical pacing differently affect left atrial function: an intra-patients evaluation. *Europace*. 2014; 16(7): 1033–1039, doi: [10.1093/europace/eut436](https://doi.org/10.1093/europace/eut436), indexed in Pubmed: [24473501](https://pubmed.ncbi.nlm.nih.gov/24473501/).
18. Fang F, Zhang Q, Chan JYS, et al. Deleterious effect of right ventricular apical pacing on left ventricular diastolic function and the impact of pre-existing diastolic disease. *Eur Heart J*. 2011; 32(15): 1891–1899, doi: [10.1093/eurheartj/ehr118](https://doi.org/10.1093/eurheartj/ehr118), indexed in Pubmed: [21531741](https://pubmed.ncbi.nlm.nih.gov/21531741/).
19. Sharma PS, Patel NR, Ravi V, et al. Clinical outcomes of left bundle branch area pacing compared to right ventricular pacing: Results from the Geisinger-Rush Conduction System Pacing Registry. *Heart Rhythm*. 2022; 19(1): 3–11, doi: [10.1016/j.hrthm.2021.08.033](https://doi.org/10.1016/j.hrthm.2021.08.033), indexed in Pubmed: [34481985](https://pubmed.ncbi.nlm.nih.gov/34481985/).
20. Andre C, Piver E, Perault R, et al. Galectin-3 predicts response and outcomes after cardiac resynchronization therapy. *J Transl Med*. 2018; 16(1): 299, doi: [10.1186/s12967-018-1675-4](https://doi.org/10.1186/s12967-018-1675-4), indexed in Pubmed: [30390680](https://pubmed.ncbi.nlm.nih.gov/30390680/).



Central illustration. The study flow-chart and the effect of right ventricular pacing and His bundle pacing on left ventricular ejection fraction after six months of pacing in intention-to-treat and as-treated analyses

Abbreviations: HBP, His bundle pacing; LVEF, left ventricular ejection fraction; PICM, pacing-induced cardiomyopathy; RVP, right ventricular myocardial pacing




Figure 1. Comparison of LVEF in the HBP and RVP groups per intention-to-treat (A) and as-treated (B) analyses

** means $P < 0.01$, *** means $P < 0.001$

Abbreviations: see [Central illustration](#)




Figure 2. Comparison of serum levels of ST2-IL, TIMP-1, MMP-9, Galectin 3, and TGF-β1 at baseline and after six months of pacing in HBP vs. RVP group per as-treated analysis

*means $P < 0.05$, **means $P < 0.01$

Abbreviations: MMP-9, matrix metalloproteinase-9; ST2-IL, suppression of tumorigenicity 2 interleukin; TGF-β1, transforming growth factor β1; TIMP-1, tissue inhibitor of metalloproteinase-1; other — see [Central illustration](#)




Figure 3. Comparison of serum levels of, Gal-3, ST2-IL, MMP-9, TIMP-1, and TFG-beta1 before implant and after six months of pacing in patients with RVP and preserved LVEF vs. declined in LVEF $\geq 5\%$

Abbreviations: see [Central illustration](#) and [Figure 2](#)




Figure 4. Receiver operating characteristic (ROC) curves of Gal-3 (A) and ST2-IL (B) in patients with and without the decline in the LVEF \geq 5% after six months of RVP

Abbreviations: see [Central illustration](#) and [Figure 2](#)

Table 1. Baseline clinical characteristics of the study population

	Intention-to-treat			As-treated		
	RVP (n = 39)	HBP (n = 53)	P-value	RVP (n = 49)	HBP (n = 43)	P-value
Age, years, mean (SD)	78 (7)	78 (8)	0.99	79 (7)	77 (8)	0.33
Male sex, n (%)	39 (80)	38 (88)	0.26	33 (85)	44 (83)	0.84
LVEF, %, mean (SD)	58 (7)	60 (5)	0.27	59 (6)	59 (4)	0.54
Arterial hypertension, n (%)	38 (97)	51 (96)	0.75	48 (98)	41 (95)	0.49
Diabetes mellitus, n (%)	16 (41)	20 (38)	0.75	18 (37)	18 (42)	0.62
CAD, n (%)	15 (38)	23 (43)	0.64	18 (37)	20 (47)	0.34
Myocardial infarction in	5 (14)	14 (27)	0.15	8 (17)	11 (26)	0.32

history, n (%)						
Spontaneous QRSd, ms) mean (SD)	126 (27)	125 (25)	0.80	126 (26)	126 (27)	0.98
Spontaneous QRS morphology, n (%)						
BBB	16 (41)	20 (38)		19 (39)	17 (40)	
Narrow (<115 ms)	12 (31)	20 (38)	0.78	16 (33)	17 (40)	0.66
NIVCD	11 (28)	13 (24)		14 (28)	9 (20)	
Pacing indication, n (%)						
AV block I. degree	5 (13)	7 (13)		6 (12)	6 (14)	
AV block II. degree	16 (41)	25 (47)	0.95	21 (43)	20 (47)	0.94
AV block III. degree	16 (41)	19 (36)		20 (41)	15 (35)	
BBB + syncope	2 (5)	2 (4)		2 (4)	2 (4)	

Abbreviations: AV block, atrioventricular block; BBB, bundle branch block; CAD, coronary artery disease; NIVCD, non-specific intraventricular conduction delay; other — see **Central illustration**

Table 2. Procedural and follow-up pacing characteristics

		Intention-to-treat			As-treated		
		RVP	HBP	P-value	RVP	HBP	P-value
Pacing thresholds (V) at 0.4 ms, mean (SD)	D1	0.7 (0.3)	1.4 (0.6)	<0.001	0.8 (0.4)	1.5 (0.6)	<0.001
	D180	0.9 (0.6)	1.7 (1.1)	0.004	1.1 (0.7)	1.7 (1.1)	0.005
	D1 vs. D180 P-value	0.35	0.11		0.40	0.21	
Ventricular sensing, mV, mean (SD)	D1	9.4 (3.5)	4.5 (3.3)	<0.001	9.3 (3.7)	3.5 (2.0)	<0.001
	D180	9.5 (2.9)	4.3 (3.2)	<0.001	9.3 (3.1)	3.2 (2.0)	<0.001
	D1 vs. D180 P-value	0.91	0.75		0.98	0.54	
Fluoroscopy time, sec, median (IQR)		242 (171; 413)	505 (270; 835)	<0.001	329 (190; 553)	399 (249; 679)	0.34
Burden of ventricular pacing after 180 days, mean (SD)		92 (18)	98 (4)	0.02	95 (17)	98 (4)	0.09
Threshold rise requiring lead revision, n (%)		2 (5)	4 (8)	0.64	4 (8)	2 (5)	0.50