



Multimorbid management in atrial fibrillation: The Polish perspective in the EHRA- PATHS study

Authors: Geraldine A Lee, Michal M Farkowski, Edward Baker, Maciej Sterliński, Isabelle C van Gelder, Rafal Dąbrowski, Lien Desteghe, Łukasz Szumowski, Jose L Merino, Ronan Collins, Michiel Rienstra, Hein Heidbuchel

Article type: Original article

Received: December 12, 2022

Accepted: February 6, 2023

Early publication date: March 16, 2023

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Multimorbid management in atrial fibrillation: The Polish perspective in the EHRA-PATHS study

Geraldine A Lee^{1*}, Michal M Farkowski^{2*}, Edward Baker^{1, 3}, Maciej Sterliński⁴, Isabelle C van Gelder⁵, Rafal Dąbrowski⁶, Lien Desteghe⁷⁻¹⁰, Łukasz Szumowski⁴, Jose L Merino¹¹, Ronan Collins¹², Michiel Rienstra⁵, Hein Heidbuchel⁷⁻⁹

¹Florence Nightingale Faculty of Nursing, Midwifery and Palliative Care, King's College London, James Clerk Maxwell Building, London, United Kingdom

²Head of Arrhythmia Unit, Department of Cardiology, Central Clinical Hospital of the Ministry of Interior and Administration, Warszawa, Poland

³Emergency Department, King's College Hospital NHS Foundation Trust, Denmark Hill, London, United Kingdom

⁴Department of Heart Arrhythmia, National Institute of Cardiology, Warszawa, Poland

⁵Department of Cardiology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

⁶Department of Coronary Disease and Cardiac Rehabilitation, National Institute of Cardiology, Warszawa, Poland

⁷Department of Cardiology, Antwerp University Hospital, Antwerp, Belgium

⁸Research Group Cardiovascular Diseases, University of Antwerp, Antwerp, Belgium

⁹Faculty of Medicine and Life Sciences, Hasselt University, Hasselt, Belgium

¹⁰Heart Center Hasselt, Jessa Hospital, Hasselt, Belgium

¹¹La Paz University Hospital, IDIPAZ, Universidad Autonoma, Madrid, Spain

¹²Tallaght University Hospital, Department of Gerontology, Trinity College Dublin, Ireland

*Both authors equally contributed to the study

Correspondence to:

Geraldine Lee, PhD, FESC,

Reader in Advanced Clinical Practice,

Florence Nightingale Faculty of Nursing,

Midwifery and Palliative Care, King's College London,

James Clerk Maxwell Building, 57 Waterloo Road, London, SE1 8WA, UK,

phone: +44 802 074 83 201,

e-mail: gerry.lee@kcl.ac.uk

WHAT'S NEW?

Compared to other European locations, the rates of specialist services and referral varied in Poland but were not significantly different to the rest of Europe. The survey showed that there were statistically higher numbers of specialised services in Poland compared to the rest of Europe for hypertension and palpitations but lower rates for sleep apnea and comprehensive geriatric care. Polish physicians seems to face more insurance and financial barriers to deliver a comprehensive care than their European colleagues. Overall, there is a need for better interdisciplinary collaboration to improving patient outcomes in all European countries.

ABSTRACT

Background: Atrial fibrillation (AF) is the most common arrhythmia and places a significant burden on individuals as well as the healthcare system. AF management requires a multidisciplinary approach in which tackling comorbidities is an important aspect.

Aims: To evaluate how multimorbidity is currently assessed and managed and to determine if interdisciplinary care is undertaken.

Methods: A 21-item online survey was undertaken over four weeks as part of the EHRA-PATHS study examining comorbidities in AF and distributed to European Heart Rhythm Association members in Europe.

Results: A total of 341 eligible responses were received of which 35 (10%) were from Polish physicians. Compared to other European locations, the rates of specialist services and referrals varied but were not significantly different. However, there were higher numbers of specialised services reported in Poland compared to the rest of Europe for hypertension (57% vs. 37%; $P = 0.02$) and palpitations/arrhythmias (63% vs. 41%; $P = 0.01$) and the rates of sleep apnea services tended to be lower (20% vs. 34%; $P = 0.10$) and comprehensive geriatric care (14% vs. 36%; $P = 0.01$). The only statistical difference between Poland and the rest of Europe in reasons for referral rates was the barrier relating to insurance and financial reasons (31% vs. 11%; $P < 0.01$, respectively).

Conclusions: There is a clear need for an integrated approach to patients with AF and associated comorbidities. Preparedness of Polish physicians to deliver such care seems to be similar to other European countries but may be hampered by financial obstacles.

Key words: atrial fibrillation, comorbidities, older people, survey

INTRODUCTION

Atrial fibrillation (AF) is the most common cardiac arrhythmia affecting approximately 33 million adults and is associated with a significant burden on healthcare systems [1]. Many patients with AF have comorbidities, including hypertension, diabetes, sleep apnoea, coronary heart disease and others, what is globally associated with increased all-cause mortality. AF is a complex long-term condition involves a multifaceted, holistic, and multidisciplinary approach. With multimorbidity defined as the presence of two or more diagnosed long-term conditions [2], in relation to AF, there is a lack of a pathway-based approach to manage AF comorbidities. One observational study identified a six-fold increased all-cause mortality risk in those with AF who had four or more comorbidities compared to those without AF comorbidities [2]. Within increasing numbers of the population diagnosed with AF and associated concomitant conditions, there is a need for new interventions to optimise outcomes using a pathway-driven approach that is systematic and standardised. Patient pathway-based interventions have been proved to be positive in other populations, but these benefits have not been consistently identified across studies and disease processes [3–6].

The EHRA-PATHS “Addressing multimorbidity in elderly atrial fibrillation patients through interdisciplinary, tailored, patient-centered care pathways” is a Horizon 2020 project coordinated by the European Heart Rhythm Association (EHRA) and the European Society of Cardiology (ESC), with 14 research collaborators from across Europe. The primary aim of EHRA-PATHS is to develop a new pathway of care for older patients (>65 years) with multimorbid AF through interdisciplinary, patient-centred and systematic approaches [7]. This survey study is one component of a work package with the objective of undertaking a clinical practice gap analysis and measuring current clinical practices including clinicians' and patients' experiences [7]. With the results from the various work packages, a patient pathway-based intervention will be developed and evaluated for the management of multimorbid AF.

In relation to the healthcare professionals and current AF comorbidity management, the study aims to capture the opinions of Polish cardiologists and electrophysiologists and allied health professionals on the current structure and interdisciplinary management of co-morbidities in patients with AF with the specific aims: (1) Evaluate how multimorbidity is currently addressed by clinicians during AF treatment to characterise the treatment structure; (2) Assess how the interdisciplinary management of multimorbid AF is currently conducted.

METHODS

This survey was developed and piloted by the research team and a multi-methods cross-sectional design using both quantitative and qualitative approaches applied. The survey aims were achieved through the following objectives: (1) Identify the specific methods used by clinicians to assess, diagnose, manage and refer multimorbid AF patients throughout Europe; (2) describe key areas of complexity in the management of multimorbid AF across Europe; and (3) highlight areas of interprofessional working to optimise health status in patients with multimorbid AF throughout Europe. The survey consisted of 21 questions including respondent characteristics, 4 questions relating to local AF referral and management practices and 10 questions relating to participants' experiences of managing multimorbid AF and a free text section for any comments.

The questionnaire was placed on the Qualtrics Survey Platform as an e-survey with a digital link to the survey sent to all EHRA members via newsletters and EHRA emails. The survey was open for 6 weeks between November 1, 2021 to December 12, 2021. It was open to physicians, registered nurses and allied healthcare professionals who work directly with patients with AF in European countries and are members of the EHRA and ESC. They were recruited through convenience sampling methods. The aim was to try and have responses from 10% of EHRA members (n = 350). Based on the eligibility criteria, all responses from outside of the EU countries were excluded from the analysis process.

This study has been registered with King's College London Research Ethics Committee under the minimal risk registration process (Ref MRA-20/21-25315).

Data Analysis (including statistical analysis)

A mixed methods approach was applied to integrate both qualitative and quantitative findings. Descriptive data analysis was conducted through the Qualtrics survey platform (2021) and comparative inferential statistics was undertaken using the statistical software, International Business Machines Corporation (IBM) Statistical Package of Social Sciences (SPSS) Version 26 for statistical analysis. Descriptive data were presented as counts and percentages. Comparisons of categorical data were calculated using chi-squared analysis, except for where the expected cell counts were ≤ 5 where a Fischer's exact test was used. Throughout, a *P*-value of <0.05 indicated statistical significance.

Qualitative data analysis of free text responses was undertaken using conventional content analysis involving both deductive and inductive reasoning with coding undertaken to identify

themes and categories within the text 8. Analysis was undertaken by EB and GL, with thematic saturation reached after approximately 200 responses and the qualitative data was managed using NVivo v.11.

RESULTS

A total of 451 responses were received, with 376 responses from 29 European countries and 75 responses submitted from outside the EU and 37 responses submitted with no data, and these were excluded from analysis as per the study eligibility criteria. A total of 341 responses were included in the data analysis with 44% of responses received from the UK, Spain, and Ireland, followed by 10% (n = 35) of responses from Poland. No statistical differences were seen between Poland and the other countries in terms of gender, number of years in their specialty and their workplace (Table 1). The Polish responses were from electrophysiologists (n = 16, 46%) and cardiologists (n=19, 54%) with no responses from allied health professionals. Responses from Polish participants reported their speciality as electrophysiologists (74%) and general cardiologists (69%) (respondents could include more than one speciality in their answer) and ranged from less than five years of experience (9%), with the majority having 20 to 30 years experience (34%).

Regarding current clinical practice for multimorbid AF management, the analysis was undertaken by comparing all responses (n = 341) to Poland (n = 35) and the rest of Europe (n = 306) (Table 2). No statistical difference was seen in the number of AF patients seen per month ($P = 0.16$) or in the proportion of patients referred to other speciality services ($P = 0.20$). In terms of specialised services available, Poland had higher numbers of referrals to hypertension (57% in Poland vs. 37% in Europe, $P = 0.02$) and arrhythmia/palpitation (63% vs. 41%, $P = 0.01$) and comprehensive geriatric assessment (14% vs. 35%, $P = 0.01$) reported. Reasons for referral rates were explored and around half reported that this was the number that needed to be referred (44%), while resourcing was cited as an issue in 23% of Polish responses. Barriers identified in relation to resources included organisational/institutional issues (57%) and a lack of integrated models of care (more than 50%). The only difference between Poland and the rest of Europe was the barrier relating to insurance and financial reasons (31% in Poland vs. 11% in Europe, $P < 0.01$).

Free-text comments were included as part of the analysis and 229 responses were completed and coding was undertaken with 56 codes identified and these were which were refined into 38 codes (coding in qualitative research involves labelling and organising the data to identify

different themes). The four identified themes highlight the lack of integrated comorbid AF management and the themes were:

- Improving access to lifestyle and health promotion interventions, including the early management of risk factors or comorbidities (this relates to risk factor modification and the need for patient education including around weight loss management and medication adherence, for example),
- Organisational restructuring to enable innovation in care provision (this includes inflexibility in the existing systems and institutional governance along with unclear pathways for managing and treating comorbidities),
- Working towards achieving an evidence-based and integrated approach to multimorbid AF care for all (achieving consensus on core components of care in the standardised practice approach with most respondents advocating for the integrated model of care as this would be expected to have the greatest impact on patient outcomes),
- Aiming for great collaboration and interdisciplinary working, especially between cardiologists and primary care/geriatrics clinicians as well as building the specialist workforce, increasing the scope of practice for nurses and allied health professionals and working with primary care clinicians.

DISCUSSION

The findings from this survey demonstrate the current issues with multimorbid AF across Europe and highlight Poland in relation to other countries: (1) Higher access to hypertension or arrhythmia specialists in the outpatient setting contrary to access to comprehensive geriatric assessment; (2) Higher impact of reimbursement/financing issues on the patients' care, 3. Apparent lower access to formalised multi-specialist AF care.

Low access to geriatricians for Polish patients is not new as according to the data presented in 2022 by the Supreme Medical Chamber, the main Polish office of the doctor's self-government, their number is more than 10-fold lower than the number of Polish cardiologists: 555 vs. 5139 (circ. 14.7/million vs. 135.9/million people), respectively [9.] This number is comparable to Denmark (15.7/million) but markedly less than France (37.3/million) or Italy (49.6 /million) [10]. The Ministry of Health had recognized that there are fewer specialist physicians within geriatrics compared to other areas and have been promoting it among graduates of medical schools for years for example by special financial incentives among

others. So far with mixed results as shown in our analysis. On the other hand, geriatrics is an independent specialization in Poland contrary to eg. Greece or Portugal where it is recognized as a competence rather than specialization [11].

Since the majority of responses came from university/teaching hospitals both in Poland and the rest of Europe there was generally a high representation of arrhythmia specialists in both cohorts. The 2016 EHRA White Book placed Poland among countries with good access to device therapy and average access to ablation [12]. Theoretically, one could extrapolate this information and assume a relatively easy access to an arrhythmia specialist for Polish patient at least in comparison to a geriatrician. There is also a potential field of professional conflict between geriatricians and cardiologists in reducing the number and doses of drugs that improve prognosis by geriatricians. Lack of reimbursement of non-vitamin K oral antagonist oral anticoagulants (NOACs) remains a challenging issue for some of the Polish patients, especially in cases of multimorbidity-derived polypharmacy and rising costs of subsequent drugs and may result in their lower prescription [13, 14]. Universal health insurance provided by a national monopolist — the National Health Fund (NHF) covers hospital bills on the basis of disease-related groups and out-patients visits on a modifiable fee-for-service basis [15]. In both cases, the overall lump sum offered by the NHF does have a cap and normally does not cover the costs of all services, and proposed tariffs are substantially lower than expected. As a result, costs incurred by health care providers that exceed this cap are in general not reimbursed by the NHS. This leads to patient queuing and waiting lists lasting up to several months or even years across all specialities. Separate specialisations have their own lump sums and separate caps. This, together with a general preference of Polish patients to be treated by a ‘specialist’, results in even longer waiting times for a specialist consultation therefore hampering most attempts of any coordinated AF care. So far, in Poland there is one real program for coordinated cardiac care with distinctive rules and financing and it is dedicated to patients with myocardial infarction [16, 17]. Its results are very promising and may lead to other programs of coordinated cardiac care [18]. Yet for now, there are reports of discrepancies and even different outcomes of AF treatment among patients living in different parts of Poland [14, 19, 20].

The survey clearly demonstrates the challenges in treating and managing AF patients with comorbidities, reflecting the findings from the main survey across Europe [21]. AF is not alone in this challenge, with previous research identifying the need for an interdisciplinary, patient-centred approach to multimorbid care that optimises health-related quality of life via the development of self-efficacy through shared health-related goal setting [3–6].

A systematic approach in assessing AF patients' multimorbidity and its impact on patient health and decision-making is warranted. The survey results suggest that this approach is the first of multiple steps needed to achieve a sustained improvement in patient health status. Organisational structures and governance are required to integrate AF and multimorbid care with greater interdisciplinary working practices. Key to this is ongoing education for both patients and clinicians considering chronic disease management and medicines optimisation as well as long-term behavioural changes in relation to associated risk factors.

Risk factor identification and management is crucial in AF and should be reviewed regularly. However, due to the lack of protocolised care, it is often unclear who is responsible for this (i.e. cardiology or primary care, for example) [22, 23]. One solution is a hospital-based AF coordination center that would support primary care physicians and hospital-based specialists in coordinating and streamlining AF care [24]. Previous pan-European studies investigating the provision of healthcare over geographically diverse areas have shown the potential impact of these variations in health inequality [25, 26].

Medication management and medicine optimisation play an important role in AF management. Previous research has identified that approximately 20% of patients with two comorbidities are prescribed between four and nine medications, with 1% prescribed 10 or more medications [27]. Primary care physicians have previously highlighted the challenge of managing polypharmacy where medications are commenced by speciality clinicians [28, 29]. The lack of a standardised approach and good communication between acute and community services has been noted and highlights the need for better collaborative partnerships [28]. Within older people, results from the STOPP-START study showed the benefits of greater interdisciplinary working between geriatricians and pharmacists in reviewing polypharmacy and complex drug regimens with the implementation of evidence-based tools [28, 30, 31].

Integrated care in AF can include several specialists, but critically the patient needs to be included in the decision-making. Previous research has highlighted that communication between clinicians and patients, and between clinicians from different disciplines, is often poor and identified a relationship between substandard communication and patient outcomes [32, 33]. Ensuring continuity of care has been shown to improve both the patient experience and the patient outcome [34, 35]. A coordinated approach to managing the older multimorbid AF is important, and there is a need to involve different specialities with a particular emphasis on gerontological expertise and communication between clinicians and patients [36]. Shared decision-making is central to optimising patient outcomes, including improving quality of life and behavioural changes relating to known AF risk factors [37–40].

Limitations

There are some limitations which need to be acknowledged. Firstly, the sample size from Polish healthcare professionals was low and may not be representative of healthcare professionals across Poland especially due to high representation of physicians based in academic/teaching hospitals. We did not collect the ages of respondents and this may be construed as a limitation. The survey was administered via EHRA and therefore does not include the opinions of those not members of EHRA. Although we captured results from many respondents across Europe, the results may not be generalisable. There was a low response rate from allied health professionals, which needs to be acknowledged but it highlights issues regarding AF care across Europe and the lack of a multidisciplinary approach.

CONCLUSION

The results of the survey highlighted the current state of clinical practice in the management of multimorbid Atrial Fibrillation in Poland and across Europe. There are clearly varying levels of specialist services available as well as evidence demonstrating the lack of a systematic approach to multimorbidity management. The respondents highlighted the need for greater collaborative working, education and improving patient self-efficacy. An integrated management of Atrial Fibrillation-related comorbidities is clearly warranted and these results will inform the next phases of the EHRA-PATHS study.

Article information

Conflict of interest: None declared.

Funding: This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 945260.

Open access: This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, which allows downloading and sharing articles with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at kardiologiapolska@ptkardio.pl.

REFERENCES

1. Hindricks G, Potpara T, Dagres N, et al. 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS): The Task Force for the diagnosis and management of atrial fibrillation of the European Society of Cardiology (ESC) Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. *Eur Heart J.* 2021; 42(5): 373–498, doi: [10.1093/eurheartj/ehaa612](https://doi.org/10.1093/eurheartj/ehaa612), indexed in Pubmed: [32860505](https://pubmed.ncbi.nlm.nih.gov/32860505/).
2. Jani BD, Nicholl BI, McQueenie R, et al. Multimorbidity and co-morbidity in atrial fibrillation and effects on survival: findings from UK Biobank cohort. *Europace.* 2018; 20(FI_3): f329–f336, doi: [10.1093/europace/eux322](https://doi.org/10.1093/europace/eux322), indexed in Pubmed: [29112751](https://pubmed.ncbi.nlm.nih.gov/29112751/).
3. Baker E, Woolley A, Xyrichis A, et al. How does the implementation of a patient pathway-based intervention in the acute care of blunt thoracic injury impact on patient outcomes? A systematic review of the literature. *Injury.* 2020; 51(8): 1733–1743, doi: [10.1016/j.injury.2020.06.002](https://doi.org/10.1016/j.injury.2020.06.002), indexed in Pubmed: [32576379](https://pubmed.ncbi.nlm.nih.gov/32576379/).
4. Roberts HC, Pickering RM, Onslow E, et al. The effectiveness of implementing a care pathway for femoral neck fracture in older people: a prospective controlled before and after study. *Age Ageing.* 2004; 33(2): 178–184, doi: [10.1093/ageing/afh063](https://doi.org/10.1093/ageing/afh063), indexed in Pubmed: [14960435](https://pubmed.ncbi.nlm.nih.gov/14960435/).
5. Devapriam J, Alexander R, Gumber R, et al. Impact of care pathway-based approach on outcomes in a specialist intellectual disability inpatient unit. *J Intellect Disabil.* 2014; 18(3): 211–220, doi: [10.1177/1744629514532453](https://doi.org/10.1177/1744629514532453), indexed in Pubmed: [24814150](https://pubmed.ncbi.nlm.nih.gov/24814150/).
6. Kwan J, Hand P, Dennis M, et al. Effects of introducing an integrated care pathway in an acute stroke unit. *Age Ageing.* 2004; 33(4): 362–367, doi: [10.1093/ageing/afh104](https://doi.org/10.1093/ageing/afh104), indexed in Pubmed: [15047573](https://pubmed.ncbi.nlm.nih.gov/15047573/).
7. Heidbuchel H, Van Gelder IC, Desteghe L. ESC and EHRA lead a path towards integrated care for multimorbid atrial fibrillation patients: the Horizon 2020 EHRA-PATHS project. *Eur Heart J.* 2022; 43(15): 1450–1452, doi: [10.1093/eurheartj/ehab672](https://doi.org/10.1093/eurheartj/ehab672), indexed in Pubmed: [34694355](https://pubmed.ncbi.nlm.nih.gov/34694355/).
8. Hsieh HF, Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res.* 2005; 15(9): 1277–1288, doi: [10.1177/1049732305276687](https://doi.org/10.1177/1049732305276687), indexed in Pubmed: [16204405](https://pubmed.ncbi.nlm.nih.gov/16204405/).
9. Chamber of Physicians and Dentists [Naczlona Izba Lekarska]: Statistical information. Available online: <https://nil.org.pl/rejstry/centralny-rejestr-lekarzy/informacje-statystyczne>. [Accessed: December 2022]. .

10. OECD (2019) Health at a glance 2019: OECD indicators (summary). OECD Publishing, Paris, 10.1787/e88a7402-en. [Accessed: 19 May 2020].
11. Soulis G, Kotovskaya Y, Bahat G, et al. Geriatric care in European countries where geriatric medicine is still emerging. *Eur Geriatr Med.* 2021; 12(1): 205–211, doi: [10.1007/s41999-020-00419-7](https://doi.org/10.1007/s41999-020-00419-7), indexed in Pubmed: [33237564](https://pubmed.ncbi.nlm.nih.gov/33237564/).
12. Raatikainen MJ, Arnar DO, Merkely B, et al. Access to and clinical use of cardiac implantable electronic devices and interventional electrophysiological procedures in the European Society of Cardiology Countries: 2016 Report from the European Heart Rhythm Association. *Europace.* 2016; 18 Suppl 3: iii1–iii79, doi: [10.1093/europace/euw244](https://doi.org/10.1093/europace/euw244), indexed in Pubmed: [27496955](https://pubmed.ncbi.nlm.nih.gov/27496955/).
13. Łodziński P, Gawalko M, Budnik M, et al. Trends in antithrombotic management of patients with atrial fibrillation. A report from the Polish part of the EURObservational Research Programme - Atrial Fibrillation General Long-Term Registry. *Pol Arch Intern Med.* 2020; 130(3): 196–205, doi: [10.20452/pamw.15157](https://doi.org/10.20452/pamw.15157), indexed in Pubmed: [31976927](https://pubmed.ncbi.nlm.nih.gov/31976927/).
14. Janion-Sadowska A, Sadowski M, Koniecznyńska M, et al. Polish regional differences in patient knowledge on atrial fibrillation and its management as well as in patterns of oral anticoagulant prescription. *Kardiol Pol.* 2019; 77(4): 437–444, doi: [10.5603/KP.a2019.0036](https://doi.org/10.5603/KP.a2019.0036), indexed in Pubmed: [30835335](https://pubmed.ncbi.nlm.nih.gov/30835335/).
15. Sowada C, Sagan A, Kowalska-Bobko I, et al. Poland health system review. *Health Syst Transit.* 2011; 13(8): 1–193, indexed in Pubmed: [22551527](https://pubmed.ncbi.nlm.nih.gov/22551527/).
16. Kubiela G, Diakowska D, Uchmanowicz I. Survival analysis of patients with acute coronary syndrome receiving comprehensive coordinated care after myocardial infarction (KOS-Zawał). *Kardiol Pol.* 2022; 80(3): 415–321, doi: [10.33963/KP.a2022.0035](https://doi.org/10.33963/KP.a2022.0035), indexed in Pubmed: [35129204](https://pubmed.ncbi.nlm.nih.gov/35129204/).
17. Jankowski P, Gašior M, Gierlotka M, et al. Coordinated care after myocardial infarction. The statement of the Polish Cardiac Society and the Agency for Health Technology Assessment and Tariff System [in Polish]. *Kardiol Pol.* 2016; 74(8): 800–811, doi: [10.5603/KP.2016.0118](https://doi.org/10.5603/KP.2016.0118), indexed in Pubmed: [27553352](https://pubmed.ncbi.nlm.nih.gov/27553352/).
18. Sagan A, Rogala M, Buszman PP, et al. Improved coordination of care after acute myocardial infarction in Poland since 2017: Promising early results. *Health Policy.* 2021; 125(5): 587–592, doi: [10.1016/j.healthpol.2021.03.010](https://doi.org/10.1016/j.healthpol.2021.03.010), indexed in Pubmed: [33832777](https://pubmed.ncbi.nlm.nih.gov/33832777/).

19. Lodziński P, Gawałko M, Kraj L, et al. District versus academic hospitals: clinical outcomes of patients with atrial fibrillation. *Pol Arch Intern Med.* 2021; 131(10), doi: [10.20452/pamw.16053](https://doi.org/10.20452/pamw.16053), indexed in Pubmed: [34213298](https://pubmed.ncbi.nlm.nih.gov/34213298/).
20. Wojdyła-Hordyńska A, Baran J, Mazurek M, et al. Results of a survey concerning atrial fibrillation ablation strategies in Poland. *Kardiol Pol.* 2020; 78(10): 974–981, doi: [10.33963/KP.15407](https://doi.org/10.33963/KP.15407), indexed in Pubmed: [32486629](https://pubmed.ncbi.nlm.nih.gov/32486629/).
21. Lee G, Baker E, Collins R, et al. The challenge of managing multimorbid atrial fibrillation: a pan-European European Heart Rhythm Association (EHRA) member survey of current management practices and clinical priorities. *Europace.* 2022; 24(12): 2004–2014, doi: [10.1093/europace/euac136](https://doi.org/10.1093/europace/euac136), indexed in Pubmed: [36036694](https://pubmed.ncbi.nlm.nih.gov/36036694/).
22. Roland M, Paddison C. Better management of patients with multimorbidity. *BMJ.* 2013; 346: f2510, doi: [10.1136/bmj.f2510](https://doi.org/10.1136/bmj.f2510), indexed in Pubmed: [23641032](https://pubmed.ncbi.nlm.nih.gov/23641032/).
23. Porter I, Davey A, Gangannagaripalli J, et al. Quality of care assessment for people with multimorbidity. *J Intern Med.* 2019; 285(3): 289–300, doi: [10.1111/joim.12881](https://doi.org/10.1111/joim.12881), indexed in Pubmed: [30719790](https://pubmed.ncbi.nlm.nih.gov/30719790/).
24. Irving G, Neves AL, Dambha-Miller H, et al. International variations in primary care physician consultation time: a systematic review of 67 countries. *BMJ Open.* 2017; 7(10): e017902, doi: [10.1136/bmjopen-2017-017902](https://doi.org/10.1136/bmjopen-2017-017902), indexed in Pubmed: [29118053](https://pubmed.ncbi.nlm.nih.gov/29118053/).
25. Jutz R. Health inequalities in Eastern Europe. Does the role of the welfare regime differ from Western Europe? *Soc Sci Med.* 2020; 267: 113357, doi: [10.1016/j.socscimed.2020.113357](https://doi.org/10.1016/j.socscimed.2020.113357), indexed in Pubmed: [32980174](https://pubmed.ncbi.nlm.nih.gov/32980174/).
26. Mackenbach JP, Valverde JR, Artnik B, et al. Trends in health inequalities in 27 European countries. *Proc Natl Acad Sci U S A.* 2018; 115(25): 6440–6445, doi: [10.1073/pnas.1800028115](https://doi.org/10.1073/pnas.1800028115), indexed in Pubmed: [29866829](https://pubmed.ncbi.nlm.nih.gov/29866829/).
27. Payne RA, Avery AJ, Duerden M, et al. Prevalence of polypharmacy in a Scottish primary care population. *Eur J Clin Pharmacol.* 2014; 70(5): 575–581, doi: [10.1007/s00228-013-1639-9](https://doi.org/10.1007/s00228-013-1639-9), indexed in Pubmed: [24487416](https://pubmed.ncbi.nlm.nih.gov/24487416/).
28. Wallace E, Salisbury C, Guthrie B, et al. Managing patients with multimorbidity in primary care. *BMJ.* 2015; 350: h176, doi: [10.1136/bmj.h176](https://doi.org/10.1136/bmj.h176), indexed in Pubmed: [25646760](https://pubmed.ncbi.nlm.nih.gov/25646760/).
29. Payne RA, Avery AJ. Polypharmacy: one of the greatest prescribing challenges in general practice. *Br J Gen Pract.* 2011; 61(583): 83–84, doi: [10.3399/bjgp11X556146](https://doi.org/10.3399/bjgp11X556146), indexed in Pubmed: [21276330](https://pubmed.ncbi.nlm.nih.gov/21276330/).

30. Avery AJ, Rodgers S, Cantrill JA, et al. A pharmacist-led information technology intervention for medication errors (PINCER): a multicentre, cluster randomised, controlled trial and cost-effectiveness analysis. *Lancet*. 2012; 379(9823): 1310–1319, doi: [10.1016/S0140-6736\(11\)61817-5](https://doi.org/10.1016/S0140-6736(11)61817-5), indexed in Pubmed: [22357106](https://pubmed.ncbi.nlm.nih.gov/22357106/).
31. Gallagher P, Ryan C, Byrne S, et al. STOPP (Screening Tool of Older Person's Prescriptions) and START (Screening Tool to Alert doctors to Right Treatment). Consensus validation. *Int J Clin Pharmacol Ther*. 2008; 46(2): 72–83, doi: [10.5414/cpp46072](https://doi.org/10.5414/cpp46072), indexed in Pubmed: [18218287](https://pubmed.ncbi.nlm.nih.gov/18218287/).
32. Tiwary A, Rimal A, Paudyal B, et al. Poor communication by health care professionals may lead to life-threatening complications: examples from two case reports. *Wellcome Open Res*. 2019; 4: 7, doi: [10.12688/wellcomeopenres.15042.1](https://doi.org/10.12688/wellcomeopenres.15042.1), indexed in Pubmed: [31448336](https://pubmed.ncbi.nlm.nih.gov/31448336/).
33. Vermeir P, Vandijck D, Degroote S, et al. Communication in healthcare: a narrative review of the literature and practical recommendations. *Int J Clin Pract*. 2015; 69(11): 1257–1267, doi: [10.1111/ijcp.12686](https://doi.org/10.1111/ijcp.12686), indexed in Pubmed: [26147310](https://pubmed.ncbi.nlm.nih.gov/26147310/).
34. Nyweide DJ, Anthony DL, Bynum JPW, et al. Continuity of care and the risk of preventable hospitalization in older adults. *JAMA Intern Med*. 2013; 173(20): 1879–1885, doi: [10.1001/jamainternmed.2013.10059](https://doi.org/10.1001/jamainternmed.2013.10059), indexed in Pubmed: [24043127](https://pubmed.ncbi.nlm.nih.gov/24043127/).
35. Saultz JW, Lochner J. Interpersonal continuity of care and care outcomes: a critical review. *Ann Fam Med*. 2005; 3(2): 159–166, doi: [10.1370/afm.285](https://doi.org/10.1370/afm.285), indexed in Pubmed: [15798043](https://pubmed.ncbi.nlm.nih.gov/15798043/).
36. Sinnott C, Mc Hugh S, Browne J, et al. GPs' perspectives on the management of patients with multimorbidity: systematic review and synthesis of qualitative research. *BMJ Open*. 2013; 3(9): e003610, doi: [10.1136/bmjopen-2013-003610](https://doi.org/10.1136/bmjopen-2013-003610), indexed in Pubmed: [24038011](https://pubmed.ncbi.nlm.nih.gov/24038011/).
37. Foster G, Taylor SJC, Eldridge SE, et al. Self-management education programmes by lay leaders for people with chronic conditions. *Cochrane Database Syst Rev*. 2007(4): CD005108, doi: [10.1002/14651858.CD005108.pub2](https://doi.org/10.1002/14651858.CD005108.pub2), indexed in Pubmed: [17943839](https://pubmed.ncbi.nlm.nih.gov/17943839/).
38. Smith SM, Wallace E, O'Dowd T, et al. Interventions for improving outcomes in patients with multimorbidity in primary care and community settings. *Cochrane Database Syst Rev*. 2021; 1(1): CD006560, doi: [10.1002/14651858.CD006560.pub4](https://doi.org/10.1002/14651858.CD006560.pub4), indexed in Pubmed: [33448337](https://pubmed.ncbi.nlm.nih.gov/33448337/).
39. Kennedy A, Reeves D, Bower P, et al. The effectiveness and cost effectiveness of a national lay-led self care support programme for patients with long-term conditions: a

pragmatic randomised controlled trial. *J Epidemiol Community Health*. 2007; 61(3): 254–261, doi: [10.1136/jech.2006.053538](https://doi.org/10.1136/jech.2006.053538), indexed in Pubmed: [17325405](https://pubmed.ncbi.nlm.nih.gov/17325405/).

40. Kennedy A, Bower P, Reeves D, et al. Implementation of self management support for long term conditions in routine primary care settings: cluster randomised controlled trial. *BMJ*. 2013; 346: f2882, doi: [10.1136/bmj.f2882](https://doi.org/10.1136/bmj.f2882), indexed in Pubmed: [23670660](https://pubmed.ncbi.nlm.nih.gov/23670660/).

Table 1. Comparative characteristics between Europe and Poland in the survey sample

N (%)	Poland (n = 35)	Europe (n = 306)	P-value
Sex			
Male	27 (77.1)	198 (64.7)	0.41
Female	7 (20.0)	102 (33.3)	
Third gender/non-binary	0	1 (0.3)	
Not disclosed	1 (2.9)	5 (1.6)	
Professional group and specialist practice area:			
Electrophysiologist	16 (45.7)	93 (30.4)	0.08
Other cardiologist	19 (54.3)	149 (48.7)	
Physician with speciality other than cardiology	0	33 (10.8)	
Nurse or allied health professional working in general cardiology	0	10 (3.3)	
Nurse or allied health professional working in electrophysiology/ arrhythmias	0	16 (5.2)	
Respondents' specialist area of interest in AF management (can choose more than 1 specialty)			
Arrhythmias/electrophysiology and devices	26 (74.3)	171 (55.9)	0.04 ^a
General cardiology	24 (68.6)	183 (59.8)	0.31
Heart failure	18 (51.4)	113 (36.9)	0.10
Valvular disease	5 (14.3)	42 (13.7)	0.93
Imaging	6 (17.1)	48 (15.7)	0.82
Interventional cardiology	2 (5.7)	33 (10.8)	0.35
Cardiovascular prevention	4 (11.4)	61 (19.9)	0.23
Congenital heart disease	3 (8.6)	16 (5.2)	0.41
Stroke	2 (5.7)	36 (11.8)	0.28
Other	0	15 (4.9)	0.18
Number of years practising in this speciality			

<5 years	3 (8.6)	53 (17.3)	0.15
5–10 years	10 (28.6)	63 (20.6)	
10–20 years	7 (20.0)	91 (29.7)	
20–30 years	12 (34.3)	83 (27.1)	
>30 years	3 (8.6)	48 (15.7)	
Hospital designation			
University hospital/academic teaching hospital	16 (45.7)	182 (59.5)	0.18
Non-academic teaching hospital	8 (22.9)	43 (14.1)	
Community or district hospital	8 (22.9)	39 (12.7)	
Specialised Heart Centre	2 (5.7)	16 (5.2)	
Other setting	1 (2.9)	26 (8.5)	

Table 2. Comparing current multimorbid AF management in Poland and more widely across Europe

N (%)	Total sample (n = 341)	Poland (n = 35)	Europe (n = 306)	P-value
Typical numbers of patients seen with AF per month				
<20	47 (13.9)	1 (2.9)	46 (15.0)	0.16
20–50	169 (49.9)	16 (45.7)	154 (50.3)	
51–100	87 (25.7)	13 (37.1)	74 (24.2)	
101–150	19 (5.6)	2 (5.7)	18 (5.9)	
>150	17 (5.0)	3 (8.6)	14 (4.6)	
What specialised outpatient services are available at your center				
Atrial fibrillation	174 (51.3)	19 (54.3)	156 (51.0)	0.71
Heart failure	249 (73.5)	23 (65.7)	224 (73.2)	0.35
Hypertension	134 (39.5)	20 (57.1)	114 (37.3)	0.02 ^a
Diabetes	177 (52.2)	19 (54.3)	158 (51.6)	0.77
Lipid	138 (40.7)	15 (42.9)	123 (40.2)	0.76
Anticoagulation	142 (41.9)	13 (37.1)	129 (42.2)	0.57
Syncope	106 (31.3)	12 (34.3)	94 (30.7)	0.68
Chest pain	146 (43.1)	13 (37.1)	134 (43.8)	0.45
Palpitations/arrhythmia/resynchronisation	148 (43.7)	22 (62.9)	124 (40.5)	0.01 ^a
Sleep apnoea	110 (32.4)	7 (20.0)	103 (33.7)	0.10
Comprehensive Geriatric Assessment (dementia, falls, frailty,	116 (34.2)	5 (14.3)	109 (35.6)	0.01 ^a

etc.) Other	17 (5.0)	1 (2.9)	23 (7.5)	0.31
What proportion of patients with comorbidities are referred to other speciality services?				
Over 80%	7 (2.1)	0	7 (2.3)	0.20
61–80%	12 (3.5)	1 (2.9)	11 (3.6)	
41–60%	55 (16.2)	10 (28.9)	45 (14.7)	
20–40%	104 (30.7)	10 (28.9)	96 (31.4)	
Less than 19%	123 (36.3)	9 (25.7)	114 (37.3)	
No response	38 (11.2)	5 (14.3)	33 (10.8)	
What is the reason for this referral rate?				
That is the number that needs referring	151 (44.5)	19 (54.3)	132 (43.1)	0.09
Resourcing issue so I need to be selective and prioritise	61 (18.0)	8 (22.9)	53 (17.3)	
There is an established process with the relevant specialties	73 (21.5)	3 (8.6)	70 (22.9)	
Other	16 (4.7)	0	17 (5.6)	
No response	38 (11.2)	5 (14.3)	34 (11.1)	
What are the barriers within your current practice which potentially impacts patient outcomes?				

Lack of integrated model of care for complex patients with AF	174 (51.3)	19 (54.3)	156 (51.0)	0.70
Lack of evidence-based guidelines	41 (12.1)	2 (5.7)	39 (12.7)	0.23
Lack of applicability of guidelines to my current practice	31 (9.1)	3 (8.6)	28 (9.2)	0.92
Lack of time	123 (36.3)	16 (45.7)	107 (35.0)	0.21
Organisational/institutional	145 (42.8)	20 (57.1)	125 (40.8)	0.06
Insurance/financial reasons	43 (12.7)	11 (31.4)	33 (10.8)	0.001 ^a
Patient adherence/compliance	126 (37.2)	12 (34.3)	115 (37.6)	0.71
Treatment related adverse events	36 (10.6)	4 (11.4)	32 (10.5)	0.85
Other	21 (6.2)	0	20 (6.5)	0.12

Abbreviations: AF, atrial fibrillation