











Association between walking pace and incident type 2 diabetes by adiposity level: A prospective cohort study from the UK Biobank

Jirapitcha Boonpor MSc^{1,2}  | Solange Parra-Soto MSc^{1,3}  | Jasunella Gore MSc¹ | Atefeh Talebi PhD¹  | Nathan Lynskey MSc¹  | Andrea Raisi MSc^{1,4}  | Paul Welsh PhD¹  | Naveed Sattar MD¹  | Jill P. Pell MD⁵ | Jason M. R. Gill PhD¹ | Stuart R. Gray PhD¹  | Frederick K. Ho PhD⁵  | Carlos A. Celis-Morales PhD^{1,6} 

¹School of Cardiovascular and Metabolic Health, University of Glasgow, Glasgow, UK

²Faculty of Public Health, Kasetsart University, Chalermphrakiat Sakon Nakhon Province Campus, Sakon Nakhon, Thailand

³Department of Nutrition and Public Health, Universidad del Bío-Bío, Chillan, Chile

⁴Center for Exercise Science and Sport, Department of Neuroscience and Rehabilitation, University of Ferrara, Ferrara, Italy

⁵School of Health and Wellbeing, University of Glasgow, Glasgow, UK

⁶Research Group on Education, Physical Activity and Health (GEEAFyS), University Católica del Maule, Talca, Chile

Correspondence

Carlos A. Celis-Morales, PhD, BHF Glasgow Cardiovascular Research Centre, School of Cardiovascular and Metabolic Health, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow G12 8TA, UK. Email: carlos.celis@glasgow.ac.uk

Funding information

Chilean Government PhD scholarship program; Medical Research Council; Northwest Regional Development Agency; Royal Thai Government Scholarship; Scottish Government; Wellcome Trust

Abstract

Aims: To investigate the combined association of adiposity and walking pace with incident type 2 diabetes.

Methods: We undertook a prospective cohort study in 194 304 White-European participants (mean age 56.5 years, 55.9% women). Participants' walking pace was self-reported as brisk, average or slow. Adiposity measures included body mass index (BMI), waist circumference (WC) and body fat percentage (BF%). Associations were investigated using Cox proportional hazard models, with a 2-year landmark analysis. A four-way decomposition analysis was used for mediation and additive interaction.

Results: The median (interquartile range) follow-up was 5.4 (4.8–6.3) years. During the follow-up period, 4564 participants developed type 2 diabetes. Compared to brisk-walking participants with normal BMI, those with obesity who walked briskly were at an approximately 10- to 12-fold higher risk of type 2 diabetes (hazard ratio [HR] 9.64, 95% confidence interval [CI] 7.24–12.84, in women; HR 11.91, 95% CI 8.80–16.12, in men), whereas those with obesity and walked slowly had an approximately 12- to 15-fold higher risk (HR 12.68, 95% CI 9.62–16.71, in women; HR 15.41, 95% CI 11.27–21.06, in men). There was evidence of an additive interaction between WC and BF% and walking pace among women, explaining 17.8% and 47.9% excess risk respectively. Obesity mediated the association in women and men, accounting for 60.1% and 44.9%, respectively.

Conclusions: Slow walking pace is a risk factor for type 2 diabetes independent of adiposity. Promoting brisk walking as well as weight management might be an effective type 2 diabetes prevention strategy given their synergistic effects.

Stuart R. Gray, Frederick K. Ho, Carlos A. Celis-Morales contributed equally and are joint senior authors.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Diabetes, Obesity and Metabolism* published by John Wiley & Sons Ltd.

KEYWORDS

body fat, gait, mediation, obesity, type 2 diabetes mellitus

1 | INTRODUCTION

Type 2 diabetes is a global public health challenge, given that 9.3% of individuals aged 20 to 79 years have known type 2 diabetes worldwide,¹ and global projections estimate that type 2 diabetes prevalence will increase to 10.9% by 2045.² Obesity is one of the most important modifiable risk factors for type 2 diabetes.^{3,4} Since obesity is a relatively stable phenotype with limited effective interventions,⁵ it is important to identify factors that might protect people with obesity from developing type 2 diabetes.

Previous studies have reported that the association of obesity with diabetes risk^{6–8} varies by physical activity level, indicating that people with obesity should consider adopting a higher level of physical activity to manage weight as well as to reduce type 2 diabetes risk.⁸ A recent study that included 117 878 participants from nine prospective studies reported that, compared to normal-weight but highly physically active individuals, those with obesity and had low physical activity levels had a 7.4 times higher risk of type 2 diabetes.⁸ However, most of the evidence is restricted to total physical activity, with limited evidence on other common forms of physical activity, such as walking pace.⁸ Usual walking pace has been identified as a strong predictor of poorer health outcomes.⁹ A recent study has suggested that walking pace could even improve predictive ability in identifying individuals at high risk of cardiovascular disease incidence, beyond that of traditional risk factors.¹⁰ Despite this, current evidence mainly focuses on stratified analysis or inclusion of adiposity as a covariate in statistical analysis.¹¹ No evidence is available on the potential additive association between walking pace combined with obesity and type 2 diabetes risk, or whether such an association differs according to different adiposity markers. Moreover, how much of the association between walking pace and type 2 diabetes is mediated by adiposity has not been addressed.

Understanding the interplay between adiposity and walking pace in type 2 diabetes risk could inform future interventions aiming to increase physical activity or reduce adiposity for type 2 diabetes prevention. The aim of this study, therefore, was to investigate the combined association of adiposity and walking pace with incident type 2 diabetes in a large prospective cohort study and to understand the extent to which the higher risk of type 2 diabetes associated with slow walking may be mediated by adiposity.

2 | MATERIALS AND METHODS

2.1 | Study population

The UK Biobank recruited >502 000 participants from the general population between 2006 and 2010 (5.5% response rate, men and women aged 37–73 years). Participants attended one of 22 assessment centres

across England, Wales and Scotland. At the baseline assessment, an electronically signed consent form and a touchscreen questionnaire were completed. Physical measurements and biological samples were also collected, including blood, urine and saliva.^{12,13} Analyses for the current study were conducted in 194 304 White-European participants from the UK Biobank cohort (age 38–71 years), who had available data records from primary care, exposures, and covariates. Participants with prevalent diabetes (type 1 or type 2 diabetes; $n = 12\,967$) or undiagnosed diabetes ($\text{HbA1c} \geq 48$ mmol/mol; $n = 1589$) at baseline were excluded from the analysis. Additionally, participants from non-White ethnic backgrounds ($n = 8390$) or who had missing data ($n = 11\,199$) were excluded from the study (Figure S1).

2.2 | Adiposity definitions

Anthropometric measurements were measured by trained research nurses using standardized protocols.¹⁴ All anthropometric measurements were taken barefoot and with light clothing. Body composition in this study were measured in a fasting state. Briefly, participants were asked not to eat or drink (except plain water) for up to 4 hours before they visited the assessment centres. They were asked to go to the physical measurement station before going to the sample collection station.¹⁴ Body weight and body fat percentage (BF%) were assessed using a Tanita BC418MA body composition analyser. Height was measured, without shoes, using the wall-mounted SECA 240 height measure. Waist circumference (WC) was measured midway between the lowest rib margin and the iliac crest, horizontally, using a non-elastic SECA 200 tape measure. For WC and BF%, sex-specific tertiles were derived and categorized as low, middle or high (46–92 cm, 78.1–101 cm and 89.1–182 cm for WC; 5%–33.8%, 23.1%–39.8% and 27.9%–69.3% for BF%). BMI was calculated from weight (kg) divided by the square of height (m). BMI was classified as normal weight (18.5–24.9 kg/m²), overweight (25–29.9 kg/m²) or obesity (≥ 30 kg/m²). Underweight was not included in the present analysis. Centrally obese WC was defined as WC ≥ 88 cm for women and ≥ 102 cm for men. High BF% was defined as >35% for women and >25% for men.¹⁵

2.3 | Walking pace assessment

Self-reported usual walking pace was collected at baseline through a touchscreen questionnaire. Participants were asked, “How would you describe your usual walking pace?”. Participants could select any of the three following answers: slow pace (defined as less than 3 miles per hour), steady average pace (defined as 3–4 miles per hour) and brisk pace (defined as more than 4 miles per hour), as described

TABLE 1 Characteristics of the study cohort, stratified by sex

Characteristics	Overall n = 194 304	Women n = 108 631 (55.9%)	Men n = 85 673 (44.1%)
Age, years (mean ± SD)	56.5 ± 8.0	56.4 ± 7.9	56.7 ± 8.1
Townsend deprivation index, n (%)			
Low deprivation	68 023 (35.0)	37 708 (34.7)	30 315 (35.4)
Middle deprivation	67 521 (34.8)	38 073 (35.1)	29 448 (34.4)
High deprivation	58 760 (30.2)	32 850 (30.2)	25 910 (30.2)
Education, n (%)			
College or university degree	73 287 (45.8)	40 073 (45.0)	33 214 (46.9)
A levels/AS levels or equivalent	21 626 (13.5)	12 759 (14.3)	8867 (12.5)
O levels/GCSEs or equivalent	41 902 (26.2)	25 682 (28.8)	16 220 (22.9)
CSEs or equivalent/NVQ or HND or HNC	23 052 (14.4)	10 534 (11.8)	12 518 (17.7)
Lifestyle			
Smoking status, n (%)			
Never	68 023 (35.0)	37 708 (34.7)	30 315 (35.4)
Previous	67 521 (34.8)	38 073 (35.1)	29 448 (34.4)
Current	58 760 (30.2)	32 850 (30.2)	25 910 (30.2)
Sleep duration, h/d (mean ± SD)	7.2 ± 1.1	7.2 ± 1.1	7.1 ± 1.0
Red meat intake, portions/week (mean ± SD)	2.1 ± 1.4	2.0 ± 1.3	2.3 ± 1.5
Process meat intake, portions/week (mean ± SD)	1.9 ± 1.1	1.6 ± 1.0	2.2 ± 1.0
Alcohol intake (mean ± SD)	2.9 ± 1.5	3.1 ± 1.5	2.6 ± 1.4
Fruit and vegetable intake, g/d (mean ± SD)	324.4 ± 185.2	346.3 ± 181.8	296.6 ± 185.8
Total sedentary time, h/d (mean ± SD)	5.0 ± 2.2	4.7 ± 2.0	5.4 ± 2.4
Walking pace, n (%)			
Brisk	80 142 (41.2)	43 901 (40.4)	36 241 (42.3)
Average	101 408 (52.2)	57 181 (52.6)	44 227 (51.6)
Slow	12 754 (6.6)	7549 (7.0)	5205 (6.1)
Adiposity			
Waist circumference, cm (mean ± SD)	89.5 ± 12.9	84.1 ± 11.9	96.3 ± 10.7
Body fat percentage (mean ± SD)	31.6 ± 8.5	36.7 ± 6.7	25.1 ± 5.6
BMI, kg/m ² (mean ± SD)	27.3 ± 4.5	27.0 ± 4.9	27.7 ± 4.0
BMI category, n (%)			
Normal weight	65 092 (33.5)	43 163 (39.7)	21 929 (25.6)
Overweight	84 812 (43.7)	41 241 (38.0)	43 571 (50.9)
Obesity	44 400 (22.9)	24 227 (22.3)	20 173 (23.6)
Multimorbidity, n (%)			
None	69 889 (36.0)	37 889 (34.9)	32 000 (37.4)
1	66 092 (34.0)	36 436 (33.5)	29 656 (34.6)
2	35 854 (18.5)	20 425 (18.8)	15 429 (18.0)
3	14 846 (7.6)	8902 (8.2)	5944 (6.9)
4	5200 (2.7)	3315 (3.1)	1885 (2.2)
≥5	2423 (1.2)	1664 (1.5)	759 (0.9)

Note: Data are presented as mean and SD for continuous variables and frequency and % for categorical variables.

Abbreviations: A levels/AS levels, Advanced/Advanced Subsidiary Levels; BMI, body mass index; CSEs, Certificate of Secondary Education; GCSE, General Certificate of Secondary Education; HNC, Higher National Certificate; HND, Higher National Diploma; NVQ, National Vocational Qualification; O levels, ordinary levels.

associations were examined using penalized cubic splines fitted in Cox proportional hazard models. The penalized spline is not as sensitive to knot numbers and placements as restricted cubic splines.^{23,24} We used

likelihood ratio tests to test nonlinearity in exposure-outcome relationships by comparing models with adiposity splines and models with linear adiposity terms.

TABLE 2 Association of walking pace and adiposity with incident type 2 diabetes

	Women				Men			
	Model 1		Model 2		Model 1		Model 2	
	HR (95% CI)	P value	HR (95% CI)	P value	HR (95% CI)	P value	HR (95% CI)	P value
Walking pace								
Brisk	1.00 (ref.)		1.00 (ref.)		1.00 (ref.)		1.00 (ref.)	
Average	1.75 (1.55; 1.96)	<0.0001	1.21 (1.08; 1.37)	0.002	1.60 (1.45; 1.75)	<0.0001	1.21 (1.10; 1.33)	<0.0001
Slow	2.52 (2.15; 2.94)	<0.0001	1.10 (0.94; 1.30)	0.242	2.05 (1.78; 2.35)	<0.0001	1.18 (1.02; 1.36)	0.024
Trend	1.59 (1.48; 1.72)	<0.0001	1.06 (0.98; 1.15)	0.160	1.46 (1.37; 1.56)	<0.0001	1.11 (1.04; 1.19)	0.002
BMI								
Normal weight	1.00 (ref.)		1.00 (ref.)		1.00 (ref.)		1.00 (ref.)	
Overweight	3.43 (2.86; 4.12)	<0.0001	3.36 (2.80; 4.02)	<0.0001	3.54 (2.95; 4.24)	<0.0001	3.48 (2.91; 4.17)	<0.0001
Obesity	10.05 (8.44; 11.97)	<0.0001	9.44 (7.91; 11.27)	<0.0001	10.12 (8.47; 12.11)	<0.0001	9.62 (8.04; 11.52)	<0.0001
Trend	3.07 (2.86; 3.30)	<0.0001	2.97 (2.76; 3.20)	<0.0001	3.01 (2.82; 3.21)	<0.0001	2.92 (2.73; 3.11)	<0.0001
WC								
Low WC	1.00 (ref.)		1.00 (ref.)		1.00 (ref.)		1.00 (ref.)	
Middle WC	3.48 (2.80; 4.33)	<0.0001	3.41 (2.74; 4.24)	<0.0001	2.46 (2.14; 2.82)	<0.0001	2.42 (2.1; 2.78)	<0.0001
High WC	12.16 (9.91; 14.91)	<0.0001	11.44 (9.32; 14.06)	<0.0001	6.53 (5.74; 7.42)	<0.0001	6.24 (5.48; 7.10)	<0.0001
Trend	3.49 (3.22; 3.78)	<0.0001	3.37 (3.11; 3.66)	<0.0001	2.59 (2.44; 2.74)	<0.0001	2.52 (2.38; 2.67)	<0.0001
BF%								
Low BF%	1.00 (ref.)		1.00 (ref.)		1.00 (ref.)		1.00 (ref.)	
Middle BF%	3.14 (2.56; 3.84)	<0.0001	3.05 (2.49; 3.74)	<0.0001	2.77 (2.39; 3.21)	<0.0001	2.72 (2.34; 3.16)	<0.0001
High BF%	7.91 (6.53; 9.59)	<0.0001	7.30 (6.01; 8.87)	<0.0001	6.47 (5.62; 7.45)	<0.0001	6.15 (5.34; 7.09)	<0.0001
Trend	2.69 (2.49; 2.90)	<0.0001	2.57 (2.38; 2.78)	<0.0001	2.47 (2.33; 2.62)	<0.0001	2.40 (2.26; 2.55)	<0.0001

Note: Data are presented as HR and 95% CI. Model 1 was adjusted for age, deprivation index, education, smoking status, fruit and vegetable, red meat and processed meat intake, alcohol consumption, total sedentary time, sleep duration and multimorbidity. Model 2 was adjusted as for Model 1 but further included walking pace when BMI, WC and BF% were the exposures and vice versa. All analyses were conducted using 2-year landmark analyses, excluding participants with type 1, type 2 diabetes, unknown diabetes and non-White ethnic background.

Abbreviations: BF%, body fat percentage; BMI, body mass index; CI, confidence interval; HR, hazard ratio; WC, waist circumference.

Statistical analyses were performed using the statistical software STATA 17 (StataCorp LP) and R v4.0.2. Significance was accepted at a P value < 0.05. The Anderson-Darling test was conducted to check the normality of variables.

2.7 | Ethics statement

The UK Biobank study was approved by the North-West Multi-Centre Research Ethics Committee (ref. 11/NW/0382 on June 17, 2011). All participants provided written informed consent to participate in the UK Biobank study. The study protocol is available online (<http://www.ukbiobank.ac.uk/>). This research has been conducted using the UK Biobank resource under application number 7155.

3 | RESULTS

In the 2-year landmark analyses and after excluding participants with prevalent diabetes, undiagnosed diabetes or non-White Europeans at

baseline, 194 304 participants with data available for incident type 2 diabetes, adiposity, walking pace and covariates were included in this study (Figure S1). The median (interquartile range) follow-up period was 5.4 (4.8-6.3) years. Over the follow-up, 4564 participants (2.3%) developed type 2 diabetes, including 1916 women (42.0%) and 2648 men (58.0%).

There was a high correlation between BMI and WC ($r = 0.81$) but a moderate to low correlation between BMI and BF% ($r = 0.57$) as well as WC and BF% ($r = 0.24$).

Table 1 shows the cohort characteristics of the men and women included in this study. In summary, the mean \pm SD age was 56.5 \pm 8.0 years, with 56.4% women. A higher proportion of the participants reported having college or university degrees and being never-smokers. Men, compared to women, reported a higher intake of red and processed meat but lower alcohol consumption and fruit and vegetable intake. A higher prevalence of overweight was observed for men compared to women (Table 1).

The associations between walking pace and incident type 2 diabetes for women and men are presented in Table 2. Compared to a brisk walking pace, women who reported either average or slow walking




FIGURE 1 Combined association between adiposity and walking pace and incident type 2 diabetes. Data are presented as hazard ratio (HR) and 95% confidence interval (CI). Analyses were adjusted for age, deprivation index, education, smoking status, fruit and vegetable intake, red meat intake, processed meat intake, alcohol consumption, total sedentary time, sleep duration and multimorbidity. All analyses were conducted using 2-year landmark analyses and excluding participants with type 1, type 2 diabetes, unknown diabetes and non-White ethnic background. A, Combined association between body mass index and walking pace and incident type 2 diabetes. B, Combined association between waist circumference (WC) and walking pace and incident type 2 diabetes. C, Combined association between body fat percentage and walking pace and incident type 2 diabetes

paces had a 75% (HR 1.75 [95% CI 1.55; 1.96]) or 2.5 times (HR 2.52 [95% CI 2.15; 2.94]) higher risk of type 2 diabetes, respectively, after adjusting for sociodemographic and lifestyle confounding factors (Model 1). When the analysis was further adjusted for adiposity, the risk of type 2 diabetes was attenuated for slow walking pace,

therefore, mediation analyses were conducted to test if adiposity mediated the association. In men, compared to a brisk walking pace, after adjusting for sociodemographic and lifestyle factors (Model 1), individuals who reported average and slow walking paces had a 60% (HR 1.60 [95% CI 1.45; 1.75]) and 2.1 times (HR 2.05 [95% CI 1.78;

TABLE 3 Four-way decomposition analysis of walking pace and adiposity with type 2 diabetes

Proportion of excess risk of slow walking due to	General obesity	Central obesity	High body fat percentage
Women			
Additive interaction (%)	1.6 (−6.3; 6.8)	17.8 (9.4; 23.8)*	47.9 (34.2; 58.0)*
Total mediation (%)	60.1 (51.8; 69.7)*	64.4 (57.2; 73.4)*	44.9 (38.2; 50.8)*
Men			
Additive interaction (%)	2.7 (−8.3; 10.3)	6.7 (−5.7; 15.2)	25.6 (12.3; 36.7)*
Total mediation (%)	44.9 (36.9; 52.3)*	44.7 (37.2; 52.9)*	35.5 (23; 42.7)*

Note: Numbers presented are percentages with 95% confidence interval. The analysis was adjusted for age, deprivation index, education, smoking status, fruit and vegetable intake, red meat intake, processed meat intake, alcohol consumption, total sedentary time, sleep duration and multimorbidity. All analyses were conducted using 2-year landmark analyses, excluding participants with type 1, type 2 diabetes, unknown diabetes and non-White ethnic background. Obesity was defined as BMI ≥ 30 kg/m²; central obesity was defined as WC ≥ 88 cm for women and ≥ 102 cm for men; BF% was defined as $>35\%$ for women and $>25\%$ for men.

Abbreviations: BF%, body fat percentage; BMI, body mass index; WC, waist circumference.

* $P < 0.0001$.

2.35]) higher risk of type 2 diabetes, respectively. The magnitude of risk of type 2 diabetes was attenuated but still significant after further adjusting for adiposity (Model 2) to a 21% (HR 1.21 [95% CI 1.10; 1.33]) and 18% (HR 1.18 [95% CI 1.02; 1.36]) higher risk, respectively. The magnitudes and directions of the associations between walking pace, adiposity and type 2 diabetes were similar when medication was added to the most adjusted model (Model 2; Tables S1).

Adiposity was positively associated with a higher risk of type 2 diabetes regardless of the adiposity marker used. Compared to normal weight as determined by BMI, both women and men who were living with overweight or obesity had a three or nine times higher risk of type 2 diabetes. For WC, women and men who were in the high WC category had an 11.4 times (HR 11.44 [95% CI 9.32; 14.06]) and 6.2 times (HR 6.24 [95% CI 5.48; 7.10]) higher risk of type 2 diabetes, respectively, compared to the lowest category. For BF%, women and men who were in the high BF% category had a 7.3 times (HR 7.30 [95% CI: 6.01; 8.87]) and 6.2 times (HR 6.15 [95% CI: 5.34; 7.09]) higher risk of type 2 diabetes, respectively, compared to the low category, as shown in Table 2.

Figure 1 shows the combined association between adiposity, walking pace, and incident type 2 diabetes. The risk of type 2 diabetes was higher when walking pace was slow. Among women, participants with obesity, central obesity or had high BF% and also reported a slow walking pace had a 12.7 times (HR 12.68 [95% CI 9.62; 16.71]), 17.9 times (HR 17.85 [95% CI 12.87; 24.78]) or 10.5 times (HR: 10.52 [95% CI 7.84; 14.10]) higher risk of developing type 2 diabetes compared to those with normal BMI, low WC or low BF% with brisk walking pace, respectively. Among men, participants with obesity, central obesity or had high BF% and also reported a slow pace had a 15.4 times (HR 15.41 [95% CI 11.27; 21.06]), 9.8 times (HR 9.82 [95% CI 7.81; 12.36]) or 10.2 times (HR 10.15 [95% CI 7.95; 12.96]) higher risk of developing type 2 diabetes compared to those with normal BMI, low WC or low BF% with brisk walking pace, respectively. Similar results were found when medication was added to the model as a covariate (Figure S2).

General and central obesity were partial mediators for the association in women, accounting for 60.1% and 64.4%, respectively. There was evidence of an additive interaction of WC and body fat with

walking pace in women; however, only body fat showed an interaction with walking pace in men (Table 3). The mediation analysis showed that BF% (Table 3) mediated 44.9% (for women) and 35.5% (for men) of the excess risk due to slow walking. Being a slow walker and having a high BF% would have a synergistic deleterious association, explaining another 47.9% and 26.5% of the excess risk for women and men, respectively. The total effect, total natural indirect effect and pure natural direct effect are shown in Table S2.

Positive nonlinear associations were found between adiposity, walking pace and incident type 2 diabetes (Figure 2). Compared to brisk walking pace, the spline for slow walking pace showed that type 2 diabetes risk increased with increasing BMI up to 33 kg/m², after which the lower risk appeared. For WC, the risk of incident type 2 diabetes for slow walking pace increased sharply with greater WC up to 103 cm, then converged. The spline for slow walking pace showed that the risk of incident type 2 diabetes seemed to increase sharply with increasing BF% up to 36%, then converge.

4 | DISCUSSION

Compared to brisk walking pace, slow walking pace was associated with a higher risk of type 2 diabetes, independent of sociodemographic and lifestyle factors among White-European people. This association was largely explained by adiposity in women but not in men. The risk of type 2 diabetes was doubled when slow walking pace was combined with obesity, high WC or BF%. These findings suggest that both walking pace and adiposity are independent risk factors for type 2 diabetes and that they have to be considered simultaneously to achieve the maximum public health benefits. Importantly, as adiposity could be a mediator between walking pace and type 2 diabetes, promoting brisk walking for weight management could have synergistic benefits for type 2 diabetes prevention. That noted, it should be clear that excess adiposity, however measured, was the dominant risk factor for type 2 diabetes so that brisk walking on its own is unlikely to be sufficient to prevent type 2 diabetes if there are other reasons, for example, poor diet, that lead to weight excess.




FIGURE 2 Penalized cubic splines for the association between walking pace, adiposity and incident type 2 diabetes. Data are presented as hazard ratios and their 95% confidence interval (CI). Analyses were adjusted for age, deprivation index, education, smoking status, fruit and vegetable intake, red meat intake, processed meat intake, alcohol consumption, total sedentary time, sleep duration and multimorbidity. All analyses were conducted using 2-year landmark analyses and excluding participants with type 1, type 2 diabetes, unknown diabetes and non-white ethnic background. BF%, body fat percentage; BMI, body mass index; WC, waist circumference

Our study is consistent with previous evidence on walking pace and the risk of type 2 diabetes. A study from the United States, conducted in 70 102 women aged 40 to 60 years, found that after adjusting for age, time period, cigarette smoking, menopausal status, parental history of

diabetes, alcohol consumption, history of hypertension, history of high cholesterol level and time spent walking per week, normal and brisk walking pace were associated with a 28% (relative risk [RR] 0.72 [95% CI 0.62; 0.85]) and 59% (RR 0.41 [95% CI 0.33; 0.52]) lower risk of type 2 diabetes,

respectively, compared to an easy walking pace.²⁵ However, after adjusting for BMI, the association for brisk walkers was slightly attenuated (RR 0.59 [95% CI 0.47; 0.73]) but remained significant.²⁵ The Health Professional's Follow-up Study, conducted in the United States in 37 918 men aged 40 to 75 years, reported that, compared to slow walkers, those who reported a normal, brisk or very brisk walking pace had a 32%, 54% and 61% lower risk of type 2 diabetes independent of BMI.²⁶

Current evidence mainly focuses on stratified analysis or including adiposity as a covariate in the statistical analysis.¹¹ However, some studies have reported the joint associations between total physical activity and BMI with type 2 diabetes risk.^{7,8} A more recent meta-analysis reported a seven times higher risk of type 2 diabetes in people with obesity and who had low physical activity compared to those with normal weight and high physical activity.⁸ Likewise, our findings showed that the combination of obesity and slow walking pace was associated with an approximately 10 times higher risk of type 2 diabetes, compared to normal weight and brisk walking pace.

Interestingly, four-way decomposition analysis in our findings showed the excess risk due to slow walking was mainly mediated through general and central obesity but had an additive interaction with body fat proportion. This indicates that people with high body fat are more vulnerable to slow walking and that increasing their walking speed during physical activity, where feasible, could potentially be effective as it could reduce body fat. Randomized controlled trials are warranted to test these hypotheses.

Our study is an observational study, so we cannot interpret causality directly. However, the association is biologically plausible. Walking, as an aerobic exercise, can improve insulin action in the muscles and liver by increasing glucose uptake acutely and during prolonged activity.²⁷ Furthermore, a previous Mendelian randomization study has suggested walking pace to be a potential causal factor for various health outcomes.²⁸ In contrast, it is well known that obesity is the strongest risk factor for type 2 diabetes.²⁹ Adiposity composition and sex hormones might explain why some of the findings differed by sex.^{30,31}

The strengths of the present study include its large number of participants, which provided a sufficient sample size to undertake the analysis. We included multiple confounding factors, which enabled the research question to be sufficiently adjusted and investigated. A 2-year landmark analysis likely reduces the risk of reverse causation impacting the results and is a strength of this study. Walking pace is a low cost equipment and is easy to administer and would therefore be relatively simple to assess in clinical practice. Although our study used self-reported usual walking pace, there is evidence that self-reported walking speed is a valuable marker of walking speed measurement and could be a marker of physical performance when walking speed is not possible to measure directly in sarcopenia and frailty.³² Self-reporting walking pace does not consider the time spent walking, which could modify the association between walking pace and obesity in type 2 diabetes. However, a previous study has shown that the association between walking pace and diabetes risk is independent of time spent walking.³³ It is well known that BMI has limitations when used to assess obesity,³ but that WC is a better predictor for evaluating the risk of type

2 diabetes.³⁴ Our study provided evidence that adiposity and fat distribution, BMI, WC or BF%, combined with a measure of physical activity such as walking pace, showed a strong relationship to type 2 diabetes risk.

The present study also has several limitations. The UK Biobank is not representative of the general population of the United Kingdom with regard to sociodemographic, physical, lifestyle and health-related characteristics. Although the UK Biobank data have been showed to have healthy volunteer selection bias, exposure-disease risk estimates can nevertheless be generalized to the broader population.³⁵ In addition, walking pace was self-reported, however, previous studies have provided robust evidence of the predictive ability of self-reported walking pace above and beyond traditional risk factors.^{9,10} We were also unable to include all ethnicities as there was insufficient statistical power to detect differences according to ethnicity. Although standardized protocols were implemented across the assessment centres, the data do not allow us to quantify the accuracy and repeatability of the methods used. Reverse causation due to the link between existing multimorbidity and walking pace cannot be ignored; however, when the analyses were adjusted for multimorbidity or medication, the associations remained similar. Although these sensitivity analyses cannot fully remove the effect of reverse causation, they provide evidence that reverse causation does not fully explain the associations observed.

To conclude, walking pace was inversely associated with the risk of type 2 diabetes for both men and women, independent of sociodemographic characteristics, lifestyle and adiposity. Slow walking pace, combined with obesity, high WC and BF%, doubled the risks of developing type 2 diabetes. Promoting brisk walking as a weight management measure might be an effective type 2 diabetes prevention strategy, given its synergistic effect.

AUTHOR CONTRIBUTIONS

Jirapitcha Boonpor, Stuart R. Gray, Frederick K. Ho and Carlos Celis-Morales contributed to the study conception and design, advised on all statistical aspects, and interpreted the data. Jirapitcha Boonpor, Solange Parra-Soto, Atefeh Talebi, Frederick K. Ho and Carlos Celis-Morales performed the statistical analyses. Jirapitcha Boonpor, Stuart R. Gray, Frederick K. Ho and Carlos Celis-Morales drafted the manuscript. All authors reviewed the manuscript and approved the final version to be published. Stuart R. Gray, Frederick K. Ho and Carlos Celis-Morales are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

ACKNOWLEDGMENTS

We are grateful to the UK Biobank participants. This research was conducted using the UK Biobank resource under application number 7155.

FUNDING INFORMATION

The UK Biobank was established by the Wellcome Trust, Medical Research Council, Department of Health, Scottish Government, and

the Northwest Regional Development Agency. It has also received funding from the Welsh Assembly Government and the British Heart Foundation. Jirapitcha Boonpor receives financial support from the Royal Thai Government Scholarship for her PhD. Solange Parra-Soto receives financial support from the Chilean Government PhD scholarship program for their PhD. The funders have no role in study design, in the collection, analysis or interpretation of data, in the report's writing, or in the decision to submit the paper for publication.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

PEER REVIEW

The peer review history for this article is available at <https://www.webofscience.com/api/gateway/wos/peer-review/10.1111/dom.15053>.

DATA AVAILABILITY STATEMENT

Data can be requested from the UK Biobank (<https://www.ukbiobank.ac.uk/>).

ORCID

Jirapitcha Boonpor  <https://orcid.org/0000-0002-9548-5690>

Solange Parra-Soto  <https://orcid.org/0000-0002-8443-7327>

Andrea Raisi  <https://orcid.org/0000-0002-0096-4534>

Paul Welsh  <https://orcid.org/0000-0002-7970-3643>

Naveed Sattar  <https://orcid.org/0000-0002-1604-2593>

Stuart R. Gray  <https://orcid.org/0000-0001-8969-9636>

Frederick K. Ho  <https://orcid.org/0000-0001-7190-9025>

Carlos A. Celis-Morales  <https://orcid.org/0000-0003-2612-3917>

REFERENCES

- International diabetes federation. IDF Diabetes atlas ninth edition. 2019 www.diabetesatlas.org
- Saeedi P, Petersohn I, Salpea P, et al. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: results from the international diabetes federation diabetes atlas, 9 (th) edition. *Diabetes Res Clin Pract.* 2019;157:107843.
- Piché M-E, Tchernof A, Després J-P. Obesity phenotypes, diabetes, and cardiovascular diseases. *Circ Res.* 2020;126(11):1477-1500.
- Verma S, Hussain ME. Obesity and diabetes: an update. *Diabetes Metab Syndr.* 2017;11(1):73-79.
- Chan RS, Woo J. Prevention of overweight and obesity: how effective is the current public health approach. *Int J Environ Res Public Health.* 2010;7(3):765-783.
- Hu G, Lindström J, Valle TT, et al. Physical activity, body mass index, and risk of type 2 diabetes in patients with Normal or impaired glucose regulation. *Arch Intern Med.* 2004;164(8):892-896.
- Weinstein AR, Sesso HD, Lee IM, et al. Relationship of physical activity vs body mass index with type 2 diabetes in women. *JAMA.* 2004;292(10):1188-1194.
- Cloostermans L, Wendel-Vos W, Doornbos G, et al. Independent and combined effects of physical activity and body mass index on the development of type 2 diabetes – a meta-analysis of 9 prospective cohort studies. *Int J Behav Nutr Phys Act.* 2015;12(1):147.
- Ganna A, Ingelsson E. 5 year mortality predictors in 498 103 UK Biobank participants: a prospective population-based study. *Lancet.* 2015;386(9993):533-540.
- Welsh CE, Celis-Morales CA, Ho FK, et al. Grip strength and walking pace and cardiovascular disease risk prediction in 406,834 UK Biobank participants. *Mayo Clin Proc.* 2020;95(5):879-888.
- Iwasaki M, Kudo A, Asahi K, et al. Fast walking is a preventive factor against new-onset diabetes mellitus in a large cohort from a Japanese general population. *Sci Rep.* 2021;11(1):716.
- Collins R. What makes UK Biobank special? *Lancet.* 2012;379(9822):1173-1174.
- Sudlow C, Gallacher J, Allen N, et al. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med.* 2015;12(3):e1001779.
- UK Biobank. UK Biobank: protocol for a large-scale prospective epidemiological resource. 2007; <http://www.ukbiobank.ac.uk/wp-content/uploads/2011/11/UK-Biobank-Protocol.pdf>. Accessed July, 2020.
- Hardman AE, Stensel DJ. Obesity and energy balance. *Physical Activity and Health: the Evidence Explained.* Vol 1. London and New York: Routledge; 2003:114-130.
- Celis-Morales CA, Gray S, Petermann F, et al. Walking pace is associated with lower risk of all-cause and cause-specific mortality. *Med Sci Sports Exerc.* 2019;51(3):472-480.
- Townsend NP, Phillimore P, Beattie A. *Health and Deprivation: Inequality and the North.* London: Croom Helm Ltd; 1988.
- VanderWeele TJ. A unification of mediation and interaction: a 4-way decomposition. *Epidemiology.* 2014;25(5):749-761.
- Tingley D, Yamamoto T, Hirose K, Keele L, Imai K. Mediation: R package for causal mediation analysis. *J Stat Softw.* 2014;59(5):1-38.
- VanderWeele TJ, Tchetgen Tchetgen EJ. Mediation analysis with time varying exposures and mediators. *J R Stat Soc Ser B Stat Methodol.* 2017;79(3):917-938.
- Boonpor J, Petermann-Rocha F, Parra-Soto S, et al. Types of diet, obesity, and incident type 2 diabetes: findings from the UK Biobank prospective cohort study. *Diabetes Obes Metab.* 2022;24(7):1351-1359.
- Shi B, Choirat C, Coull BA, VanderWeele TJ, Valeri L. CMAverse: a suite of functions for reproducible causal mediation analyses. *Epidemiology.* 2021;32(5):e20-e22.
- Govindarajulu US, Malloy EJ, Ganguli B, Spiegelman D, Eisen EA. The comparison of alternative smoothing methods for fitting non-linear exposure-response relationships with cox models in a simulation study. *Int J Biostat.* 2009;5(1):Article 2.
- Ho, FK, Cole TJ. Non-linear predictor outcome associations. *BMJ Medicine,* 2023;2(1):e000396. doi:10.1136/bmjmed-2022-000396
- Hu FB, Sigal RJ, Rich-Edwards JW, et al. Walking compared with vigorous physical activity and risk of type 2 diabetes in women a prospective study. *JAMA.* 1999;282(15):1433-1439.
- Hu FB, Leitzmann MF, Stampfer MJ, Colditz GA, Willett WC, Rimm EB. Physical activity and television watching in relation to risk for type 2 diabetes mellitus in men. *Arch Intern Med.* 2001;161(12):1542-1548.
- Colberg SR, Sigal RJ, Yardley JE, et al. Physical activity/exercise and diabetes: a position Statement of the American Diabetes Association. *Diabetes Care.* 2016;39(11):2065-2079.
- Timmins IR, Zaccardi F, Nelson CP, Franks PW, Yates T, Dudbridge F. Genome-wide association study of self-reported walking pace suggests beneficial effects of brisk walking on health and survival. *Commun Biol.* 2020;3(1):634.
- Carbone S, Del Buono MG, Ozemek C, Lavie CJ. Obesity, risk of diabetes and role of physical activity, exercise training and cardiorespiratory fitness. *Prog Cardiovasc Dis.* 2019;62(4):327-333.
- Mauvais-Jarvis F. Epidemiology of gender differences in diabetes and obesity. In: Mauvais-Jarvis F, ed. *Sex and Gender Factors Affecting Metabolic Homeostasis, Diabetes and Obesity.* Cham: Springer International Publishing; 2017:3-8.

31. Sattar N. Type 2 diabetes-related sex differences in cardiovascular risk: reasons, ramifications, and clinical realities. *Eur Heart J*. 2020; 41(13):1354-1356.
32. Syddall HE, Westbury LD, Cooper C, Sayer AA. Self-reported walking speed: a useful marker of physical performance among community-dwelling older people? *J Am Med Dir Assoc*. 2015;16(4):323-328.
33. Boonpor J, Ho FK, Gray SR, Celis-Morales CA. Association of Self-reported Walking Pace with Type 2 diabetes incidence in the UK Biobank prospective cohort study. *Mayo Clin Proc*. 2022;97(9):1631-1640.
34. Schulze MB, Thorand B, Fritsche A, et al. Body adiposity index, body fat content and incidence of type 2 diabetes. *Diabetologia*. 2012; 55(6):1660-1667.
35. Fry A, Littlejohns TJ, Sudlow C, et al. Comparison of sociodemographic and health-related characteristics of UK Biobank participants with those of the general population. *Am J Epidemiol*. 2017;186(9):1026-1034.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Boonpor J, Parra-Soto S, Gore J, et al. Association between walking pace and incident type 2 diabetes by adiposity level: A prospective cohort study from the UK Biobank. *Diabetes Obes Metab*. 2023;25(7):1900-1910. doi:[10.1111/dom.15053](https://doi.org/10.1111/dom.15053)