

LEVEL SPACING STATISTICS FOR THE MULTI-DIMENSIONAL QUANTUM HARMONIC OSCILLATOR: ALGEBRAIC CASE

ALAN HAYNES, ROLAND ROEDER

ABSTRACT. We study the statistical properties of the spacings between neighboring energy levels for the multi-dimensional quantum harmonic oscillator that occur in a window $[E, E + \Delta E)$ of fixed width ΔE as E tends to infinity. This regime provides a notable exception to the Berry-Tabor Conjecture from Quantum Chaos and, for that reason, it was studied extensively by Berry and Tabor in their seminal paper from 1977. We focus entirely on the case that the (ratios of) frequencies $\omega_1, \omega_2, \dots, \omega_d$ together with 1 form a basis for an algebraic number field Φ of degree $d + 1$, allowing us to use tools from algebraic number theory. This special case was studied by Dyson, Bleher, Bleher-Homma-Ji-Roeder-Shen, and others. Under a suitable rescaling, we prove that the distribution of spacings behaves asymptotically quasiperiodically in $\log E$. We also prove that the distribution of ratios of neighboring spacings behaves asymptotically quasiperiodically in $\log E$. The same holds for the distribution of finite words in the finite alphabet of rescaled spacings.

Mathematically, our work is a higher dimensional version of the Steinhaus Conjecture (Three Gap Theorem) involving the fractional parts of a linear form in more than one variable, and it is of independent interest from this perspective.

1. INTRODUCTION

We study the statistical properties of spacings (also called gaps and nearest neighbor distances) between neighboring energy levels for the multi-dimensional quantum harmonic oscillator in the limit as the energy tends to infinity. This is a fundamental problem from physics whose motivations date back to the origins of the study of Quantum Chaos.

Our main techniques are from algebraic number theory and to make these techniques accessible we will focus entirely on the case that the (ratios of) frequencies $\omega_1, \omega_2, \dots, \omega_d$ together with 1 form a basis for an algebraic number field Φ of degree $d + 1$. (Here the dimension of the harmonic oscillator is $d + 1$.) This is a rather special case, however it is also very important because of the emphasis placed on it by previous works of Berry and Tabor [2], Dyson [14], Bleher [4], and others.

Mathematically, the problem reduces to studying the fractional parts of a linear form in more than one variable, which is a topic of independent interest in number theory. Study of this topic provides multi-dimensional analogs and extensions of the famous Three Gap Theorem (also called the Steinhaus Conjecture) of Sós [26], Surányi [27], and Świerczkowski [28].

In order to reach readers from both of these perspectives we will now describe the motivations, background, and also state our results in both contexts.

1.1. Physical Perspective. Given a quantum-mechanical system (or equivalently a Hamiltonian H), one of the central problems from Quantum Chaos is to study the distribution of spacings between neighboring energy levels in the limit as the energy E tends to infinity. One typically

Date: July 12, 2021.

Key words and phrases. Quantum Harmonic Oscillator, Energy Level Statistics, Steinhaus problem, Three Gap Theorem.

MSC 2020: 81Q50, 11J13, 11J17.

This is the author's manuscript of the article published in final edited form as:

considers all energy levels in a window

$$E \leq e_1 < e_2 < \dots < e_\ell < E + \Delta E,$$

where the width ΔE is fixed and one takes the limit as $E \rightarrow \infty$. In this way, the ordered energy levels $e_1(E) < e_2(E) < \dots < e_{\ell(E)}(E)$ all depend on the energy $E > 0$ as does the number $\ell \equiv \ell(E)$ of them. One is interested in the statistics of the distances between neighboring energy levels

$$\delta_i(E) := e_{i+1}(E) - e_i(E) \quad \text{for } 1 \leq i \leq \ell(E) - 1,$$

and their fluctuations in the limit as $E \rightarrow \infty$. As E increases, the size of these spacings decrease. However, one can multiply by a factor $\lambda(E)$ in order to produce normalized spacings

$$\bar{\delta}_i(E) := \lambda(E)\delta_i(E) = \lambda(E)(e_{i+1}(E) - e_i(E)) \quad \text{for } 1 \leq i \leq \ell(E) - 1,$$

where $\lambda(E)$ is chosen so that the normalized spacings $\bar{\delta}_i(E)$ have average length equal to 1. This allows one to ask whether the normalized spacings are described by a limiting law and to ask how that law depends on the nature of the system (Hamiltonian H) being considered. We have:

Conjecture of Berry and Tabor (1977) [2]:

If the classical dynamical system associated to the Hamiltonian H is integrable, then the limiting distribution of the normalized spacings for quantum energy levels is governed by a Poisson law.

Conjecture of Bohigas, Giannoni, and Schmit (1984) [8]:

If the classical dynamical system associated to the Hamiltonian H is chaotic, then the limiting distribution of the normalized spacings for the quantum energy levels is governed by the eigenvalue statistics of one of the three standard ensembles of random matrices, GOE, GUE, or GSE.

For more information on these conjectures and various exciting extensions of them to other contexts we recommend [19, 6].

As discussed in the paper of Berry and Tabor [2], the multi-dimensional quantum harmonic oscillator serves as a notable exception to these conjectures. Even though the classical multi-dimensional harmonic oscillator is completely integrable, the asymptotic behavior of the normalized spacings distributions is not governed by a Poisson law because the energy level contours are flat. Rather, it depends dramatically on the arithmetic properties of the ratio of frequencies. A considerable portion of the paper by Berry and Tabor is devoted to this situation. This was followed by the works of Dyson [14], Boshernitzan [9, 10], Bleher [4, 5], and others.

The multi-dimensional quantum harmonic oscillator is given by the Hamiltonian

$$H = - \sum_{j=1}^{d+1} \frac{\hbar^2}{2m} \frac{\partial^2}{\partial x_j^2} + \sum_{j=1}^{d+1} \frac{k_j}{2} x_j^2.$$

Applying Schrödinger's equation, the quantum energy levels of the system are determined by $d+1$ non-negative integers, m_0, \dots, m_d , and they are of the form

$$e = e_0 + m_0\alpha_0 + \dots + m_d\alpha_d,$$

where $\alpha_0, \dots, \alpha_d$ are positive real numbers depending on the spring constant \mathbf{k} and the mass m . We have

$$e = e_0 + \alpha_0(m_0 + m_1\omega_1 + \dots + m_d\omega_d),$$

where $\omega_i = \alpha_i/\alpha_0 > 0$. We will refer to $\omega_1, \dots, \omega_d$ the *ratios of frequencies*.

It is convenient to set $\Delta E = \alpha_0$ and to perform a linear change of variables to the energy scale, letting $\mathcal{E} = (E - e_0)/\alpha_0$, $\varepsilon = (e - e_0)/\alpha_0$, and hence $\Delta\mathcal{E} = 1$. Then, the problem becomes to find spacings between each of the neighboring energy levels

$$\varepsilon = m_0 + m_1\omega_1 + \dots + m_d\omega_d$$

lying between a given \mathcal{E} and $\mathcal{E} + 1$ as $\mathcal{E} \rightarrow \infty$. All of our results will be described under this change of variables.

The case $d = 1$ corresponds to a two-dimensional harmonic oscillator and it was carefully studied by Bleher [4, 5], using methods involving continued fractions. He proved in [5, Theorem 1.5] that for a generic ratio of the frequencies $\omega \equiv \omega_1$ there is no limiting distribution of normalized spacings. Meanwhile in the special case that the ratio of frequencies is the golden mean $\omega = (\sqrt{5}-1)/2$, Bleher explicitly describes how the distribution of normalized spacings depends on \mathcal{E} ; see [4, Theorem 3]. From these formulae one can see that the distribution behaves periodically in $\log \mathcal{E}$, asymptotically as $\mathcal{E} \rightarrow \infty$. In other words, although there is no limit of the distribution of normalized spacings, the asymptotic behavior remains rather simple.

Later it was proved by Bleher-Homma-Ji-Roeder-Shen in [7, Theorem 1.6] that this phenomenon carries over to the three-dimensional quantum harmonic oscillator when the ratios of frequencies satisfy that $1, \omega_1, \omega_2$ form a basis for a cubic algebraic number field Φ that has one fundamental unit. Rather than using continued fractions, which do not generalize to higher dimensions, basic techniques from algebraic number theory were used to prove that the distribution of normalized spacings behaves asymptotically quasi-periodically in $\log \mathcal{E}$ as $\mathcal{E} \rightarrow \infty$.

In this paper we build on these previous results to prove:

Theorem A. *Suppose that the ratios of frequencies satisfy that $1, \omega_1, \dots, \omega_d \in \mathbb{R}$ form a \mathbb{Q} -basis for an algebraic number field Φ . We then have:*

- (1) **Uniform Labeling:** *There is a finite set $\mathcal{S} := \{s_1, \dots, s_J\} \subset \Phi$ and a rescaling factor $u(\mathcal{E}) > 1$ such that for any $\mathcal{E} > 1$ the rescaled spacings satisfy:*

$$\bar{\delta}_i(\mathcal{E}) := u(\mathcal{E})(\varepsilon_{i+1}(\mathcal{E}) - \varepsilon_i(\mathcal{E})) \in \mathcal{S} \quad \text{for } 1 \leq i \leq \ell(\mathcal{E}) - 1.$$

- (2) **Quasiperiodicity:** *For each $1 \leq j \leq J$ let $p_j(\mathcal{E})$ denote the proportion of the rescaled spacings $\bar{\delta}_i(\mathcal{E})$ such that $\bar{\delta}_i(\mathcal{E}) = s_j \in \mathcal{S}$. Then, there are integers $0 \leq k, r \leq d + 1$, a Lipschitz continuous function*

$$g : \mathbb{T}^k \times [0, 1]^r \rightarrow P := \left\{ (p_1, \dots, p_J) : \sum p_j = 1, p_j \geq 0 \right\},$$

angles $\boldsymbol{\theta} = (\theta_1, \dots, \theta_k) \in \mathbb{T}^k$, rates $\boldsymbol{\beta} = (\beta_1, \dots, \beta_r) \in \mathbb{R}^r$, and $0 < \alpha < 1$ such that

$$(p_1(\mathcal{E}), \dots, p_J(\mathcal{E})) = g(\boldsymbol{\theta} \log \mathcal{E}, \{\boldsymbol{\beta} \log \mathcal{E}\}) + \mathcal{O}(\alpha^{\log \mathcal{E}}).$$

Here, we use the notation

$$\{\boldsymbol{\beta} \log \mathcal{E}\} := (\{\beta_1 \log \mathcal{E}\}, \dots, \{\beta_r \log \mathcal{E}\}) \in [0, 1]^r,$$

where $\{x\}$ denotes the fractional part of a real number x .

Remark 1. *Note that $\mathcal{E} \mapsto (\boldsymbol{\theta} \log \mathcal{E}, \{\boldsymbol{\beta} \log \mathcal{E}\})$ describes a linear flow in $\log \mathcal{E}$ time on the “generalized annulus” $\mathbb{T}^k \times [0, 1]^r$, i.e. a quasiperiodic motion. This is why we say that the rescaled spacings distribution depends quasiperiodically on $\log \mathcal{E}$, asymptotically as $\mathcal{E} \rightarrow \infty$.*

Remark 2. *Note also that the rescaling factor $u(\mathcal{E})$ is adapted to the number field Φ and therefore it might not precisely equal the normalization factor $\lambda(\mathcal{E})$. I.e. the rescaled spacings may not be*

normalized to have average length exactly equal to 1. This is why we refer to $u(\mathcal{E})$ as the “rescaling factor” rather than the “normalizing factor”.

For many quantum systems it is difficult to determine the normalization factor $\lambda(E)$. In 2007 it was proposed by Oganesyanyan and Huse [23] that instead of studying the statistics of the normalized spacings $\bar{\delta}_i(E)$ for $2 \leq i \leq \ell(\mathcal{E})$ one can study the ratios of neighboring spacings

$$\rho_i(\mathcal{E}) := \frac{\delta_i(\mathcal{E})}{\delta_{i-1}(\mathcal{E})} \quad \text{for } 2 \leq i \leq \ell(\mathcal{E}) - 1.$$

(Note that in [23] they take the reciprocal of $\rho_i(E)$ in the case that $\rho_i(E) > 1$, but we will not do that.) The conjectures of Berry and Tabor and of Bohigas, Giannoni, and Schmit can then be re-phrased in terms of ratios. See, for example, [1].

Considering the ratios of neighboring spacings is also quite suitable in our setting because the ambiguity associated to the choice of rescaling factor $u(\mathcal{E})$ that is described in Remark 2 is eliminated. We have:

Theorem B. *Suppose that the ratios of frequencies satisfy that $1, \omega_1, \dots, \omega_d \in \mathbb{R}$ form a \mathbb{Q} -basis for an algebraic number field Φ . We then have:*

- (1) **Uniform Set of Ratios:** *There is a finite set $\mathcal{R} := \{r_1, \dots, r_J\} \subset \Phi$ such that for any $\mathcal{E} > 1$ the ratios of neighboring spacings satisfy:*

$$\rho_i(\mathcal{E}) := \frac{\delta_i(\mathcal{E})}{\delta_{i-1}(\mathcal{E})} \in \mathcal{R} \quad \text{for } 2 \leq i \leq \ell(\mathcal{E}) - 1.$$

- (2) **Quasiperiodicity:** *For each $1 \leq j \leq J$ let $p_j(\mathcal{E})$ denote the proportion of the ratios of neighboring spacings $\rho_i(\mathcal{E})$ such that $\rho_i(\mathcal{E}) = r_j \in \mathcal{R}$. Then, there are integers $0 \leq k, r \leq d + 1$, a Lipschitz continuous function*

$$h : \mathbb{T}^k \times [0, 1]^r \rightarrow P := \left\{ (p_1, \dots, p_J) : \sum p_j = 1, p_j \geq 0 \right\},$$

angles $\boldsymbol{\theta} = (\theta_1, \dots, \theta_k) \in \mathbb{T}^k$, rates $\boldsymbol{\beta} = (\beta_1, \dots, \beta_r) \in \mathbb{R}^r$, and $0 < \alpha < 1$ such that

$$(p_1(\mathcal{E}), \dots, p_J(\mathcal{E})) = h(\boldsymbol{\theta} \log \mathcal{E}, \{\boldsymbol{\beta} \log \mathcal{E}\}) + \mathcal{O}(\alpha^{\log \mathcal{E}}).$$

Remark 3. *Theorem B does not immediately follow from Theorem A since the ratios of neighboring spacings are considered rather than the ratios between arbitrary pairs of spacings. However, in Section 5 we will prove a generalization of Theorem A which implies Theorem B.*

In order to prove Theorems A and B we need a way to compute all of the energy levels ε occurring in the window $\mathcal{E} \leq \varepsilon < \mathcal{E} + 1$ for a give choice of $\mathcal{E} > 0$. For any vector $\mathbf{m} = (m_1, \dots, m_d)$ of non-negative integers such that $\mathbf{m} \cdot \boldsymbol{\omega} < \mathcal{E} + 1$, there is exactly one integer $m_0 \geq 0$ that forces

$$\mathcal{E} \leq \varepsilon := m_0 + m_1 \omega_1 + \dots + m_d \omega_d < \mathcal{E} + 1.$$

This allows us to reduce the problem modulo 1, considering differences between the fractional parts of the numbers $m_1 \omega_1 + \dots + m_d \omega_d$ determined by integer vectors $\mathbf{m} \in R(t)$. Here, $R(t)$ is the homothetic expansion of

$$R = \{ \mathbf{v} \in \mathbb{R}^n : v_1 \omega_1 + \dots + v_d \omega_d < 1 \quad \text{and} \quad v_i \geq 0 \quad \text{for } 1 \leq i \leq d \}$$

by a factor of $t = \mathcal{E} + 1$ about the origin. Therefore, Theorems A and B will be special cases of Theorems A' and B' that are stated in the next subsection.

1.2. Mathematical Perspective. Having provided the physical context and statements of our results, we will now rephrase them purely in the mathematical context.

Suppose that $d \in \mathbb{N}$ and that $\boldsymbol{\omega} = (\omega_1, \dots, \omega_d) \in \mathbb{R}^d$. Let R be a bounded, convex region in \mathbb{R}^d with non-empty interior, and for $t \geq 1$ let $R(t)$ denote the homothetic dilation of R by a factor of t . Throughout the paper we will refer to $t \geq 1$ as the “scale”. Let $\{x\}$ denote the fractional part of a real number x , set $M(t) = R(t) \cap \mathbb{Z}^d$, and write the elements of the set

$$Y(t) := \{\{\mathbf{m} \cdot \boldsymbol{\omega}\} : \mathbf{m} \in M(t)\}$$

in order as

$$0 \leq y_1(t) \leq \dots \leq y_{|M(t)|}(t) < 1.$$

For each value of $i = 1, \dots, |M(t)| - 1$, let

$$\delta_i(t) = y_{i+1}(t) - y_i(t),$$

and let $D(t)$ be the number of distinct elements of the set $\{\delta_i(t)\}_{i=1}^{|M(t)|-1}$. Finally, let

$$\Delta_1(t) < \dots < \Delta_{D(t)}(t)$$

be the ordered sequence of these distinct elements. We may also write Δ_i for $\Delta_i(t)$ and, following [7], we refer to the quantities δ_i and Δ_i as *spacings*.

The classical Three Gap Theorem (also called the Steinhaus Conjecture and the Three Distance Theorem) states that if $d = 1$, then $D(t) \leq 3$ for all t . This was first proved in 1957 by Sós [26], in 1958 by Surányi [27], and in 1959 by Świerczkowski [28]. In the case when $d \geq 2$, estimating the size of $D(t)$ is a more difficult problem. It was known to Geelen and Simpson (attributed by them to Holzman in [16, Section 4]) that, in the case when $d = 2$, if R is a square with sides parallel to the coordinate axes and if $1, \omega_1$, and ω_2 are \mathbb{Q} -linearly dependent, then

$$(1) \quad \sup_{t \geq 1} D(t) < \infty.$$

A proof of this, as well as an extension to $d \geq 3$, is given in [18, Section 4]. A problem attributed to Erdős is to determine for what values of $\boldsymbol{\omega}$ the quantity $D(t)$ remains bounded. It was first speculated that the condition that the numbers $1, \omega_1, \dots, \omega_d$ be \mathbb{Q} -linearly dependent is necessary in order for $D(t)$ to remain bounded. However, the situation is more subtle.

A vector $\boldsymbol{\omega} \in \mathbb{R}^d$ is called *Diophantine* with exponent γ if there is some positive number K such that for all nonzero vectors $\mathbf{m} \in \mathbb{Z}^d$, we have

$$(2) \quad |\mathbf{m} \cdot \boldsymbol{\omega}| \geq \frac{K}{|\mathbf{m}|^\gamma}, \text{ where } |\mathbf{m}| = \sqrt{m_1^2 + \dots + m_d^2}.$$

It follows from the Minkowski’s Theorem that $\gamma \geq d$. We call $\boldsymbol{\omega}$ *badly approximable* if $\gamma = d$. The set of badly approximable $\boldsymbol{\omega}$ has zero Lebesgue measure, but it is known by work of Jarník [21] and Wolfgang Schmidt [25] to be a subset of \mathbb{R}^d of Hausdorff dimension d .

Boshernitzan-Dyson Theorem. *If $\boldsymbol{\omega} \in \mathbb{R}^d$ is badly approximable, then (1) holds.*

This result was not published by Boshernitzan and Dyson, but a proof can be found in [7].

In the other direction, it was proved in [18] that when $d \geq 2$, for almost all choices of $\boldsymbol{\omega} \in \mathbb{R}^d$,

$$(3) \quad \sup_{t \geq 1} D(t) = \infty.$$

The proof given in [18] uses ergodic theory in spaces of unimodular lattices in \mathbb{R}^{d+1} . Part of the interest in this problem lies in the fact that, for $d \geq 2$, if (3) holds, then

$$\liminf_{n \rightarrow \infty} n \|n\omega_1\| \cdots \|n\omega_d\| = 0,$$

where $\|\cdot\|$ denotes the distance to the nearest integer [18, Theorem 3]. In other words, if (3) holds then the generalized Littlewood conjecture is true for ω . The converse of this statement, however, is not true.

We also remark that, as another twist in the above mentioned problem of Erdős, it turns out that there do exist vectors $\omega \in \mathbb{R}^d$ that are not badly approximable but for which (1) holds. An explicit example is given in [3] for $d = 2$.

Now let us focus on the situation when $1, \omega_1, \dots, \omega_d \in \mathbb{R}$ form a \mathbb{Q} -basis for an algebraic number field Φ of degree $d+1$. In this case it is known by work of Perron [24] that ω is badly approximable, so it follows from the Boshernitzan-Dyson Theorem that (1) holds. (In fact, Dyson originally proved the theorem in this special case.) Actually, more is true, as demonstrated by the following result.

Theorem 4. [7, Theorem 1.6] *If $1, \omega_1, \dots, \omega_d \in \mathbb{R}$ form a \mathbb{Q} -basis for an algebraic number field Φ , then there exists a finite set*

$$\mathcal{S} = \{s_1, \dots, s_J\} \subseteq \Phi$$

such that every spacing Δ_i has the form us_j for some unit u in the ring of integers \mathbb{Z}_Φ and some $s_j \in \mathcal{S}$.

Throughout this paper we will denote by \mathbb{Z}_Φ^\times the group of units in the ring of integers \mathbb{Z}_Φ of an algebraic number field Φ .

Consider the one dimensional case $d = 1$. As $t \geq 1$ increases, additional points $y_j(t)$ are added one-by-one. Most of the time, this results in one of the large-sized spacings being split into one of the mid-sized spacings and one of the small-sized spacings. This continues until each of the large-sized spacings is split, at which point one renames the medium-sized spacings to be “large” and the small-sized spacings to be “medium” and then repeats the process, splitting each new large-sized spacing into a new medium-sized spacing and a (truly new) small-sized spacing. Therefore, as t increases, the proportions of spacings occurring that are deemed to be “small”, “medium”, and “large” depend in an organized way on t , which can be made precise using the theory of continued fractions.

The purpose of this paper is to describe analogous behavior in the far more complicated situation when $d \geq 2$, at least in the algebraic case when $1, \omega_1, \dots, \omega_d \in \mathbb{R}$ form a \mathbb{Q} -basis for an algebraic number field Φ . We will prove in Part (1) of Theorem A', below, a stronger version of Theorem 4 that allows us to *uniformly* label the spacings $\Delta_1(t), \dots, \Delta_{D(t)}(t)$ using the elements of a finite set $\mathcal{S} \subset \Phi$. We can then prove in Part (2) of Theorem A' a description of the time t evolution of the proportion of spacings realizing these labels from \mathcal{S} . We will also prove a similar theorem about the frequencies with which the ratios of neighboring spacings occur, see Theorem B' below.

Theorem A'. *Suppose that $1, \omega_1, \dots, \omega_d \in \mathbb{R}$ form a \mathbb{Q} -basis for an algebraic number field Φ and let $r \geq 1$ denote the rank of \mathbb{Z}_Φ^\times . We then have:*

- (1) **Uniform Labeling:** *There is a finite set $\mathcal{S} := \{s_1, \dots, s_J\} \subset \Phi$ and a rescaling factor $u(t) \in \mathbb{Z}_\Phi^\times$ such that for any $t > 1$ we have*

$$\overline{\delta}_i(t) := u(t)\delta_i(t) \in \mathcal{S} \quad \text{for } 1 \leq i \leq |M(t)| - 1.$$

We will refer to the $\{\overline{\delta}_i(t)\}$ as the rescaled spacings.

- (2) **Quasiperiodicity:** *For each $1 \leq j \leq J$ let $p_j(t)$ denote the proportion of the rescaled spacings $\overline{\delta}_i(t)$ such that $\overline{\delta}_i(t) = s_j \in \mathcal{S}$. Then, there is an integer $0 \leq k \leq d+1$, a Lipschitz continuous function*

$$g : \mathbb{T}^k \times [0, 1]^r \rightarrow P := \left\{ (p_1, \dots, p_J) : \sum p_j = 1, p_j \geq 0 \right\},$$

angles $\boldsymbol{\theta} = (\theta_1, \dots, \theta_k) \in \mathbb{T}^k$, rates $\boldsymbol{\beta} = (\beta_1, \dots, \beta_r) \in \mathbb{R}^r$, and $0 < \alpha < 1$ such that

$$(p_1(t), \dots, p_J(t)) = g(\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\}) + \mathcal{O}(\alpha^{\log t}).$$

Here, we use the notation

$$\{\boldsymbol{\beta} \log t\} := (\{\beta_1 \log t\}, \dots, \{\beta_r \log t\}) \in [0, 1]^r,$$

where $\{x\}$ denotes the fractional part of a real number x .

Note that $t \mapsto (\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\})$ describes a linear flow in $\log t$ time on the ‘‘generalized annulus’’ $\mathbb{T}^k \times [0, 1]^r$, i.e. a quasiperiodic motion. Therefore, Theorem A’ asserts that the frequencies at which the rescaled spacings $s_1 < s_2 < \dots < s_J$ occur, as a function of t , depend quasiperiodically on $\log t$, as $t \rightarrow \infty$. To understand some of the details of this dependence, before delving into the complexities of the proof, first note there are many ways to choose a basis $\{\epsilon_1, \dots, \epsilon_r\}$ for a finite index subgroup of \mathbb{Z}_Φ^\times , none of which in general can be assumed to be canonical. All of the parameters in the statement of the theorem depend on this choice of basis and, once it has been made, they are explicitly computable. The units $u(t)$ will turn out to be determined by

$$u(t)^{-1} = \epsilon_1^{\lfloor \beta_1 \log t \rfloor} \dots \epsilon_r^{\lfloor \beta_r \log t \rfloor},$$

for a suitable choice of $\boldsymbol{\beta}$, and this in turn determines \mathcal{S} . The Minkowski embedding of Φ into \mathbb{R}^{d+1} (see next section for details) allows us to view multiplication by $u(t)^{-1}$ in Φ as a linear transformation $U(t)$ on \mathbb{R}^{d+1} . By using the Jordan decomposition of a complex matrix defined using a particular choice of matrix logarithm for $U(t)$, we are able to understand the time evolution of $U(t)$ as $t \rightarrow \infty$, and to prove that it is governed completely by two types of generalized eigenspaces: one dimensional spaces corresponding to purely imaginary eigenvalues $2\pi i\theta_1, \dots, 2\pi i\theta_k$ (of which there is at least one), and spaces with eigenvalues whose real parts are negative. This gives k and $\boldsymbol{\theta}$, and α is determined by the maximum of the real parts of the eigenvalues which are not purely imaginary (if there are any). Finally, the function g is defined explicitly in Section 4.3 and equation (23), it is straightforward to compute and depends only on \mathcal{S} and the region R .

Remark 5. *Under the additional hypothesis that Φ has one fundamental unit (which restricts the theorem to quadratic fields and cubic fields with a complex embedding), a preliminary version of Theorem A’ was proved in [7, Thms 1.5 and 1.6]. The term involving rate $\beta \equiv \beta_1$ does not appear in those theorems because they are expressed at a sequence of times $t_n = \eta e^{n/\beta}$ that is chosen to make the sequence $\{\beta \log t_n\}$ constant. When $r > 1$ such a choice is not typically possible. However, several aspects of the proofs from [7, Thms 1.5 and 1.6] will play an important role in our proof of Theorem A’.*

The labeling of spacings by elements of \mathcal{S} may seem ad hoc because it depends on the construction of $u(t)$ from the proof of Theorem A’, Part (1). By adjusting the choice of $u(t)$ we could easily change the set of labels \mathcal{S} . For this reason, it may be more natural to consider the ratios between the neighboring spacings.

Theorem B’. *Suppose that $1, \omega_1, \dots, \omega_d \in \mathbb{R}$ form a \mathbb{Q} -basis for an algebraic number field Φ and let $r \geq 1$ denote the rank of \mathbb{Z}_Φ^\times . We then have:*

- (1) **Uniform Set of Ratios:** *There is a finite set $\mathcal{R} := \{r_1, \dots, r_J\} \subset \Phi$ such that for any $t > 1$ the ratios of neighboring spacings satisfy:*

$$\rho_i(t) := \frac{\delta_i(t)}{\delta_{i-1}(t)} \in \mathcal{R} \quad \text{for } 2 \leq i \leq |M(t)| - 1.$$

- (2) **Quasiperiodicity:** For each $1 \leq j \leq J$ let $p_j(t)$ denote the proportion of the ratios of neighboring spacings $\rho_i(t)$ such that $\rho_i(t) = r_j \in \mathcal{R}$. Then, there is an integer $0 \leq k \leq d+1$, a Lipschitz continuous function

$$h : \mathbb{T}^k \times [0, 1]^r \rightarrow P := \left\{ (p_1, \dots, p_J) : \sum p_j = 1, p_j \geq 0 \right\},$$

angles $\boldsymbol{\theta} = (\theta_1, \dots, \theta_k) \in \mathbb{T}^k$, rates $\boldsymbol{\beta} = (\beta_1, \dots, \beta_r) \in \mathbb{R}^r$, and $0 < \alpha < 1$ such that

$$(p_1(t), \dots, p_J(t)) = h(\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\}) + \mathcal{O}(\alpha^{\log t}).$$

Remark 6. Theorem B' does not immediately follow from Theorem A' because only the ratios of neighboring spacings are considered, rather than the ratios between arbitrary pairs of spacings. However, in Section 5 we will state and prove a stronger version of Theorem A', namely Theorem C, about the frequencies with which a given word $s_{j_1} s_{j_2} \dots s_{j_l}$ appears among the consecutive rescaled spacings. Theorem B' then follows immediately from Theorem C using words of length $l = 2$.

Question 1. Let $\boldsymbol{\omega} \in \mathbb{R}^d$ and suppose Statement (1) from Theorem B' holds for the ratios of spacings determined by $\boldsymbol{\omega}$. Must $1, \omega_1, \dots, \omega_d$ be a \mathbb{Q} -basis for an algebraic number field Φ ?

1.3. Plan for the paper. As explained at the end of Section 1.1, Theorems A and B are direct consequences of Theorems A' and B'.

Our proof of Part (1) of Theorem A' is an application of transference principles from Diophantine approximation, together with well known results from algebraic number theory. In Section 2 we will review some of these results, and in Section 3 we will present the proof of Part (1) of Theorem A'. Our proof of Part (2) of Theorem A' uses many results from the proofs of [7, Thms 1.5 and 1.6] combined with several new ideas that are needed when $\mathbb{Z}_{\Phi}^{\times}$ has rank $r > 1$. It is presented in Section 4. In Section 5 we prove Theorem C, the generalization of Theorem A' mentioned in Remark 6. Since Theorem B' is an immediate corollary to Theorem C this will also complete the proof of Theorem B'. Finally, in Section 6 we work out the details of Theorems A' for a particular example (a totally real cubic field) which highlights the computational aspects and importance of many of the steps in our proofs.

1.4. Acknowledgments. The second author thanks Pavel Bleher for introducing him to this subject and for many interesting conversations about it. We thank Evgeny Mukhin and Vitaly Tarasov for providing us with the proof of Lemma 12, which plays a crucial role in our paper. The second author also thanks Aneesh Dasgupta for interesting conversations about this subject. The work of the first author was supported by NSF grant DMS-2001248. The work of the second author was supported by NSF grant DMS-1348589.

2. NOTATION AND PRELIMINARY RESULTS

For $x \in \mathbb{R}$, we write $\{x\}$ for the fractional part of x and $\|x\|$ for the distance from x to the nearest integer. For $d \in \mathbb{N}$ and $\boldsymbol{x} \in \mathbb{R}^d$, we write $|\boldsymbol{x}|$ for the Euclidean norm of \boldsymbol{x} .

Results from Diophantine approximation known as transference principles (see [11, Section V, Theorem VI]) imply that $\boldsymbol{\omega} \in \mathbb{R}^d$ is badly approximable if and only if there exists a constant $K' > 0$ with the property that, for any $t \geq 1$, any ball of diameter K'/t^d in $[0, 1)$ contains a point of the set

$$\{\{\boldsymbol{m} \cdot \boldsymbol{\omega}\} : \boldsymbol{m} \in M(t)\}.$$

If $1, \omega_1, \dots, \omega_d$ form a \mathbb{Q} -basis for an algebraic number field of degree $d+1$ over \mathbb{Q} , then the results from [24] mentioned in the introduction imply that $\boldsymbol{\omega}$ is badly approximable. By the transference

principle just cited, we thus have for all $t \geq 1$ and for all i that

$$(4) \quad \Delta_i \leq \frac{K'}{t^d}.$$

Next we summarize some basic facts from algebraic number theory, proofs of which can be found in [29, Chapters 1, 3]. As above, suppose that Φ is an algebraic number field of degree $d + 1$ over \mathbb{Q} . There are $d + 1$ distinct embeddings of Φ into \mathbb{C} , and the non-real complex embeddings come in complex conjugate pairs. Suppose there are r_1 real embeddings and $2r_2$ complex embeddings, and write $\sigma_1, \dots, \sigma_{r_1}$ for the real embeddings and σ_i, σ_{i+r_2} , with $r_1 < i \leq r_1 + r_2$, for each pair of complex conjugate embeddings. Identifying \mathbb{C} with \mathbb{R}^2 , we define a map $\sigma : \Phi \rightarrow \mathbb{R}^{d+1}$ by

$$\sigma(\alpha) = (\sigma_1(\alpha), \dots, \sigma_{r_1+r_2}(\alpha)).$$

This map is injective, and the set

$$\Gamma = \sigma(\mathbb{Z}_\Phi)$$

is called the Minkowski embedding of the ring of integers of Φ into \mathbb{R}^{d+1} . It is a discrete subgroup of \mathbb{R}^{d+1} , and the quotient \mathbb{R}^{d+1}/Γ has a measurable fundamental domain of finite volume. In other words, Γ is a lattice in \mathbb{R}^{d+1} .

Let $k \in \mathbb{Z}$ be chosen so that for each $1 \leq j \leq d$ we have $k\omega_j \in \mathbb{Z}_\Phi$. Notice that

$$(5) \quad \mathbb{M}_\Phi := \{\alpha \in \Phi : k\alpha \in \mathbb{Z}_\Phi\}$$

is a \mathbb{Z}_Φ module that contains $\omega_1, \dots, \omega_d$. In particular, for any $t \geq 1$ each spacing $\Delta_i(t) \in \mathbb{M}_\Phi$ and moreover $x\Delta_i(t) \in \mathbb{M}_\Phi$ for any $x \in \mathbb{Z}_\Phi$. The image

$$(6) \quad \Gamma' = \sigma(\mathbb{M}_\Phi)$$

is a lattice because $\Gamma' = \frac{1}{k}\Gamma$, with Γ a lattice.

Next, let \mathbb{Z}_Φ^\times denote the group of multiplicative units of \mathbb{Z}_Φ . By the Dirichlet unit theorem, this group has rank $r_1 + r_2 - 1$. Consider the map $\varphi : \mathbb{Z}_\Phi^\times \rightarrow \mathbb{R}^{r_1+r_2}$ defined by

$$(7) \quad \varphi(u) = (\log |\sigma_1(u)|, \dots, \log |\sigma_{r_1+r_2}(u)|).$$

This map is well defined, since $|\sigma_i(u)| \neq 0$. The norm of any unit is ± 1 , so the image of φ is contained in the hyperplane in $\mathbb{R}^{r_1+r_2}$ with equation

$$(8) \quad x_1 + \dots + x_{r_1} + 2x_{r_1+1} + \dots + 2x_{r_1+r_2} = 0.$$

Furthermore, the image of φ is a lattice in this hyperplane (see the proof of [29, Theorem 11]). All of these facts will be useful to us in what follows.

3. PROOF OF PART (1) OF THEOREM A' (UNIFORM LABELING)

Suppose without loss of generality that $t \geq 1$. Since $\omega \in \mathbb{R}^d$ we know that $r_1 \geq 1$, so let us assume that $\sigma_1 : \Phi \rightarrow \mathbb{R}$ is the trivial embedding which maps each number ω_j to itself. Each spacing $\Delta_i(t)$ has the form

$$\Delta_i(t) = (\mathbf{m} - \mathbf{m}') \cdot \omega,$$

for some $\mathbf{m}, \mathbf{m}' \in M(t)$. It follows from this and (4) that there exists a constant $C > 0$, which does not depend on t , with the property that for each spacing Δ_i , we have that

$$|\sigma_1(\Delta_i)| \leq \frac{C}{t^d}.$$

Also, since $|\mathbf{m} - \mathbf{m}'|$ is bounded by a constant times t , and the maps σ_j are homomorphisms, it immediately follows that

$$|\sigma_j(\Delta_i)| \leq Ct.$$

for $2 \leq j \leq r_1 + r_2$.

Since $\varphi(\mathbb{Z}_\Phi^\times)$ is a lattice in the hyperplane defined by (8), there is a constant $C' > 0$ with the property that, for any point $\mathbf{x} \in \mathbb{R}^{r_1+r_2}$ satisfying (8), there is an element of $\varphi(\mathbb{Z}_\Phi^\times)$ in the ball of radius C' centered at \mathbf{x} . Using the fact that $r_1 + 2r_2 = d + 1$, we apply this observation with

$$(9) \quad \mathbf{x} = (d \log t, -\log t, \dots, -\log t).$$

We thus deduce that there is a unit $u = u(t)$ with the properties that

$$|\sigma_1(u)| \leq e^{C'} t^d$$

and, for $2 \leq j \leq r_1 + r_2$, that

$$|\sigma_j(u)| \leq \frac{e^{C'}}{t}.$$

It follows that $\sigma(u\Delta_i)$ is a point of the set Γ' from (6), which lies in a cube \mathcal{C} of side length $2Ce^{C'}$ centered at the origin in \mathbb{R}^{d+1} . Let \mathcal{S} be the collection of all elements of \mathbb{M}_Φ (defined in (5)) whose images under σ lie in \mathcal{C} . Since Γ' is a lattice and σ is injective, the set \mathcal{S} is finite. The statement of the theorem thus follows. \square (Part (1) of Theorem A')

4. PROOF OF PART (2) OF THEOREM A' (QUASIPERIODICITY)

Several times in this section we will need to refer to the multiplicative inverse of $u(t)$ and thus we will denote it by $u_1 \equiv u_1(t) := u(t)^{-1}$.

Let $\mathbf{n} : \Phi \rightarrow \mathbb{Q}^{d+1}$ denote the expansion of an element of Φ in terms of the basis $1, \omega_1, \dots, \omega_d$. That is, for any $\alpha \in \Phi$,

$$\mathbf{n}(\alpha) = (n_0, n_1, \dots, n_d) \quad \text{iff} \quad \alpha = n_0 + n_1\omega_1 + \dots + n_d\omega_d.$$

Let $\mathbf{m} : \Phi \rightarrow \mathbb{Q}^d$ be the ‘‘truncated expansion’’ of α given by

$$\mathbf{m}(\alpha) = (n_1, \dots, n_d) \quad \text{if} \quad \mathbf{n}(\alpha) = (n_0, n_1, \dots, n_d).$$

The preliminary versions of Part (2) of Theorem A' from [7] are proved in three steps:

- (1) Describing the proportions of spacings at a given scale $t \geq 1$ in terms of suitable partitions of $M(t)$ and R ,
- (2) Relating these partitions to $\frac{\mathbf{n}(u_1(t))}{t}$, the normalized expansion of $u_1(t)$ in the basis $1, \omega_1, \dots, \omega_d$, and
- (3) Analysis of the asymptotic behavior of $\frac{\mathbf{n}(u_1(t))}{t}$ as $t \rightarrow \infty$.

Steps (1) and (2) carry over directly to our setting. We will describe them in Sections 4.1 and 4.3, with a discussion of any necessary adaptations. However, Step (3) requires some new ideas, which we present in Section 4.4, thus completing the proof of Part (2) of Theorem A'. (Section 4.2 presents some lemmas that are needed in Sections 4.3 and 4.4.)

4.1. Partitions of $M(t)$ and R . For any $1 \leq j \leq J$, let $Y_j(t)$ be the set of numbers $y_i(t)$ such that $\delta_i(t) = y_{i+1}(t) - y_i(t) = s_j u_1(t)$. Here, $u_1(t) = u(t)^{-1}$, where $u(t)$ is the unit from Part (1) of Theorem A' and s_j is an element of the finite set \mathcal{S} , which has been ordered so that $s_1 < \dots < s_J$.

Let $M_j(t)$ be the set of vectors $\mathbf{m} \in M(t)$ such that $\{\mathbf{m} \cdot \boldsymbol{\omega}\} \in Y_j(t)$. Up to the single point corresponding to the largest element $y_{\ell(t)}(t) \in Y(t)$ we have that

$$\bigsqcup_{j=1}^J M_j(t) = M(t),$$

so, by a slight abuse of notation, we will call $\{M_j(t)\}$ a partition of $M(t)$. We refer the reader to [7, Fig. 1] for an explicit example. We conclude that at scale $t \geq 1$ the proportion of times that the rescaled spacing $s_j \in \mathcal{S}$ occurs is

$$p_j(t) = \frac{|M_j(t)|}{|M(t)| - 1}.$$

We will need the following result from [7].

Proposition 7 (Prop. 5.1 from [7]). *For any $1 \leq j \leq J$ let*

$$\mathbf{v}_j(t) = \mathbf{m}(s_j u_1(t)).$$

Then we have

$$(10) \quad M_j(t) = [M(t) \cap (M(t) - \mathbf{v}_j(t))] \setminus \bigcup_{i=1}^{j-1} (M(t) - \mathbf{v}_i(t)),$$

where $\Omega + \mathbf{v}$ is defined to be $\{\mathbf{u} + \mathbf{v} : \mathbf{u} \in \Omega\}$ for any $\Omega \subset \mathbb{R}^d$ and $\mathbf{v} \in \mathbb{R}^d$.

Remark 8. *Our formula for $M_j(t)$ above has minus signs where the analogous formula in [7] has plus signs. The reason is that we define $\delta_i(t) = y_{i+1}(t) - y_i(t)$ while $y_i(t) - y_{i-1}(t)$ is used in [7].*

Denote the power set of our region R by $\mathcal{P}(R)$ and let

$$P : (\mathbb{R}^d)^J \rightarrow \mathcal{P}(R)^J$$

be the mapping which sends the J -tuple of vectors $\mathbf{v} = (\mathbf{v}_1, \dots, \mathbf{v}_J)$ to the J -tuple $(P_1(\mathbf{v}), \dots, P_J(\mathbf{v}))$ of subsets of R where,

$$P_j(\mathbf{v}) = [R \cap (R - \mathbf{v}_j)] \setminus \bigcup_{i=1}^{j-1} (R - \mathbf{v}_i).$$

Proposition 9 (Prop. 5.2 from [7]). *If*

$$(11) \quad \mathbf{v} := \mathbf{v}(t) = \left(\frac{\mathbf{m}(s_1 u_1(t))}{t}, \dots, \frac{\mathbf{m}(s_J u_1(t))}{t} \right),$$

then

$$(12) \quad p_j(t) - \frac{\text{volume}(P_j(\mathbf{v}))}{\text{volume}(R)} = \frac{|M_j(t)|}{|M(t)| - 1} - \frac{\text{volume}(P_j(\mathbf{v}))}{\text{volume}(R)} = \mathcal{O}\left(\frac{1}{t}\right).$$

The proof of [7, Prop. 5.4] uses the estimate that for any convex $\Omega \subset \mathbb{R}^2$ we have

$$\text{area}(t\Omega) - |t\Omega \cap \mathbb{Z}^2| = \mathcal{O}(t).$$

This estimate adapts to work for a convex region $\Omega \subset \mathbb{R}^d$, with the area becoming volume and the error becoming $\mathcal{O}(t^{d-1})$. This is the only change needed to adapt the proof of [7, Prop. 5.4] to the present setting.

Proposition 10 (Prop. 5.3 from [7]). *The function P is Lipschitz continuous with respect to the infinity norm on $(\mathbb{R}^d)^J$ and the metric*

$$d(P^{(1)}, P^{(2)}) = \sum_{j=1}^J \text{vol}(P_j^{(1)} \Delta P_j^{(2)})$$

on J -tuples of subsets of R , where Δ denotes the symmetric difference of sets.

The only change needed to adapt the proof of Prop. 5.3 from [7] from dimension two to dimension $d \geq 2$ is that one needs to remark that the polynomial $Q(x) = \left(1 + \frac{x}{a}\right)^d - 1$ which appears in that proof is Lipschitz on the interval $[0, 1]$.

4.2. Matrix representation of multiplication by elements of Φ and their logarithms. Multiplication by any non-zero $a \in \Phi$ corresponds to an invertible linear mapping from Φ to itself.

Lemma 11. *For any two non-zero $a, b \in \Phi$ the $(d+1) \times (d+1)$ dimensional matrices A and B representing multiplication by a and b in terms of the basis $1, \omega_1, \dots, \omega_d$ commute.*

Proof. This is an immediate consequence of commutativity $ab = ba$ in Φ . □

Combined with Lemma 11 the following lemma plays a crucial role in our proof.

Lemma 12. *For any $k \geq 1$ let A_1, \dots, A_k be a commuting collection of invertible $n \times n$ matrices. Then, there exist $n \times n$ matrices L_1, \dots, L_k such that*

$$(13) \quad A_j = e^{L_j}$$

for each $1 \leq j \leq k$ and such that all k matrices L_1, \dots, L_k commute.

As usual, the matrix exponential in (13) is interpreted using the standard power series for e^z . The matrices L_1, \dots, L_k are called *logarithms* of A_1, \dots, A_k ; see, for example [17, Sec. 2.3]. Although it seems that this lemma should be well-known we could not find a suitable reference. We thank Evgeny Mukhin and Vitaly Tarasov for providing us with the following proof.

Proof. We claim that there is an invertible $n \times n$ matrix P such that for every $1 \leq j \leq k$ we have $A_j = PD_jP^{-1}$ with each D_j a block diagonal matrix

$$D_j = \text{diag}(B_{j,1}, \dots, B_{j,m}),$$

with the size of the blocks independent of $1 \leq j \leq k$, and each block having the form

$$B_{j,\ell} = \lambda_{j,\ell} \mathbf{I} + N_{j,\ell},$$

where $\lambda_{j,\ell} \in \mathbb{C} \setminus \{0\}$ and $N_{j,\ell}$ is a nilpotent matrix for each $1 \leq j \leq k$ and $1 \leq \ell \leq m$. Here, \mathbf{I} denotes the identity matrix of the appropriate dimension.

Let us first see how this claim yields the desired result. Notice that it suffices to find commuting logarithms of the block diagonal matrices D_j , for $1 \leq j \leq k$, because the desired L_1, \dots, L_k will then be obtained by conjugating by P . Moreover, matrix exponentials respect block-diagonal structure, so it suffices to find mutually commuting matrix logarithms for k commuting matrices of the form

$$M_j = \lambda_j \mathbf{I} + N_j,$$

where N_j is nilpotent for each $1 \leq j \leq k$. One can do this using the Mercator series to define

$$\mathcal{L}_j := \log \lambda_j \mathbf{I} - \sum_{m=1}^{\infty} \frac{1}{m} \left(\frac{-N_j}{\lambda_j} \right)^m,$$

for each $1 \leq j \leq k$. (One can choose any complex logarithm $\log \lambda_j$ that one likes.) The series converge because they terminate in finitely many steps, since each N_j is nilpotent. Moreover $\mathcal{L}_1, \dots, \mathcal{L}_k$ commute because the N_1, \dots, N_k commute.

Now we are left to establish the claim from the beginning of the proof. Existence of the matrix P that simultaneously conjugates the A_1, \dots, A_k to the desired block diagonal form is a generalization of the well-known fact that commuting diagonalizable matrices are simultaneously diagonalizable. We sketch it here in the case $k = 2$, leaving the straightforward generalization to larger k to the reader.

By the Jordan decomposition we can write \mathbb{R}^n as a direct sum of generalized eigenspaces of A_1 . Let V be any one of the generalized eigenspaces of A_1 , corresponding to eigenvalue λ . By definition, it consists of the vectors in \mathbb{R}^n in the kernel of $(A_1 - \lambda I)^\ell$ for some $\ell \geq 1$. On V the matrix A_1 is conjugate to $\lambda I + N$ for some nilpotent matrix N , by the Jordan form.

Commutativity of A_1 and A_2 implies that $A_2(V)$ is a subspace of V . We can therefore decompose V into generalized eigenspaces of A_2 and, for any such subspace W of V , commutativity of A_1 and A_2 implies $A_1 W$ is a subspace of W . In particular $N(W)$ is a subspace of W .

This proves that \mathbb{R}^n can be decomposed into a direct sum of spaces that are simultaneously generalized eigenspaces of A_1 and A_2 , on each of which A_1 is conjugate to $\lambda I + N_1$ and A_2 is conjugate to $\mu I + N_2$ for some $\lambda, \mu \in \mathbb{C} \setminus \{0\}$ and nilpotent matrices N_1 and N_2 . \square

4.3. Reduction to analysis of normalized expansions of $u_1(t)$. Here we reduce the proof of Part (2) of Theorem A' to the proof of the following theorem.

Theorem 13. *Under the hypotheses of Theorem A' there is an integer $0 \leq k \leq d + 1$, a Lipschitz continuous function*

$$g_3 : \mathbb{T}^k \times [0, 1]^r \rightarrow \mathbb{R}^{d+1},$$

angles $\boldsymbol{\theta} = (\theta_1, \dots, \theta_k) \in \mathbb{T}^k$, "rates" $\boldsymbol{\beta} = (\beta_1, \dots, \beta_r) \in \mathbb{R}^r$, and $0 < \alpha < 1$ such that

$$\frac{\mathbf{n}(u_1(t))}{t} = g_3(\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\}) + \mathcal{O}(\alpha^{\log t}).$$

Proof of Part (2) of Theorem A' supposing Theorem 13. Let us first summarize what Propositions 7-10 achieve. Let

$$g_1 : (\mathbb{R}^d)^J \rightarrow \mathbb{R}^J$$

be given by

$$g_1(\mathbf{v}) = \left(\frac{\text{volume}(P_1(\mathbf{v}))}{\text{volume}(R)}, \dots, \frac{\text{volume}(P_J(\mathbf{v}))}{\text{volume}(R)} \right).$$

Then, g_1 is Lipschitz continuous and satisfies that for any $t \geq 1$

$$g_1(\mathbf{v}(t)) - (p_1(t), \dots, p_J(t)) = \mathcal{O}(t^{-1}),$$

where $\mathbf{v}(t)$ is given by (11).

We now claim that there is a linear function

$$g_2 : \mathbb{R}^{d+1} \rightarrow (\mathbb{R}^d)^J$$

such that

$$\mathbf{v}(t) = g_2 \left(\frac{\mathbf{n}(u_1(t))}{t} \right).$$

For any $1 \leq j \leq J$ let S_j denote the matrix expressing multiplication by $s_j \in \mathcal{S}$ in terms of the basis $1, \omega_1, \dots, \omega_d$. For any $1 \leq j \leq J$ we let the j -th component $\mathbf{v}_j(t)$ of $\mathbf{v}(t)$ be the projection of

$$\frac{\mathbf{n}(s_j u_1(t))}{t} = S_j \frac{\mathbf{n}(u_1(t))}{t} \in \mathbb{R}^{d+1}$$

onto its last d components.

Therefore, given the function $g_3 : \mathbb{T}^k \times [0, 1]^r \rightarrow \mathbb{R}^{d+1}$ whose existence is asserted by Theorem 13, we can let $g = g_1 \circ g_2 \circ g_3$ so that

$$\begin{aligned} g(\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\}) &= g_1 \left(g_2 \left(\frac{\mathbf{n}(u_1(t))}{t} + \mathcal{O}(\alpha^{\log t}) \right) \right) \\ &= g_1 \left(\mathbf{v}(t) + \mathcal{O}(\alpha^{\log t}) \right) \\ &= (p_1(t), \dots, p_J(t)) + \mathcal{O}(t^{-1}) + \mathcal{O}(\alpha^{\log t}), \end{aligned}$$

with the last equality using that g_1 is Lipschitz. Since $t^{-1} = (1/e)^{\log t}$ this proves the claim. \square

4.4. Analysis of normalized expansions of $u_1(t)$ as $t \rightarrow \infty$. We will now use results from linear algebra to finish the proof of Theorem 13. We begin by deriving an explicit formula for $u_1(t) = u(t)^{-1}$. Let $r = r_1 + r_2 - 1$ denote the rank of the unit group $\mathbb{Z}_{\Phi}^{\times}$ and let $\epsilon_1, \dots, \epsilon_r$ be a basis for a finite index subgroup of the multiplicative group $\mathbb{Z}_{\Phi}^{\times}$. Recall that the image of $\mathbb{Z}_{\Phi}^{\times}$ under the mapping φ given in (7) forms a lattice in the hyperplane $H \subset \mathbb{R}^{r_1+r_2}$ defined by (8). Therefore we can use the coordinate system

$$(y_1, \dots, y_r) \mapsto y_1 \varphi(\epsilon_1) + \dots + y_r \varphi(\epsilon_r)$$

on H . In these coordinates the image under ϕ of the group generated by $\epsilon_1, \dots, \epsilon_r$ becomes the integer lattice \mathbb{Z}^r . The path $\mathbf{x}(t)$ defined in (9) becomes

$$(14) \quad \mathbf{x}(t) = -\mathbf{w} \log t$$

for some suitable non-zero vector $\mathbf{w} = (w_1, \dots, w_r)$. We can then use

$$u_1(t) = u(t)^{-1} = \epsilon_1^{\lfloor w_1 \log t \rfloor} \dots \epsilon_r^{\lfloor w_r \log t \rfloor}$$

as the unit in Part (1) of Theorem A'.

Let $U(t)$ denote the matrix representing multiplication by $u_1(t)$ in the basis $1, \omega_1, \dots, \omega_d$. We have

$$U(t) = E_1^{\lfloor w_1 \log t \rfloor} \dots E_r^{\lfloor w_r \log t \rfloor},$$

where E_1, \dots, E_r are the matrices representing multiplication by the units $\epsilon_1, \dots, \epsilon_r$.

Let us approximate $U(t)$ by a continuous version. According to Lemmas 11 and 12 we can choose logarithms L_1, \dots, L_r of the matrices E_1, \dots, E_r in a way that they all commute. Let

$$(15) \quad \tilde{U}(t) = e^{w_1(\log t)L_1 + \dots + w_r(\log t)L_r} = e^{(\log t)L},$$

where

$$L := w_1 L_1 + \dots + w_r L_r.$$

We remark that, since L_1, \dots, L_r commute, we also have that

$$(16) \quad \tilde{U}(t) = E_1^{w_1 \log t} \dots E_r^{w_r \log t},$$

since the definition of the real power of a matrix gives $E_j^{w_j \log t} := e^{w_j \log t L_j}$ for $1 \leq j \leq r$. Note that making a different choice of matrix logarithm can lead to a different value of the real power of a matrix, just like for the real power of a real number. However, we have fixed our choices of logarithms once and for all when we defined (15).

For any $t \geq 1$, again using the commutativity of L_1, \dots, L_r , we have that

$$(17) \quad U(t) = A(t) \tilde{U}(t)$$

where

$$(18) \quad A(t) = e^{-\{w_1 \log t\}L_1 \cdots - \{w_r \log t\}L_r}.$$

We think of $A(t)$ as the “multiplicative error” between $U(t)$ and our continuous approximation $\tilde{U}(t)$. It ranges over a compact subset of the space of invertible $(d+1) \times (d+1)$ matrices.

Remark 14. *The product $A(t)\tilde{U}(t)$ is a real matrix, because $U(t)$ is. However, the matrices $A(t)$ and $\tilde{U}(t)$ are not (necessarily) real because the logarithms L_1, \dots, L_r are not necessarily real.*

To prove Theorem 13 we must estimate

$$\frac{\mathbf{n}(u_1(t))}{t} = \frac{1}{t}U(t) \mathbf{e}_1 = A(t) \left(\frac{1}{t}\tilde{U}(t) \mathbf{e}_1 \right),$$

where $\mathbf{e}_1 = (1, 0, \dots, 0)^T$. To do this we write

$$\frac{1}{t}\tilde{U}(t) = e^{-(\log t)I} e^{(\log t)L} = e^{(\log t)(L-I)},$$

where I denotes the $(d+1) \times (d+1)$ identity matrix.

Lemma 15. *The real part of every eigenvalue of $L - I$ is non-positive and there exist eigenvalues whose real part is 0. In the Jordan canonical form for $L - I$, each of the purely imaginary eigenvalues corresponds to a 1×1 (i.e. trivial) Jordan block.*

Proof. If $a, b : [1, \infty) \rightarrow \mathbb{R}$ are functions, we will use the asymptotic notation $a(t) \asymp b(t)$ to denote that there exist constants $C_1, C_2 > 0$ such that for every $t \geq 1$ we have

$$C_1 \leq \frac{a(t)}{b(t)} \leq C_2.$$

Recall from Section 2 that, because ω is badly approximable, the transference principle implies that there is a $K' > 0$ such that each spacing satisfies

$$\Delta_i(t) \leq \frac{K'}{t^d}.$$

Focusing on the smallest spacing, Part (1) of Theorem A' implies that $\Delta_1(t) = s_j(t)u_1(t)$ for some $s_j(t)$ in the finite set \mathcal{S} . This gives

$$\Delta_1(t) = (1, \omega_1, \dots, \omega_d) S_j(t) U(t) \mathbf{e}_1 \leq \frac{K'}{t^d},$$

where $S_j(t)$ is the matrix representing multiplication by $s_j(t)$ in the basis $\{1, \omega_1, \dots, \omega_d\}$. Since the vector $(\omega_1, \dots, \omega_d)$ is badly approximable, we find that there is a constant $C_1 > 0$ such that

$$|S_j(t) U(t) \mathbf{e}_1| > C_1 t$$

for every $t \geq 1$.

Also observe that, since

$$\Delta_1(t) = \{(\mathbf{m}_1 - \mathbf{m}_2) \cdot \omega\}$$

for some $\mathbf{m}_1, \mathbf{m}_2 \in M(t)$, there is a constant $C_2 > 0$ such that for every $t \geq 1$ we have

$$|S_j(t) U(t) \mathbf{e}_1| < C_2 t.$$

In summary, we have

$$|S_j(t) U(t) \mathbf{e}_1| \asymp t.$$

Since the $S_j(t)$ range over a finite set of invertible matrices (corresponding to multiplication by elements of the finite set S) we conclude that

$$(19) \quad |U(t) \mathbf{e}_1| \asymp t.$$

For each $1 \leq i \leq d$, let W_i denote multiplication by ω_i in the basis $\{1, \omega_1, \dots, \omega_d\}$. We then have

$$|U(t) \mathbf{e}_{i+1}| = |U(t) W_i \mathbf{e}_1| = |W_i U(t) \mathbf{e}_1| \asymp t.$$

The last equality holds because of Lemma 11, and the last assertion follows from (19) because W_i is invertible.

Finally, since $U(t) = A(t)\tilde{U}(t)$ with $A(t)$ given by (18) and hence varying over a compact set of invertible matrices, we have

$$|\tilde{U}(t) \mathbf{e}_i| \asymp t.$$

for each $1 \leq i \leq d+1$. In other words, this gives that for each $1 \leq i \leq d+1$ we have

$$(20) \quad \left| e^{\log t(L-1)} \mathbf{e}_i \right| = \left| \frac{1}{t} \tilde{U}(t) \mathbf{e}_i \right| \asymp 1.$$

Now we will use the Jordan decomposition $L - I = PJP^{-1}$, where P is an invertible matrix and J is in Jordan form. It follows from the power series definition of the matrix exponential that

$$e^{(\log t)(L-1)} = P e^{(\log t)J} P^{-1}.$$

Since P is invertible, (20) implies for each $1 \leq i \leq d+1$ that we have

$$(21) \quad \left| e^{(\log t)J} P^{-1} \mathbf{e}_i \right| \asymp 1.$$

Let us consider the upper left Jordan block, which we suppose is $k \times k$:

$$J_1 = \begin{bmatrix} \lambda & 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & \lambda & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & \lambda & 1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & \lambda \end{bmatrix} = \lambda I + N,$$

where N is the nilpotent matrix whose only non-zero entries are ones on the ‘‘super-diagonal’’ of J_1 . It satisfies $N^k = 0$. Since I commutes with every matrix we have

$$e^{(\log t)J_1} = e^{(\log t)\lambda I} e^{(\log t)N} = \text{diag}(e^{\log t\lambda}, \dots, e^{\log t\lambda}) p((\log t)N)$$

where

$$p(x) = 1 + x + \frac{x^2}{2!} + \cdots + \frac{x^{k-1}}{(k-1)!}.$$

From this, one can check that the $(1, k)$ entry of $e^{(\log t)J_1}$ equals

$$\frac{1}{(k-1)!} (\log t)^{k-1} e^{\log t\lambda}$$

and that all other entries have moduli that are smaller, asymptotically as $t \rightarrow \infty$, by at least a factor of $\log t$.

Because $\mathbf{e}_1, \dots, \mathbf{e}_{d+1}$ form a basis for \mathbb{R}^{d+1} there exists $1 \leq j \leq d+1$ such that the k -th entry of $P^{-1} \mathbf{e}_j$ is non-zero. It follows from the previous paragraph that

$$|\mathbf{e}_1 \cdot (e^{(\log t)J} P^{-1} \mathbf{e}_j)| \asymp (\log t)^{k-1} e^{\log t \operatorname{Re}(\lambda)}.$$

Combined with (21) this implies that $\operatorname{Re}(\lambda) \leq 0$ and that if $\operatorname{Re}(\lambda) = 0$ then $k = 1$.

By permuting the Jordan blocks of J we find that the same holds for every other Jordan block. Finally, existence of at least one purely imaginary eigenvalue is needed for the lower bound implied by (21) to hold. \square

Proof of Theorem 13. By Lemma 15, we can write $L - I = PJP^{-1}$ with J being block-diagonal of the form

$$J = \operatorname{diag}(2\pi i\theta_1, \dots, 2\pi i\theta_k, J_{k+1}, \dots, J_\ell)$$

with $\theta_1, \dots, \theta_k \in \mathbb{R}$ for some $1 \leq k \leq d+1$ and with the blocks J_{k+1}, \dots, J_ℓ all corresponding to eigenvalues with real parts less than some $\gamma < 0$. Note that at this step we have selected the angles $\boldsymbol{\theta} = (\theta_1, \dots, \theta_k)$ which are asserted to exist in the statement of the theorem.

Consider the $(d+1) \times (d+1)$ diagonal matrix:

$$\hat{J} := \operatorname{diag}(2\pi i\theta_1, \dots, 2\pi i\theta_k, 0, \dots, 0).$$

Let $e^\gamma < \alpha < 1$. Then, it follows from the calculations of exponentials of Jordan blocks in the end of the proof of Lemma 15 that for any $t \geq 1$ we have

$$(22) \quad e^{(\log t)J} - e^{(\log t)\hat{J}} = \mathcal{O}(\alpha^{\log t}).$$

Here we mean that the modulus of each corresponding component of the difference is $\mathcal{O}(\alpha^{\log t})$.

Define $g_3 : \mathbb{T}^k \times [0, 1]^r \rightarrow \mathbb{R}^{d+1}$ by

$$(23) \quad g_3(\boldsymbol{\psi}, \mathbf{x}) := \operatorname{Re} \left(e^{-x_1 L_1 \dots - x_r L_r} P \operatorname{diag}(e^{2\pi i\psi_1}, \dots, e^{2\pi i\psi_k}, 0, \dots, 0) P^{-1} \mathbf{e}_1 \right).$$

Clarifications:

- (1) Here, as usual, we denote the angles $\boldsymbol{\psi} \in \mathbb{T}^k$ by their lifts in \mathbb{R}^k . However the formula clearly only depends on the angles themselves.
- (2) The Re denotes that we are taking the real part of each component of the resulting vector.
- (3) This function is differentiable, hence Lipschitz.

If we define our rates by $\boldsymbol{\beta} = \mathbf{w}$ (see (14)) then we have

$$g_3(\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\}) = \operatorname{Re} \left(A(t) P e^{\log t \hat{J}} P^{-1} \mathbf{e}_1 \right).$$

Finally, observe that

$$\frac{\mathbf{n}(u_1(t))}{t} = A(t) \left(\frac{1}{t} \tilde{U}(t) \mathbf{e}_1 \right) = A(t) e^{\log t(L-1)} \mathbf{e}_1 = A(t) P e^{\log t \hat{J}} P^{-1} \mathbf{e}_1.$$

Since $A(t)$ ranges over a compact set of matrices, the result follows from (22) and the fact that $\frac{\mathbf{n}(u_1(t))}{t}$ is real. \square

5. QUASIPERIODICITY OF FINITE WORDS AND PROOF OF THEOREM B'

We will show that a relatively simple modification of the proof of Theorem A' yields the following stronger statement.

Theorem C. *Suppose that $1, \omega_1, \dots, \omega_d \in \mathbb{R}$ form a \mathbb{Q} -basis for an algebraic number field Φ and let $r \geq 1$ denote the rank of \mathbb{Z}_Φ^\times . Let \mathcal{S} be the finite set given by Part (1) of Theorem A'.*

For each choice of $1 \leq j_0, \dots, j_l \leq J$ let $p_{j_0, \dots, j_l}(t)$ denote the proportion of the points from $\{y_i(t) : 1 \leq i \leq \ell(t) - l - 1\}$ such that

$$(24) \quad \begin{aligned} \delta_i(t) &= y_{i+1}(t) - y_i(t) = s_{j_0} u(t)^{-1}, \\ \delta_{i+1}(t) &= y_{i+2}(t) - y_{i+1}(t) = s_{j_1} u(t)^{-1}, \\ &\vdots \\ \delta_{i+l-1}(t) &= y_{i+l}(t) - y_{i+l-1}(t) = s_{j_{l-1}} u(t)^{-1}, \quad \text{and} \\ \delta_{i+l}(t) &= y_{i+l+1}(t) - y_{i+l}(t) = s_{j_l} u(t)^{-1}. \end{aligned}$$

Equivalently, $p_{j_0, \dots, j_l}(t)$ is the proportion of the points $y_i(t)$ such that the sequence of $l+1$ consecutive rescaled spacings starting at $y_i(t)$ forms the word $s_{j_0} s_{j_1} \dots s_{j_l}$.

Then, there is an integer $0 \leq k \leq d+1$, a Lipschitz continuous function

$$h : \mathbb{T}^k \times [0, 1]^r \rightarrow P := \left\{ (p_{1, \dots, 1}, \dots, p_{J, \dots, J}) : \sum p_{j_0, \dots, j_l} = 1, p_{j_0, \dots, j_l} \geq 0 \right\},$$

angles $\boldsymbol{\theta} = (\theta_1, \dots, \theta_k) \in \mathbb{T}^k$, rates $\boldsymbol{\beta} = (\beta_1, \dots, \beta_r) \in \mathbb{R}^r$, and $0 < \alpha < 1$ such that

$$(p_{1, \dots, 1, 1}(t), p_{1, \dots, 1, 2}(t), \dots, p_{J, \dots, J, J}(t)) = h(\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\}) + \mathcal{O}(\alpha^{\log t}).$$

Proof. The only changes to the proof of Theorem A' that are necessary are adaptations to Section 4.1 about the partitions.

For any length $l+1$ word $j_0 j_1 \dots j_l \in J^{l+1}$ let $Y_{j_0 j_1 \dots j_l}(t)$ be the subset of those

$$\{y_i(t) : 1 \leq i \leq \ell(t) - l - 1\}$$

such that (24) holds. Let $M_{j_0 j_1 \dots j_l}(t)$ be the set of vectors $\mathbf{m} \in M(t)$ such that $\{\mathbf{m} \cdot \boldsymbol{\omega}\} \in Y_{j_0 j_1 \dots j_l}(t)$. These sets form a partition of $M(t)$ up to the l points corresponding to $\{y_{\ell(t)-l}, \dots, y_{\ell(t)}\}$. As in the proof of Theorem A' we have for any word $j_0 j_1 \dots j_l$ that

$$p_{j_0 j_1 \dots j_l}(t) = \frac{|M_{j_0 j_1 \dots j_l}(t)|}{|M(t)| - l}.$$

Let us first consider the case of words of length two. As in Proposition 7, for any $1 \leq j \leq J$ let

$$\mathbf{v}_j(t) = \mathbf{m}(s_j u_1(t)).$$

Then, we claim that

$$M_{j_0 j_1}(t) = M_{j_0}(t) \cap (M_{j_1}(t) - \mathbf{v}_{j_0}(t)).$$

To see this, notice that if $\mathbf{m}(y_i(t)) = \mathbf{v}$ then to have $\delta_i(t) = s_{j_0} u(t)^{-1}$ we need to have $\mathbf{v} \in M_{j_0}(t)$ and in order to have $\delta_{i+1}(t) = s_{j_1} u(t)^{-1}$ we need to have $\mathbf{v} + \mathbf{v}_{j_0}(t) \in M_{j_1}(t)$. Here, we are using that since $\delta_i(t) = s_{j_0} u(t)^{-1}$ we have $\mathbf{m}(y_{i+1}(t)) = \mathbf{v} + \mathbf{v}_{j_0}(t)$.

A simple induction yields the following formula in the general case:

$$M_{j_0 j_1 \dots j_l}(t) = M_{j_0}(t) \cap (M_{j_1}(t) - \mathbf{v}_{j_0}(t)) \cap \dots \cap (M_{j_l}(t) - \mathbf{v}_{j_0}(t) - \mathbf{v}_{j_1}(t) - \dots - \mathbf{v}_{j_{l-1}}(t)).$$

In each of these formulae the sets $M_j(t)$ are defined as in Section 4.1 and given by (10).

Denote the power set of our region R by $\mathcal{P}(R)$ and let

$$\mathcal{P} : (\mathbb{R}^d)^J \rightarrow \mathcal{P}(R)^{J^l}$$

be the mapping which sends $\mathbf{v} = (\mathbf{v}_1, \dots, \mathbf{v}_J) \in (\mathbb{R}^d)^J$ to $(\mathcal{P}_{11\dots 1}(\mathbf{v}), \dots, \mathcal{P}_{JJ\dots J}(\mathbf{v})) \in \mathcal{P}(R)^{J^l}$, where,

$$\mathcal{P}_{j_0 j_1 \dots j_l}(\mathbf{v}) = P_{j_0}(\mathbf{v}) \cap (P_{j_1}(\mathbf{v}) - \mathbf{v}_{j_0}) \cap \dots \cap (P_{j_l}(\mathbf{v}) - \mathbf{v}_{j_0} - \mathbf{v}_{j_1} - \dots - \mathbf{v}_{j_{l-1}}).$$

Here,

$$P_j(\mathbf{v}) = [R \cap (R - \mathbf{v}_j)] \setminus \bigcup_{i=1}^{j-1} (R - \mathbf{v}_i),$$

is the same formula as from Section 4.1.

We then claim that the appropriately generalized versions of Propositions 9 and 10, expressed in terms of the definitions and formulae above for $M_{j_0 j_1 \dots j_l}(t)$ and $\mathcal{P}_{j_0 j_1 \dots j_l}(\mathbf{v})$, also hold in this context. We leave this to the reader to check. The remainder of the proof holds exactly as the proof of Theorem A' \square

Theorem B' is now an immediate consequence of Theorem C in the case of words of length two. \square (Theorem B')

6. WORKED EXAMPLE: A TOTALLY REAL CUBIC FIELD

In this section we will work out the details of our quasiperiodicity theorem (Part (2) of Theorem A') in a particular example which highlights many of the important steps in its proof. For our example we take a totally real cubic field of smallest discriminant, for which we can use the table in [13] to identify a pair of generators for a finite index subgroup of the group of units in \mathbb{Z}_Φ^\times . We also take the region R to be the half open unit square, so that $R(t) = [0, t)^2$ and $M(t) = \mathbb{Z}^2 \cap [0, t)^2$.

Let ω_1 be the smallest real root of the cubic polynomial $f(x) = x^3 - 7x^2 + 14x - 7$, and let $\omega_2 = \omega_1^2$. Then $1, \omega_1$, and ω_2 form a \mathbb{Q} -basis for the algebraic number field $\Phi = \mathbb{Q}(\omega_1)$ of degree $d + 1 = 3$ over \mathbb{Q} , and the ring of integers of Φ is $\mathbb{Z}_\Phi = \mathbb{Z}[\omega_1]$ (see [13]).

All three of the roots of $f(x)$ are real and positive, so let us list them as $0 < \alpha_1 < \alpha_2 < \alpha_3$ (note that $\omega_1 = \alpha_1$). For each $1 \leq i \leq 3$ let σ_i be the embedding of Φ into \mathbb{R} which maps ω_1 to α_i . It is clear that $r_1 = 3$ and $2r_2 = 0$, so the rank of the group of units is $r_1 + r_2 - 1 = 2$. Now we let $\epsilon_1, \epsilon_2 \in \Phi$ be defined by

$$(25) \quad \epsilon_1 = 2 - 4\alpha_1 + \alpha_1^2 \quad \text{and} \quad \epsilon_2 = -5 + 5\alpha_1 - \alpha_1^2.$$

It follows from [13] that ϵ_1 and ϵ_2 generate a finite index subgroup of the group of units of \mathbb{Z}_Φ .

Next, following (14), we let $\beta = (\beta_1, \beta_2)$ be determined by

$$\begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} = \begin{pmatrix} \log |\epsilon_1| & \log |\epsilon_2| \\ \log |\sigma_2(\epsilon_1)| & \log |\sigma_2(\epsilon_2)| \end{pmatrix}^{-1} \begin{pmatrix} -2 \\ 1 \end{pmatrix},$$

so that

$$\beta \approx (1.96080, -0.70061),$$

and for $t \geq 1$ we set

$$u_1(t) = \epsilon_1^{\lfloor \beta_1 \log t \rfloor} \epsilon_2^{\lfloor \beta_2 \log t \rfloor}.$$

Note for later that the fact the ϵ_1 and ϵ_2 are units also gives that

$$\beta_1 \log |\sigma_3(\epsilon_1)| + \beta_2 \log |\sigma_3(\epsilon_2)| = -\beta_1 \log |\epsilon_1 \sigma_2(\epsilon_1)| - \beta_2 \log |\epsilon_2 \sigma_2(\epsilon_2)| = 1.$$

The first claim of Theorem A' is that there is a finite set $S \subseteq \Phi$ with the property that, for any $t \geq 1$, every spacing $\Delta_i(t)$ has the form $u_1(t)s$ for some $s \in S$. In order to identify such a set, we must first derive an upper bound for $\Delta_i(t)$. As indicated by our proof above, we will do this using Cassels's transference principle. First notice that if $(m_1, m_2) \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}$ satisfies $|m_1|, |m_2| \leq t$, and if $m_0 \in \mathbb{Z}$ is chosen so that

$$\|\mathbf{m} \cdot \boldsymbol{\omega}\| = m_0 + m_1 \omega_1 + m_2 \omega_2,$$

then we have that

$$|m_0| \leq |m_1\omega_1 + m_2\omega_2| + 1/2 \leq (1/2 + \alpha_1 + \alpha_1^2)t.$$

This in turn gives for $j = 2$ and 3 that

$$|m_0 + m_1\alpha_j + m_2\alpha_j^2| \leq (1/2 + \alpha_1 + \alpha_1^2 + \alpha_j + \alpha_j^2)t,$$

and it follows that

$$\|\mathbf{m} \cdot \boldsymbol{\omega}\| = \frac{\text{Norm}(m_0 + m_1\alpha_1 + m_2\alpha_1^2)}{|m_0 + m_1\alpha_2 + m_2\alpha_2^2| \cdot |m_0 + m_1\alpha_3 + m_2\alpha_3^2|} \geq \frac{1}{Kt^2},$$

with

$$K = (1/2 + \alpha_1 + \alpha_1^2 + \alpha_2 + \alpha_2^2)(1/2 + \alpha_1 + \alpha_1^2 + \alpha_3 + \alpha_3^2).$$

The inhomogeneous transference principle [11, Section V, Theorem VI] then implies that, for any $t \geq 1$ and for any real number γ , there is an integer solution $(n_1, n_2) \in \mathbb{Z}^2$ to the inequality

$$\|\mathbf{n} \cdot \boldsymbol{\omega} - \gamma\| \leq \frac{\lfloor K \rfloor + 1}{2Kt^2}$$

satisfying

$$|n_1|, |n_2| \leq \left(\frac{\lfloor K \rfloor + 1}{2} \right) t.$$

Rescaling, we conclude that there is always an integer solution $(n_1, n_2) \in \mathbb{Z}^2$ to the inequality

$$\|\mathbf{n} \cdot \boldsymbol{\omega} - \gamma\| \leq \frac{(\lfloor K \rfloor + 1)^3}{8Kt^2},$$

with $|n_1|, |n_2| \leq t$. It follows that, for $t \geq 1$, every spacing $\Delta_i(t)$ must satisfy the inequality

$$(26) \quad \Delta_i(t) \leq \frac{(\lfloor K \rfloor + 1)^3}{4Kt^2}.$$

Since every spacing $\Delta_i(t)$ must be of the form $\mathbf{m} \cdot \boldsymbol{\omega}$ for some $\mathbf{m} \in M(t)$, from the above discussion we also have for $j = 2$ and 3 that

$$\sigma_j(\Delta_i(t)) \leq (1/2 + \alpha_1 + \alpha_1^2 + \alpha_j + \alpha_j^2)t.$$

Next, with help from the math software Sage, we find that

$$\begin{aligned} |u_1(t)^{-1}| &= |\epsilon_1|^{-\lfloor \beta_1 \log t \rfloor} |\epsilon_2|^{-\lfloor \beta_2 \log t \rfloor} \\ &= t^2 \exp(\{\beta_1 \log t\} \log |\epsilon_1| + \{\beta_2 \log t\} \log |\epsilon_2|) \\ &\leq |\epsilon_2| t^2, \end{aligned}$$

that

$$\begin{aligned} |\sigma_2(u_1(t)^{-1})| &= |\sigma_2(\epsilon_1)|^{-\lfloor \beta_1 \log t \rfloor} |\sigma_2(\epsilon_2)|^{-\lfloor \beta_2 \log t \rfloor} \\ &= t^{-1} \exp(\{\beta_1 \log t\} \log |\sigma_2(\epsilon_1)| + \{\beta_2 \log t\} \log |\sigma_2(\epsilon_2)|) \\ &\leq \frac{|\sigma_2(\epsilon_1)| |\sigma_2(\epsilon_2)|}{t}, \end{aligned}$$

and that

$$\begin{aligned} |\sigma_3(u_1(t)^{-1})| &= |\sigma_3(\epsilon_1)|^{-\lfloor \beta_1 \log t \rfloor} |\sigma_3(\epsilon_2)|^{-\lfloor \beta_2 \log t \rfloor} \\ &= t^{-1} \exp(\{\beta_1 \log t\} \log |\sigma_3(\epsilon_1)| + \{\beta_2 \log t\} \log |\sigma_3(\epsilon_2)|) \\ &\leq \frac{|\sigma_3(\epsilon_1)|}{t}. \end{aligned}$$

This means that the Minkowski embedding $\sigma(u_1(t)^{-1}\Delta_i)$ of $u_1(t)^{-1}\Delta_i$ into \mathbb{R}^3 is a point of the lattice $\Gamma = \sigma(\mathbb{Z}_\Phi)$ which lies in the box

$$[-K_1, K_1] \times [-K_2, K_2] \times [-K_3, K_3],$$

with

$$\begin{aligned} K_1 &= \frac{(\lfloor K \rfloor + 1)^3 |\epsilon_2|}{4K}, \\ K_2 &= (1/2 + \alpha_1 + \alpha_1^2 + \alpha_2 + \alpha_2^2) |\sigma_2(\epsilon_1)| |\sigma_2(\epsilon_2)|, \text{ and} \\ K_3 &= (1/2 + \alpha_1 + \alpha_1^2 + \alpha_3 + \alpha_3^2) |\sigma_3(\epsilon_1)|. \end{aligned}$$

Therefore, for our finite set \mathcal{S} we may take the collection of all points of Γ which lie in this box. Unfortunately, here there is a bit of a disappointment. The volume of the box defined above is approximately 11,034,177, while a fundamental domain for Γ has volume 7. This means that the number of lattice points in the box is close to 10^6 . While it is not computationally infeasible to find and list all of these points, further computations of the areas of the regions from Proposition 9 become unwieldy. They are also somewhat unenlightening, because most of the regions end up being empty- in all cases we have computed, which includes all $t \leq 300$, there are no more than 10 distinct spacings. However, we can still continue further to explore the quasiperiodic behavior of the function g_3 from the statement of Theorem 13.

The linear transformations E_1 and E_2 of Γ determined by multiplication by ϵ_1 and ϵ_2 in Φ (with respect to the basis $\sigma(1), \sigma(\omega_1), \sigma(\omega_2)$ of Γ) are given by

$$E_1 = \begin{pmatrix} 2 & 7 & 21 \\ -4 & -12 & -35 \\ 1 & 3 & 9 \end{pmatrix} \quad \text{and} \quad E_2 = \begin{pmatrix} -5 & -7 & -14 \\ 5 & 9 & 21 \\ -1 & -2 & -5 \end{pmatrix}.$$

These matrices commute, and they are diagonalizable, therefore they are simultaneously diagonalizable. Explicitly, let

$$\lambda_1 = 2 - \alpha_1, \quad \lambda_2 = 2 - 4\alpha_1 + \alpha_1^2, \quad \lambda_3 = -5 + 5\alpha_1 - \alpha_1^2,$$

let

$$D_1 = \text{diag}(\lambda_1, \lambda_2, \lambda_3), \quad D_2 = \text{diag}(\lambda_2, \lambda_3, \lambda_1),$$

and let

$$Q = \begin{pmatrix} 1 & 1 & 1 \\ -3 + 2\alpha_1 - (3/7)\alpha_1^2 & -\alpha_1 + (1/7)\alpha_1^2 & -1 - \alpha_1 + (2/7)\alpha_1^2 \\ 1 - (5/7)\alpha_1 + (1/7)\alpha_1^2 & (1/7)\alpha_1 & (4/7)\alpha_1 - (1/7)\alpha_1^2 \end{pmatrix}.$$

Then we have for $i = 1$ and 2 that

$$E_i = QD_iQ^{-1}.$$

Noting that $\lambda_1 > 0$ while $\lambda_2, \lambda_3 < 0$, and choosing a branch of the logarithm which includes both the positive and negative real axes, a pair of commuting logarithms of E_1 and E_2 is given by

$$\begin{aligned} L_1 &= Q \text{diag}(\log \lambda_1, \log |\lambda_2| + i\pi, \log |\lambda_3| + i\pi) Q^{-1}, \text{ and} \\ L_2 &= Q \text{diag}(\log |\lambda_2| + i\pi, \log |\lambda_3| + i\pi, \log \lambda_1) Q^{-1}. \end{aligned}$$

Now, following the proof of Theorem 13, we take $L = \beta_1 L_1 + \beta_2 L_2$. In this case, again using Sage, we have that

$$L - I = PJP^{-1}$$

with

$$P \approx \begin{pmatrix} -0.52319 & -0.48157 & 0.82671 \\ 0.83239 & 0.83647 & -0.55556 \\ -0.18274 & -0.26156 & 0.08893 \end{pmatrix}$$

and

$$J \approx \text{diag}(6.16003 i, -2.20103 i, -3.00000 + 3.95900 i).$$

These numbers have been computed using (complex) double float precision, but for readability we have rounded them to five digits. This means that $k = 2$ in the statement of Theorems A' and 13, and that

$$\boldsymbol{\theta} \approx \left(\frac{6.16003}{2\pi}, \frac{-2.20103}{2\pi} \right).$$

Finally, with g_3 defined as in (23), we have that

$$\frac{\mathbf{n}(u_1(t))}{t} = g_3(\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\}) + O(\alpha^{\log t}),$$

for any $\alpha > e^{-3}$. Below is a table comparing the actual values of $\mathbf{n}(u_1(t))$ with the approximate values given by $tg_3(\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\})$. The values of t have been sampled along the sequence $\lfloor 10^{i/2} \rfloor$.

i	$t = \lfloor 10^{i/2} \rfloor$	$\mathbf{n}(u_1(t))$	$tg_3(\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\})$
1	3	(-5, 8, -2)	(-4.80194, 7.86690, -1.97869)
2	10	(-3, 4, 0)	(-3.02177, 4.01463, -0.00234)
3	31	(-41, 68, -18)	(-40.99761, 67.99839, -17.99974)
4	100	(186, -308, 81)	(186.00012, -308.00008, 81.00001)
5	316	(-20, 74, -63)	(-20.00001, 74.00001, -63.00000)
6	1000	(424, -609, 61)	(424.00000, -609.00000, 61.00000)

Consistent with our observations above, this data indicates that the error in approximating $\mathbf{n}(u_1(t))$ by $tg_3(\boldsymbol{\theta} \log t, \{\boldsymbol{\beta} \log t\})$ is roughly on the order of magnitude of $1/t^2$. In conclusion, this is an example in which the frequencies with which the elements of \mathcal{S} appear in Theorem A' are determined quasiperiodically by a linear flow on a two dimensional torus with flow direction determined by $\boldsymbol{\theta}$ and a linear flow on $[0, 1]^2$ with flow direction determined in $\boldsymbol{\beta}$.

REFERENCES

- [1] YY Atas, E Bogomolny, O Giraud, and G Roux. Distribution of the ratio of consecutive level spacings in random matrix ensembles. *Physical review letters*, 110(8):084101, 2013.
- [2] Michael Victor Berry and Michael Tabor. Level clustering in the regular spectrum. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 356(1686):375–394, 1977.
- [3] Valérie Berthé and Dong Han Kim. Some constructions for the higher-dimensional three-distance theorem. *Acta Arith.*, 184(4):385–411, 2018.
- [4] P. M. Bleher. “The Energy Level Spacing for Two Harmonic Oscillators with Golden Mean Ratio of Frequencies.” *Journal of Statistical Physics Belgium*, 61 (3-4): 869-876, November 1990.
- [5] P. M. Bleher. “The Energy Level Spacing for Two Harmonic Oscillators with Generic Ratio of Frequencies.” *Journal of Statistical Physics Belgium*, 63 (1-2): 261-283, April 1991.
- [6] Pavel M. Bleher, Freeman J. Dyson, and Joel L. Lebowitz. Non-Gaussian energy level statistics for some integrable systems. *Phys. Rev. Lett.*, 71(19):3047–3050, 1993.

- [7] P. M. Bleher, Y. Homma, L. L. Ji, R. K. W. Roeder, J. D. Shen: *Nearest neighbor distances on a circle: multidimensional case*, J. Stat. Phys. 146 (2012), no. 2, 446–465.
- [8] Oriol Bohigas, Marie-Joya Giannoni, and Charles Schmit. Characterization of chaotic quantum spectra and universality of level fluctuation laws. *Physical review letters*, 52(1):1, 1984.
- [9] M. D. Boshernitzan: Letter to F. J. Dyson, December 9, 1991.
- [10] M. D. Boshernitzan: Letter to F. J. Dyson, May 20, 1992.
- [11] J. W. S. Cassels: *An introduction to Diophantine approximation*, Cambridge Tracts in Mathematics and Mathematical Physics, No. 45. Cambridge University Press, New York, 1957.
- [12] N. Chevallier: *Three distance theorem and grid graph*, Discrete Math. 223 (2000), no. 1-3, 355-362.
- [13] T. Cusick, L. Schoenfeld: *A table of fundamental pairs of units in totally real cubic fields*, Math. Comp. 48 (1987), no. 177, 147–158.
- [14] F. J. Dyson: *Nearest neighbor distances on a circle*, IAS Preprint IASSNS-HEP-92/27.
- [15] A. S. Fraenkel, R. Holzman: *Gaps problems for integer part and fractional part sequences*, J. Number Theory 50 (1995), 66–86.
- [16] J. F. Geelen, R. J. Simpson: *A two-dimensional Steinhaus theorem*, Australas. J. Combin. 8 (1993), 169–197.
- [17] B. Hall: *Lie groups, Lie algebras, and representations. An elementary introduction.*, Graduate Texts in Mathematics, 222. Springer-Verlag, New York, 2003.
- [18] A. Haynes, J. Marklof: *Higher dimensional Steinhaus and Slater problems via homogeneous dynamics*, Ann. Sci. Ec. Norm. Super. 53 (2020) 537–557.
- [19] Dennis A. Hejhal, Joel Friedman, Martin C. Gutzwiller, and Andrew M. Odlyzko, editors. *Emerging applications of number theory*, volume 109 of *The IMA Volumes in Mathematics and its Applications*. Springer-Verlag, New York, 1999. Papers from the IMA Summer Program held at the University of Minnesota, Minneapolis, MN, July 15–26, 1996.
- [20] R. Horn, C. Johnson: *Matrix Analysis*, Cambridge University Press, (1985) ISBN 978-0-521-38632-6
- [21] V. Jarník: *Zur metrischen Theorie der diophantischen Approximationen*, Puce mar. fiz. 36 (1928), 91–106.
- [22] J. Marklof, A. Strömbergsson: *The three gap theorem and the space of lattices*, Amer. Math. Monthly 124 (2017), no. 8, 741–745.
- [23] Vadim Oganesyan and David A Huse. Localization of interacting fermions at high temperature. *Physical review b*, 75(15):155111, 2007.
- [24] O. Perron: *Über diophantische Approximationen*, Math. Ann. 83 (1921), no. 1-2, 77–84.
- [25] W. M. Schmidt: *Badly approximable systems of linear forms*, J. Number Theory 1 (1969), 139–154.
- [26] V. T. Sós: *On the theory of diophantine approximations I*, Acta Math. Acad. Sci. Hungar. 8 (1957), 461–472.
- [27] J. Surányi: *Über die Anordnung der Vielfachen einer reellen Zahl mod 1*, Ann. Univ. Sci. Budapest Eötvös Sect. Math. 1 (1958) 107–111.
- [28] S. Świerczkowski: *On successive settings of an arc on the circumference of a circle*, Fund. Math. 46 (1959), 187–189.
- [29] H. P. F. Swinnerton-Dyer: *A brief guide to algebraic number theory*, London Mathematical Society Student Texts, 50, Cambridge University Press, Cambridge, 2001.

AH: Department of Mathematics,
 University of Houston,
 Houston, TX, United States.

haynes@math.uh.edu

RR: Department of Mathematical Sciences
Indiana University–Purdue University Indianapolis,
Indianapolis, IN, United States.
rroeder@math.iupui.edu