



## Article

# Views from women and maternity care professionals on routine discussion of previous trauma in the perinatal period: A qualitative evidence synthesis

Cull, Joanne, Thomson, Gill, Downe, Soo, Fine, Michelle and Topalidou, Anastasia

Available at <https://clock.uclan.ac.uk/46134/>

*Cull, Joanne orcid iconORCID: 0000-0001-8990-154X, Thomson, Gill orcid iconORCID: 0000-0003-3392-8182, Downe, Soo orcid iconORCID: 0000-0003-2848-2550, Fine, Michelle and Topalidou, Anastasia orcid iconORCID: 0000-0003-0280-6801 (2023) Views from women and maternity care professionals on routine discussion of previous trauma in the perinatal period: A qualitative evidence synthesis. PLoS One .*

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.1101/2022.10.25.22281395>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

# PLOS ONE

## Views from women and maternity care professionals on routine discussion of previous trauma in the perinatal period: A qualitative evidence synthesis --Manuscript Draft--

<b>Manuscript Number:</b>	PONE-D-22-29747R1
<b>Article Type:</b>	Research Article
<b>Full Title:</b>	Views from women and maternity care professionals on routine discussion of previous trauma in the perinatal period: A qualitative evidence synthesis
<b>Short Title:</b>	Discussing previous trauma in maternity care: a qualitative evidence synthesis
<b>Corresponding Author:</b>	Joanne Joanne Cull University of Central Lancashire Preston, UNITED KINGDOM
<b>Keywords:</b>	Trauma-informed care; Adverse Childhood Experiences; perinatal mental health; Psychosocial screening; Psychological trauma; childhood sexual abuse
<b>Abstract:</b>	<p><b>Background</b> Over a third of pregnant women (around 250,000) each year in the United Kingdom have experienced trauma such as domestic abuse, childhood trauma or sexual assault. These experiences can have a long-term impact on women's mental and physical health. This global qualitative evidence synthesis explores the views of women and maternity care professionals on routine discussion of previous trauma in the perinatal period.</p> <p><b>Methods</b> Systematic database searches (MEDLINE, EMBASE, CINAHL Plus, APA PsycINFO and Global Index Medicus) were conducted in July 2021 and updated in April 2022. The quality of each study was assessed using the Critical Appraisal Skills Programme. We thematically synthesised the data and assessed confidence in findings using GRADE-CERQual.</p> <p><b>Results</b> We included 25 papers, from five countries, published between 2001 and 2022. All the studies were conducted in high-income countries; therefore findings cannot be applied to low- or middle-income countries. Confidence in most of the review findings was moderate or high.</p> <p>The findings are presented in six themes. These themes described how women and clinicians felt trauma discussions were valuable and worthwhile, provided there was adequate time and appropriate referral pathways. However, women often found being asked about previous trauma to be unexpected and intrusive, and women with limited English faced additional challenges. Many pregnant women were unaware of the extent of the trauma they have suffered, or its impact on their lives. Before disclosing trauma, women needed to have a trusting relationship with a clinician; even so, some women chose not to share their histories. Hearing trauma disclosures could be distressing for clinicians.</p> <p><b>Conclusion</b> Discussions of previous trauma should be undertaken when women want to have the discussion, when there is time to understand and respond to the needs and concerns of each individual, and when there are effective resources available for follow up if needed. Continuity of carer should be considered a key feature of routine trauma discussion, as many women will not disclose their histories to a stranger. All women should be provided with information about the impact of trauma and how to independently access support in the event of non-disclosures. Care providers need support to carry out these discussions.</p>
<b>Order of Authors:</b>	<p>Joanne Cull</p> <p>Gill Thomson</p> <p>Soo Downe</p> <p>Michelle Fine</p> <p>Anastasia Topalidou</p>

<b>Opposed Reviewers:</b>	
<b>Response to Reviewers:</b>	Please see 'response to reviewers' document attached.
<b>Additional Information:</b>	
<b>Question</b>	<b>Response</b>
<p><b>Financial Disclosure</b></p> <p>Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the <a href="#">submission guidelines</a> for detailed requirements. View published research articles from <a href="#">PLOS ONE</a> for specific examples.</p> <p>This statement is required for submission and <b>will appear in the published article</b> if the submission is accepted. Please make sure it is accurate.</p> <p><b>Unfunded studies</b> Enter: <i>The author(s) received no specific funding for this work.</i></p> <p><b>Funded studies</b> Enter a statement with the following details:</p> <ul style="list-style-type: none"> <li>• Initials of the authors who received each award</li> <li>• Grant numbers awarded to each author</li> <li>• The full name of each funder</li> <li>• URL of each funder website</li> <li>• Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript?</li> <li>• <b>NO</b> - Include this sentence at the end of your statement: <i>The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.</i></li> <li>• <b>YES</b> - Specify the role(s) played.</li> </ul> <p>* typeset</p>	<p>Joanne Cull is funded by a National Institute for Health Research (NIHR) Wellbeing of Women Doctoral Fellowship (grant number NIHR301525). This paper presents independent research funded by the National Institute for Health Research (NIHR) and the charity Wellbeing of Women. The views expressed are those of the authors and not necessarily those of Wellbeing of Women, the NHS, the NIHR or the Department of Health and Social Care. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.</p> <p><a href="https://www.nihr.ac.uk/">https://www.nihr.ac.uk/</a>  <a href="https://www.wellbeingofwomen.org.uk/">https://www.wellbeingofwomen.org.uk/</a></p>
<p><b>Competing Interests</b></p> <p>Use the instructions below to enter a competing interest statement for this submission. On behalf of all authors, disclose any <a href="#">competing interests</a> that could be perceived to bias this work—acknowledging all financial support and any other relevant financial or non-</p>	<p>The authors have declared that no competing interests exist.</p>

financial competing interests.

This statement is **required** for submission and **will appear in the published article** if the submission is accepted. Please make sure it is accurate and that any funding sources listed in your Funding Information later in the submission form are also declared in your Financial Disclosure statement.

View published research articles from [PLOS ONE](#) for specific examples.

**NO authors have competing interests**

Enter: *The authors have declared that no competing interests exist.*

**Authors with competing interests**

Enter competing interest details beginning with this statement:

*I have read the journal's policy and the authors of this manuscript have the following competing interests: [insert competing interests here]*

\* typeset

**Ethics Statement**

Enter an ethics statement for this submission. This statement is required if the study involved:

- Human participants
- Human specimens or tissue
- Vertebrate animals or cephalopods
- Vertebrate embryos or tissues
- Field research

Write "N/A" if the submission does not require an ethics statement.

General guidance is provided below. Consult the [submission guidelines](#) for

N/A

detailed instructions. **Make sure that all information entered here is included in the Methods section of the manuscript.**

#### Format for specific study types

##### Human Subject Research (involving human participants and/or tissue)

- Give the name of the institutional review board or ethics committee that approved the study
- Include the approval number and/or a statement indicating approval of this research
- Indicate the form of consent obtained (written/oral) or the reason that consent was not obtained (e.g. the data were analyzed anonymously)

##### Animal Research (involving vertebrate animals, embryos or tissues)

- Provide the name of the Institutional Animal Care and Use Committee (IACUC) or other relevant ethics board that reviewed the study protocol, and indicate whether they approved this research or granted a formal waiver of ethical approval
- Include an approval number if one was obtained
- If the study involved *non-human primates*, add *additional details* about animal welfare and steps taken to ameliorate suffering
- If anesthesia, euthanasia, or any kind of animal sacrifice is part of the study, include briefly which substances and/or methods were applied

##### Field Research

Include the following details if this study involves the collection of plant, animal, or other materials from a natural setting:

- Field permit number
- Name of the institution or relevant body that granted permission

##### Data Availability

Authors are required to make all data underlying the findings described fully available, without restriction, and from the time of publication. PLOS allows rare

Yes - all data are fully available without restriction

exceptions to address legal and ethical concerns. See the [PLOS Data Policy](#) and [FAQ](#) for detailed information.

A Data Availability Statement describing where the data can be found is required at submission. Your answers to this question constitute the Data Availability Statement and **will be published in the article**, if accepted.

**Important:** Stating 'data available on request from the author' is not sufficient. If your data are only available upon request, select 'No' for the first question and explain your exceptional situation in the text box.

Do the authors confirm that all data underlying the findings described in their manuscript are fully available without restriction?

**Describe where the data may be found in full sentences. If you are copying our sample text, replace any instances of XXX with the appropriate details.**

- If the data are **held or will be held in a public repository**, include URLs, accession numbers or DOIs. If this information will only be available after acceptance, indicate this by ticking the box below. For example: *All XXX files are available from the XXX database (accession number(s) XXX, XXX).*
- If the data are all contained **within the manuscript and/or Supporting Information files**, enter the following: *All relevant data are within the manuscript and its Supporting Information files.*
- If neither of these applies but you are able to provide **details of access elsewhere**, with or without limitations, please do so. For example:

*Data cannot be shared publicly because of [XXX]. Data are available from the XXX Institutional Data Access / Ethics Committee (contact via XXX) for researchers who meet the criteria for*

This is a systematic review; references for the included papers are given in the manuscript.

*access to confidential data.*

*The data underlying the results presented in the study are available from (include the name of the third party and contact information or URL).*

- This text is appropriate if the data are owned by a third party and authors do not have permission to share the data.

\* typeset

Additional data availability information:

Dear Sir / Madam,

Thank you for considering our submission of this global qualitative evidence synthesis which explores routine discussion of previous trauma within maternity care.

Increasing recognition of the prevalence and long-term impact of trauma such as sexual abuse and violence has led to calls for universal screening for previous trauma within maternity care. However, research to date has failed to show clear evidence of benefit, and there are concerns that this sensitive conversation could be re-traumatising for women, increase unnecessary safeguarding referrals or stigmatise women with a history of trauma.

This synthesis is the first to bring together the views of women and maternity care professionals on routine discussion of previous trauma in the perinatal period, and we believe the findings will be of great interest and relevance to readers of PLOS ONE. Supporting open science, the paper is uploaded as a preprint in medRxiv at <https://doi.org/10.1101/2022.10.25.22281395>.

Yours faithfully,

Joanne Cull, National Institute for Health Research Wellbeing of Women Doctoral Fellow,  
Professor Gill Thomson, Professor Soo Downe, Distinguished Professor Michelle Fine,  
and Dr Anastasia Topalidou



## Title

Full title: Views from women and maternity care professionals on routine discussion of previous trauma in the perinatal period: A qualitative evidence synthesis

## Authors

Joanne Cull<sup>1\*</sup>

Roles: Conceptualisation, Formal analysis, Investigation, Methodology, Writing - original draft preparation, Writing - review and editing

Gill Thomson<sup>1</sup>

Roles: Formal analysis, Methodology, Supervision, Writing - review and editing

Soo Downe<sup>1</sup>

Roles: Supervision, Writing - review and editing

Michelle Fine<sup>2</sup>

Roles: Supervision, Writing - review and editing

Anastasia Topalidou<sup>1</sup>

Roles: Supervision, Writing - review and editing

---

<sup>1</sup> School of Community Health and Midwifery, University of Central Lancashire, Preston, England

<sup>2</sup> Public Science Project, The Graduate Center, City University of New York, New York, USA

\* Corresponding author. Email: [jcull@uclan.ac.uk](mailto:jcull@uclan.ac.uk) (JC). Orcid iD 0000-0001-8990-154X

# Abstract

## Background

Over a third of pregnant women (around 250,000) each year in the United Kingdom have experienced trauma such as domestic abuse, childhood trauma or sexual assault. These experiences can have a long-term impact on women's mental and physical health. This global qualitative evidence synthesis explores the views of women and maternity care professionals on routine discussion of previous trauma in the perinatal period.

## Methods

Systematic database searches (MEDLINE, EMBASE, CINAHL Plus, APA PsycINFO and Global Index Medicus) were conducted in July 2021 and updated in April 2022. The quality of each study was assessed using the Critical Appraisal Skills Programme. We thematically synthesised the data and assessed confidence in findings using GRADE-CERQual.

## Results

We included 25 papers, from five countries, published between 2001 and 2022. All the studies were conducted in high-income countries; therefore findings cannot be applied to low- or middle-income countries. Confidence in most of the review findings was moderate or high.

The findings are presented in six themes. These themes described how women and clinicians felt trauma discussions were valuable and worthwhile, provided there was adequate time and appropriate referral pathways. However, women often found being asked about previous trauma to be unexpected and intrusive, and women with limited English faced additional challenges. Many pregnant women were unaware of the extent of the trauma they have suffered, or its impact on their lives. Before disclosing trauma, women needed to have a trusting relationship with a clinician; even so, some women chose not to share their histories. Hearing trauma disclosures could be distressing for clinicians.

## **Conclusion**

Discussions of previous trauma should be undertaken when women want to have the discussion, when there is time to understand and respond to the needs and concerns of each individual, and when there are effective resources available for follow up if needed. Continuity of carer should be considered a key feature of routine trauma discussion, as many women will not disclose their histories to a stranger. All women should be provided with information about the impact of trauma and how to independently access support in the event of non-disclosures. Care providers need support to carry out these discussions.

# Introduction

Over a third of pregnant women (~250,000) each year in the United Kingdom (UK) have experienced significant trauma such as violence or sexual abuse in childhood or adulthood [1]. Exposure to trauma can have a severe and prolonged impact on mental health, physical health, and health seeking behaviours [2-4]. Internationally, preventing trauma and reducing its impact is a public health priority [5].

Some women who have experienced trauma will have recovered from their experiences at the time of pregnancy, while others begin the pregnancy with unresolved trauma which negatively affects their mental health and the maternal-infant bond [6, 7]. Women who have experienced trauma may find the perinatal period particularly challenging and find that aspects of maternity care, such as clinical procedures, trigger memories of their previous abuse [8].

An increased understanding of the long-term effects of trauma exposure on both mother and baby have led to calls for universal screening within maternity care [9]. The American College of Obstetricians and Gynecologists recommend that women's healthcare providers screen all women for current and past trauma [10]. In the UK, whilst National Institute for Health and Care Excellence guidelines do not address routine enquiry about previous trauma, anecdotally a number of maternity hospitals have introduced it at the initial midwife appointment. However, concerns have also been raised that this could be re-traumatising for women [11], increase unnecessary or unwarranted safeguarding referrals [12], or stigmatise women with a history of adverse events [13].

It is evident that routinely discussing prior trauma with pregnant women requires careful consideration and sensitivity to ensure these conversations create value rather than cause harm. To understand whether routine trauma discussion should be carried out, and if so how it should be done, we aimed to collate and synthesise all relevant qualitative studies to capture women's and maternity care professionals' views of this issue.

We use the term 'routine' to indicate raising the issue of previous trauma with all women accessing maternity services, as opposed to selectively for women who the healthcare professional suspects may have suffered trauma. The term 'routine' in this context does not negate the need for personalised care. We have chosen to use the term 'discussion' rather than 'enquiry' as discussion leaves open the possibility that trauma could be raised in a way that does not involve a direct question.

The review question was 'What are the views of women and maternity care professionals on routine discussion of previous trauma in the perinatal period?'. The study was registered in PROSPERO with the reference number CRD42021247160 [14]. The review was informed by guidance produced by the Cochrane Effective Practice and Organisation of Care group on carrying out and reporting qualitative evidence syntheses [15-17].

# Methods

## Reflexive note

We considered our pre-existing beliefs on routine trauma discussion and how these might influence the design and conduct of the review [18].

JC is a midwife and doctoral student. She was unsure at the outset whether routine trauma discussion was beneficial to women. She believed that for some groups of women, who face discrimination based on their class, race, immigration status or a range of other factors, disclosure of trauma could be harmful and increase the risk of unnecessary or unwanted safeguarding or mental health involvement. SD is a midwife with 18 years of clinical practice and a maternity care researcher, and held similar beliefs to JC. GT has a psychology academic background, and a long history of undertaking research with women who have experienced perinatal mental health problems. Her views were that conversations that were trauma-informed were important to enable needs-led care. AT is a researcher in the field of maternal and neonatal care, and her views were that an appropriate supporting model of care is needed to enable trauma-informed conversations, that will be beneficial to women. MF teaches critical psychology in the US and is a visiting professor in South Africa. With years of participatory work with women in and out of prison, highly marginalized young people and most recently Muslim American youth, MF is interested in collective and individual trauma as sites of wounds and creativity, memory and activism, a rich and painful source of knowledge and wisdom.

A 'research collective' comprising experts by lived experience, from the voluntary sector and healthcare professionals is supporting the doctoral study of which this review is part. Involvement of the collective in the design of the review helped minimise the risk that our pre-understandings would influence the review. Further, the range of backgrounds and views on the review team also helped enhance the rigor of the analysis.

#### **Search strategy and selection criteria**

We searched the databases MEDLINE, CINAHL Plus, EMBASE, APA Psycinfo, and Global Index Medicus using the search terms 'trauma-informed' and 'trauma informed'.

We developed the search strategy by reviewing key relevant papers we were already aware of and examining relevant systematic reviews on PROSPERO. Each of the key papers referred to 'trauma-informed' in their keywords, title or abstract. We trialled searching using this term on Medline and duplicated it using a variety of different terms such as trauma-sensitive, trauma-focused, and trauma-responsive. These alternative terms had very few hits and had always been used in addition to trauma-informed. We therefore decided to only use the search terms 'trauma-informed' or 'trauma informed' and hand select relevant papers that concerned the care of women in the perinatal period.

Forward and backward citation tracking and key author searches for studies included in the review were carried out to identify additional relevant studies. Searches were conducted in July 2021 and updated in April 2022.

Only qualitative studies and qualitative aspects of mixed methods studies were included in the review. Studies not based in a maternity setting, or which did not include women in the perinatal period (defined for this purpose as pregnancy and up to one year after birth) were excluded. The review was focused on previous psychological trauma: this could include all past trauma or specific types such as adverse childhood experiences or sexual abuse but excluded studies with participants who have experienced physical trauma such as injury. Studies published at any date, and in any language were eligible for inclusion in the review.

Two reviewers (GT and JC) independently screened 20% of papers at title and abstract stage using Rayyan blind screening. The level of agreement was set at 95%. As the reviewers achieved 100% agreement, the remaining 80% of papers were reviewed by JC. At the second stage, all papers were blind screened by two reviewers (GT and JC), with any differences of opinion about inclusion resolved through discussion.



### **Data extraction and management**

Papers eligible for inclusion were uploaded to the software program MAXQDA Plus 2020 for data extraction, analysis, and thematic synthesis. We developed a standardised data extraction form using Excel and piloted it prior to beginning data collection. JC used the form to record basic contextual and methodological information about each study, including bibliographic information, country of study, setting, study design, data collection, participants' characteristics, data analysis methods and key themes. To reduce bias and errors, GT independently extracted data from 20% of studies, and JC and GT compared results, resolving disagreements through dialogue.

### **Appraisal of the methodological quality of included studies**

JC assessed the quality of each included paper using the Critical Appraisal Skills Programme (CASP) quality assessment checklist for qualitative studies [19]. The CASP checklist comprises 10 questions: 1 mark was allocated to each question if the criterion was met. The overall quality of each study was categorised as 'strong' (score 8-10/10; minimal methodological issues), 'adequate' (score 5-7/10; no major methodological issues) or 'weak' (0-4/10; major methodological issues). Studies scoring 'weak' were excluded on quality grounds to ensure that the credibility or trustworthiness of the review findings was not compromised by including studies with important methodological limitations [15]. GT independently quality assessed 20% of included studies, and JC and GT compared results, resolving disagreement through re-examination and discussion. As suggested by Carroll and Booth [20] post hoc sensitivity analyses were carried out to

assess the impact on the review findings of excluding a study on quality grounds. Quality ratings contributed to the GRADE-CERQual assessments (described below).

When we were quality appraising each study, we also reflected on broader issues of the integration and inclusion of women's voices, and these additional considerations are reflected in the discussion.

### **Data synthesis**

Data were synthesised thematically using the method developed by Thomas and Harden [21]. First, the findings of each study were inductively coded on a line-by-line basis. The codes were then organised into related areas, constructing 'descriptive' themes (summaries of findings). Finally, the descriptive themes were organised into analytical themes. This work was led by JC, and, at all stages, emerging concepts were shared, discussed, and refined with the review team.

### **Assessment of confidence in the review findings**

The GRADE 'Confidence in the Evidence from Reviews of Qualitative research' (GRADE-CERQual) approach was used to assess confidence in the synthesised findings [22-27]. This approach facilitates explicit and transparent assessment of whether the findings from a qualitative synthesis reasonably represent the phenomenon of interest. Each summary of findings (descriptive theme) was assessed in terms of methodological limitations, coherence, adequacy of data, and relevance; with the four assessments contributing to an overall assessment of confidence. The summaries of findings and associated CERQual assessment of confidence are presented in an Evidence Profile and Summary of Qualitative Findings table.

### **Results of the search**

Overall, 3,888 papers were identified after removal of duplicates, of which 25 met the inclusion criteria for the study. The PRISMA flow chart can be found at Figure 1. Three papers were excluded due to quality [28-30]. Post hoc examination of the three excluded papers indicated that their inclusion in the review would not have altered the final themes.

**<insert figure 1 here>**

### **Description of the studies**

The characteristics and quality of the 25 studies included in the review are summarised in Table 1. The included studies were published between 2001 and 2022. In terms of

study setting, 12 of the studies were carried out in Australia, nine in the United States, two in Sweden, and one each in England and Canada. Routine trauma discussion was explored from the perspective of women in thirteen of the papers, eight looked at the perspective of healthcare professionals, and the remaining four papers looked at both viewpoints. Most data were collected by individual interviews, focus groups and/or surveys. The studies represented the views of 1602 women and 286 healthcare professionals and experts from the voluntary sector.

**<Insert Table 1 here>**

#### **Assessment of methodological strengths and limitations**

Of the 25 included papers, sixteen were assessed as methodologically strong and nine as adequate. As discussed above, a further three papers were assessed as methodologically weak and were excluded from the review. A lack of reflexivity was noted across most of the studies. Other common methodological weaknesses included insufficient information about data analysis, the lack of a clear statement of findings, and minimal discussion of ethical issues.

#### **Confidence in the review findings**

Table 2 shows the Summary of Findings and CERQual rating for each summary of finding. Confidence in most of the review findings was moderate or high, reflecting the quality and quantity of the studies included in the review. Each summary of finding was mapped to

an analytical theme and these themes are discussed in the next section. All studies that met the inclusion criteria were from high-income countries. Because these findings were applicable to the UK, we did not downgrade the CERQual ratings for income status of the countries. However, the findings are not applicable to low- and middle- income countries.

<Insert Table 2 here>

## Findings

Six analytical themes were identified relating to women's and maternity care providers' views and experiences of routine trauma discussion. The first theme *'I did not know how to say it, and no-one asked me'* considers whether maternity care providers should ask women about previous trauma. *'A real whitefella way to start'* explores standardised compared to more unstructured ways of asking about trauma. In the theme *'You say it is confidential... but you are going to report me'*, fear of judgement as a barrier to disclosure, and the importance of trust and relationships in trauma discussions is highlighted. The theme *'I'm not quite sure what is going on, but I feel really vulnerable'* calls attention to the intensity of the perinatal period, which is often challenging but also has the potential for healing and growth. *Embedding trauma in routine practice* explores barriers and facilitators to successful implementation. Finally, *'You go home and it's playing on your mind as you're cooking'* considers the impact on care providers of hearing trauma disclosures.

**'I did not know how to say it, and no-one asked me': should maternity care providers ask women about previous trauma?**

This theme explores whether women and maternity care providers feel that routine trauma discussion should take place.

Participants in 14 studies expressed that they felt routine trauma discussion is acceptable and worthwhile [31-44]. One of the participants in the study undertaken by Montgomery, Seng and Chang (2021) [36] proposed:

*"It might have just put a thought in my head, even if it wasn't something that I shared with anybody, it might have just put a thought in my head which might have been useful at some point" [36].*

Some women felt it was difficult to broach the subject of previous trauma and would not have disclosed unless the clinician raised the issue:

*"At the time, I could not and did not tell the healthcare professionals of my survivor status. I did not know how to say it, and no one asked me" [36].*

The overall finding that women accepted routine trauma discussion masks several complexities and contradictions. In eight studies, women reported feeling unprepared for

the discussion and found it intrusive [32, 33, 35-39, 42]. This was reflected by a woman who was interviewed in the study by Millar et al. (2021) [35]:

*“Like it doesn’t feel good when you first meet someone, and they just start like trying to jump into your life. Like they know you. I hate that”* [35].

Some participants spoke of how they would have welcomed forewarning of the discussion:

*“I think they could have told me what they were going to ask before I even arrived for my appointment. I had no idea that was what was coming”* [39].

In 11 studies, professionals reported that they felt routine trauma discussion was worthwhile [32, 34, 45-53]. A clinician taking part in the study by Flanagan et al. (2018) [47] reflected that while *‘most of the time (the) screen is negative’*, when finding the individual who had faced previous trauma, *‘you’re so glad you did’* [47].

Participants in the study by Kohlkoff et al. (2021) felt there were three key benefits of routine trauma discussion: identify women at higher risk of mental health problems or family violence, increase referrals to appropriate support services, and provide support and education [47]. However, in seven studies, women felt that trauma discussions should only take place if clinicians had enough time to respond to disclosures and could

provide or refer into appropriate support [32, 38, 39, 41-44]. Having disclosed prior trauma, some women expected that they would be treated more sensitively and that other care providers would be aware of their history, and were aggravated when this wasn't the case [39, 41]:

*“Why don't they take the extra time just to read over [my file] and if they have any more questions about it then they can ask. If it's already there then why bother... it is really frustrating”* [39].

Maternity care providers similarly spoke of their support for routine trauma discussion as contingent on having sufficient time and appropriate referral pathways [32, 34, 45-48, 50-53]. Without good quality support services, clinicians were reluctant to discuss prior trauma, fearing this would open a 'Pandora's box' of issues they were unable to deal with:

*“We see perinatal depression and anxiety but this is a continuum of social disadvantage and intergenerational trauma. We have super complicated patients with so many problems. Where do we fit mental health in where there are so few resources to respond properly?”* [32].

#### **'A real whitefella way to start': standardisation and tick-boxes in trauma discussion**

This theme explores how maternity care providers should raise the issue of previous trauma. Participants in several studies discussed limitations in the use of questionnaires to raise the issue of previous trauma [31, 32, 37, 45, 54]. Participants suggested that closed



questions (for example ‘in the last year, have you experienced...’) can prevent disclosures [31, 37] This Aboriginal participant in a study by Carlin, Atkinson and Marley (2019) [32] proposed:

*“You talk about things because they are important to talk about not cause they happened one week ago! It is a real whitefella way to start. It’s like you’re in or you’re out. You see that hey? Like what happens if it was a bit longer, then the lady might think oh no, it’s not important, I won’t talk about that”* [31].

Mendel, Sperlich and Fava (2021) [54] investigated the use of the Adverse Childhood Experiences questionnaire (ACE-10) within maternity research. The researchers found that the questionnaire contains confusing and ambiguous questions, excludes important traumatic events in childhood (such as the death of a parent) and fails to ascertain the severity or duration of the traumatic experience. For example, one participant in the study had an ACE score of one out of ten but had suffered extensive abuse over 12 years of her childhood, resulting in seven miscarriages. The authors concluded that completion of the questionnaire might not give a true representation of the extent of trauma the woman has suffered.

Women who took part in the study by Carlin et al. (2020) [32] felt that direct questions could cause women to disengage, and that broad, gentle questions were the best approach to ask about difficult experiences. The voluntary sector experts and healthcare professionals who participated in the study by Chamberlain et al. (2020) [45] noted that

direct questions about trauma can be problematic, because avoiding thinking about trauma experiences can be a way of coping. Instead, indirect methods of gentle communication were preferred, asked by *“someone trusted – this sort of information will naturally become evident so the trusted person can gently empathise and draw attention to, as opposed to ask directly and abruptly”* [45]. Another participant noted that clinicians should ask *‘slowly, gently and only where there is the possibility of being able to ‘hold a space’ and deal appropriately with the answer’* [45].

Carlin et al. (2020) [32] explored clinicians’ views of a questionnaire-based approach versus questionnaire plus narrative. Several participants reported that they only used the questionnaire, describing the narrative section as ‘aspirational’ due to time constraints and concerns that it positioned them as a counsellor. However, those who did use the narrative approach felt it enhanced rapport with women and that women understood the limits around the assistance they could provide:

*“Generally I think women are keen to share some of their problems with us as nurses even though we cannot solve these issues as such but we can listen, we can advise them where to seek help and how we can assist as a support for some of their problems.”* [32].

#### **‘You say it is confidential... but you are going to report me’: the importance of trust**

This theme concerns the importance of trust and relationships to women who have suffered trauma. In twelve studies, fear of judgement was reported as a reason for non-

disclosure [31, 32, 36-38, 40-45, 51]. This included a general fear of being perceived as a bad parent, and specific concerns that their child would be removed from their care; *“You guys are bound by law [to report certain things]... You say it is confidential... but you are going to report me”* [43]. Concerns about confidentiality were raised by participants in several studies, as was the misconception that the abused becomes the abuser. One mother stated:

*“Speaking from personal experience, I felt at times that my past trauma was being used to assess the likelihood I would harm my own child, rather than as a means of identifying what support I might need as an individual”* [38].

Participants in 13 studies highlighted the importance of a trusting relationship, built through multiple encounters, as a prerequisite for trauma disclosure [31-33, 35-40, 42, 44, 45, 52] A mother from the study by Millar et al. undertaken in 2021 [35] suggested:

*“I think if I had a relationship, then yes I [would disclose trauma history]. But with the amount of time I was seeing them, no. ‘Cause I was always seeing someone different”* [35].

Participants in five studies proposed that trauma should not be discussed at the first appointment, but at a later appointment, enabling a relationship to be built first [35, 38, 42, 45, 52]. This woman chose to disclose later in pregnancy:

*“When she gave me the initial, you know, the history form... when I saw ‘were you abused?’ I said no. There was no way I was going to tell her”* [40].

Although maternity care providers agreed that women were more likely to disclose prior trauma after a relationship with the clinician had been established, some felt it was appropriate to raise trauma at the first consultation, seeing this as the start of an ongoing conversation:

*“If... they’re not opening up... they’ll go home and think about it and reflect on that and they may come back the next time and open up a bit more. It’s just opening the door, isn’t it?”* [51].

In twelve studies, participants described the importance of how the clinician asked about previous trauma [31-34, 37-41, 43-45]. Desired attributes consistently included kindness, friendliness, sensitivity, a non-judgemental attitude, respect, care, and compassion. One woman who took part in the study carried out by Sobel et al. (2018) [41] expressed that *“I opened up to my midwife because I felt comfortable with her. That’s it”* [41] Conversely, this participant in Seng et al.’s study (2002) [40] felt the need to change care provider:

*“The doctor was kind of cold, not personable at all, and those feelings [emotional memory of being abused, shame, vulnerability, nakedness] would come back to me in his office, and I found myself crying at every visit”* [40].

For women with limited English, non-verbal signals like smiling and a relaxed manner were vital in inspiring trust: *“She make me like not scared because she smile a lot, her smiling and the way she spoke was really helpful”* [39].

Participants proposed that trauma discussions should be held in private, comfortable and welcoming surroundings [33, 38, 43, 45]. The ideal combination of a trusted care provider and warm environment is summed up by a woman who took part in Gokhale et al.’s study (2020) [33]:

*“The only way my health care providers can help me with my trauma is every time I come, make me feel like I’m at home. Make me feel comfortable. Make me feel safe and make me feel like I have nothing to worry about”*.

Participants in three studies felt that if the conversation was handled badly, routine trauma discussion could impact upon future health care access and experiences [36, 38, 39]. One respondent to the survey carried out by Olsen, Galloway and Guthman (2021) [38] proposed:

*“If it isn’t asked about in a sensitive way under the right circumstances, it could feel really intrusive or could be so upsetting or off-putting that someone could avoid needed health care entirely”* [38]

Participants in thirteen studies reported that they would not disclose previous trauma to the healthcare professional looking after them, although not all studies explored the reasons for this [32, 33, 35-41, 44-46, 51]. Mule et al. (2021) carried out a survey asking women whether they had chosen to fully disclose their histories during antenatal psychosocial assessment: 161 women responded that they had not, and completed an open-ended question giving their reasons. The researchers found there were a range of reasons, including lack of trust of the person asking, fear of judgement, use of closed-ended questions and lack of time, but also simply privacy: they did not want to share this information [37]. Similarly, a participant in the study by Olsen, Galloway and Guthman (2021) proposed that *“some people may not be ready”* [38] while a woman taking part in the study by Gokhale et al. (2020) suggested that *“it’s not easy speaking up about situations like that and a lot of people don’t because they don’t feel comfortable or they don’t feel like they could trust people enough to do that”* [33]. This implies that even within a trusting relationship, some women who would benefit from support will choose not to share their histories. Accordingly, participants in the study by Seng et al. (2002) [40] proposed that care providers should assume women are trauma survivors if they display signs or symptoms of trauma, irrespective of whether they have disclosed.

**‘I’m not quite sure what is going on, but I feel really vulnerable’: the intensity of the perinatal period**

This theme explores experiences of the perinatal period for women who have suffered trauma. In seven of the studies, some participants reported feeling that there was no connection between their trauma histories and their current wellbeing and pregnancy [33, 36, 37, 39, 40, 42, 45]:

*“It’s not really affecting me now... my main concern is getting through the pregnancy, not worrying about my past stuff” [39].*

Some women wanted to focus on the pregnancy and a positive future and felt discussion of trauma could trigger distressing feelings. This was the case even when the perinatal period could be expected to bring up strong emotions, such as for this woman whose infant had been murdered:

*“Cause when you come to the visit you want to hear stuff about your baby. You don’t want to keep dwelling on this that happened in the past and you trying to have a happy moment.” [33].*

Some women had not foreseen that their pregnancy would be so difficult: *“It’s hard to put into words because I’m not quite sure what is going on, but I feel really super vulnerable” [40].* Even women who appeared to be far along in recovery and living happy lives were often unprepared for the intensity of the perinatal period:

*“I was really looking forward to the cuddling time with the baby and breastfeeding... I didn’t expect this whole other ugliness” [40].*

Women commonly felt a loss of control over their body, due both to the pregnancy and a sense of powerlessness within maternity care. Vaginal examinations, birth, or even seemingly benign clinical procedures such as blood pressure measurement caused

flashbacks to abuse. Some women feared bodily exposure during labour and birth, with a participant in the study by Sobel et al. (2018) [41] reporting:

*“I was so concerned with being covered up... I would have been devastated [by a vaginal delivery]. I did not know how I was going to keep my clothes on and have a baby”* [41]

Until pregnancy, some women were not fully aware of the trauma they had suffered. Seng et al. (2002) [40] explored this issue in detail through narrative interviews with 15 women who had suffered childhood sexual trauma and subsequently accessed maternity care. At the time of the pregnancy, four of the fifteen women had only a vague understanding that they had been subjected to abuse. Participants described indications in their thoughts and behaviour of the effects of trauma, such as fleeting flashbacks, suicidal intentions, and extreme promiscuity:

*“I realized [later] there were pieces that had been floating around for a long time that I wouldn’t acknowledge prior to [postpartum]”* [40].

Because they had not fully admitted it to themselves, these women were not able to disclose the abuse to their healthcare providers:

*“I knew early that I was not going to deliver vaginally. I knew in my head that I was not going there. So that piece I connected...I don’t know that I drew a real direct line because of how vulnerable I felt. I wasn’t probably ready to acknowledge that...So it was knowing and not knowing at the same time”* [40].



Participants talked about the potential for post-traumatic growth in the perinatal period [31, 40, 41, 46]. However, this was not always an easy process, as described by this participant in the study by Seng et al. (2002) [40]:

*“I kind of knew in some way it was affecting me, but I just couldn’t connect the dots ever... but when I got pregnant it all just came out, came clear, and it was hard, and I’m grateful... and I think it’s going to help me grow past it and deal with it... but pregnancy is enough to deal with.”* [40].

#### **Embedding trauma discussion in routine practice**

This theme investigates how trauma discussion can be introduced and explores barriers and facilitators to successful implementation.

Care providers in four studies reported that they quickly adapted to routine trauma discussion, and found it feasible within their workloads [47, 51-53].

*“I just think it’s the initial getting used to... just even logging into it, and doing all of that was a hassle when I first started. It’s “Oh, this is all so hard.” But it’s so simple now, because we’re used to it... it’s like anything,... any tool that you use over and over again, it becomes more simple”* [53].

Participants in seven studies discussed the challenges of ensuring all women are asked about prior trauma [46-48, 50-53]. Care providers taking part in studies by Choi and Seng (2014) and Stenson, Sidenvall and Heimer (2005) spoke of having preconceived notions of who might or might not have suffered trauma: *“If you have a feeling something isn’t quite right, then it’s easier. But just when you don’t think...”* [52]. Participants in the study undertaken by Schmied et al. (2020) felt *“they’re just not appropriate questions to ask”* [51], or avoided asking the question where a woman was felt to be ‘keeping her distance’. Describing a successful implementation of routine trauma discussion, this participant in Reilly et al.’s study (2020) proposed that key individuals were helpful in getting more reluctant staff on board:

*“Even if you have people who are sceptical, if you are enthusiastic, and the clients get enthusiastic and really feel cared for, that automatically rolls over to the staff members that are sitting on the sidelines and saying “I don’t know if all this is necessary”... They start seeing that it is making a positive impact on people’s lives...”* [50].

Participants in the study by Stenson, Sidenvall and Heimer (2005) [52] found it difficult to remember to raise the issue, resulting in some women not being asked about prior trauma. Documentation of abuse in hand-held notes could be a confidentiality risk, but several people taking part in the study suggested a check box in the records, to act as an aide memoire, and ensure the discussion is accorded the same importance as other issues [52]. Participants in Kohlhoff’s study (2021) [48] proposed setting up a flag on the system to ensure effective information sharing among the maternity team.

Partner presence at appointments could influence the discussion of previous trauma [46, 48, 49, 52]. This was felt by some participants in the study by Stenson, Sidenvall and Heimer (2005) to be a modern phenomenon, with one care provider proposing “*this generation of couples, the husband’s there the whole time*” [52]. Whilst a participant in Kohlhoff et al.’s study (2021) [48] noted that “*pregnant women can be quite vulnerable and anxious, and it is good to have that support person with them*”, another pointed out:

*“They’re not going to be able to divulge anything while their partner’s there, especially if their partner doesn’t know about it, and sometimes that is the case”* [48].

Trauma discussions were more difficult for women with limited English [39, 52, 53]. Women often did not want to disclose sensitive issues through an interpreter, and where family members or partners were acting as interpreters, this provided a further barrier to disclosure. In the study by Stenson, Sidenvall and Heimer (2005) [52], staff reported that partners were often asked to sit in the waiting room while this discussion took place: few partners insisted on being present, but the midwives sometimes felt in a difficult position where the woman had limited English:

*“I would really prefer a professional interpreter but on most occasions the men say ‘no’. They want to do the interpreting”* [52].

**'You go home and it's playing on your mind as you're cooking': the impact on care providers of hearing trauma disclosures**

Hearing trauma disclosures could be challenging for clinicians [34, 46, 48, 52, 55]. Mollart, Newing and Foureur (2009) [55] explored this issue in detail in their Australian study in which they carried out focus groups with 18 midwives who undertook routine discussion of prior trauma. The midwife participants in the study reported that the cumulative, complex disclosures they heard affected them emotionally and impacted on their home and work life:

*"You go home and it's playing on your mind as you're cooking. I don't know how long it usually goes on for, probably till you get that next bad case" [55].*

Some participants in the study reported that they continued to think about trauma disclosures after work:

*"Sometimes I've gone home and actually worried about people, then you've got to remember that they told me this today and they've been living with this for how long? Just keep telling yourself that" [55].*

For some, this impacted on their family life:

*"For me, I explode at home, I don't explode here [at work] because I know that no-one would put up with that kind of behaviour. But I do it to my kids, and that's not very good." [55].*

Marley et al. (2017) interviewed healthcare providers about their experiences of routine trauma discussion with Aboriginal women in Western Australia. A participant in the study expressed: *“At the end of the day, it’s hard not to want to neck a bottle of wine to cope with [hearing their stories]”* [34]. The use of unhealthy coping strategies, such as excess alcohol, was echoed by a participant in Mollart, Newing and Foureur (2020):

*“I can debrief 10, 20, 30 times, and the information is still with me, and I don’t know where to channel that sometimes. Sometimes you channel that into things that are probably not appropriate.”* [55].

Participants in studies undertaken by Marley et al. (2017) [34] and Mollart, Newing and Foureur (2009) [55] felt clinical supervision is vital for midwives carrying out routine trauma discussion, and those with no access to supervision expressed resentment. However, not all care providers who were offered supervision chose to participate in it:

*“The only way I know how to deal with it is I talk to colleagues... even though sometimes when you talk to colleagues, you know they’re thinking about the booking-in they had, they’re only half listening and you haven’t actually been heard”* [55].

Stenson, Sidenvall and Heimer (2005) [52] was the only study to raise the issue of care providers who have experienced trauma themselves. The researchers carried out five focus groups with 21 midwives in Sweden to explore their views on discussing prior and current violence with women. The researchers noted that none of the participants spontaneously raised the issue. When the moderator brought it up *‘several admitted that they had not considered the possibility that colleagues might have been subjected to*

*abuse*" [52]. The group discussed whether midwives could act professionally in this situation and concluded that they could, but it might be difficult if they are still in a violent relationship. The group did not explore whether discussion of previous trauma might be more challenging for midwives with personal experience in this area, or how they could be supported.

## **Discussion**

This qualitative evidence synthesis found that although many women feel positive about routine discussion of previous trauma, this simple statement masks complexity. Women often find the conversation unexpected and intrusive; they expect their care to change after trauma disclosure and are disappointed if it does not; and women with limited English face additional challenges. The findings of the review suggest that many pregnant women are unaware of the extent of the trauma they have suffered, or its impact on their lives. This highlights the difficulty of discussing trauma with women who will be at very different stages of recovery.

The review underscores the importance of an established relationship and trust in trauma discussions. While midwife-led continuity models of care have been shown to be beneficial for mothers and babies, New Zealand is the only country to have achieved this at scale, and the national target of implementation of continuity of carer in the UK has now been removed [56-58]. Notwithstanding this, our review found that many women will

not disclose previous trauma in the absence of a trusting relationship and will consequently fail to receive the care they need. Some women will choose not to disclose their histories; this is problematic because in the included studies, women who disclosed trauma were generally provided with information and follow-up care, while women who did not disclose trauma were not. Consideration should therefore be given to providing all women with information about the impact of trauma and the means to independently access support should they wish to do so.

Our review highlights specific areas of training which are necessary for maternity care professionals. The focus of trauma discussions should not be eliciting specific details of past trauma, but rather ascertaining what resources, support, and / or adjustments to the care plan would be helpful. It is important that women are forewarned of the discussion, including any limits to confidentiality and that disclosure of past trauma is entirely voluntary. The review found that partner presence can be a barrier to trauma discussions. National Institute for Health and Care Excellence guidance (2021) [59] states that maternity care providers should ensure women are asked about domestic abuse in a private, one-to-one discussion; we propose that this also applies to discussion of previous trauma. For both women who have suffered trauma and professionals, support for routine discussion of previous trauma was found to be contingent on adequate time to explore these complex issues. This is a serious caveat in maternity systems which were already understaffed and have been further impacted by the COVID-19 pandemic, both in the UK and internationally [60]. Description of the discussion as easy and quick by some clinician participants in the included studies

implies an unknowingly 'performative' approach to routine trauma discussion in which professionals feel they have given women the opportunity to disclose, but women may not feel comfortable disclosing or get the support they need.

This performative approach could also be a protective mechanism for clinicians who feel trauma discussions are emotionally difficult. Midwifery work carries a heavy emotional burden, with midwives suffering significantly higher levels of burnout, stress, anxiety, and depression than the general population [61-63]. Added to this, the review found that hearing women's histories of sexual assault, childhood abuse, and domestic violence can profoundly affect clinicians. None of the studies in the review explored whether discussion of previous trauma might be more challenging for midwives who have personally suffered trauma, or how they could be supported. There is a need for an appropriate support and supervision model which allows midwives the opportunity to explore the challenging situations they are exposed to. Reflective supervision would enable clinicians experiencing vicarious traumatisation or those whose own trauma experience have been triggered through their interactions with survivors to receive the support they need.

Women's perceptions and experiences of trauma discussion might be affected by characteristics such as class, ethnicity, or immigration status, but it was not possible to differentiate and draw distinct themes for different groups of participants. For example, the review found with high confidence that women are fearful of disclosing their histories



in case it raises concerns that they are unable to safely care for their child: it may be that this is a greater barrier for women from population groups who experience disproportionately high levels of child welfare involvement [64]. Women with insecure immigration status are highly vulnerable to violence and abuse [65], but were excluded from almost all of the studies by default as their inclusion criteria required that participants could speak English: this is an area where further research is needed.

In 2014, the US Substance Abuse and Mental Health Services administration (SAMHSA) [66] published a seminal conceptual document about trauma-informed care, a model of service delivery which aims to develop an environment in which people who have experienced trauma feel safe and can build trust with their healthcare provider. This framework has been further developed and tailored to the perinatal period by others including Seng and Taylor (2015) [67], Sperlich et al. (2017) [68] and Law et al. (2021) [9]. SAMHSA (2014) [66] propose that trauma-informed approaches are underpinned by four assumptions: staff realise the prevalence of trauma and its impact on behaviour; they are able to recognise the signs of trauma, including by trauma screening; the organisation's policies and procedures take into consideration the experiences of trauma among service users and staff; and active steps are taken to resist re-traumatisation of clients and staff by ensuring psychological safety. Within a trauma-informed care framework, implementation of routine screening for previous trauma should be considered within the context of much broader changes to services, including staff training, continuity of carer, emotional support for staff, and service evaluation [9, 66-68]. Further research is needed

to ascertain what resources and support are helpful to women who have suffered trauma, bearing in mind that this must focus on the preferences and needs of the individual.

### **Strengths and limitations of the review**

This synthesis is the first to bring together the views of women and maternity care professionals on routine discussion of previous trauma in the perinatal period. We adopted an inclusive approach to the search, with broad search terms and multiple search strategies used to ensure no key articles were missed. The findings are strengthened by the large number of participants. We minimised the risk of over or under interpretation of the data through explicit positionality, reflexivity, and discussion.

However, the synthesis has limitations. All studies that met the inclusion criteria were from high-income countries, therefore the findings cannot be applied to women in low- or middle-income settings. Further, there were no studies set in Asia, and only in three in Europe. Research is needed on the optimal ways of addressing previous trauma in pregnancy and the perinatal period for different populations, in line with the Sustainable Development Goals and guidance by the European Parliament that sexual and reproductive health services should consider the needs of women who have suffered sexual or gender-based violence [69, 70].

## **Conclusion**

The review provides insight into the barriers and facilitators of women in high-income countries sharing their trauma histories, and clinicians asking about previous trauma. Areas for future research are highlighted. Discussions of previous trauma are complex and require careful consideration and sensitivity. While implementation of continuity of carer is no longer a target in maternity services in the UK, many women will not disclose previous trauma in the absence of a trusting relationship. As some women will not disclose their histories, consideration should be given to providing all women with information about the impact of trauma, and means of independently accessing support. These findings also highlight the need for time to undertake authentically sensitive and tailored discussions about trauma, and for appropriate support and supervision for care providers. There is a need for routine trauma discussions to be co-developed with childbearing women and the midwives who will be initiating the conversation, to ensure that women receive the support they need, and the wellbeing of care providers is protected.

## **Acknowledgements**

We would like to thank the EMPATHY study Research Collective for generously sharing their knowledge and expertise. Members of the collective are: Dr Laura Abbott, Juliet Albert, Kirsty Armstrong, Jill Benjoya Miller, Dr Emma Brooks, Dr Geraldine Butcher, Jo Doherty, Amber Jackson, Isobel Martin, Dr Elsa Montgomery, Sam Pointon, Sarah-Jayne

Pomeroy, Erjola Sadria, Gill Skene, Memuna Sowe, Dr Kim Thomas, and Lucy Warwick-Guasp.

## References

1. Crown Prosecution Service. Violence Against Women and Girls Report. England: 2019.
2. Bellis M, Hughes K, Hardcastle K, Ashton K, Ford K, Quigg Z, et al. The impact of adverse childhood experiences on health service use across the life course using a retrospective cohort study. *J Health Serv Res Policy*. 2017;22(3):168-77.
3. Bellis MA, Hughes K, Ford K, Ramos Rodriguez G, Sethi D, Passmore J. Life course health consequences and associated annual costs of adverse childhood experiences across Europe and North America: a systematic review and meta-analysis. *The Lancet Public Health*. 2019;4(10):e517-28.
4. Hughes K, Bellis MA, Hardcastle KA, Sethi D, Butchart A, Mikton C, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *The Lancet Public Health*. 2017;2(8):e356-66.
5. Sara G, Lappin J. Childhood trauma: psychiatry's greatest public health challenge? *The Lancet Public Health*. 2017;2(7):e300-1.
6. Muzik M, Ads M, Bonham C, Lisa Rosenblum K, Broderick A, Kirk R. Perspectives on trauma-informed care from mothers with a history of childhood maltreatment: A qualitative study. *Child Abuse Neglect*. 2013;37(12):1215-24.
7. Seng JS, D'Andrea W, Ford JD. Complex Mental Health Sequelae of Psychological Trauma Among Women in Prenatal Care. *Psychological trauma: theory, research, practice and policy*. 2014;6(1):41-9.
8. Montgomery E, Chang Y, Youssef A, Ivan F, Maxted F, Cheadle S, et al. *Pregnancy, Birth and Parenthood after Childhood Sexual Abuse*. The Survivors Trust. 2019.
9. Law C, Wolfenden L, Sperlich M, Taylor J. A good practice guide to support implementation of trauma-informed care in the perinatal period. NHS England and NHS Improvement; 2021.

10. American College of Obstetricians and Gynecologists. Caring for Patients Who Have Experienced Trauma: ACOG Committee Opinion Summary, Number 825. *Obstet Gynecol.* 2021;137(4):757-8.
11. Ford K, Hughes K, Hardcastle K, Di Lemma LCG, Davies AR, Edwards S, et al. The evidence base for routine enquiry into adverse childhood experiences: A scoping review. *Child Abuse & Neglect.* 2019;91:131-46.
12. Underwood E. California has begun screening for early childhood trauma, but critics urge caution. *Science (American Association for the Advancement of Science).* 2020.
13. Racine N, Killam T, Madigan S. Trauma-Informed Care as a Universal Precaution: Beyond the Adverse Childhood Experiences Questionnaire. *JAMA Pediatrics.* 2020;174(1):5-6.
14. Cull J, Thomson G, Downe S, Fine M, Topalidou A. What are the views of women and maternity care professionals on routine discussion of previous trauma in the perinatal period? A qualitative evidence synthesis. PROSPERO CRD42021247160. . 2021.
15. Glenton C, Bohren M, Downe S, Paulson E, Lewin S, On behalf of Effective Practice and Organisation of Care, (EPOC). EPOC Qualitative Evidence Synthesis: Protocol and review template. Version 1.3. EPOC Resources for review authors. Oslo: Norwegian Institute of Public Health; 2022 b.
16. Glenton C, Lewin S, Downe S, Paulsen E, Munabi-Babigumira S, Johansen M, et al. Qualitative Evidence Syntheses Within Cochrane Effective Practice and Organisation of Care: Developing a Template and Guidance. *International Journal of Qualitative Methods.* 2021;20:16094069211041959.
17. Glenton C, Lewin S, Downe S, Paulsen E, Munabi-Babigumira S, Agarwal S, et al. Cochrane Effective Practice and Organisation of Care (EPOC) Qualitative Evidence Syntheses, Differences From Reviews of Intervention Effectiveness and Implications for Guidance. *International Journal of Qualitative Methods.* 2022 a;21:16094069211061950.
18. Walsh D, Downe S. Appraising the quality of qualitative research. *Midwifery.* 2006 Jun;22(2):108-19.
19. CASP Qualitative Research Checklist [Internet].; 2018 [cited 2.1.2020 2020]. Available from: <https://casp-uk.net/wp-content/uploads/2018/01/CASP-Qualitative-Checklist-2018.pdf>.
20. Carroll, C., and Booth, A. Quality assessment of qualitative evidence for systematic review and synthesis: Is it meaningful, and if so, how should it be performed? . *Research Synthesis Methods.* 2015;6(2):149-54.

21. Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Medical Research Methodology*. 2008;8(1):45.
22. Lewin S, Bohren M, Rashidian A, Munthe-Kaas H, Glenton C, Colvin CJ, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 2: how to make an overall CERQual assessment of confidence and create a Summary of Qualitative Findings table. *Implementation Science*. 2018 b;13(1):10.
23. Lewin S, Booth A, Glenton C, Munthe-Kaas H, Rashidian A, Wainwright M, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings: introduction to the series. *Implementation Science*. 2018 a;13(2).
24. Munthe-Kaas H, Bohren MA, Glenton C, Lewin S, Noyes J, Tunçalp Ö, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 3: how to assess methodological limitations. *Implementation Science*. 2018;13(1):9.
25. Colvin CJ, Garside R, Wainwright M, Munthe-Kaas H, Glenton C, Bohren MA, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 4: how to assess coherence. *Implementation Science*. 2018;13(1):13.
26. Glenton C, Carlsen B, Lewin S, Munthe-Kaas H, Colvin CJ, Tunçalp Ö, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 5: how to assess adequacy of data. *Implementation Science*. 2018;13(1):14.
27. Noyes J, Booth A, Lewin S, Carlsen B, Glenton C, Colvin CJ, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 6: how to assess relevance of the data. *Implementation Science*. 2018 b;13(1):4.
28. Matthey S, White T, Phillips J, Taouk R, Chee TT, Barnett B. Acceptability of routine antenatal psychosocial assessments to women from English and non-English speaking backgrounds. *Archives of Women's Mental Health*. 2005;8(3):171-80.
29. Rollans M, Schmied V, Kemp L, Meade T. 'We just ask some questions...' the process of antenatal psychosocial assessment by midwives. *Midwifery*. 2013 a;29(8):935-42.
30. Rollans M, Kohlhoff J, Meade T, Kemp L, Schmied V. Partner involvement: negotiating the presence of partners in psychosocial assessment as conducted by midwives and child and family health nurses. *Infant Ment Health J*. 2016;37(3):302-12.
31. Carlin E, Atkinson D, Marley JV. 'Having a Quiet Word': Yarning with Aboriginal Women in the Pilbara Region of Western Australia about Mental Health and Mental Health Screening during the Perinatal Period. *International Journal of Environmental Research and Public Health*. 2019;16(21):4253.

32. Carlin E, Spry E, Atkinson D, Marley JV. Why validation is not enough: Setting the scene for the implementation of the Kimberley Mum's Mood Scale. *PLOS ONE*. 2020;15(6):e0234346.
33. Gokhale P, Young MR, Williams MN, Reid SN, Tom LS, O'Brian CA, et al. Refining Trauma-Informed Perinatal Care for Urban Prenatal Care Patients with Multiple Lifetime Traumatic Exposures: A Qualitative Study. *Journal of Midwifery and Women's Health*. 2020;65(2):224-30.
34. Marley JV, Kotz J, Engelke C, Williams M, Stephen D, Coutinho S, et al. Validity and Acceptability of Kimberley Mum's Mood Scale to Screen for Perinatal Anxiety and Depression in Remote Aboriginal Health Care Settings. *PLOS ONE*. 2017;12(1):e0168969.
35. Millar HC, Lorber S, Vandermorris A, Thompson G, Thompson M, Allen L, et al. "No, you need to explain what you are doing": Obstetric care experiences and preferences of adolescent mothers with a history of childhood trauma. *J Pediatr Adolesc Gynecol*. 2021;34(4):538-45.
36. Montgomery E, Seng JS, Chang Y. Co-production of an e-resource to help women who have experienced childhood sexual abuse prepare for pregnancy, birth, and parenthood. *BMC Pregnancy & Childbirth*. 2021 Jan 07;21(1):30.
37. Mule V, Reilly NM, Schmied V, Kingston D, Austin MV. Why do some pregnant women not fully disclose at comprehensive psychosocial assessment with their midwife? *Women and Birth*. 2021;35(1):80-6.
38. Olsen JM, Galloway EG, Guthman PL. Exploring women's perspectives on prenatal screening for adverse childhood experiences. *Public Health Nurs*. 2021 Nov;38(6):997-1008.
39. Rollans M, Schmied V, Kemp L, Meade T. Digging over that old ground: an Australian perspective of women's experience of psychosocial assessment and depression screening in pregnancy and following birth. *BMC Womens Health*. 2013;13(1):18.
40. Seng JS, Sparbel KJH, Low LK, Killion C. Abuse-related posttraumatic stress and desired maternity care practices: women's perspectives. *J Midwifery Womens Health*. 2002;47(5):360-70.
41. Sobel L, O'Rourke-Suchoff D, Holland E, Remis K, Resnick K, Perkins R, et al. Pregnancy and Childbirth After Sexual Trauma: Patient Perspectives and Care Preferences. *Obstetrics & Gynecology*. 2018;132(6):1461-8.
42. Stenson K, Saarinen H, Heimer G, Sidenvall B. Women's attitudes to being asked about exposure to violence. *Midwifery*. 2001;17(1):2-10.

43. White A, Danis M, Gillece J. Abuse survivor perspectives on trauma inquiry in obstetrical practice. *Arch Womens Ment Health*. 2016;19(2):423-7.
44. Willey SM, Blackmore RP, Gibson-Helm ME, Ali R, Boyd LM, McBride J, et al. "If you don't ask ... you don't tell": Refugee women's perspectives on perinatal mental health screening. *Women and Birth*. 2020;33(5):e429-37.
45. Chamberlain C, Gee G, Gartland D, Mensah FK, Mares S, Clark Y, et al. Community Perspectives of Complex Trauma Assessment for Aboriginal Parents: 'Its Important, but How These Discussions Are Held Is Critical'. *Front Psychol*. 2020 Sep 15;11:2014.
46. Choi KR, Seng JS. Pilot for Nurse-Led, Interprofessional In-Service Training on Trauma-Informed Perinatal Care. *J Contin Educ Nurs*. 2015 Nov;46(11):515-21.
47. Flanagan T, Alabaster A, McCaw B, Stoller N, Watson C, Young-Wolff K. Feasibility and Acceptability of Screening for Adverse Childhood Experiences in Prenatal Care. *Journal of Women's Health*. 2018;27(7):903-11.
48. Kohlhoff J, Cibralic S, Tooke S, Hickinbotham R, Knox C, Roach V, et al. Health professional perspectives on an antenatal mental health screening program in a private hospital. *Aust N Z J Obstet Gynaecol*. 2021;61(6):891-7.
49. Preis H, Whitney C, Kocis C, Lobel M. Saving time, signaling trust: Using the PROMOTE self-report screening instrument to enhance prenatal care quality and therapeutic relationships. *PEC Innovation*. 2022;1:100030.
50. Reilly N, Brake E, Kalra H, Austin M. Insights into implementation of routine depression screening and psychosocial assessment in a private hospital setting: A qualitative study. *Aust N Z J Obstet Gynaecol*. 2020;60(3):419-24.
51. Schmied V, Reilly N, Black E, Kingston D, Talcevska K, Mule V, et al. Opening the door: midwives' perceptions of two models of psychosocial assessment in pregnancy- a mixed methods study. *BMC Pregnancy and Childbirth*. 2020;20(1):451.
52. Stenson K, Sidenvall B, Heimer G. Midwives' experiences of routine antenatal questioning relating to men's violence against women. *Midwifery*. 2005;21(4):311-21.
53. Willey SM, Gibson-Helm ME, Finch TL, East CE, Khan NN, Boyd LM, et al. Implementing innovative evidence-based perinatal mental health screening for women of refugee background. *Women and Birth*. 2020;33(3):e245-55.
54. Mendel WE, Sperlich M, Fava NM. "Is there anything else you would like me to know?": Applying a trauma-informed approach to the administration of the adverse childhood experiences questionnaire. *J Community Psychol*. 2021;49(5):1079-99.



55. Mollart L, Newing C, Foureur M. Midwives' emotional wellbeing: Impact of conducting a Structured Antenatal Psychosocial Assessment (SAPSA). *Women and Birth*. 2009;22(3):82-8.
56. Sandall J, Soltani H, Gates S, Shennan A, Devane D. Midwife-led continuity models versus other models of care for childbearing women. *Cochrane Database of Systematic Reviews*. 2016(4).
57. Bradford BF, Wilson AN, Portela A, McConville F, Fernandez Turienzo C, Homer CSE. Midwifery continuity of care: A scoping review of where, how, by whom and for whom? *PLOS Global Public Health*. 2022;2(10):e0000935.
58. May R, Dunkley-Bent J, Jolly M. Midwifery Continuity of Carer. NHS England; 21.9.22 21.9.22.
59. National Institute for Health and Care Excellence. Antenatal Care: NICE guideline NG201. 2021.
60. Cordey S, Moncrieff G, Cull J, Sarian A, the ASPIRE-COVID 19 CG. 'There's only so much you can be pushed': Magnification of the maternity staffing crisis by the 2020/21 COVID-19 pandemic. *BJOG: Int J Obstet Gy*. 2022;129(8):1408-9.
61. NHS England. Better Births: Improving outcomes of maternity services in England. England: NHS England; 2016.
62. Hunter B, Henley J, Fenwick J, Sidebotham M, Pallant J. Work, Health and Emotional Lives of Midwives in the United Kingdom: The UK WHELM study. Cardiff University: School of Healthcare Sciences; 2018.
63. Sidhu R, Su B, Shapiro KR, Stoll K. Prevalence of and factors associated with burnout in midwifery: A scoping review. *European Journal of Midwifery*. 2020;4.
64. Mason W, Morris K, Webb C, Daniels B, Featherstone B, Bywaters P, et al. Toward Full Integration of Quantitative and Qualitative Methods in Case Study Research: Insights From Investigating Child Welfare Inequalities. *Journal of Mixed Methods Research*. 2020;14(2):164-83.
65. Long J, Wertans E, Harper K, Brennan D, Harvey H, Allen R, et al. Femicide Census: UK Femicides, 2009-2018. 2020.
66. SAMHSA (U.S. Department of Health and Human Services Substance Abuse and Mental Health Services Administration). SAMHSA's Concept of Trauma and Guidance for a Trauma-Informed Approach. 2014.

67. Seng J, Taylor J, editors. Trauma informed care in the perinatal period. Edinburgh: Dunedin Academic Press Ltd.; 2015.

68. Sperlich M, Seng JS, Li Y, Taylor J, BradburyJones C. Integrating trauma-informed care into maternity care practice: Conceptual and practical issues. J Midwifery Womens Health. 2017;62(6):661-72.

69. UN General Assembly. Transforming our world: the 2030 Agenda for Sustainable Development. 2015.

70. Sexual and reproductive health and rights in the EU, in the frame of women's health, European Parliament, (2021).

## **Supporting information**

S1 PRISMA 2020 Checklist

Table 1. Characteristics of included studies

Study no.	Authors	Year	Country	Study design	Participants (number / type)	CASP Quality assessment rating	Focus of paper
1	Carlin, Atkinson and Marley	2019	Australia	Yarning - conversational process involving telling of stories and development of knowledge	15 Aboriginal women	Strong	Women's perspectives
2	Carlin et al.	2020	Australia	Health professionals - online survey or semi-structured interview. Aboriginal women - in depth interviews	18 health professionals 10 Aboriginal women	Strong	Women and clinician perspectives
3	Chamberlain et al.	2020	Australia	Stakeholder workshop	57 key stakeholders, with extensive experience working with Aboriginal families.	Strong	Women and clinician perspectives
4	Choi and Seng	2014	United States	Semi-structured telephone interviews	20 perinatal care providers	Strong	Clinician perspectives
5	Flanagan et al.	2018	United States	Childbearing women - ACE questionnaire plus telephone interview. Clinicians - surveys and focus groups	210 childbearing women; 26 clinicians	Adequate	Women and clinician perspectives
6	Gokhale et al.	2020	United States	Semi-structured interviews and completion of trauma history questionnaire	30 pregnant women	Strong	Women's perspectives
7	Kohlhoff et al.	2021	Australia	Focus groups and semi-structured interviews	Nine midwives, two obstetricians, and one nephrologist	Adequate	Clinician perspectives
8	Marley et al.	2017	Australia	Childbearing women - questionnaire. Clinicians - questionnaire and follow-up interview	81 women; 9 study personnel	Adequate	Women and clinician perspectives
9	Mendel, Sperlich, and Fava	2021	United States	Semi structured interviews	99 first time mothers	Adequate	Women's perspectives
10	Millar et al.	2021	Canada	Questionnaire consisting of ACE-10 plus open- and closed-ended questions; semi-structured interviews	Questionnaire - 29 adolescent mothers Follow-up interview - 5 mothers	Strong	Women's perspectives
11	Mollart, Newing and Foureur	2009	Australia	Focus group interviews	18 midwives from 2 study sites	Strong	Clinician perspectives
12	Montgomery, Seng and Chang	2021	England	Focus groups, interviews, and an online survey	2 focus groups, 2 interviews, 29 responses to online survey	Strong	Women's perspectives
13	Mule et al.	2021	Australia	Open-ended question giving reason for	161 childbearing women	Adequate	Women's perspectives


				non-disclosure of trauma history			
14	Olsen, Galloway and Guthman	2021	United States	Online survey with quantitative and open-ended questions	154 women	Strong	Women's perspectives
15	Reilly et al.	2020	Australia	Semi-structured interviews	3 midwives, 3 obstetricians, 2 managers, 1 mental health worker	Strong	Clinician perspectives
16	Rollans et al.	2013	Australia	Qualitative ethnographic study – observation of antenatal and postnatal appointments plus face to face interviews.	34 observed antenatally; 20 of the same women who were observed during postnatal visit; 31 antenatal interviews, 29 postnatal interviews	Strong	Women's perspectives
17	Schmied et al.	2020	Australia	Survey before introduction of new psychosocial assessment, second survey following implementation; focus groups	First survey - 26 midwives, second survey - 27 midwives (9 midwives completed both). Focus groups - 16 midwives	Strong	Clinician perspectives
18	Seng et al.	2002	United States	Narrative interviews	15 childbearing women	Strong	Women's perspectives
19	Sobel et al.	2018	United States	Semi-structured interviews	20 women with history of sexual trauma; 10 without	Adequate	Women's perspectives
20	Stenson et al.	2001	Sweden	Open-ended written / telephone question about abuse screening in pregnancy	879 women	Adequate	Women's perspectives
21	Stenson, Sidenvall and Heimer	2005	Sweden	Focus groups	21 midwives in 5 focus groups	Adequate	Clinician perspectives
22	White, Danis and Gillece	2015	United States	Focus group	6 women	Strong	Women's perspectives
23	Willey et al.	2020 a	Australia	Focus groups and semi-structured interviews	24 healthcare professionals	Strong	Clinician perspectives
24	Willey et al.	2020 b	Australia	Focus group and semi-structured interviews	22 women who were refugees, 5 women who were migrants	Strong	Women's perspectives
25	Preis et al.	2022	United States	Focus groups and semi-structured interviews	22 healthcare professionals	Adequate	Clinician's perspectives

Table 2. Summary of Findings and CERQual ratings

Review finding	Relevant studies (study numbers as per Table 1)	CERQual assessment of confidence in the evidence	Theme
Women feel positively about routine trauma discussion	14 studies (1,2,6,8,10,12,13,14,16,18,19,20,22,24)	Low	'I did not know how to say it, and no-one asked me': should maternity care providers ask women about previous trauma?
Some women find routine trauma discussion invasive and unexpected	8 studies (2,6,13,14,16,20)	High	
Maternity care providers feel routine trauma discussion is valuable	11 studies (2,3,4,5,7,8,15,17,21,23,25)	Moderate	
Support for routine trauma discussion is contingent on adequate time and resources	16 studies (2,3,4,5,7,8,14,15,16,17,19,20,21,22,23,24)	High	
Women favour a broad, conversational approach to discussing trauma	5 studies (1,2,9,12,13)	Very low	'A real whitefella way to start': standardisation and tickboxes in trauma discussion
Women who have suffered trauma want relationship-based care	6 studies (1,6,10,12,14,18)	Moderate	
Choice and control is important to women	6 studies (1,10,12,14,18,19)	High	
Women want further therapeutic support	7 studies (1,3,6,10,14,22,24)	Moderate	
Women fear judgement if they disclose their histories	12 studies (1,2,3,12,13,14,17,18,19,20,22,24)	High	'You say it is confidential... but you are going to report me': the importance of trust
Relationships are a critical prerequisite to trauma disclosure	13 studies (1,2,3,6,10,12,13,14,16,18,20,21,24)	High	
The manner of the person asking and the environment are also important	12 studies (1,2,3,6,8,13,14,16,18,19,22,24)	High	
If not handled sensitively, trauma discussion could affect future health care access and experiences	3 studies (12,14,16)	Low	
Some women will choose not to disclose previous trauma	13 studies (2,3,4,6,10,12,13,14,16,17,18,19,24)	High	

Some women feel their previous experiences are irrelevant to their current pregnancy	7 studies (3,6,12,13,16,18,20)	Moderate	'I'm not quite sure what is going on, but I feel really vulnerable': the intensity of the perinatal period
The perinatal period can be intense and challenging	10 studies (1,2,4,10,12,14,18,19,20,24)	High	
Not all women were fully aware of the extent or impact of the trauma they had suffered	10 studies (1,2,3,4,6,12,13,14,18,22)	High	
The perinatal period carries potential for healing and growth	7 studies (1,4,6,8,12,18,19)	Moderate	
Embedding trauma discussion in routine practice is challenging	7 studies (4,5,7,15,17,21,23)	Moderate	'Heavens! I forgot it!': challenges to embedding trauma discussion in routine practice
Partner presence can be a barrier to trauma discussion	2 studies (7,21)	Very low	
Women with limited English face additional challenges in discussing trauma	3 studies (16,23,24)	Low	
Hearing trauma disclosures can be distressing for maternity care providers	5 studies (4,7,8,11,21)	Low	'You go home and it's playing on your mind as you're cooking': the impact on care providers of hearing trauma disclosures

Figure 1: PRISMA flow chart





Click here to access/download  
**Supporting Information**  
S1 Checklist .docx





## **Title**

Full title: Views from women and maternity care professionals on routine discussion of previous trauma in the perinatal period: A qualitative evidence synthesis

## **Authors**

Joanne Cull<sup>1\*</sup>

Roles: Conceptualisation, Formal analysis, Investigation, Methodology, Writing - original draft preparation, Writing - review and editing

Gill Thomson<sup>1</sup>

Roles: Formal analysis, Methodology, Supervision, Writing - review and editing

Soo Downe<sup>1</sup>

Roles: Supervision, Writing - review and editing

Michelle Fine<sup>2</sup>

Roles: Supervision, Writing - review and editing

Anastasia Topalidou<sup>1</sup>

Roles: Supervision, Writing - review and editing

---

<sup>1</sup> School of Community Health and Midwifery, University of Central Lancashire, Preston, England

<sup>2</sup> Public Science Project, The Graduate Center, City University of New York, New York, USA

\* Corresponding author. Email: [jcull@uclan.ac.uk](mailto:jcull@uclan.ac.uk) (JC). Orcid iD 0000-0001-8990-154X

# **Abstract**

## **Background**

Over a third of pregnant women (around 250,000) each year in the United Kingdom have experienced trauma such as domestic abuse, childhood trauma or sexual assault. These experiences can have a long-term impact on women's mental and physical health. This global qualitative evidence synthesis explores the views of women and maternity care professionals on routine discussion of previous trauma in the perinatal period.

## **Methods**

Systematic database searches (MEDLINE, EMBASE, CINAHL Plus, APA PsycINFO and Global Index Medicus) were conducted in July 2021 and updated in April 2022. The quality of each study was assessed using the Critical Appraisal Skills Programme. We thematically synthesised the data and assessed confidence in findings using GRADE-CERQual.

## **Results**

We included 25 papers, from five countries, published between 2001 and 2022. All the studies were conducted in high-income countries; therefore findings cannot be applied to

low- or middle-income countries. Confidence in most of the review findings was moderate or high.

The findings are presented in six themes. These themes described how women and clinicians felt trauma discussions were valuable and worthwhile, provided there was adequate time and appropriate referral pathways. However, women often found being asked about previous trauma to be unexpected and intrusive, and women with limited English faced additional challenges. Many pregnant women were unaware of the extent of the trauma they have suffered, or its impact on their lives. Before disclosing trauma, women needed to have a trusting relationship with a clinician; even so, some women chose not to share their histories. Hearing trauma disclosures could be distressing for clinicians.

## **Conclusion**

Discussions of previous trauma should be undertaken when women want to have the discussion, when there is time to understand and respond to the needs and concerns of each individual, and when there are effective resources available for follow up if needed. Continuity of carer should be considered a key feature of routine trauma discussion, as many women will not disclose their histories to a stranger. All women should be provided with information about the impact of trauma and how to independently access support in the event of non-disclosures. Care providers need support to carry out these discussions.



# Introduction

Over a third of pregnant women (~250,000) each year in the United Kingdom (UK) have experienced significant trauma such as violence or sexual abuse in childhood or adulthood [1]. Exposure to trauma can have a severe and prolonged impact on mental health, physical health, and health seeking behaviours [2-4]. Internationally, preventing trauma and reducing its impact is a public health priority [5].

Some women who have experienced trauma will have recovered from their experiences at the time of pregnancy, while others begin the pregnancy with unresolved trauma which negatively affects their mental health and the maternal-infant bond [6, 7]. Women who have experienced trauma may find the perinatal period particularly challenging and find that aspects of maternity care, such as clinical procedures, trigger memories of their previous abuse [8].

An increased understanding of the long-term effects of trauma exposure on both mother and baby have led to calls for universal screening within maternity care [9]. The American College of Obstetricians and Gynecologists recommend that women's healthcare providers screen all women for current and past trauma [10]. In the UK, whilst National Institute for Health and Care Excellence guidelines do not address routine enquiry about previous trauma, anecdotally a number of maternity hospitals have introduced it at the initial midwife appointment. However, concerns have also been raised that this could be re-traumatising for women [11], increase unnecessary or unwarranted safeguarding referrals [12], or stigmatise women with a history of adverse events [13].

It is evident that routinely discussing prior trauma with pregnant women requires careful consideration and sensitivity to ensure these conversations create value rather than cause harm. To understand whether routine trauma discussion should be carried out, and if so how it should be done, ~~how best to support this work,~~ we aimed to collate and synthesise all relevant qualitative studies to capture women's and maternity care professionals' views of this issue. ~~routine discussion of previous trauma in the perinatal period. We aimed to illuminate how best to support women and professionals in having routine trauma conversations.~~

We use the term 'routine' to indicate raising the issue of previous trauma with all women accessing maternity services, as opposed to selectively for women who the healthcare professional suspects may have suffered trauma. The term 'routine' in this context does not negate the need for personalised care. We have chosen to use the term 'discussion' rather than 'enquiry' as discussion leaves open the possibility that trauma could be raised in a way that does not involve a direct question.

The review question was 'What are the views of women and maternity care professionals on routine discussion of previous trauma in the perinatal period?'. The study was registered in PROSPERO with the reference number CRD42021247160 [14]. The review was informed by guidance produced by the Cochrane Effective Practice and Organisation of Care group on carrying out and reporting qualitative evidence syntheses [15-17].

## Methods

### Reflexive note

We considered our pre-existing beliefs on routine trauma discussion and how these might influence the design and conduct of the review [18].

JC is a midwife and doctoral student. She was unsure at the outset whether routine trauma discussion was beneficial to women. She believed that for some groups of women, who face discrimination based on their class, race, immigration status or a range of other factors, disclosure of trauma could be harmful and increase the risk of unnecessary or unwanted safeguarding or mental health involvement. SD is a midwife with 18 years of clinical practice and a maternity care researcher, and held similar beliefs to JC. GT has a psychology academic background, and a long history of undertaking research with women who have experienced perinatal mental health problems. Her views were that conversations that were trauma-informed were important to enable needs-led care. AT is a researcher in the field of maternal and neonatal care, and her views were that an appropriate supporting model of care is needed to enable trauma-informed conversations, that will be beneficial to women. MF teaches critical psychology in the US and is a visiting professor in South Africa. With years of participatory work with women in and out of prison, highly marginalized young people and most recently Muslim American youth, MF is interested in collective and individual trauma as sites of wounds and creativity, memory and activism, a rich and painful source of knowledge and wisdom.

A 'research collective' comprising experts by lived experience, from the voluntary sector and healthcare professionals is supporting the doctoral study of which this review is part. Involvement of the collective in the design of the review helped minimise the risk that our pre-understandings would influence the review. Further, the range of backgrounds and views on the review team also helped enhance the rigor of the analysis.

## **Search strategy and selection criteria**

We searched the databases MEDLINE, CINAHL Plus, EMBASE, APA Psycinfo, and Global Index Medicus using the search terms 'trauma-informed' and 'trauma informed'.

We developed the search strategy by reviewing key relevant papers we were already aware of and examining relevant systematic reviews on PROSPERO. Each of the key papers referred to 'trauma-informed' in their keywords, title or abstract. We trialled searching using this term on Medline and duplicated it using a variety of different terms such as trauma-sensitive, trauma-focused, and trauma-responsive. These alternative terms had very few hits and had always been used in addition to trauma-informed. We therefore decided to only use the search terms 'trauma-informed' or 'trauma informed' and hand select relevant papers that concerned the care of women in the perinatal period.



Forward and backward citation tracking and key author searches for studies included in the review were carried out to identify additional relevant studies. Searches were conducted in July 2021 and updated in April 2022.

Only qualitative studies and qualitative aspects of mixed methods studies were included in the review. Studies not based in a maternity setting, or which did not include women in the perinatal period (defined for this purpose as pregnancy and up to one year after birth) were excluded. The review was focused on previous psychological trauma: this could include all past trauma or specific types such as adverse childhood experiences or sexual abuse but excluded studies with participants who have experienced physical trauma such as injury. Studies published at any date, and in any language were eligible for inclusion in the review.

Two reviewers (GT and JC) independently screened 20% of papers at title and abstract stage using Rayyan blind screening. The level of agreement was set at 95%. As the reviewers achieved 100% agreement, the remaining 80% of papers were reviewed by JC. At the second stage, all papers were blind screened by two reviewers (GT and JC), with any differences of opinion about inclusion resolved through discussion.

## Data extraction and management

Papers eligible for inclusion were uploaded to the software program MAXQDA Plus 2020 for data extraction, analysis, and thematic synthesis. We developed a standardised data extraction form using Excel and piloted it prior to beginning data collection. JC used the form to record basic contextual and methodological information about each study, including bibliographic information, country of study, setting, study design, data collection, participants' characteristics, data analysis methods and key themes. To reduce bias and errors, GT independently extracted data from 20% of studies, and JC and GT compared results, resolving disagreements through dialogue.

## Appraisal of the methodological quality of included studies

JC assessed the quality of each included paper using the Critical Appraisal Skills Programme (CASP) quality assessment checklist for qualitative studies [19]. The CASP checklist comprises 10 questions: 1 mark was allocated to each question if the **criteria** was met. ~~This enabled comparison between papers and reviewers.~~ The overall quality of each study was categorised as 'strong' (score 8-10/10; minimal methodological issues), 'adequate' (score 5-7/10; no major methodological issues) or 'weak' (0-4/10; major methodological issues). Studies scoring 'weak' were excluded on quality grounds to ensure that the credibility or trustworthiness of the review findings was not compromised

by including studies with important methodological limitations [15]. GT independently quality assessed 20% of included studies, and JC and GT compared results, resolving disagreement through re-examination and discussion. As suggested by Carroll and Booth [20] post hoc sensitivity analyses were carried out to assess the impact on the review findings of excluding a study on quality grounds. Quality ratings contributed to the GRADE-CERQual assessments (described below).

When we were quality appraising each study, we also reflected on broader issues of the integration and inclusion of women's voices, and these additional considerations are reflected in the discussion.

## **Data synthesis**

Data were synthesised thematically using the method developed by Thomas and Harden [21]. First, the findings of each study were inductively coded on a line-by-line basis. The codes were then organised into related areas, constructing 'descriptive' themes (summaries of findings). Finally, the descriptive themes were organised into analytical themes. This work was led by JC, and, at all stages, emerging concepts were shared, discussed, and refined with the review team.

## **Assessment of confidence in the review findings**

The GRADE 'Confidence in the Evidence from Reviews of Qualitative research' (GRADE-CERQual) approach was used to assess confidence in the synthesised findings [22-27]. This approach facilitates explicit and transparent assessment of whether the findings from a qualitative synthesis reasonably represent the phenomenon of interest. Each summary of findings (descriptive theme) was assessed in terms of methodological limitations, coherence, adequacy of data, and relevance; with the four assessments contributing to an overall assessment of confidence. The summaries of findings and associated CERQual assessment of confidence are presented in an Evidence Profile and Summary of Qualitative Findings table.

## **Results of the search**

Overall, 3,888 papers were identified after removal of duplicates, of which 25 met the inclusion criteria for the study. The PRISMA flow chart can be found at Figure 1. Three papers were excluded due to quality [28-30]. Post hoc examination of the three excluded papers indicated that their inclusion in the review would not have altered the final themes.

**<insert figure 1 here>**

## **Description of the studies**

The characteristics and quality of the 25 studies included in the review are summarised in Table 1. The included studies were published between 2001 and 2022. In terms of study setting, 12 of the studies were carried out in Australia, nine in the United States, two in Sweden, and one each in England and Canada. Routine trauma discussion was explored from the perspective of women in thirteen of the papers, eight looked at the perspective of healthcare professionals, and the remaining four papers looked at both viewpoints. Most data were collected by individual interviews, focus groups and/or surveys. The studies represented the views of 1602 women and 286 healthcare professionals and experts from the voluntary sector.

<Insert Table 1 here>

## **Assessment of methodological strengths and limitations**

Of the 25 included papers, sixteen were assessed as methodologically strong and nine as adequate. As discussed above, a further three papers were assessed as methodologically weak and were excluded from the review. A lack of reflexivity was noted across most of the studies. Other common methodological weaknesses included insufficient information about data analysis, the lack of a clear statement of findings, and minimal discussion of ethical issues.

## Confidence in the review findings

Table 2 shows the Summary of Findings and CERQual rating for each summary of finding. Confidence in most of the review findings was moderate or high, reflecting the quality and quantity of the studies included in the review. Each summary of finding was mapped to an analytical theme and these themes are discussed in the next section. All studies that met the inclusion criteria were from high-income countries. Because these findings were applicable to the UK, we did not downgrade the CERQual ratings for income status of the countries. However, the findings are not applicable to low- and middle- income countries.

<Insert Table 2 here>

## Findings

Six analytical themes were identified relating to women's and maternity care providers' views and experiences of routine trauma discussion. The first theme '*I did not know how to say it, and no-one asked me*' considers whether maternity care providers should ask women about previous trauma. '*A real whitefella way to start*' explores standardised compared to more unstructured ways of asking about trauma. In the theme '*You say it is confidential... but you are going to report me*', fear of judgement as a barrier to disclosure, and the importance of trust and relationships in trauma discussions is highlighted. The theme '*I'm not quite sure what is going on, but I feel really vulnerable*' calls attention to

the intensity of the perinatal period, which is often challenging but also has the potential for healing and growth. *Embedding trauma in routine practice* explores barriers and facilitators to successful implementation. ~~*Heavens! I forgot it!* looks at the challenges of embedding trauma in routine practice.~~ Finally, *'You go home and it's playing on your mind as you're cooking'* considers the impact on care providers of hearing trauma disclosures.

## **'I did not know how to say it, and no-one asked me': should maternity care providers ask women about previous trauma?**

This theme explores whether women and maternity care providers feel that routine trauma discussion should take place.

Participants in 14 studies expressed that they felt routine trauma discussion is acceptable and worthwhile [31-44]. One of the participants in the study undertaken by Montgomery, Seng and Chang (2021) [36] proposed:

*"It might have just put a thought in my head, even if it wasn't something that I shared with anybody, it might have just put a thought in my head which might have been useful at some point"* [36].

Some women felt it was difficult to broach the subject of previous trauma and would not have disclosed unless the clinician raised the issue:

*“At the time, I could not and did not tell the healthcare professionals of my survivor status. I did not know how to say it, and no one asked me” [36].*

The overall finding that women accepted routine trauma discussion masks several complexities and contradictions. In eight studies, women reported feeling unprepared for the discussion and found it intrusive [32, 33, 35-39, 42]. This was reflected by a woman who was interviewed in the study by Millar et al. (2021) [35]:

*“Like it doesn’t feel good when you first meet someone, and they just start like trying to jump into your life. Like they know you. I hate that” [35].*

Some participants spoke of how they would have welcomed forewarning of the discussion:

*“I think they could have told me what they were going to ask before I even arrived for my appointment. I had no idea that was what was coming” [39].*

In 11 studies, professionals reported that they felt routine trauma discussion was worthwhile [32, 34, 45-53]. A clinician taking part in the study by Flanagan et al. (2018)



[47] reflected that while *'most of the time (the) screen is negative'*, when finding the individual who had faced previous trauma, *'you're so glad you did'* [47].

Participants in the study by Kohlkoff et al. (2021) felt there were three key benefits of routine trauma discussion: identify women at higher risk of mental health problems or family violence, increase referrals to appropriate support services, and provide support and education [47]. However, in seven studies, women felt that trauma discussions should only take place if clinicians had enough time to respond to disclosures and could provide or refer into appropriate support [32, 38, 39, 41-44]. Having disclosed prior trauma, some women expected that they would be treated more sensitively and that other care providers would be aware of their history, and were aggravated when this wasn't the case [39, 41]:

*"Why don't they take the extra time just to read over [my file] and if they have any more questions about it then they can ask. If it's already there then why bother... it is really frustrating"* [39].

Maternity care providers similarly spoke of their support for routine trauma discussion as contingent on having sufficient time and appropriate referral pathways [32, 34, 45-48, 50-53]. Without good quality support services, clinicians were reluctant to discuss prior trauma, fearing this would open a 'Pandora's box' of issues they were unable to deal with:

*“We see perinatal depression and anxiety but this is a continuum of social disadvantage and intergenerational trauma. We have super complicated patients with so many problems. Where do we fit mental health in where there are so few resources to respond properly?” [32].*

## **‘A real whitefella way to start’: standardisation and tick-boxes in trauma discussion**

This theme explores how maternity care providers should raise the issue of previous trauma. Participants in several studies discussed limitations in the use of questionnaires to raise the issue of previous trauma [31, 32, 37, 45, 54]. Participants suggested that closed questions (for example ‘in the last year, have you experienced...’) can prevent disclosures [31, 37] This Aboriginal participant in a study by Carlin, Atkinson and Marley (2019) [32] proposed:

*“You talk about things because they are important to talk about not cause they happened one week ago! It is a real whitefella way to start. It’s like you’re in or you’re out. You see that hey? Like what happens if it was a bit longer, then the lady might think oh no, it’s not important, I won’t talk about that” [31].*

Mendel, Sperlich and Fava (2021) [54] investigated the use of the Adverse Childhood Experiences questionnaire (ACE-10) within maternity research. The researchers found

that the questionnaire contains confusing and ambiguous questions, excludes important traumatic events in childhood (such as the death of a parent) and fails to ascertain the severity or duration of the traumatic experience. For example, one participant in the study had an ACE score of one out of ten but had suffered extensive abuse over 12 years of her childhood, resulting in seven miscarriages. The authors concluded that completion of the questionnaire might not give a true representation of the extent of trauma the woman has suffered.

Women who took part in the study by Carlin et al. (2020) [32] felt that direct questions could cause women to disengage, and that broad, gentle questions were the best approach to ask about difficult experiences. The voluntary sector experts and healthcare professionals who participated in the study by Chamberlain et al. (2020) [45] noted that direct questions about trauma can be problematic, because avoiding thinking about trauma experiences can be a way of coping. Instead, indirect methods of gentle communication were preferred, asked by *“someone trusted – this sort of information will naturally become evident so the trusted person can gently empathise and draw attention to, as opposed to ask directly and abruptly”* [45]. Another participant noted that clinicians should ask *‘slowly, gently and only where there is the possibility of being able to ‘hold a space’ and deal appropriately with the answer’* [45].

Carlin et al. (2020) [32] explored clinicians’ views of a questionnaire-based approach versus questionnaire plus narrative. Several participants reported that they only used the

questionnaire, describing the narrative section as 'aspirational' due to time constraints and concerns that it positioned them as a counsellor. However, those who did use the narrative approach felt it enhanced rapport with women and that women understood the limits around the assistance they could provide:

*"Generally I think women are keen to share some of their problems with us as nurses even though we cannot solve these issues as such but we can listen, we can advise them where to seek help and how we can assist as a support for some of their problems."* [32].

## **'You say it is confidential... but you are going to report me': the importance of trust**

This theme concerns the importance of trust and relationships to women who have suffered trauma. In twelve studies, fear of judgement was reported as a reason for non-disclosure [31, 32, 36-38, 40-45, 51]. This included a general fear of being perceived as a bad parent, and specific concerns that their child would be removed from their care; *"You guys are bound by law [to report certain things]... You say it is confidential... but you are going to report me"* [43]. Concerns about confidentiality were raised by participants in several studies, as was the misconception that the abused becomes the abuser. One mother stated:

*"Speaking from personal experience, I felt at times that my past trauma was being used to assess the likelihood I would harm my own child, rather than as a means of identifying what support I might need as an individual"* [38].

Participants in 13 studies highlighted the importance of a trusting relationship, built through multiple encounters, as a prerequisite for trauma disclosure [31-33, 35-40, 42, 44, 45, 52] A mother from the study by Millar et al. undertaken in 2021 [35] suggested:

*“I think if I had a relationship, then yes I [would disclose trauma history]. But with the amount of time I was seeing them, no. ‘Cause I was always seeing someone different”* [35].

Participants in five studies proposed that trauma should not be discussed at the first appointment, but at a later appointment, enabling a relationship to be built first [35, 38, 42, 45, 52]. This woman chose to disclose later in pregnancy:

*“When she gave me the initial, you know, the history form... when I saw ‘were you abused?’ I said no. There was no way I was going to tell her”* [40].

Although maternity care providers agreed that women were more likely to disclose prior trauma after a relationship with the clinician had been established, some felt it was appropriate to raise trauma at the first consultation, seeing this as the start of an ongoing conversation:

*“If... they’re not opening up... they’ll go home and think about it and reflect on that and they may come back the next time and open up a bit more. It’s just opening the door, isn’t it?”* [51].

In twelve studies, participants described the importance of how the clinician asked about previous trauma [31-34, 37-41, 43-45]. Desired attributes consistently included kindness, friendliness, sensitivity, a non-judgemental attitude, respect, care, and compassion. One woman who took part in the study carried out by Sobel et al. (2018) [41] expressed that *“I opened up to my midwife because I felt comfortable with her. That’s it”* [41] Conversely, this participant in Seng et al.’s study (2002) [40] felt the need to change care provider:

*“The doctor was kind of cold, not personable at all, and those feelings [emotional memory of being abused, shame, vulnerability, nakedness] would come back to me in his office, and I found myself crying at every visit”* [40].

For women with limited English, non-verbal signals like smiling and a relaxed manner were vital in inspiring trust: *“She make me like not scared because she smile a lot, her smiling and the way she spoke was really helpful”* [39].

Participants proposed that trauma discussions should be held in **private, comfortable and welcoming surroundings** ~~a safe, private, and homely environment~~ [33, 38, 43, 45]. The ideal combination of a trusted care provider and warm environment is summed up by a woman who took part in Gokhale et al.’s study (2020) [33]:

*“The only way my health care providers can help me with my trauma is every time I come, make me feel like I’m at home. Make me feel comfortable. Make me feel safe and make me feel like I have nothing to worry about”.*

Participants in three studies felt that if the conversation was handled badly, routine trauma discussion could impact upon future health care access and experiences [36, 38, 39]. One respondent to the survey carried out by Olsen, Galloway and Guthman (2021) [38] proposed:

*“If it isn’t asked about in a sensitive way under the right circumstances, it could feel really intrusive or could be so upsetting or off-putting that someone could avoid needed health care entirely” [38]*

Participants in thirteen studies reported that they would not disclose previous trauma to the healthcare professional looking after them, although not all studies explored the reasons for this [32, 33, 35-41, 44-46, 51]. Mule et al. (2021) carried out a survey asking women whether they had chosen to fully disclose their histories during antenatal psychosocial assessment: 161 women responded that they had not, and completed an open-ended question giving their reasons. The researchers found there were a range of reasons, including lack of trust of the person asking, fear of judgement, use of closed-ended questions and lack of time, but also simply privacy: they did not want to share this information [37]. Similarly, a participant in **the study by** Olsen, Galloway and Guthman (2021) proposed that *“some people may not be ready”* [38] while a woman taking part in

the study by Gokhale et al. (2020) suggested that *“it’s not easy speaking up about situations like that and a lot of people don’t because they don’t feel comfortable or they don’t feel like they could trust people enough to do that”* [33]. This implies that even within a trusting relationship, some women who would benefit from support will choose not to share their histories. Accordingly, participants in the study by Seng et al. (2002) [40] proposed that care providers should assume women are trauma survivors if they display signs or symptoms of trauma, irrespective of whether they have disclosed.

## **‘I’m not quite sure what is going on, but I feel really vulnerable’: the intensity of the perinatal period**

This theme explores experiences of the perinatal period for women who have suffered trauma. In seven of the studies, **some participants women** reported feeling that there was no connection between their trauma histories and their current wellbeing and pregnancy [33, 36, 37, 39, 40, 42, 45]:

*“It’s not really affecting me now... my main concern is getting through the pregnancy, not worrying about my past stuff”* [39].

Some women wanted to focus on the pregnancy and a positive future and felt discussion of trauma could trigger distressing feelings. This was the case even when the perinatal period could be expected to bring up strong emotions, such as for this woman whose infant had been murdered:



*“Cause when you come to the visit you want to hear stuff about your baby. You don’t want to keep dwelling on this that happened in the past and you trying to have a happy moment.” [33].*

Some women had not foreseen that their pregnancy would be so difficult: *“It’s hard to put into words because I’m not quite sure what is going on, but I feel really super vulnerable”* [40]. Even women who appeared to be far along in recovery and living happy lives were often unprepared for the intensity of the perinatal period:

*“I was really looking forward to the cuddling time with the baby and breastfeeding... I didn’t expect this whole other ugliness”* [40].

Women commonly felt a loss of control over their body, due both to the pregnancy and a sense of powerlessness within maternity care. Vaginal examinations, birth, or even seemingly benign clinical procedures such as blood pressure measurement caused flashbacks to abuse. Some women feared bodily exposure during labour and birth, with a participant in the study by Sobel et al. (2018) [41] reporting:

*“I was so concerned with being covered up... I would have been devastated [by a vaginal delivery]. I did not know how I was going to keep my clothes on and have a baby”* [41]

Until pregnancy, some women were not fully aware of the trauma they had suffered. Seng et al. (2002) [40] explored this issue in detail through narrative interviews with 15 women who had suffered childhood sexual trauma and subsequently accessed maternity care. At the time of the pregnancy, four of the fifteen women had only a vague understanding that they had been subjected to abuse. Participants described indications in their thoughts and behaviour of the effects of trauma, such as fleeting flashbacks, suicidal intentions, and extreme promiscuity:

*“I realized [later] there were pieces that had been floating around for a long time that I wouldn’t acknowledge prior to [postpartum]” [40].*

Because they had not fully admitted it to themselves, these women were not able to disclose the abuse to their healthcare providers:

*“I knew early that I was not going to deliver vaginally. I knew in my head that I was not going there. So that piece I connected...I don’t know that I drew a real direct line because of how vulnerable I felt. I wasn’t probably ready to acknowledge that...So it was knowing and not knowing at the same time” [40].*

Participants talked about the potential for post-traumatic growth in the perinatal period [31, 40, 41, 46]. However, this was not always an easy process, as described by this participant in the study by Seng et al. (2002) [40]:

*“I kind of knew in some way it was affecting me, but I just couldn’t connect the dots ever... but when I got pregnant it all just came out, came clear, and it was hard, and I’m*

*grateful... and I think it's going to help me grow past it and deal with it... but pregnancy is enough to deal with."* [40].

## ***'Heavens! I forgot it!': challenges to Embedding trauma discussion in routine practice***

This theme investigates how trauma discussion can be introduced and explores barriers and facilitators to successful implementation. ~~including practical steps to aid implementation.~~

Care providers in four studies reported that they quickly adapted to routine trauma discussion, and found it feasible within their workloads [47, 51-53].

*"I just think it's the initial getting used to... just even logging into it, and doing all of that was a hassle when I first started. It's "Oh, this is all so hard." But it's so simple now, because we're used to it... it's like anything,... any tool that you use over and over again, it becomes more simple"* [53].

Participants in seven studies discussed the challenges of ensuring all women are asked about prior trauma [46-48, 50-53]. Care providers taking part in studies by Choi and Seng (2014) and Stenson, Sidenvall and Heimer (2005) spoke of having preconceived notions

of who might or might not have suffered trauma: *“If you have a feeling something isn’t quite right, then it’s easier. But just when you don’t think...”* [52]. Participants in the study undertaken by Schmied et al. (2020) felt *“they’re just not appropriate questions to ask”* [51], or avoided asking the question where a woman was felt to be ‘keeping her distance’. Describing a successful implementation of routine trauma discussion, this participant in Reilly et al.’s study (2020) proposed that key individuals were helpful in getting more reluctant staff on board:

*“Even if you have people who are sceptical, if you are enthusiastic, and the clients get enthusiastic and really feel cared for, that automatically rolls over to the staff members that are sitting on the sidelines and saying “I don’t know if all this is necessary”... They start seeing that it is making a positive impact on people’s lives...”* [50].

Participants in the study by Stenson, Sidenvall and Heimer (2005) [52] found it difficult to remember to raise the issue, resulting in some women not being asked about prior trauma. Documentation of abuse in hand-held notes could be a confidentiality risk, but several people taking part in the study suggested a check box in the records, to act as an aide memoire, and ensure the discussion is accorded the same importance as other issues [52]. Participants in Kohlhoff’s study (2021) [48] proposed setting up a flag on the system to ensure effective information sharing among the maternity team.

Partner presence at appointments could influence the discussion of previous trauma [46, 48, 49, 52]. This was felt by some participants in the study by Stenson, Sidenvall and Heimer (2005) to be a modern phenomenon, with one care provider proposing *“this*

*generation of couples, the husband's there the whole time" [52]. Whilst a participant in Kohlhoff et al.'s study (2021) [48] noted that "pregnant women can be quite vulnerable and anxious, and it is good to have that support person with them", another pointed out: "They're not going to be able to divulge anything while their partner's there, especially if their partner doesn't know about it, and sometimes that is the case" [48].*

Trauma discussions were more difficult for women with limited English [39, 52, 53]. Women often did not want to disclose sensitive issues through an interpreter, and where family members or partners were acting as interpreters, this provided a further barrier to disclosure. In the study by Stenson, Sidenvall and Heimer (2005) [52], staff reported that partners were often asked to sit in the waiting room while this discussion took place: few partners insisted on being present, but the midwives sometimes felt in a difficult position where the woman had limited English:

*"I would really prefer a professional interpreter but on most occasions the men say 'no'. They want to do the interpreting" [52].*

# **‘You go home and it’s playing on your mind as you’re cooking’: the impact on care providers of hearing trauma disclosures**

Hearing trauma disclosures could be challenging for clinicians [34, 46, 48, 52, 55]. Mollart, Newing and Foureur (2009) [55] explored this issue in detail in their Australian study in which they carried out focus groups with 18 midwives who undertook routine discussion of prior trauma. The midwife participants in the study reported that the cumulative, complex disclosures they heard affected them emotionally and impacted on their home and work life:

*“You go home and it’s playing on your mind as you’re cooking. I don’t know how long it usually goes on for, probably till you get that next bad case” [55].*

Some participants in the study reported that they continued to think about trauma disclosures after work:

*“Sometimes I’ve gone home and actually worried about people, then you’ve got to remember that they told me this today and they’ve been living with this for how long? Just keep telling yourself that” [55].*

For some, this impacted on their family life:

*“For me, I explode at home, I don’t explode here [at work] because I know that no-one would put up with that kind of behaviour. But I do it to my kids, and that’s not very good.”* [55].

Marley et al. (2017) interviewed healthcare providers about their experiences of routine trauma discussion with Aboriginal women in Western Australia. A participant in the study expressed: *“At the end of the day, it’s hard not to want to neck a bottle of wine to cope with [hearing their stories]”* [34]. The use of unhealthy coping strategies, such as excess alcohol, was echoed by a participant in Mollart, Newing and Foureur (2020):

*“I can debrief 10, 20, 30 times, and the information is still with me, and I don’t know where to channel that sometimes. Sometimes you channel that into things that are probably not appropriate.”* [55].

Participants in studies undertaken by Marley et al. (2017) [34] and Mollart, Newing and Foureur (2009) [55] felt clinical supervision is vital for midwives carrying out routine trauma discussion, and those with no access to supervision expressed resentment. However, not all care providers who were offered supervision chose to participate in it:

*“The only way I know how to deal with it is I talk to colleagues... even though sometimes when you talk to colleagues, you know they’re thinking about the booking-in they had, they’re only half listening and you haven’t actually been heard”* [55].

Stenson, Sidenvall and Heimer (2005) [52] was the only study to raise the issue of care providers who have experienced trauma themselves. The researchers carried out five

focus groups with 21 midwives in Sweden to explore their views on discussing prior and current violence with women. The researchers noted that none of the participants spontaneously raised the issue. When the moderator brought it up ‘*several admitted that they had not considered the possibility that colleagues might have been subjected to abuse*’ [52]. The group discussed whether midwives could act professionally in this situation and concluded that they could, but it might be difficult if they are still in a violent relationship. The group did not explore whether discussion of previous trauma might be more challenging for midwives with personal experience in this area, or how they could be supported.

## Discussion

This qualitative evidence synthesis found that although many women feel positive about routine discussion of previous trauma, this simple statement masks complexity. Women often find the conversation unexpected and intrusive; they expect their care to change after trauma disclosure and are disappointed if it does not; and women with limited English face additional challenges. The findings of the review suggest that many pregnant women are unaware of the extent of the trauma they have suffered, or its impact on their lives. This highlights the difficulty of discussing trauma with women who will be at very different stages of recovery.



The review underscores the importance of an established relationship and trust in trauma discussions. While midwife-led continuity models of care have been shown to be beneficial for mothers and babies, New Zealand is the only country to have achieved this at scale, and the national target of implementation of continuity of carer in the UK has now been removed [56-58]. Notwithstanding this, our review found that many women will not disclose previous trauma in the absence of a trusting relationship and will consequently fail to receive the care they need. Some women will choose not to disclose their histories; this is problematic because in the included studies, women who disclosed trauma were generally provided with information and follow-up care, while women who did not disclose trauma were not. Consideration should therefore be given to providing all women with information about the impact of trauma and the means to independently access support should they wish to do so.

Our review highlights specific areas of training which are necessary for maternity care professionals. The focus of trauma discussions should not be eliciting specific details of past trauma, but rather ascertaining what resources, support, and / or adjustments to the care plan would be helpful. It is important that women are forewarned of the discussion, including any limits to confidentiality and that disclosure of past trauma is entirely voluntary. The review found that partner presence can be a barrier to trauma discussions. National Institute for Health and Care Excellence guidance (2021) [59] states that maternity care providers should ensure women are asked about domestic abuse in a private, one-to-one discussion; we propose that this also applies to discussion of previous trauma. For both women who have suffered trauma and

professionals, support for routine discussion of previous trauma was found to be contingent on adequate time to explore these complex issues. This is a serious caveat in maternity systems which were already understaffed and have been further impacted by the COVID-19 pandemic, both in the UK and internationally [60]. Description of the discussion as easy and quick by some clinician participants in the included studies implies an unknowingly 'performative' approach to routine trauma discussion in which professionals feel they have given women the opportunity to disclose, but women may not feel comfortable disclosing or get the support they need.

This performative approach could also be a protective mechanism for clinicians who feel trauma discussions are emotionally difficult. Midwifery work carries a heavy emotional burden, with midwives suffering significantly higher levels of burnout, stress, anxiety, and depression than the general population [61-63]. Added to this, the review found that hearing women's histories of sexual assault, childhood abuse, and domestic violence can profoundly affect clinicians. None of the studies in the review explored whether discussion of previous trauma might be more challenging for midwives who have personally suffered trauma, or how they could be supported. There is a need for an appropriate support and supervision model which allows midwives the opportunity to explore the challenging situations they are exposed to. Reflective supervision would enable clinicians experiencing vicarious traumatisation or those whose own trauma experience have been triggered through their interactions with survivors to receive the support they need.

Women's perceptions and experiences of trauma discussion might be affected by characteristics such as class, ethnicity, or immigration status, but it was not possible to differentiate and draw distinct themes for different groups of participants. For example, the review found with high confidence that women are fearful of disclosing their histories in case it raises concerns that they are unable to safely care for their child: it may be that this is a greater barrier for women from population groups who experience disproportionately high levels of child welfare involvement [64]. Women with insecure immigration status are highly vulnerable to violence and abuse [65], but were excluded from almost all of the studies by default as their inclusion criteria required that participants could speak English: this is an area where further research is needed.

In 2014, the US Substance Abuse and Mental Health Services administration (SAMHSA) [66] published a seminal conceptual document about trauma-informed care, **a model of service delivery which aims to develop an environment in which people who have experienced trauma feel safe and can build trust with their healthcare provider.** This framework has been further developed and tailored to the perinatal period by others including Seng and Taylor (2015) [67], Sperlich et al. (2017) [68] and Law et al. (2021) [9]. SAMHSA (2014) [66] **propose that trauma-informed approaches are underpinned by four assumptions: staff realise the prevalence of trauma and its impact on behaviour; they are able to recognise the signs of trauma, including by trauma screening; the organisation's policies and procedures take into consideration the experiences of trauma among service**

users and staff; and active steps are taken to resist re-traumatisation of clients and staff by ensuring psychological safety. Within a trauma-informed care framework, implementation of routine screening for previous trauma should be considered within the context of much broader changes to services, including staff training, continuity of carer, emotional support for staff, and service evaluation [9, 66-68]. Law et al. (2021) proposed that to be trauma-informed, organisations must co-produce services with trauma survivors [9]. We suggest that the clinicians expected to carry out trauma discussions must also be involved in developing trauma discussion frameworks, to ensure that they are given the training and support they need. Further research is needed to ascertain what resources and support are helpful to women who have suffered disclose trauma, bearing in mind that this must focus on the preferences and needs of the individual.

## **Strengths and limitations of the review**

This synthesis is the first to bring together the views of women and maternity care professionals on routine discussion of previous trauma in the perinatal period. We adopted an inclusive approach to the search, with broad search terms and multiple search strategies used to ensure no key articles were missed. The findings are strengthened by the large number of participants. We minimised the risk of over or under interpretation of the data through explicit positionality, reflexivity, and discussion.

However, the synthesis has limitations. All studies that met the inclusion criteria were from high-income countries, therefore the findings cannot be applied to women in low- or middle-income settings. Further, there were no studies set in Asia, and only in three in Europe. Research is needed on the optimal ways of addressing previous trauma in pregnancy and the perinatal period for different populations, in line with the Sustainable Development Goals and guidance by the European Parliament that sexual and reproductive health services should consider the needs of women who have suffered sexual or gender-based violence [69, 70]

## **Conclusion**

The review provides insight into the barriers and facilitators of women in high-income countries sharing their trauma histories, and clinicians asking about previous trauma. Areas for future research are highlighted. Discussions of previous trauma are complex and require careful consideration and sensitivity. While implementation of continuity of carer is no longer a target in maternity services in the UK, many women will not disclose previous trauma in the absence of a trusting relationship. As some women will not disclose their histories, consideration should be given to providing all women with information about the impact of trauma, and means of independently accessing support. These findings also highlight the need for time to undertake authentically sensitive and tailored discussions about trauma, and for appropriate support and supervision for care providers. There is a need for routine trauma discussions to be co-

developed with childbearing women and the midwives who will be initiating the conversation, to ensure that women receive the support they need, and the wellbeing of care providers is protected.

## Acknowledgements

We would like to thank the EMPATHY study Research Collective for generously sharing their knowledge and expertise. Members of the collective are: Dr Laura Abbott, Juliet Albert, Kirsty Armstrong, Jill Benjoya Miller, **Dr** Emma Brooks, Dr Geraldine Butcher, Jo Doherty, Amber Jackson, Isobel Martin, **Dr** Elsa Montgomery, Sam Pointon, Sarah-Jayne Pomeroy, Erjola Sadria, Gill Skene, Memuna Sowe, **Dr** Kim Thomas, and Lucy Warwick-Guasp.

## References

1. Crown Prosecution Service. Violence Against Women and Girls Report. England: 2019.
2. Bellis M, Hughes K, Hardcastle K, Ashton K, Ford K, Quigg Z, et al. The impact of adverse childhood experiences on health service use across the life course using a retrospective cohort study. *J Health Serv Res Policy*. 2017;22(3):168-77.
3. Bellis MA, Hughes K, Ford K, Ramos Rodriguez G, Sethi D, Passmore J. Life course health consequences and associated annual costs of adverse childhood experiences across Europe and North America: a systematic review and meta-analysis. *The Lancet Public Health*. 2019;4(10):e517-28.
4. Hughes K, Bellis MA, Hardcastle KA, Sethi D, Butchart A, Mikton C, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *The Lancet Public Health*. 2017;2(8):e356-66.

5. Sara G, Lappin J. Childhood trauma: psychiatry's greatest public health challenge? *The Lancet Public Health*. 2017;2(7):e300-1.
6. Muzik M, Ads M, Bonham C, Lisa Rosenblum K, Broderick A, Kirk R. Perspectives on trauma-informed care from mothers with a history of childhood maltreatment: A qualitative study. *Child Abuse Neglect*. 2013;37(12):1215-24.
7. Seng JS, D'Andrea W, Ford JD. Complex Mental Health Sequelae of Psychological Trauma Among Women in Prenatal Care. *Psychological trauma: theory, research, practice and policy*. 2014;6(1):41-9.
8. Montgomery E, Chang Y, Youssef A, Ivan F, Maxted F, Cheadle S, et al. *Pregnancy, Birth and Parenthood after Childhood Sexual Abuse*. The Survivors Trust. 2019.
9. Law C, Wolfenden L, Sperlich M, Taylor J. A good practice guide to support implementation of trauma-informed care in the perinatal period. NHS England and NHS Improvement; 2021.
10. American College of Obstetricians and Gynecologists. *Caring for Patients Who Have Experienced Trauma: ACOG Committee Opinion Summary, Number 825*. *Obstet Gynecol*. 2021;137(4):757-8.
11. Ford K, Hughes K, Hardcastle K, Di Lemma LCG, Davies AR, Edwards S, et al. The evidence base for routine enquiry into adverse childhood experiences: A scoping review. *Child Abuse & Neglect*. 2019;91:131-46.
12. Underwood E. California has begun screening for early childhood trauma, but critics urge caution. *Science (American Association for the Advancement of Science)*. 2020.
13. Racine N, Killam T, Madigan S. Trauma-Informed Care as a Universal Precaution: Beyond the Adverse Childhood Experiences Questionnaire. *JAMA Pediatrics*. 2020;174(1):5-6.
14. Cull J, Thomson G, Downe S, Fine M, Topalidou A. What are the views of women and maternity care professionals on routine discussion of previous trauma in the perinatal period? A qualitative evidence synthesis. PROSPERO CRD42021247160. . 2021.
15. Glenton C, Bohren M, Downe S, Paulson E, Lewin S, On behalf of Effective Practice and Organisation of Care, (EPOC). *EPOC Qualitative Evidence Synthesis: Protocol and review template*. Version 1.3. EPOC Resources for review authors. Oslo: Norwegian Institute of Public Health; 2022 b.
16. Glenton C, Lewin S, Downe S, Paulsen E, Munabi-Babigumira S, Johansen M, et al. *Qualitative Evidence Syntheses Within Cochrane Effective Practice and Organisation of*

Care: Developing a Template and Guidance. *International Journal of Qualitative Methods*. 2021;20:16094069211041959.

17. Glenton C, Lewin S, Downe S, Paulsen E, Munabi-Babigumira S, Agarwal S, et al. Cochrane Effective Practice and Organisation of Care (EPOC) Qualitative Evidence Syntheses, Differences From Reviews of Intervention Effectiveness and Implications for Guidance. *International Journal of Qualitative Methods*. 2022 a;21:16094069211061950.

18. Walsh D, Downe S. Appraising the quality of qualitative research. *Midwifery*. 2006 Jun;22(2):108-19.

19. CASP Qualitative Research Checklist [Internet].; 2018 [cited 2.1.2020 2020]. Available from: <https://casp-uk.net/wp-content/uploads/2018/01/CASP-Qualitative-Checklist-2018.pdf>.

20. Carroll, C., and Booth, A. Quality assessment of qualitative evidence for systematic review and synthesis: Is it meaningful, and if so, how should it be performed? . *Research Synthesis Methods*. 2015;6(2):149-54.

21. Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Medical Research Methodology*. 2008;8(1):45.

22. Lewin S, Bohren M, Rashidian A, Munthe-Kaas H, Glenton C, Colvin CJ, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 2: how to make an overall CERQual assessment of confidence and create a Summary of Qualitative Findings table. *Implementation Science*. 2018 b;13(1):10.

23. Lewin S, Booth A, Glenton C, Munthe-Kaas H, Rashidian A, Wainwright M, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings: introduction to the series. *Implementation Science*. 2018 a;13(2).

24. Munthe-Kaas H, Bohren MA, Glenton C, Lewin S, Noyes J, Tunçalp Ö, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 3: how to assess methodological limitations. *Implementation Science*. 2018;13(1):9.

25. Colvin CJ, Garside R, Wainwright M, Munthe-Kaas H, Glenton C, Bohren MA, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 4: how to assess coherence. *Implementation Science*. 2018;13(1):13.

26. Glenton C, Carlsen B, Lewin S, Munthe-Kaas H, Colvin CJ, Tunçalp Ö, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 5: how to assess adequacy of data. *Implementation Science*. 2018;13(1):14.



27. Noyes J, Booth A, Lewin S, Carlsen B, Glenton C, Colvin CJ, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings—paper 6: how to assess relevance of the data. *Implementation Science*. 2018 b;13(1):4.
28. Matthey S, White T, Phillips J, Taouk R, Chee TT, Barnett B. Acceptability of routine antenatal psychosocial assessments to women from English and non-English speaking backgrounds. *Archives of Women's Mental Health*. 2005;8(3):171-80.
29. Rollans M, Schmied V, Kemp L, Meade T. 'We just ask some questions...' the process of antenatal psychosocial assessment by midwives. *Midwifery*. 2013 a;29(8):935-42.
30. Rollans M, Kohlhoff J, Meade T, Kemp L, Schmied V. Partner involvement: negotiating the presence of partners in psychosocial assessment as conducted by midwives and child and family health nurses. *Infant Ment Health J*. 2016;37(3):302-12.
31. Carlin E, Atkinson D, Marley JV. 'Having a Quiet Word': Yarning with Aboriginal Women in the Pilbara Region of Western Australia about Mental Health and Mental Health Screening during the Perinatal Period. *International Journal of Environmental Research and Public Health*. 2019;16(21):4253.
32. Carlin E, Spry E, Atkinson D, Marley JV. Why validation is not enough: Setting the scene for the implementation of the Kimberley Mum's Mood Scale. *PLOS ONE*. 2020;15(6):e0234346.
33. Gokhale P, Young MR, Williams MN, Reid SN, Tom LS, O'Brian CA, et al. Refining Trauma-Informed Perinatal Care for Urban Prenatal Care Patients with Multiple Lifetime Traumatic Exposures: A Qualitative Study. *Journal of Midwifery and Women's Health*. 2020;65(2):224-30.
34. Marley JV, Kotz J, Engelke C, Williams M, Stephen D, Coutinho S, et al. Validity and Acceptability of Kimberley Mum's Mood Scale to Screen for Perinatal Anxiety and Depression in Remote Aboriginal Health Care Settings. *PLOS ONE*. 2017;12(1):e0168969.
35. Millar HC, Lorber S, Vander Morris A, Thompson G, Thompson M, Allen L, et al. "No, you need to explain what you are doing": Obstetric care experiences and preferences of adolescent mothers with a history of childhood trauma. *J Pediatr Adolesc Gynecol*. 2021;34(4):538-45.
36. Montgomery E, Seng JS, Chang Y. Co-production of an e-resource to help women who have experienced childhood sexual abuse prepare for pregnancy, birth, and parenthood. *BMC Pregnancy & Childbirth*. 2021 Jan 07;21(1):30.

37. Mule V, Reilly NM, Schmied V, Kingston D, Austin MV. Why do some pregnant women not fully disclose at comprehensive psychosocial assessment with their midwife? *Women and Birth*. 2021;35(1):80-6.
38. Olsen JM, Galloway EG, Guthman PL. Exploring women's perspectives on prenatal screening for adverse childhood experiences. *Public Health Nurs*. 2021 Nov;38(6):997-1008.
39. Rollans M, Schmied V, Kemp L, Meade T. Digging over that old ground: an Australian perspective of women's experience of psychosocial assessment and depression screening in pregnancy and following birth. *BMC Womens Health*. 2013;13(1):18.
40. Seng JS, Sparbel KJH, Low LK, Killion C. Abuse-related posttraumatic stress and desired maternity care practices: women's perspectives. *J Midwifery Womens Health*. 2002;47(5):360-70.
41. Sobel L, O'Rourke-Suchoff D, Holland E, Remis K, Resnick K, Perkins R, et al. Pregnancy and Childbirth After Sexual Trauma: Patient Perspectives and Care Preferences. *Obstetrics & Gynecology*. 2018;132(6):1461-8.
42. Stenson K, Saarinen H, Heimer G, Sidenvall B. Women's attitudes to being asked about exposure to violence. *Midwifery*. 2001;17(1):2-10.
43. White A, Danis M, Gillece J. Abuse survivor perspectives on trauma inquiry in obstetrical practice. *Arch Womens Ment Health*. 2016;19(2):423-7.
44. Willey SM, Blackmore RP, Gibson-Helm ME, Ali R, Boyd LM, McBride J, et al. "If you don't ask ... you don't tell": Refugee women's perspectives on perinatal mental health screening. *Women and Birth*. 2020;33(5):e429-37.
45. Chamberlain C, Gee G, Gartland D, Mensah FK, Mares S, Clark Y, et al. Community Perspectives of Complex Trauma Assessment for Aboriginal Parents: 'Its Important, but How These Discussions Are Held Is Critical'. *Front Psychol*. 2020 Sep 15;11:2014.
46. Choi KR, Seng JS. Pilot for Nurse-Led, Interprofessional In-Service Training on Trauma-Informed Perinatal Care. *J Contin Educ Nurs*. 2015 Nov;46(11):515-21.
47. Flanagan T, Alabaster A, McCaw B, Stoller N, Watson C, Young-Wolff K. Feasibility and Acceptability of Screening for Adverse Childhood Experiences in Prenatal Care. *Journal of Women's Health*. 2018;27(7):903-11.
48. Kohlhoff J, Cibralic S, Tooke S, Hickinbotham R, Knox C, Roach V, et al. Health professional perspectives on an antenatal mental health screening program in a private hospital. *Aust N Z J Obstet Gynaecol*. 2021;61(6):891-7.

49. Preis H, Whitney C, Kocis C, Lobel M. Saving time, signaling trust: Using the PROMOTE self-report screening instrument to enhance prenatal care quality and therapeutic relationships. *PEC Innovation*. 2022;1:100030.
50. Reilly N, Brake E, Kalra H, Austin M. Insights into implementation of routine depression screening and psychosocial assessment in a private hospital setting: A qualitative study. *Aust N Z J Obstet Gynaecol*. 2020;60(3):419-24.
51. Schmied V, Reilly N, Black E, Kingston D, Talcevska K, Mule V, et al. Opening the door: midwives' perceptions of two models of psychosocial assessment in pregnancy- a mixed methods study. *BMC Pregnancy and Childbirth*. 2020;20(1):451.
52. Stenson K, Sidenvall B, Heimer G. Midwives' experiences of routine antenatal questioning relating to men's violence against women. *Midwifery*. 2005;21(4):311-21.
53. Willey SM, Gibson-Helm ME, Finch TL, East CE, Khan NN, Boyd LM, et al. Implementing innovative evidence-based perinatal mental health screening for women of refugee background. *Women and Birth*. 2020;33(3):e245-55.
54. Mendel WE, Sperlich M, Fava NM. "Is there anything else you would like me to know?": Applying a trauma-informed approach to the administration of the adverse childhood experiences questionnaire. *J Community Psychol*. 2021;49(5):1079-99.
55. Mollart L, Newing C, Foureur M. Midwives' emotional wellbeing: Impact of conducting a Structured Antenatal Psychosocial Assessment (SAPSA). *Women and Birth*. 2009;22(3):82-8.
56. Sandall J, Soltani H, Gates S, Shennan A, Devane D. Midwife-led continuity models versus other models of care for childbearing women. *Cochrane Database of Systematic Reviews*. 2016(4).
57. Bradford BF, Wilson AN, Portela A, McConville F, Fernandez Turienzo C, Homer CSE. Midwifery continuity of care: A scoping review of where, how, by whom and for whom? *PLOS Global Public Health*. 2022;2(10):e0000935.
58. May R, Dunkley-Bent J, Jolly M. Midwifery Continuity of Carer. NHS England; 21.9.22 21.9.22.
59. National Institute for Health and Care Excellence. Antenatal Care: NICE guideline NG201. 2021.
60. Cordey S, Moncrieff G, Cull J, Sarian A, the ASPIRE-COVID 19 CG. 'There's only so much you can be pushed': Magnification of the maternity staffing crisis by the 2020/21 COVID-19 pandemic. *BJOG: Int J Obstet Gy*. 2022;129(8):1408-9.

61. NHS England. Better Births: Improving outcomes of maternity services in England. England: NHS England; 2016.
62. Hunter B, Henley J, Fenwick J, Sidebotham M, Pallant J. Work, Health and Emotional Lives of Midwives in the United Kingdom: The UK WHELM study. Cardiff University: School of Healthcare Sciences; 2018.
63. Sidhu R, Su B, Shapiro KR, Stoll K. Prevalence of and factors associated with burnout in midwifery: A scoping review. *European Journal of Midwifery*. 2020;4.
64. Mason W, Morris K, Webb C, Daniels B, Featherstone B, Bywaters P, et al. Toward Full Integration of Quantitative and Qualitative Methods in Case Study Research: Insights From Investigating Child Welfare Inequalities. *Journal of Mixed Methods Research*. 2020;14(2):164-83.
65. Long J, Wertans E, Harper K, Brennan D, Harvey H, Allen R, et al. Femicide Census: UK Femicides, 2009-2018. 2020.
66. SAMHSA (U.S. Department of Health and Human Services Substance Abuse and Mental Health Services Administration). SAMHSA's Concept of Trauma and Guidance for a Trauma-Informed Approach. 2014.
67. Seng J, Taylor J, editors. Trauma informed care in the perinatal period. Edinburgh: Dunedin Academic Press Ltd.; 2015.
68. Sperlich M, Seng JS, Li Y, Taylor J, BradburyJones C. Integrating trauma-informed care into maternity care practice: Conceptual and practical issues. *J Midwifery Womens Health*. 2017;62(6):661-72.
69. UN General Assembly. Transforming our world: the 2030 Agenda for Sustainable Development. 2015.
70. Sexual and reproductive health and rights in the EU, in the frame of women's health, European Parliament, (2021).

## Response to reviewers

### Reviewer #1

Reviewer comment	Response
<p>I appreciate the opportunity to review this well-written review paper on routine discussion of previous trauma in the perinatal period. The methodology is well articulated, and it is a strength that the authors have included a reflexive note for the authors. I offer the following queries and suggestions primarily centering on the results and discussion sections.</p>	<p>Sincere thanks for taking the time to review our paper and for your well thought through and helpful suggestions to improve it.</p>
<p>For the first theme of “‘I did not know how to say it, and no-one asked me’: should maternity care providers ask women about previous trauma?”, the authors note that although very many of the included studies’ participants found routine inquiry acceptable, that there are issues that are masking some complexities and contradictions. Some of these complexities are elucidated in this theme and other themes – notably the “‘You say it is confidential... but you are going to report me’: the importance of trust” theme. While I agree with the authors that routine inquiry would be better positioned in the context of a trusted relationship and with continuity of care, give that this is difficult to establish with the rise of the “hospitalist” system of care, it will be difficult to realize this ideal in the short term. Therefore, this underscores the need for more training on the part of perinatal professionals, in trauma-informed care, but also in trauma inquiry. The authors do acknowledge this in the discussion briefly, but I think it bears more discussion. Specifically, the authors might add that routine trauma inquiry should NOT be about eliciting specific details of past trauma, and should rather be present-focused in that clinicians should be communicating that they are trying to ascertain what the present needs are and how they might be able to assist the</p>	<p>Thank you for these comments which we entirely agree with. We have expanded on the discussion section to address your points.</p>

<p>patient in the here and now with helpful resources and/or adjustments to the care plan that take the survivor’s specific needs into consideration. I also might add that there should be a preamble prior to inquiry that a) normalizes that you are asking all patients about this, b) that sharing such information is entirely voluntary, c) that any limits to confidentiality are fully explained, and that d) the point of the inquiry should be as mentioned above, present-focused, and that e) should only be done when there is adequacy of time to do the inquiry in a not-rushed manner, and that f) resources and supports are readily at hand and offered to all patients regardless of disclosure (as the authors have already alluded to).</p>	
<p>p. 26 – note that “homely” in American English means “unattractive.” Perhaps use “homey” or some other word?</p>	<p>Thank you - this is helpful!</p> <p>I have replaced the sentence with: Participants proposed that trauma discussions should be held in private, comfortable and welcoming surroundings</p>
<p>The theme title: ““Heavens! I forgot it!”: challenges to embedding trauma discussion in routine practice” could be better explained in the description of the theme. Also, I am not sure this quote is emblematic of the overall theme category – since this is where not only workload issues are covered but also where the authors are discussing partner presence at appointments as well as challenges related to language interpretation. Also, there are strategies that those working in IPV contexts employ to elicit trauma exposures despite the presence of the partner – perhaps include more information about this here or in the discussion?</p>	<p>We have expanded this section and changed the title to “Embedding trauma discussion in routine practice” to make it more applicable to the section contents.</p> <p>Within the discussion, we have noted the importance of ensuring the care provider can talk with the woman privately.</p>
<p>I am glad that the authors also bring up the role of supervision – which should go hand in hand with training and support for clinicians. I feel this could be highlighted even more, especially the need for reflective supervision which would allow support for clinicians who</p>	<p>We have expanded on the findings section related to care providers emotional wellbeing and the discussion section on supervision.</p>

<p>experience vicarious traumatization or triggers of their own trauma experiences through their interactions with survivor clients.</p>	
<p>Finally, the authors rightly call for systems change and the need for trauma-informed approaches. I feel this could also be augmented in the paper, perhaps by alluding not only to the Law et al 2021 guide but also to other published works regarding TIC in the perinatal period across provider contexts. I think there is opportunity here to provide more specificity as to how TIC might foster improvement in routine inquiry process, especially regarding the systems change the authors are calling for – what is really needed is a trauma informed continuum of responsiveness, of which inquiry is one piece (See Sperlich et al., 2017 below, as well as some other likely relevant resources):</p> <p>Kuehn, B. M. (2020). Trauma-informed care may ease patient fear, clinician burnout. <i>JAMA</i>, 323(7), 595-597.</p> <p>Kuzma, E. K., Pardee, M., &amp; Morgan, A. (2020). Implementing patient-centered trauma-informed care for the perinatal nurse. <i>The Journal of Perinatal &amp; Neonatal Nursing</i>, 34(4), E23-E31.</p> <p>Mosley, E. A., &amp; Lanning, R. K. (2020). Evidence and guidelines for trauma-informed doula care. <i>Midwifery</i>, 83, 102643.</p> <p>Nagle-Yang, S., Sachdeva, J., Zhao, L. X., Shenai, N., Shirvani, N., Worley, L. L., ... &amp; Byatt, N. (2022). Trauma-Informed Care for Obstetric and Gynecologic Settings. <i>Maternal and Child Health Journal</i>, 1-8.</p> <p>Seng, J. (2015). <i>Trauma informed care in the perinatal period</i>. Dunedin Academic Press Ltd.</p> <p>Sperlich, M., Seng, J. S., Li, Y., Taylor, J., &amp; Bradbury- Jones, C. (2017). Integrating trauma- informed care into maternity care practice: conceptual and practical issues.</p>	<p>We have expanded the discussion of trauma-informed approaches.</p>

<p>Journal of Midwifery &amp; Women's Health, 62(6), 661-672.</p> <p>Ward, L. G. (2020). Trauma-informed perinatal healthcare for survivors of sexual violence. The Journal of Perinatal &amp; Neonatal Nursing, 34(3), 199-202.</p>	
--	--

## Reviewer #2

Reviewer comment	Response
<p>This paper reports a qualitative evidence synthesis on routine discussion of previous trauma in the perinatal period and considers views from both women and maternity care professionals, which is a strength of the review. This is an important issue that is relevant to an international audience. The paper is well written, and a sensitive approach is evident. Overall, a thorough, systematic review is described, which incorporates a reflexive note regarding pre-existing beliefs of the authors on routine trauma discussion. The authors recognise the strengths and limitations of their review. I recommend that it is accepted subject to some minor revisions as detailed below.</p>	<p>Many thanks indeed for taking the time to review this paper and offering suggestions for its improvement. We are very grateful for your insight.</p>
<p>P11, first 2 paragraphs – ‘To understand how best to support this work’ an explicit statement about which work would be helpful. As written, the statement implies that routine trauma conversations are needed, and the aim supports this assumption (‘how best to support women and professionals in having routine trauma conversations’). However, the review question, set out in the paragraph below, suggests that women and maternity care professionals will not necessarily be in favour of routine discussion, and this is something the review will investigate. I am concerned that these paragraphs could be</p>	<p>Thank you for highlighting this. We propose the following amendment:</p> <p>To understand whether routine trauma discussion should be carried out, and if so how it should be done, we aimed to collate and synthesise all relevant qualitative studies to capture women’s and maternity care professionals’ views of this issue.</p>



<p>interpreted as bias by the authors, which is not evident in the rest of the paper.</p>	
<p>P13, more information on search terms would be helpful. The authors indicate that they used broad search terms, but the only terms mentioned are 'trauma-informed' and 'trauma informed'. Were any others used?</p>	<p>We have added a further paragraph in this section for clarity:</p> <p>We developed the search strategy by reviewing key relevant papers we were already aware of and examining relevant systematic reviews on PROSPERO. Each of the key papers referred to 'trauma-informed' in their keywords, title or abstract. We trialed searching using this term on Medline and duplicated it using a variety of different terms such as trauma-sensitive, trauma-focused, and trauma-responsive. These alternative terms had very few hits and had always been used in addition to trauma-informed. We therefore decided to only use the search terms 'trauma-informed' or 'trauma informed' and to hand select relevant papers that concerned the care of women in the perinatal period.</p>
<p>P14, 4th line from bottom of page – 'This enabled comparison between papers and reviewers.' I found this comment confusing here and think it should be deleted as an explanation of comparison between papers and reviewers is provided on the following page</p>	<p>Comment deleted</p>
<p>P27 'a participant in Olsen...', P32 'Participants in Marley et al. [34] and Mollart...' Please could reference to these three studies be amended to say, 'in the study/studies by'. This feels more respectful and is a construct used elsewhere.</p>	<p>Now amended, I appreciate your comment and agree that this wording is better</p>
<p>P27-28 'women reported feeling that there was no connection between trauma history and current pregnancy'. Please could this statement be qualified. As written, this could be misinterpreted as suggesting that all</p>	<p>Now qualified as follows, "in seven of the studies, some participants reported feeling..."</p>

<p>women in these 7 studies reported there was no connection between their trauma history and current wellbeing, which is not the case. Sometimes it is a small minority of women who hold this view.</p>	
<p>P30-31 'Heavens! I forgot it!' This section feels very brief. The introductory sentence indicates that the theme will investigate how trauma discussion can be introduced and include practical steps to aid implementation. I do not recognise this description in the theme. Rather than indicating how trauma discussion can be introduced, it just indicates feasibility. It raises a couple of potential challenges but does not seem to include practical steps to implementation. This section would benefit from review by the authors.</p>	<p>We have expanded this section and changed the title to make it more applicable to the section contents.</p>
<p>P32, the last theme is also very brief. I am wondering if there is any more relevant data in the papers that could be added to the section. If not, it would be helpful if the lack of data on this important issue could be addressed in the discussion.</p>	<p>We have expanded on the findings section related to care providers emotional wellbeing.</p>
<p>P35, 'Further research is needed to ascertain what support is helpful to women who disclose trauma'. I would find a further exploration of this statement helpful. As presented, it implies that it will be possible to identify specific support for women who disclose trauma. I suspect that this will be dependent on individual women and how far through their journey to recovery they are.</p>	<p>We have developed this phrase further:</p> <p>Further research is needed to ascertain what resources and support are helpful to women who have suffered trauma, bearing in mind that this must focus on the preferences and needs of the individual.</p>
<p>I have some concerns about the idea of 'routine' in relation to this work, as in the title and throughout the paper. It belies the complexity of working with people who have</p>	<p>We have added a paragraph in the introduction to clarify our thoughts around this:</p> <p>We use the term 'routine' to indicate raising the issue of previous trauma with all women</p>

<p>experienced trauma and the need for responsive care</p>	<p>accessing maternity services, as opposed to selectively for women who the healthcare professional suspects may have suffered trauma. The term 'routine' in this context does not negate the need for personalised care. We have chosen to use the term 'discussion' rather than 'enquiry' as discussion leaves open the possibility that trauma could be raised in a way that does not involve a direct question.</p>
<p>P14 Line 3 of section Appraisal of the methodological quality of included studies: 1 mark was allocated to each question if the criteria was met. Should be 'criterion'</p>	<p>Now amended</p>
<p>When in-text citations include mention of authors, there is a lack of consistency as to whether dates are included. Please could this be addressed.</p>	<p>Now amended, thank you</p>
<p>Table 1, country column – Sometimes the authors refer to United States and sometimes to America. Consistency would be helpful</p>	<p>Now amended, thank you for highlighting this</p>