

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/158604/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Shams, Mudassir, Kausar, Nasreen, Alayyash, Khulud, Al-Shamiri, Mohammed M., Arif, Nayyab and Ismail, Rashad 2023. Semi-analytical scheme for solving intuitionistic fuzzy system of differential equations. IEEE Access 11 , pp. 33205-33223. 10.1109/ACCESS.2023.3241482 file

Publishers page: <http://dx.doi.org/10.1109/ACCESS.2023.3241482>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



RESEARCH ARTICLE

Semi-Analytical Scheme for Solving Intuitionistic Fuzzy System of Differential Equations

MUDASSIR SHAMS¹, NASREEN KAUSAR^{1,2}, KHULUD ALAYYASH³,
MOHAMMED M. AL-SHAMIRI^{4,5}, NAYYAB ARIF¹,
AND RASHAD ISMAIL^{4,5}

¹Department of Mathematics and Statistics, Riphah International University, Islamabad 44000, Pakistan

²Department of Mathematics, Faculty of Arts and Sciences, Yildiz Technical University, Esenler, 34220 Istanbul, Turkey

³Department of Mathematics, Cardiff University, CF10 3AT Cardiff, U.K.

⁴Department of Mathematics, Faculty of Science and Arts, Mahayl Assir, King Khalid University, Abha 62529, Saudi Arabia

⁵Department of Mathematics and Computers, Faculty of Sciences, Ibb University, Ibb, Yemen

Corresponding author: Rashad Ismail (rashadif@gmail.com)

This work was supported by the Deanship of Scientific Research at King Khalid University through Small Groups Project under Grant R.G.P.1/277/43.

ABSTRACT The aim of this article is to implement the Generalized Modified Adomian Decomposition Method to compute the semi-numerical solution of the linear system of intuitionistic fuzzy initial value problems. Here, we consider the initial values as generalized trapezoidal intuitionistic fuzzy numbers. The technique is applied to brine tanks problem and coupled mass spring systems. Theoretically, different approaches to solving a system of generalized trapezoidal intuitionistic fuzzy differential equations are discussed in this study under the presumption that the coefficients of the system of the differential equations are associated to generalized trapezoidal intuitionistic fuzzy numbers. The approximate results are compared with exact solutions which shows good efficiency. The corresponding graphs at different levels of uncertainty show the example's numerical outcomes. The graphical representations further demonstrate the effectiveness and accuracy of the proposed method in comparison to existing semi-numerical methods in the literature.

INDEX TERMS Fuzzy set, fuzzy number, generalized trapezoidal intuitionistic fuzzy number, system of fuzzy differential equation, analytical technique, engineering applications.

I. INTRODUCTION

System of differential equation plays a significant role in modeling and studying many naturally occurring phenomenon such as population models, economic models, friction model, bacteria culture model, predator-prey model, weight loss and oil production model, bank account and drug concentration problem, human immunodeficiency virus (HIV) model. In classical set theory, the variables or parameters are taken as crisp numbers. But in actual case, these variables or parameters are usually uncertain or vague. So, these variables may be considered as a fuzzy numbers. In other words, to overcome uncertainty we use

fuzzy numbers. So the system of differential equations are converted to system of fuzzy differential equations (SFDE).

The concept of fuzzy set theory was firstly introduced by Zadeh in 1965, as the extension of classical set theory [1]. The concept of fuzzy set theory has been applied to various fields of science and engineering to handle vagueness and uncertainty. In 1987, Kandel and Byatt [2] introduced the fuzzy differential equations. The fuzzy differential equations have been applied in numerous daily life problems [3], [4], [5]. Vasavi et al. [6] discussed fuzzy differential for cooling problems. Devi and Ganesan used fuzzy differential equations in modelling electric circuit problem [7]. Ahmad et al. [8] studied a mathematical method to find the solution of fuzzy integro differential equations. Sadeghi et al. [9] studied the system of fuzzy differential equation. Buckley et al find the solution of system of first

The associate editor coordinating the review of this manuscript and approving it for publication was Longzhi Yang¹⁰.

order linear fuzzy differential equations by extension principle [10]. Hashemi et al. find the series solution of SFDE [11]. In 1986, Atanassov [12] introduced an extension of fuzzy set theory known as intuitionistic fuzzy set. The intuitionistic fuzzy set [13] not only provides the information about membership values but also the non-membership values respectively, and so that the sum of both values is less than one. Intuitionistic fuzzy differential equations are being studied widely and being used in various fields of Physics, Chemistry, Biology as well as among other fields of science and engineering. Melliani and Chadli obtained the approximate and numerical solutions of intuitionistic fuzzy differential equations with linear differential operators [14], [15]. Gulzar et al. worked on fuzzy algebra [16], [17], [18]. Akin and Bayeg [19], [20] studied a method to find general solution of second order intuitionistic fuzzy differential equation and to solve the system of intuitionistic fuzzy differential equations with intuitionistic fuzzy initial values. Mondal and Roy [21], [22], [23] studied the generalized intuitionistic fuzzy Laplace transform method and to solve the system of differential equations with initial value as triangular intuitionistic fuzzy number. Saw et al introduced a method for solving system of linear intuitionistic fuzzy equations [24].

The Adomian Decomposition Method (ADM) which is a semi analytical method was first presented by Adomian in 1980's [25], [26]. This method is very efficient in finding the solutions of differential equations, algebraic equations as well as integral equations. In this article, we will propose the Generalized Modified Adomian Decomposition Method (GMADM) to find the solutions of the system of linear intuitionistic fuzzy differential equations with initial values as generalized trapezoidal intuitionistic fuzzy number. This modification was proposed by Wazwaz [27]. He presented a reliable modification to the ADM. In this modification Wazwaz divides the original function into two parts, one part assigned to the initial term of the series and the other to the second term. This modification results in a different series being generated. The efficiency of this method depends only on the choice of the parts into which the original function is to be divided.

First order system of fuzzy differential equations is important among all the fuzzy differential equations. There are many approaches to solve the SFDEs. Buckley and Feuring [28] solving the linear system of first order ordinary differential equations with fuzzy initial conditions by extension principle using triangular fuzzy number. The geometric approach is developed by Gasilova et al. [29] and series solution is developed by Hashemi et al. [30]. Mondal and Roy [31] studied strong and weak solution of first order homogeneous intuitionistic fuzzy differential equation, subsequently and studied system of differential equation in literature. Melliani, et al. [32] discussed the existence and uniqueness of the solution of the intuitionistic fuzzy differential equation and its system using the analytical technique. Therefore, finding an efficient and accurate algorithm for investigating

FIE has been one the hot areas of research in recent time. To achieve these goals, various methods and procedures were used to handle differentia equations, using triangular fuzzy number, for details, see [9], [33].

In this study, motivated by the aforementioned work, we solve the system of differential equations using a GMADM and a more generalized fuzzy system of differential equations, namely a trapezoidal intuitionistic fuzzy system of intuitionistic differential equations.

The main contributions of this research work are summarized below.

- GMADM is used to solve a system of differential equations using initial conditions as a Generalized trapezoidal intuitionistic fuzzy number.
- In order to solve a system of fuzzy intuitionistic differential equations that have not before been explored, the computational complexity of the suggested GMADM is discussed.
- Applications of system of Generalized trapezoidal intuitionistic fuzzy differential equations in mechanical engineering are taken into consideration in a Generalized trapezoidal intuitionistic fuzzy environment.
- Computational tools are used to evaluate the effectiveness and applicability of the suggested analytical scheme.

This paper is organized as follows: In section II, we recall some basic definitions which we will use in further sections. In sections III, we introduced our proposed method. In section IV, the efficiency of this method has been illustrated by applications. In the last section, we give conclusions.

II. PRELIMINARIES

In this section, the fundamental definitions of fuzzy set and intuitionistic fuzzy set are presented.

Definition 1 [34]: Let $\overset{\circ}{U}$ be the largest set under consideration then $\overset{\star}{F}$ be a subset of $\overset{\circ}{U}$ is said to be a fuzzy set if it is defined as:

$$\mu_{\overset{\star}{F}}(\hat{u}) = \overset{\circ}{U} \longrightarrow [0, 1],$$

defines the degree of membership of an element $\hat{u} \in \overset{\circ}{U}$ to the set $\overset{\star}{F}$ which is a subset of $\overset{\circ}{U}$.

Definition 2 [34]: α -cut of a fuzzy set $\overset{\star}{F}$ is a crisp set $\overset{\star}{F}_{\alpha}$ which is defined as:

$$\overset{\star}{F}_{\alpha} = \{(\hat{u} \mid \mu_{\overset{\star}{F}}(\hat{u}) \geq \alpha : \hat{u} \in \overset{\circ}{U})\}.$$

Definition 3 [23]: If $\overset{\star}{F}$ is a fuzzy set then height of a fuzzy set is denoted by $h(\overset{\star}{F})$ and is defined as the largest membership function obtained by any element in that set i.e.,

$$h(\overset{\star}{F}) = \sup \mu_{\overset{\star}{F}}(\hat{u}).$$

Definition 4 [19]: Let $\overset{\circ}{U}$ be a nonempty finite set of real numbers, then an intuitionistic fuzzy set $\overset{\star}{I}$ on $\overset{\circ}{U}$ is:

$$\overset{\star}{I} = \{(\hat{u}, \mu_{\overset{\star}{I}}(\hat{u}), \nu_{\overset{\star}{I}}(\hat{u})) : \hat{u} \in \overset{\circ}{U}\},$$

where the functions,

$$\mu_{\overset{\star}{I}}(\hat{u}) = \overset{\circ}{U} \rightarrow [0, 1],$$

$$\nu_{\overset{\star}{I}}(\hat{u}) = \overset{\circ}{U} \rightarrow [0, 1],$$

define the degree of membership and degree of non-membership respectively, of an element $\hat{u} \in \overset{\circ}{U}$ to the set $\overset{\star}{I}$ which is a subset of $\overset{\circ}{U}$, and for every $\hat{u} \in \overset{\circ}{U}$, the

$$0 \leq \mu_{\overset{\star}{I}}(\hat{u}) + \nu_{\overset{\star}{I}}(\hat{u}) \leq 1,$$

condition must be satisfied.

Definition 5 [23]: An intuitionistic fuzzy set $\overset{\star}{I}$ is said to be normal if there exists an $\hat{u}_0 \in \overset{\circ}{U}$, such that $\mu_{\overset{\star}{I}}(\hat{u}_0) = 1$ so $\nu_{\overset{\star}{I}}(\hat{u}_0) = 0$.

Definition 6 [23]: An intuitionistic fuzzy set $\overset{\star}{I}$ is said to be convex set for the membership function if it satisfy the following condition:

$$\begin{aligned} \mu_{\overset{\star}{I}}(\hat{u})(\eta\hat{u} + (1 - \eta)\hat{s}) &\geq \min(\mu_{\overset{\star}{I}}(\hat{u}), \mu_{\overset{\star}{I}}(\hat{s})); \\ \forall \hat{u}, \hat{s} \in \overset{\circ}{U}, \eta \in [0, 1]. \end{aligned}$$

Definition 7 [23]: An intuitionistic fuzzy set $\overset{\star}{I}$ is said to be concave set for the non-membership function if it satisfy the following condition:

$$\begin{aligned} \nu_{\overset{\star}{I}}(\hat{u})(\eta\hat{u} + (1 - \eta)\hat{s}) &\geq \max(\nu_{\overset{\star}{I}}(\hat{u}), \nu_{\overset{\star}{I}}(\hat{s})); \\ \forall \hat{u}, \hat{s} \in \overset{\circ}{U}, \eta \in [0, 1]. \end{aligned}$$

A. GENERALIZED INTUITIONISTIC FUZZY NUMBER

Definition 8 [35]: A generalized intuitionistic fuzzy number

$$\overset{\star}{T} = \langle (s_1^{\star}, s_2^{\star}, s_3^{\star}, s_4^{\star}; \nu_A); (t_1^{\star}, t_2^{\star}, t_3^{\star}, t_4^{\star}; \nu_B) \rangle$$

is said to be generalized trapezoidal intuitionistic fuzzy number (GTIFN) (as shown in Figure 1) if its membership and non-membership functions are defined as follows:

$$\mu_{\overset{\star}{T}}(\hat{u}) = \begin{cases} \nu_A \left(\frac{\hat{u} - s_1^{\star}}{s_2^{\star} - s_1^{\star}} \right), & s_1^{\star} \leq \hat{u} \leq s_2^{\star}, \\ \nu_A, & s_2^{\star} \leq \hat{u} \leq s_3^{\star}, \\ \nu_A \left(\frac{s_4^{\star} - \hat{u}}{s_4^{\star} - s_3^{\star}} \right), & s_3^{\star} \leq \hat{u} \leq s_4^{\star}, \\ 0, & \text{otherwise,} \end{cases} \dots$$

$$\nu_{\overset{\star}{T}}(\hat{u}) = \begin{cases} \frac{(t_2^{\star} - \hat{u}) + \nu_B(t_1^{\star} - \hat{u})}{t_2^{\star} - t_1^{\star}}, & t_1^{\star} \leq \hat{u} \leq t_2^{\star}, \\ \nu_B, & t_2^{\star} \leq \hat{u} \leq t_3^{\star}, \\ \frac{(\hat{u} - t_3^{\star}) + \nu_B(t_4^{\star} - \hat{u})}{t_4^{\star} - t_3^{\star}}, & t_3^{\star} \leq \hat{u} \leq t_4^{\star}, \\ 1, & \text{otherwise,} \end{cases}$$




FIGURE 1. Generalized trapezoidal intuitionistic fuzzy number.

where $t_1^{\star} \leq s_1^{\star} \leq t_2^{\star} \leq s_2^{\star} \leq s_3^{\star} \leq t_3^{\star} \leq s_4^{\star} \leq t_4^{\star}$, $0 \leq \nu_A, \nu_B \leq 1$ and $0 < \nu_A + \nu_B \leq 1$.

Definition 9 [36], [37]: (α, β) -cut set of a generalized trapezoidal intuitionistic fuzzy number $\overset{\star}{T} = \langle (s_1^{\star}, s_2^{\star}, s_3^{\star}, s_4^{\star}; \nu_A); (t_1^{\star}, t_2^{\star}, t_3^{\star}, t_4^{\star}; \nu_B) \rangle$ is a crisp subset of $\overset{\circ}{U}$ which is defined as:

$$\begin{aligned} \overset{\star}{T}(\alpha, \beta) &= \{(\hat{u}, \mu_{\overset{\star}{T}}(\hat{u}), \nu_{\overset{\star}{T}}(\hat{u})) : \hat{u} \in U, \mu_{\overset{\star}{T}}(\hat{u}) \geq \alpha, \nu_{\overset{\star}{T}}(\hat{u}) \geq \beta\} \\ &= \{[T_1(\alpha), T_2(\alpha)]; [T_1(\beta), T_2(\beta)]\}, \end{aligned}$$

where $\alpha + \beta < 1$, $\alpha \in [0, \nu_A]$ and $\beta \in [\nu_B, 1]$.

Figure 1: Shows membership and non-membership function of generalized trapezoidal intuitionistic fuzzy number.

B. ARITHMETIC OPERATIONS ON GTIFNs

Definition 10 [37], [38]: Let $\overset{\star}{T}_1 = \langle (s_1^{\star}, s_2^{\star}, s_3^{\star}, s_4^{\star}; \nu_{A1}); (t_1^{\star}, t_2^{\star}, t_3^{\star}, t_4^{\star}; \nu_{B1}) \rangle$ and $\overset{\star}{T}_2 = \langle (u_1^{\star}, u_2^{\star}, u_3^{\star}, u_4^{\star}; \nu_{B1}); (v_1^{\star}, v_2^{\star}, v_3^{\star}, v_4^{\star}; \nu_{\star}) \rangle$ be two GTIFNs and ϖ be a real number. Then

- $$\begin{aligned} \overset{\star}{T}_1 + \overset{\star}{T}_2 &= \langle (s_1^{\star} + u_1^{\star}, s_2^{\star} + u_2^{\star}, s_3^{\star} + u_3^{\star}, s_4^{\star} + u_4^{\star}; \min\{\nu_{A1}, \nu_{B1}\}); \\ & (t_1^{\star} + v_1^{\star}, t_2^{\star} + v_2^{\star}, t_3^{\star} + v_3^{\star}, t_4^{\star} + v_4^{\star}; \max\{\nu_{A2}, \nu_{B2}\}) \rangle. \end{aligned}$$
- $$\begin{aligned} \overset{\star}{T}_1 - \overset{\star}{T}_2 &= \langle (s_1^{\star} - u_4^{\star}, s_2^{\star} - u_3^{\star}, s_3^{\star} - u_2^{\star}, s_4^{\star} - u_1^{\star}; \min\{\nu_{A1}, \nu_{B1}\}); \\ & (t_1^{\star} - v_4^{\star}, t_2^{\star} - v_3^{\star}, t_3^{\star} - v_2^{\star}, t_4^{\star} - v_1^{\star}; \max\{\nu_{A2}, \nu_{B2}\}) \rangle. \end{aligned}$$

- $$\begin{aligned} \overset{\star}{T}_1 \times \overset{\star}{T}_2 = & \langle (\overset{\star}{s}_1 \overset{\star}{u}_1, \overset{\star}{s}_2 \overset{\star}{u}_2, \overset{\star}{s}_3 \overset{\star}{u}_3, \overset{\star}{s}_4 \overset{\star}{u}_4; \min\{v_{A_1}, v_{B_1}\}); \\ & (t_1 \overset{\star}{v}_1, t_2 \overset{\star}{v}_2, t_3 \overset{\star}{v}_3, t_4 \overset{\star}{v}_4; \max\{v_{A_2}, v_{B_2}\}) \rangle, \end{aligned}$$

where $\overset{\star}{T}_1 > 0, \overset{\star}{T}_2 > 0$.

- $$\begin{aligned} \overset{\star}{T}_1 \times \overset{\star}{T}_2 = & \langle (\overset{\star}{s}_1 \overset{\star}{u}_4, \overset{\star}{s}_2 \overset{\star}{u}_3, \overset{\star}{s}_3 \overset{\star}{u}_2, \overset{\star}{s}_4 \overset{\star}{u}_1; \min\{v_{A_1}, v_{B_1}\}); \\ & (t_1 \overset{\star}{v}_4, t_2 \overset{\star}{v}_3, t_3 \overset{\star}{v}_2, t_4 \overset{\star}{v}_1; \max\{v_{A_2}, v_{B_2}\}) \rangle, \end{aligned}$$

where $\overset{\star}{T}_1 < 0, \overset{\star}{T}_2 > 0$.

- $$\begin{aligned} \overset{\star}{T}_1 \times \overset{\star}{T}_2 = & \langle (\overset{\star}{s}_4 \overset{\star}{u}_4, \overset{\star}{s}_3 \overset{\star}{u}_3, \overset{\star}{s}_2 \overset{\star}{u}_2, \overset{\star}{a}_1 \overset{\star}{u}_1; \min\{v_{A_1}, v_{B_1}\}); \\ & (t_4 \overset{\star}{v}_4, t_3 \overset{\star}{v}_3, t_2 \overset{\star}{v}_2, t_1 \overset{\star}{v}_1; \max\{v_{A_2}, v_{B_2}\}) \rangle, \end{aligned}$$

where $\overset{\star}{T}_1 < 0, \overset{\star}{T}_2 < 0$.

- $$\begin{aligned} \overset{\star}{T}_1 \div \overset{\star}{T}_2 = & \langle (\frac{\overset{\star}{s}_1}{\overset{\star}{u}_4}, \frac{\overset{\star}{s}_2}{\overset{\star}{u}_3}, \frac{\overset{\star}{s}_3}{\overset{\star}{u}_2}, \frac{\overset{\star}{s}_4}{\overset{\star}{u}_1}; \min\{v_{A_1}, v_{B_1}\}); \\ & (\frac{t_1}{\overset{\star}{v}_4}, \frac{t_2}{\overset{\star}{v}_3}, \frac{t_3}{\overset{\star}{v}_2}, \frac{t_4}{\overset{\star}{v}_1}; \max\{v_{A_2}, v_{B_2}\}) \rangle, \end{aligned}$$

where $\overset{\star}{T}_2 > 0$.

- $$\begin{aligned} \varpi \overset{\star}{T}_1 = & \langle (\varpi \overset{\star}{s}_1, \varpi \overset{\star}{s}_2, \varpi \overset{\star}{s}_3, \varpi \overset{\star}{s}_4; \min\{v_{A_1}, v_{B_1}\}); \\ & (\varpi t_1, \varpi t_2, \varpi t_3, \varpi t_4; \max\{v_{A_2}, v_{B_2}\}) \rangle, \end{aligned}$$

where $\varpi > 0$.

- $$\begin{aligned} \varpi \overset{\star}{T}_1 = & \langle (\varpi \overset{\star}{s}_4, \varpi \overset{\star}{s}_3, \varpi \overset{\star}{s}_2, \varpi \overset{\star}{s}_1; \min\{v_{A_1}, v_{B_1}\}); \\ & (\varpi t_4, \varpi t_3, \varpi t_2, \varpi t_1; \max\{v_{A_2}, v_{B_2}\}) \rangle, \end{aligned}$$

where $\varpi < 0$.

III. THE GENERALIZED MODIFIED ADOMIAN DECOMPOSITION METHOD

Let us consider the system of intuitionistic fuzzy differential equations with linear differential operator as follows:

$$\left. \begin{aligned} \overset{\boxtimes}{L} \overset{\star}{x}(t) + \overset{\boxtimes}{R} \overset{\star}{x}(t) + \overset{\boxtimes}{R} \overset{\star}{y}(t) + N(t, \overset{\star}{x}(t), \overset{\star}{y}(t)) &= \overset{\star}{g}(t), \\ \overset{\boxtimes}{L} \overset{\star}{y}(t) + \overset{\boxtimes}{R} \overset{\star}{x}(t) + \overset{\boxtimes}{R} \overset{\star}{y}(t) + N(t, \overset{\star}{x}(t), \overset{\star}{y}(t)) &= \overset{\star}{h}(t), \end{aligned} \right\} \quad (1)$$

where $\overset{\boxtimes}{L}$ is the highest order linear differential operator, $\overset{\boxtimes}{R}$ is the remaining part of the linear differential operator, N may be linear or nonlinear function of $t, \overset{\star}{x}(t)$ and $\overset{\star}{y}(t)$, $\overset{\star}{g}(t)$ and $\overset{\star}{h}(t)$ are non-homogeneous terms. Here, in this case we take N as a linear function of $\overset{\star}{x}(t), \overset{\star}{y}(t)$ and t . Taking (α, β) -cut of (1), we get (2), as shown at the bottom of the page.

From (2), we obtain the following equations:

$$\left. \begin{aligned} \overset{\boxtimes}{L} \overset{\star}{x}_1(t, \alpha) + \overset{\boxtimes}{R} \overset{\star}{x}_1(t, \alpha) + \overset{\boxtimes}{R} \overset{\star}{y}_1(t, \alpha) + \\ N_1(t, \overset{\star}{x}_1(t, \alpha), \overset{\star}{y}_1(t, \alpha)) &= \overset{\star}{g}_1(t, \alpha), \\ \overset{\boxtimes}{L} \overset{\star}{y}_1(t, \alpha) + \overset{\boxtimes}{R} \overset{\star}{x}_1(t, \alpha) + \overset{\boxtimes}{R} \overset{\star}{y}_1(t, \alpha) + \\ N_1(t, \overset{\star}{x}_1(t, \alpha), \overset{\star}{y}_1(t, \alpha)) &= \overset{\star}{h}_1(t, \alpha). \end{aligned} \right\} \quad (3)$$

$$\left. \begin{aligned} \overset{\boxtimes}{L} \overset{\star}{x}_2(t, \alpha) + \overset{\boxtimes}{R} \overset{\star}{x}_2(t, \alpha) + \overset{\boxtimes}{R} \overset{\star}{y}_2(t, \alpha) + \\ N_2(t, \overset{\star}{x}_2(t, \alpha), \overset{\star}{y}_2(t, \alpha)) &= \overset{\star}{g}_2(t, \alpha), \\ \overset{\boxtimes}{L} \overset{\star}{y}_2(t, \alpha) + \overset{\boxtimes}{R} \overset{\star}{x}_2(t, \alpha) + \overset{\boxtimes}{R} \overset{\star}{y}_2(t, \alpha) + \\ N_2(t, \overset{\star}{x}_2(t, \alpha), \overset{\star}{y}_2(t, \alpha)) &= \overset{\star}{h}_2(t, \alpha). \end{aligned} \right\} \quad (4)$$

$$\left. \begin{aligned} \overset{\boxtimes}{L} \overset{\star}{x}_1(t, \beta) + \overset{\boxtimes}{R} \overset{\star}{x}_1(t, \beta) + \overset{\boxtimes}{R} \overset{\star}{y}_1(t, \beta) + \\ N_1(t, \overset{\star}{x}_1(t, \beta), \overset{\star}{y}_1(t, \beta)) &= \overset{\star}{g}_1(t, \beta), \\ \overset{\boxtimes}{L} \overset{\star}{y}_1(t, \beta) + \overset{\boxtimes}{R} \overset{\star}{x}_1(t, \beta) + \overset{\boxtimes}{R} \overset{\star}{y}_1(t, \beta) + \\ N_1(t, \overset{\star}{x}_1(t, \beta), \overset{\star}{y}_1(t, \beta)) &= \overset{\star}{h}_1(t, \beta). \end{aligned} \right\} \quad (5)$$

$$\left. \begin{aligned} \overset{\boxtimes}{L}([\overset{\star}{x}_1(t, \alpha), \overset{\star}{x}_2(t, \alpha)]; [\overset{\star}{x}_1(t, \beta), \overset{\star}{x}_2(t, \beta)]) + \overset{\boxtimes}{R}([\overset{\star}{x}_1(t, \alpha), \overset{\star}{x}_2(t, \alpha)]; \\ [\overset{\star}{x}_1(t, \beta), \overset{\star}{x}_2(t, \beta)]) + \overset{\boxtimes}{R}([\overset{\star}{y}_1(t, \alpha), \overset{\star}{y}_2(t, \alpha)]; [\overset{\star}{y}_1(t, \beta), \overset{\star}{y}_2(t, \beta)]) \\ + ([N_1(t, \overset{\star}{x}_1(t, \alpha), \overset{\star}{y}_1(t, \alpha)), N_2(t, \overset{\star}{x}_2(t, \alpha), \overset{\star}{y}_2(t, \alpha))]; \\ [N_1(t, \overset{\star}{x}_1(t, \beta), \overset{\star}{y}_1(t, \beta)), N_2(t, \overset{\star}{x}_2(t, \beta), \overset{\star}{y}_2(t, \beta))]) \\ = ([\overset{\star}{g}_1(t, \alpha), \overset{\star}{g}_2(t, \alpha)]; [\overset{\star}{g}_1(t, \beta), \overset{\star}{g}_2(t, \beta)]), \\ \overset{\boxtimes}{L}([\overset{\star}{y}_1(t, \alpha), \overset{\star}{y}_2(t, \alpha)]; [\overset{\star}{y}_1(t, \beta), \overset{\star}{y}_2(t, \beta)]) + \overset{\boxtimes}{R}([\overset{\star}{x}_1(t, \alpha), \overset{\star}{x}_2(t, \alpha)]; \\ [\overset{\star}{x}_1(t, \beta), \overset{\star}{x}_2(t, \beta)]) + \overset{\boxtimes}{R}([\overset{\star}{y}_1(t, \alpha), \overset{\star}{y}_2(t, \alpha)]; [\overset{\star}{y}_1(t, \beta), \overset{\star}{y}_2(t, \beta)]) \\ + ([N_1(t, \overset{\star}{x}_1(t, \alpha), \overset{\star}{y}_1(t, \alpha)), N_2(t, \overset{\star}{x}_2(t, \alpha), \overset{\star}{y}_2(t, \alpha))]; \\ [N_1(t, \overset{\star}{x}_1(t, \beta), \overset{\star}{y}_1(t, \beta)), N_2(t, \overset{\star}{x}_2(t, \beta), \overset{\star}{y}_2(t, \beta))]) \\ = ([\overset{\star}{h}_1(t, \alpha), \overset{\star}{h}_2(t, \alpha)]; [\overset{\star}{h}_1(t, \beta), \overset{\star}{h}_2(t, \beta)]). \end{aligned} \right\} \quad (2)$$

$$\left. \begin{aligned} & \mathbb{L}^* \dot{x}_2(t, \beta) + \mathbb{R}^* \dot{x}_2(t, \beta) + \mathbb{R}^* \dot{y}_2(t, \beta) + \\ & N_1(t, \dot{x}_2(t, \beta), \dot{y}_2(t, \beta)) = \dot{g}_2(t, \beta), \\ & \mathbb{L}^* \dot{y}_2(t, \beta) + \mathbb{R}^* \dot{x}_2(t, \beta) + \mathbb{R}^* \dot{y}_2(t, \beta) + \\ & N_1(t, \dot{x}_2(t, \beta), \dot{y}_2(t, \beta)) = \dot{h}_2(t, \beta). \end{aligned} \right\} \quad (6)$$

Applying the \mathbb{L}^{-1} operator on both sides of (3), (4), (5) and (6), we get;

$$\left. \begin{aligned} \dot{x}_1(t, \alpha) &= \Psi_1(t, \alpha) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{x}_1(t, \alpha)) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{y}_1(t, \alpha)) \\ & - \mathbb{L}^{-1} (N_1(t, \dot{x}_1(t, \alpha), \dot{y}_1(t, \alpha))) + \mathbb{L}^{-1} (g_1(t, \alpha)), \\ \dot{y}_1(t, \alpha) &= \Phi_1(t, \alpha) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{x}_1(t, \alpha)) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{y}_1(t, \alpha)) \\ & - \mathbb{L}^{-1} (N_1(t, \dot{y}_1(t, \alpha))) + \mathbb{L}^{-1} (h_1(t, \alpha)). \end{aligned} \right\} \quad (7)$$

$$\left. \begin{aligned} \dot{x}_2(t, \alpha) &= \Psi_2(t, \alpha) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{x}_2(t, \alpha)) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{y}_2(t, \alpha)) \\ & - \mathbb{L}^{-1} (N_2(t, \dot{x}_2(t, \alpha), \dot{y}_2(t, \alpha))) + \mathbb{L}^{-1} (g_2(t, \alpha)), \\ \dot{y}_2(t, \alpha) &= \Phi_2(t, \alpha) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{x}_2(t, \alpha)) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{y}_2(t, \alpha)) \\ & - \mathbb{L}^{-1} (N_2(t, \dot{x}_2(t, \alpha), \dot{y}_2(t, \alpha))) + \mathbb{L}^{-1} (h_2(t, \alpha)), \end{aligned} \right\} \quad (8)$$

$$\left. \begin{aligned} \dot{x}_1(t, \beta) &= \Psi_1(t, \beta) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{x}_1(t, \beta)) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{y}_1(t, \beta)) \\ & - \mathbb{L}^{-1} (N_1(t, \dot{x}_1(t, \beta), \dot{y}_1(t, \beta))) + \mathbb{L}^{-1} (g_1(t, \beta)), \\ \dot{y}_1(t, \beta) &= \Phi_1(t, \beta) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{x}_1(t, \beta)) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{y}_1(t, \beta)) \\ & - \mathbb{L}^{-1} (N_1(t, \dot{x}_1(t, \beta), \dot{y}_1(t, \beta))) + \mathbb{L}^{-1} (h_1(t, \beta)), \end{aligned} \right\} \quad (9)$$

$$\left. \begin{aligned} \dot{x}_2(t, \beta) &= \Psi_2(t, \beta) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{x}_2(t, \beta)) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{y}_2(t, \beta)) \\ & - \mathbb{L}^{-1} (N_2(t, \dot{x}_2(t, \beta), \dot{y}_2(t, \beta))) + \mathbb{L}^{-1} (g_2(t, \beta)), \\ \dot{y}_2(t, \beta) &= \Phi_2(t, \beta) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{x}_2(t, \beta)) - \mathbb{L}^{-1} (\mathbb{R}^* \dot{y}_2(t, \beta)) \\ & - \mathbb{L}^{-1} (N_2(t, \dot{x}_2(t, \beta), \dot{y}_2(t, \beta))) + \mathbb{L}^{-1} (h_2(t, \beta)), \end{aligned} \right\} \quad (10)$$

where,

$$\left. \begin{aligned} \Psi_i(t, \alpha) &= \mathbb{L}^{-1} \Psi_i(t, \alpha) = 0, i = 1, 2 \\ \Phi_i(t, \alpha) &= \mathbb{L}^{-1} \Phi_i(t, \alpha) = 0, i = 1, 2 \\ \Psi_i(t, \beta) &= \mathbb{L}^{-1} \Psi_i(t, \beta) = 0, i = 1, 2 \\ \Phi_i(t, \beta) &= \mathbb{L}^{-1} \Phi_i(t, \beta) = 0, i = 1, 2 \end{aligned} \right\}$$

the above functions are found by using the initial conditions. Now by using the GMADM the solutions of the (7), (8), (9) and (10), can be expressed in the form of an infinite series for the unknown functions as follows:

$$\left. \begin{aligned} \dot{x}_1(t, \alpha) &= \sum_{n=0}^{\infty} \dot{x}_{1n}(t, \alpha), \\ \dot{y}_1(t, \alpha) &= \sum_{n=0}^{\infty} \dot{y}_{1n}(t, \alpha), \end{aligned} \right\} \quad (11)$$

$$\left. \begin{aligned} \dot{x}_2(t, \alpha) &= \sum_{n=0}^{\infty} \dot{x}_{2n}(t, \alpha), \\ \dot{y}_2(t, \alpha) &= \sum_{n=0}^{\infty} \dot{y}_{2n}(t, \alpha), \end{aligned} \right\} \quad (12)$$

$$\left. \begin{aligned} \dot{x}_1(t, \beta) &= \sum_{n=0}^{\infty} \dot{x}_{1n}(t, \beta), \\ \dot{y}_1(t, \beta) &= \sum_{n=0}^{\infty} \dot{y}_{1n}(t, \beta), \end{aligned} \right\} \quad (13)$$

$$\left. \begin{aligned} \dot{x}_2(t, \beta) &= \sum_{n=0}^{\infty} \dot{x}_{2n}(t, \beta), \\ \dot{y}_2(t, \beta) &= \sum_{n=0}^{\infty} \dot{y}_{2n}(t, \beta), \end{aligned} \right\} \quad (14)$$

Using (11), (12), (13) and (14), into (7), (8), (9) and (10), we have:

$$\left. \begin{aligned} \sum_{n=0}^{\infty} \dot{x}_{1n}(t, \alpha) &= \Psi_1(t, \alpha) - \mathbb{L}^{-1} (\mathbb{R}^* \sum_{n=0}^{\infty} \dot{x}_{1n}(t, \alpha)) \\ & - \mathbb{L}^{-1} (\mathbb{R}^* \sum_{n=0}^{\infty} \dot{y}_{1n}(t, \alpha)) - \mathbb{L}^{-1} (N_1(t, \sum_{n=0}^{\infty} \dot{x}_{1n}(t, \alpha), \\ & \sum_{n=0}^{\infty} \dot{y}_{1n}(t, \alpha))) + \mathbb{L}^{-1} (g_1(t, \alpha)), \end{aligned} \right\} \quad (15)$$

$$\left. \begin{aligned} \sum_{n=0}^{\infty} \dot{y}_{1n}(t, \alpha) &= \Phi_1(t, \alpha) - \mathbb{L}^{-1} (\mathbb{R}^* \sum_{n=0}^{\infty} \dot{x}_{1n}(t, \alpha)) \\ & - \mathbb{L}^{-1} (\mathbb{R}^* \sum_{n=0}^{\infty} \dot{y}_{1n}(t, \alpha)) - \mathbb{L}^{-1} (N_1(t, \sum_{n=0}^{\infty} \dot{x}_{1n}(t, \alpha), \\ & \sum_{n=0}^{\infty} \dot{y}_{1n}(t, \alpha))) + \mathbb{L}^{-1} (h_1(t, \alpha)), \end{aligned} \right\}$$

$$\left. \begin{aligned} \sum_{n=0}^{\infty} \dot{x}_{2n}(t, \alpha) &= \Psi_2(t, \alpha) - \mathbb{L}^{-1} (\mathbb{R}^* \sum_{n=0}^{\infty} \dot{x}_{2n}(t, \alpha)) \\ & - \mathbb{L}^{-1} (\mathbb{R}^* \sum_{n=0}^{\infty} \dot{y}_{2n}(t, \alpha)) - \mathbb{L}^{-1} (N_2(t, \sum_{n=0}^{\infty} \dot{x}_{2n}(t, \alpha), \\ & \sum_{n=0}^{\infty} \dot{y}_{2n}(t, \alpha))) + \mathbb{L}^{-1} (g_2(t, \alpha)), \end{aligned} \right\} \quad (16)$$

$$\left. \begin{aligned} \sum_{n=0}^{\infty} \dot{y}_{2n}(t, \alpha) &= \Phi_2(t, \alpha) - \mathbb{L}^{-1} (\mathbb{R}^* \sum_{n=0}^{\infty} \dot{x}_{2n}(t, \alpha)) \\ & - \mathbb{L}^{-1} (\mathbb{R}^* \sum_{n=0}^{\infty} \dot{y}_{2n}(t, \alpha)) - \mathbb{L}^{-1} (N_2(t, \sum_{n=0}^{\infty} \dot{x}_{2n}(t, \alpha), \\ & \sum_{n=0}^{\infty} \dot{y}_{2n}(t, \alpha))) + \mathbb{L}^{-1} (h_2(t, \alpha)), \end{aligned} \right\}$$

$$\left. \begin{aligned}
 \sum_{n=0}^{\infty} \star x_{1n}(t, \beta) &= \Psi_1(t, \beta) - \bar{L}^{-1} \left(\bar{R} \sum_{n=0}^{\infty} \star x_{1n}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \sum_{n=0}^{\infty} \star y_{1n}(t, \beta) \right) - \bar{L}^{-1} \left(N_1(t, \sum_{n=0}^{\infty} \star x_{1n}(t, \beta), \right. \\
 &\quad \left. \sum_{n=0}^{\infty} \star y_{1n}(t, \beta) \right) + \bar{L}^{-1} \left(\star g_1(t, \beta) \right), \\
 \sum_{n=0}^{\infty} \star y_{1n}(t, \beta) &= \Phi_1(t, \beta) - \bar{L}^{-1} \left(\bar{R} \sum_{n=0}^{\infty} \star x_{1n}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \sum_{n=0}^{\infty} \star y_{1n}(t, \beta) \right) - \bar{L}^{-1} \left(N_1(t, \sum_{n=0}^{\infty} \star x_{1n}(t, \beta), \right. \\
 &\quad \left. \sum_{n=0}^{\infty} \star y_{1n}(t, \beta) \right) + \bar{L}^{-1} \left(\star h_1(t, \beta) \right), \\
 \sum_{n=0}^{\infty} \star x_{2n}(t, \beta) &= \Psi_2(t, \beta) - \bar{L}^{-1} \left(\bar{R} \sum_{n=0}^{\infty} \star x_{2n}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \sum_{n=0}^{\infty} \star y_{2n}(t, \beta) \right) - \bar{L}^{-1} \left(N_2(t, \sum_{n=0}^{\infty} \star x_{2n}(t, \beta), \right. \\
 &\quad \left. \sum_{n=0}^{\infty} \star y_{2n}(t, \beta) \right) + \bar{L}^{-1} \left(\star g_2(t, \beta) \right), \\
 \sum_{n=0}^{\infty} \star y_{2n}(t, \beta) &= \Phi_2(t, \beta) - \bar{L}^{-1} \left(\bar{R} \sum_{n=0}^{\infty} \star x_{2n}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \sum_{n=0}^{\infty} \star y_{2n}(t, \beta) \right) - \bar{L}^{-1} \left(N_1(t, \sum_{n=0}^{\infty} \star x_{2n}(t, \beta), \right. \\
 &\quad \left. \sum_{n=0}^{\infty} \star y_{2n}(t, \beta) \right) + \bar{L}^{-1} \left(\star h_2(t, \beta) \right).
 \end{aligned} \right\} (17)$$

$$\left. \begin{aligned}
 \star x_{20}(t, \alpha) &= \Psi_2(t, \alpha), \\
 \star y_{20}(t, \alpha) &= \Phi_2(t, \alpha), \\
 \star x_{21}(t, \alpha) &= \bar{L}^{-1} \left(\star g_2(t, \alpha) \right) - \bar{L}^{-1} \left(\bar{R} \star x_{20}(t, \alpha) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \star y_{20}(t, \alpha) \right) - \bar{L}^{-1} \left(N_2(t, \star x_{20}(t, \alpha), \star y_{20}(t, \alpha)) \right), \\
 \star y_{21}(t, \alpha) &= \bar{L}^{-1} \left(\star h_2(t, \alpha) \right) - \bar{L}^{-1} \left(\bar{R} \star x_{20}(t, \alpha) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \star y_{20}(t, \alpha) \right) - \bar{L}^{-1} \left(N_2(t, \star x_{20}(t, \alpha), \star y_{20}(t, \alpha)) \right), \\
 \star x_{2k+1}(t, \alpha) &= -\bar{L}^{-1} \left(\bar{R} \star x_{2k}(t, \alpha) \right) - \bar{L}^{-1} \left(\bar{R} \star y_{2k}(t, \alpha) \right) \\
 -\bar{L}^{-1} \left(N_2(t, \star x_{2k}(t, \alpha), \star y_{2k}(t, \alpha)) \right), k \geq 1, \\
 \star y_{2k+1}(t, \alpha) &= -\bar{L}^{-1} \left(\bar{R} \star x_{2k}(t, \alpha) \right) - \bar{L}^{-1} \left(\bar{R} \star y_{2k}(t, \alpha) \right) \\
 -\bar{L}^{-1} \left(N_2(t, \star x_{2k}(t, \alpha), \star y_{2k}(t, \alpha)) \right), k \geq 1, \\
 \star x_{10}(t, \beta) &= \Psi_1(t, \beta), \\
 \star y_{10}(t, \beta) &= \Phi_1(t, \beta), \\
 \star x_{11}(t, \beta) &= \bar{L}^{-1} \left(\star g_1(t, \beta) \right) - \bar{L}^{-1} \left(\bar{R} \star x_{10}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \star y_{10}(t, \beta) \right) - \bar{L}^{-1} \left(N_1(t, \star x_{10}(t, \beta), \star y_{10}(t, \beta)) \right), \\
 \star y_{11}(t, \beta) &= \bar{L}^{-1} \left(\star h_1(t, \beta) \right) - \bar{L}^{-1} \left(\bar{R} \star x_{10}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \star y_{10}(t, \beta) \right) - \bar{L}^{-1} \left(N_1(t, \star x_{10}(t, \beta), \star y_{10}(t, \beta)) \right), \\
 \star x_{1k+1}(t, \beta) &= -\bar{L}^{-1} \left(\bar{R} \star x_{1k}(t, \beta) \right) - \bar{L}^{-1} \left(\bar{R} \star y_{1k}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(N_1(t, \star x_{1k}(t, \beta), \star y_{1k}(t, \beta)) \right), k \geq 1, \\
 \star y_{1k+1}(t, \beta) &= -\bar{L}^{-1} \left(\bar{R} \star x_{1k}(t, \beta) \right) - \bar{L}^{-1} \left(\bar{R} \star y_{1k}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(N_1(t, \star x_{1k}(t, \beta), \star y_{1k}(t, \beta)) \right), k \geq 1,
 \end{aligned} \right\} (20)$$

According to the GMADM the recursive relation for the (15), (16),(17) and (18), is as follows:

$$\left. \begin{aligned}
 \star x_{10}(t, \alpha) &= \Psi_1(t, \alpha), \\
 \star y_{10}(t, \alpha) &= \Phi_1(t, \alpha), \\
 \star x_{11}(t, \alpha) &= \bar{L}^{-1} \left(\star g_1(t, \alpha) \right) - \bar{L}^{-1} \left(\bar{R} \star x_{10}(t, \alpha) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \star y_{10}(t, \alpha) \right) - \bar{L}^{-1} \left(N_1(t, \star x_{10}(t, \alpha), \star y_{10}(t, \alpha)) \right), \\
 \star y_{11}(t, \alpha) &= \bar{L}^{-1} \left(\star h_1(t, \alpha) \right) - \bar{L}^{-1} \left(\bar{R} \star x_{10}(t, \alpha) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \star y_{10}(t, \alpha) \right) - \bar{L}^{-1} \left(N_1(t, \star x_{10}(t, \alpha), \star y_{10}(t, \alpha)) \right), \\
 \star x_{1k+1}(t, \alpha) &= -\bar{L}^{-1} \left(\bar{R} \star x_{1k}(t, \alpha) \right) - \bar{L}^{-1} \left(\bar{R} \star y_{1k}(t, \alpha) \right) \\
 -\bar{L}^{-1} \left(N_1(t, \star x_{1k}(t, \alpha), \star y_{1k}(t, \alpha)) \right), k \geq 1, \\
 \star y_{1k+1}(t, \alpha) &= -\bar{L}^{-1} \left(\bar{R} \star x_{1k}(t, \alpha) \right) - \bar{L}^{-1} \left(\bar{R} \star y_{1k}(t, \alpha) \right) \\
 -\bar{L}^{-1} \left(N_1(t, \star x_{1k}(t, \alpha), \star y_{1k}(t, \alpha)) \right), k \geq 1, \\
 \star x_{20}(t, \beta) &= \Psi_2(t, \beta), \\
 \star y_{20}(t, \beta) &= \Phi_2(t, \beta), \\
 \star x_{21}(t, \beta) &= \bar{L}^{-1} \left(\star g_2(t, \beta) \right) - \bar{L}^{-1} \left(\bar{R} \star x_{20}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \star y_{20}(t, \beta) \right) - \bar{L}^{-1} \left(N_2(t, \star x_{20}(t, \beta), \star y_{20}(t, \beta)) \right), \\
 \star y_{21}(t, \beta) &= \bar{L}^{-1} \left(\star h_2(t, \beta) \right) - \bar{L}^{-1} \left(\bar{R} \star x_{20}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(\bar{R} \star y_{20}(t, \beta) \right) - \bar{L}^{-1} \left(N_2(t, \star x_{20}(t, \beta), \star y_{20}(t, \beta)) \right), \\
 \star x_{2k+1}(t, \beta) &= -\bar{L}^{-1} \left(\bar{R} \star x_{2k}(t, \beta) \right) - \bar{L}^{-1} \left(\bar{R} \star y_{2k}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(N_2(t, \star x_{2k}(t, \beta), \star y_{2k}(t, \beta)) \right), k \geq 1, \\
 \star y_{2k+1}(t, \beta) &= -\bar{L}^{-1} \left(\bar{R} \star x_{2k}(t, \beta) \right) - \bar{L}^{-1} \left(\bar{R} \star y_{2k}(t, \beta) \right) \\
 -\bar{L}^{-1} \left(N_2(t, \star x_{2k}(t, \beta), \star y_{2k}(t, \beta)) \right), k \geq 1.
 \end{aligned} \right\} (19)$$

The n th term approximation to the solution is defined as follows:

$$\left. \begin{aligned} \varphi_{1n}(t, \alpha) &= \sum_{i=0}^{n-1} \overset{*}{x}_{1i}(t, \alpha), \\ \phi_{1n}(t, \alpha) &= \sum_{i=0}^{n-1} \overset{*}{y}_{1i}(t, \alpha), \\ \varphi_{2n}(t, \alpha) &= \sum_{i=0}^{n-1} \overset{*}{x}_{2i}(t, \alpha), \\ \phi_{2n}(t, \alpha) &= \sum_{i=0}^{n-1} \overset{*}{y}_{2i}(t, \alpha), \\ \varphi_{1n}(t, \beta) &= \sum_{i=0}^{n-1} \overset{*}{x}_{1i}(t, \beta), \\ \phi_{1n}(t, \beta) &= \sum_{i=0}^{n-1} \overset{*}{y}_{1i}(t, \beta), \\ \varphi_{2n}(t, \beta) &= \sum_{i=0}^{n-1} \overset{*}{x}_{2i}(t, \beta), \\ \phi_{1n}(t, \beta) &= \sum_{i=0}^{n-1} \overset{*}{y}_{1i}(t, \beta). \end{aligned} \right\}$$

Hence,

$$\left\{ \begin{aligned} \left\{ \lim_{n \rightarrow \infty} (\varphi_{1n}(t, \alpha)), \lim_{n \rightarrow \infty} \phi_{1n}(t, \alpha) \right\} &= \{\overset{*}{x}_1(t, \alpha), \overset{*}{y}_1(t, \alpha)\}, \\ \left\{ \lim_{n \rightarrow \infty} (\varphi_{2n}(t, \alpha)), \lim_{n \rightarrow \infty} \phi_{2n}(t, \alpha) \right\} &= \{\overset{*}{x}_2(t, \alpha), \overset{*}{y}_2(t, \alpha)\}, \\ \left\{ \lim_{n \rightarrow \infty} (\varphi_{1n}(t, \beta)), \lim_{n \rightarrow \infty} \phi_{1n}(t, \beta) \right\} &= \{\overset{*}{x}_1(t, \beta), \overset{*}{y}_1(t, \beta)\}, \\ \left\{ \lim_{n \rightarrow \infty} (\varphi_{2n}(t, \beta)), \lim_{n \rightarrow \infty} \phi_{2n}(t, \beta) \right\} &= \{\overset{*}{x}_2(t, \beta), \overset{*}{y}_2(t, \beta)\}. \end{aligned} \right.$$

Raza et al. [39], Ray [40], Zo'bi et al. [41], and many others discuss the convergence of the ADM or Decomposition method and GMADM.

A. STABILITY OF GMADM

When the solution produced by a technique is unaffected by small changes in the inputs and parameters and when it is expected that changes in the parameters carried on by impacts in equations and conditions, the method is said to be stable. By giving examples and analyzing the stability of the GMADM in this study, we suggested contrasting the GMADM with other existing methods i.e., ADM and Taylor series method (TSM).

B. APPLICATIONS

Example 1 Brine Tanks Problem [42], [43]: Two tanks A and B are connected by pipes. Tank A contains 100 gal of brine and tank B contains 200 gal of brine. Through one pipe solution is pumped from first tank to the second at 30 gal/min. Through the other solution is pumped at the rate of 10 gal/min from the second tank to the first, and the brine in tank B flows out at 20 gal/min. If $\overset{*}{x}(t)$ and $\overset{*}{y}(t)$ denotes the amount of salt in tanks A and B respectively, then what will be the salt content in each tank at any time t . It is to be noted that at time $t = 0$

there is $\langle(98, 99, 100, 101, 0.7); (97, 99, 100, 102; 0.2)\rangle$ lb salt in tank A and $\langle(50, 51, 52, 53; 0.7); (49, 51, 52, 54; 0.2)\rangle$ lb in tank B.

The system of intuitionistic fuzzy differential equation related to above problem is as follows:

$$\left. \begin{aligned} \frac{d\overset{*}{x}(t, \alpha)}{dt} &= \frac{\overset{*}{y}}{20} - \frac{3\overset{*}{x}}{10}, \\ \frac{d\overset{*}{y}(t, \alpha)}{dt} &= \frac{3\overset{*}{x}}{10} - \frac{3\overset{*}{y}}{20}. \end{aligned} \right\} \tag{23}$$

with initial conditions,

$$\left\{ \begin{aligned} \overset{*}{x}(0) &= \langle(98, 99, 100, 101, 0.7); (97, 99, 100, 102; 0.2)\rangle, \\ \overset{*}{y}(0) &= \langle(50, 51, 52, 53; 0.7); (49, 51, 52, 54; 0.2)\rangle. \end{aligned} \right. \tag{24}$$

By taking (α, β) -cut of (24) and (24), we obtain the following equations:

$$\left\{ \begin{aligned} \frac{d\overset{*}{x}(t, \alpha)}{dt} &= \frac{\overset{*}{y}}{20} - \frac{3\overset{*}{x}}{10}, \overset{*}{x}_1(0, \alpha) = 1.429\alpha + 98, \\ \frac{d\overset{*}{y}(t, \alpha)}{dt} &= \frac{3\overset{*}{x}}{10} - \frac{3\overset{*}{y}}{20}, \overset{*}{y}_1(0, \alpha) = 1.429\alpha + 50, \end{aligned} \right. \tag{25}$$

$$\left\{ \begin{aligned} \frac{d\overset{*}{x}(t, \alpha)}{dt} &= \frac{\overset{*}{y}}{20} - \frac{3\overset{*}{x}}{10}, \overset{*}{x}_2(0, \alpha) = -1.429\alpha + 101, \\ \frac{d\overset{*}{y}(t, \alpha)}{dt} &= \frac{3\overset{*}{x}}{10} - \frac{3\overset{*}{y}}{20}, \overset{*}{y}_2(0, \alpha) = -1.429\alpha + 53, \end{aligned} \right. \tag{26}$$

$$\left\{ \begin{aligned} \frac{d\overset{*}{x}(t, \beta)}{dt} &= \frac{\overset{*}{y}}{20} - \frac{3\overset{*}{x}}{10}, \overset{*}{x}_1(0, \beta) = -2.5\beta + 99, \\ \frac{d\overset{*}{y}(t, \beta)}{dt} &= \frac{3\overset{*}{x}}{10} - \frac{3\overset{*}{y}}{20}, \overset{*}{y}_1(0, \beta) = -2.5\beta + 51, \end{aligned} \right. \tag{27}$$

$$\left\{ \begin{aligned} \frac{d\overset{*}{x}(t, \beta)}{dt} &= \frac{\overset{*}{y}}{20} - \frac{\overset{*}{x}}{20}, \overset{*}{x}_2(0, \beta) = -2.5\beta + 99.5, \\ \frac{d\overset{*}{y}(t, \beta)}{dt} &= \frac{\overset{*}{x}}{20} - \frac{\overset{*}{y}}{20}, \overset{*}{y}_2(0, \beta) = 2.5\beta - 51.5. \end{aligned} \right. \tag{28}$$

Here $\overset{*}{L} = \frac{d}{dt}$ and by taking $\overset{*}{L}^{-1} (\cdot) = \int_0^t (\cdot) dt$ on both sides of (25), (26), (27) and (28), and using the initial conditions we obtain;

$$\left\{ \begin{aligned} \overset{*}{x}_1(t, \alpha) &= \int_0^t (-0.30\overset{*}{x}_1(u, \alpha) + 0.05\overset{*}{y}_1(u, \alpha))du \\ &\quad + 1.429\alpha + 98, \\ \overset{*}{y}_1(t, \alpha) &= \int_0^t (-0.150\overset{*}{y}_1(u, \alpha) + 0.30\overset{*}{x}_1(u, \alpha))du \\ &\quad + 1.429\alpha + 50, \end{aligned} \right. \tag{29}$$

$$\left\{ \begin{aligned} \overset{*}{x}_2(t, \alpha) &= \int_0^t (-0.30\overset{*}{x}_2(u, \alpha) + 0.05\overset{*}{y}_2(u, \alpha))du \\ &\quad - 1.429\alpha + 101, \\ \overset{*}{y}_2(t, \alpha) &= \int_0^t (-0.15\overset{*}{y}_2(u, \alpha) + 0.30\overset{*}{x}_2(u, \alpha))du \\ &\quad - 1.429\alpha + 53, \end{aligned} \right. \tag{30}$$

$$\left\{ \begin{aligned} \overset{*}{x}_1(t, \beta) &= \int_0^t (-0.30\overset{*}{x}_1(u, \beta) + 0.05\overset{*}{y}_1(u, \beta))du \\ &\quad + 99 - 2.5\beta, \\ \overset{*}{y}_1(t, \beta) &= \int_0^t (-0.15\overset{*}{y}_1(u, \beta) + 0.30\overset{*}{x}_1(u, \beta))du \\ &\quad + 51 - 2.5\beta, \end{aligned} \right. \tag{31}$$

$$\left\{ \begin{aligned} \overset{*}{x}_2(t, \beta) &= \int_0^t (-0.05\overset{*}{x}_2(u, \beta) + 0.05\overset{*}{y}_2(u, \beta))du \\ &\quad + 2.5\beta + 99.5, \\ \overset{*}{y}_2(t, \beta) &= \int_0^t (-0.05\overset{*}{y}_2(u, \beta) + 0.05\overset{*}{x}_2(u, \beta))du \\ &\quad + 0.25\beta + 0.65. \end{aligned} \right. \tag{32}$$

Now by using GMADM the solution of (29), (30), (31) and (32), can be expressed as;

$$\left\{ \begin{array}{l} \begin{array}{l} \overset{*}{x}_{1_0}(t, \alpha) = 1.429\alpha + 98, \\ \overset{*}{y}_{1_0}(t, \alpha) = 1.429\alpha + 50, \\ \overset{*}{x}_{1_1}(t, \alpha) = -0.357250\alpha t - 26.90t, \\ \overset{*}{y}_{1_1}(t, \alpha) = 0.214350\alpha t + 21.90t, \\ \overset{*}{x}_{1_{k+1}}(t, \alpha) = \int_0^t (-0.30\overset{*}{x}_{1_k}(u, \alpha) \\ + 0.05\overset{*}{y}_{1_k}(u, \alpha))du, k \geq 1, \\ \overset{*}{y}_{1_{k+1}}(t, \alpha) = \int_0^t (-0.15\overset{*}{y}_{1_k}(u, \alpha) \\ + 0.30\overset{*}{x}_{1_k}(u, \alpha))du, k \geq 1. \end{array} \end{array} \right. \quad (33)$$

$$\left\{ \begin{array}{l} \begin{array}{l} \overset{*}{x}_{2_0}(t, \alpha) = -1.429\alpha + 101, \\ \overset{*}{y}_{2_0}(t, \alpha) = -1.429\alpha + 50, \\ \overset{*}{x}_{2_1}(t, \alpha) = 0.357250\alpha t - 27.650t, \\ \overset{*}{y}_{2_1}(t, \alpha) = -0.214350\alpha t + 22.350t, \\ \overset{*}{x}_{2_{k+1}}(t, \alpha) = \int_0^t (-0.30\overset{*}{x}_{2_k}(u, \alpha) \\ + 0.05\overset{*}{y}_{2_k}(u, \alpha))du, k \geq 1, \\ \overset{*}{y}_{2_{k+1}}(t, \alpha) = \int_0^t (-0.15\overset{*}{y}_{2_k}(u, \alpha) \\ + 0.30\overset{*}{x}_{2_k}(u, \alpha))du, k \geq 1. \end{array} \end{array} \right. \quad (34)$$

$$\left\{ \begin{array}{l} \begin{array}{l} \overset{*}{x}_{1_0}(t, \beta) = 99 - 2.5\beta, \\ \overset{*}{y}_{1_0}(t, \beta) = 51 - 2.5\beta, \\ \overset{*}{x}_{1_1}(t, \beta) = -27.150t + 0.6250\beta t, \\ \overset{*}{y}_{1_1}(t, \beta) = 22.050t - 0.3750\beta t, \\ \overset{*}{x}_{1_{k+1}}(t, \beta) = \int_0^t (-0.30\overset{*}{x}_{1_k}(u, \beta) \\ + 0.05\overset{*}{y}_{1_k}(u, \beta))du, k \geq 1, \\ \overset{*}{y}_{1_{k+1}}(t, \beta) = \int_0^t (-0.15\overset{*}{y}_{1_k}(u, \beta) \\ + 0.30\overset{*}{x}_{1_k}(u, \beta))du, k \geq 1. \end{array} \end{array} \right. \quad (35)$$

$$\left\{ \begin{array}{l} \begin{array}{l} \overset{*}{x}_{2_0}(t, \beta) = 2.5\beta + 99.5, \\ \overset{*}{y}_{2_0}(t, \beta) = 2.5\beta + 51.5, \\ \overset{*}{x}_{2_1}(t, \beta) = -0.6250\beta t - 27.2750t, \\ \overset{*}{y}_{2_1}(t, \beta) = 0.3750\beta t + 22.1250t, \\ \overset{*}{x}_{2_{k+1}}(t, \beta) = \int_0^t (-0.30\overset{*}{x}_{2_k}(u, \beta) \\ + 0.05\overset{*}{y}_{2_k}(u, \beta))du, k \geq 1, \\ \overset{*}{y}_{2_{k+1}}(t, \beta) = \int_0^t (-0.15\overset{*}{y}_{2_k}(u, \beta) \\ + 0.30\overset{*}{x}_{2_k}(u, \beta))du, k \geq 1. \end{array} \end{array} \right. \quad (36)$$

By solving (33), (34), (35) and (36), we get the approximate solution after four iterations as follows:

$$\begin{aligned} & (\overset{*}{x}_1(t, \alpha), \overset{*}{y}_1(t, \alpha)) \\ &= (1.429\alpha + 98 - 0.357250\alpha t - 26.90t \\ &+ 0.0589462500t^2\alpha + 4.582500000t^2 \\ &- 0.007055687500\alpha t^3 - 0.5528750000t^3 \\ &+ 0.0006463992188\alpha t^4 + 0.05074218750t^4, \\ &1.429\alpha + 50 + 0.2143500000\alpha t + 21.90000000t \\ &- 0.06966375000\alpha t^2 - 5.677500000t^2 \end{aligned}$$

$$\begin{aligned} & + 0.009377812500\alpha t^3 + 0.7421250000t^3 \\ &- 0.0008808445312\alpha t^4 - 0.06929531250t^4), \\ & (\overset{*}{x}_2(t, \alpha), \overset{*}{y}_2(t, \alpha)) \\ &= (-1.429\alpha + 101 + 0.357250\alpha t - 27.650t \\ &- 0.05894625000\alpha t^2 + 4.706250000t^2 \\ &+ 0.007055687499\alpha t^3 - 0.5676874999t^3 \\ &- 0.0006463992188\alpha t^4 + 0.05209921875t^4, \\ &- 1.429\alpha + 53 - 0.2143500000\alpha t + 22.35000000t \\ &+ 0.06966375000\alpha t^2 - 5.823750000t^2 \\ &- 0.009377812500\alpha t^3 + 0.7618125000t^3 \\ &+ 0.0008808445312\alpha t^4 - 0.07114453125t^4), \\ & (\overset{*}{x}_1(t, \beta), \overset{*}{y}_1(t, \beta)) \\ &= (99 - 2.5\beta - 27.150t + 0.6250\beta t \\ &+ 4.623750t^2 - 0.1031250000\beta t^2 \\ &- 0.5578125000t^3 + 0.01234375000\beta t^3 \\ &+ 0.05119453125t^4 - 0.001130859375\beta t^4 \\ &, 51 - 2.5\beta + 22.050t - 0.3750\beta t \\ &- 5.726250000t^2 + 0.1218750000\beta t^2 \\ &+ 0.7486875000t^3 - 0.01640625000\beta t^3 \\ &- 0.06991171875t^4 + 0.001541015625\beta t^4), \\ & (\overset{*}{x}_2(t, \beta), \overset{*}{y}_2(t, \beta)) \\ &= (2.5\beta + 99.5 - 0.6250\beta t - 27.2750t \\ &+ 0.1031250\beta t^2 + 4.644375000t^2 \\ &- 0.01234375000\beta t^3 - 0.5602812500t^3 \\ &+ 0.001130859375\beta t^4 + 0.05142070312t^4, \\ &2.5\beta + 51.5 + 0.3750\beta t + 22.1250t \\ &- 0.1218750000\beta t^2 - 5.750625000t^2 \\ &+ 0.01640625000\beta t^3 + 0.7519687500t^3 \\ &- 0.001541015625\beta t^4 - 0.07021992188t^4). \end{aligned}$$

In Table 1, $\overset{*}{x}_1(t, \alpha)$, $\overset{*}{x}_2(t, \alpha)$, $\overset{*}{y}_1(t, \alpha)$ and $\overset{*}{y}_2(t, \alpha)$ represents approximate solution of the membership functions of the Example 1 for $\alpha \in [0, 0.7]$ and $\overset{*}{x}_1(t, \beta)$, $\overset{*}{x}_2(t, \beta)$, $\overset{*}{y}_1(t, \beta)$ and $\overset{*}{y}_2(t, \beta)$ represents approximate solution of non-membership function of the Example 1 for $\beta \in [0.2, 1.0]$.

The following is the mathematical and exact solution to Example 1 using the classical method, as shown in Figure 2(a,b) and Table 2:

$$\begin{aligned} & (\overset{*}{x}_1(t, \alpha), \overset{*}{y}_1(t, \alpha)) \\ &= \left(-\frac{1}{34000}(-17 + \sqrt{17})(92000 + 1429\alpha \right. \\ &- 6000\sqrt{17})e^{\frac{1}{40}(-9 + \sqrt{17})t} \\ &+ \frac{1}{34000}(17 + \sqrt{17})(92000 + 1429\alpha \\ &+ 6000\sqrt{17})e^{-\frac{1}{40}(9 + \sqrt{17})t}, \end{aligned}$$

TABLE 1. Illustrates the approximation to the generalized trapezoidal intuitionistic fuzzy initial value problem from Example 1 for t=1.

Table 1: Approximate solution of example1 at t = 1.									
α	$\tilde{x}_1^*(t, \alpha)$	$\tilde{y}_1^*(t, \alpha)$	$\tilde{x}_2^*(t, \alpha)$	$\tilde{y}_2^*(t, \alpha)$	β	$\tilde{x}_1^*(t, \beta)$	$\tilde{y}_1^*(t, \beta)$	$\tilde{x}_2^*(t, \beta)$	$\tilde{y}_2^*(t, \beta)$
0	75.18	57.90	77.54	61.06	0.2	75.57	58.42	76.75	60.00
0.1	75.29	58.05	77.43	60.90	0.3	75.38	58.16	76.95	60.27
0.2	75.41	58.20	77.32	60.75	0.4	75.18	57.90	77.15	60.53
0.3	75.52	58.35	77.20	60.60	0.5	74.98	57.63	77.34	60.79
0.4	75.63	58.50	77.09	60.45	0.6	74.79	57.37	77.54	61.06
0.5	75.74	58.65	76.98	60.30	0.7	74.59	57.11	77.73	61.32
0.6	75.85	58.80	76.87	60.15	0.8	74.39	56.84	77.93	61.58
0.7	75.97	58.95	76.75	60.00	0.9	74.20	56.58	78.13	61.84
					1.0	74.00	56.32	78.33	62.11

TABLE 2. Illustrates the exact solution to the generalized trapezoidal intuitionistic fuzzy initial value problem from Example 3 for t=1.

Table 2: Exact solution of example1 at t = 1.									
α	$\tilde{x}_1(t, \alpha)$	$\tilde{y}_1(t, \alpha)$	$\tilde{x}_2(t, \alpha)$	$\tilde{y}_2(t, \alpha)$	β	$\tilde{x}_1(t, \beta)$	$\tilde{y}_1(t, \beta)$	$\tilde{x}_2(t, \beta)$	$\tilde{y}_2(t, \beta)$
0	74.78	51.00	77.52	70.35	0.2	75.57	67.39	76.75	69.03
0.1	74.89	51.15	77.41	70.13	0.3	75.38	67.13	76.95	69.38
0.2	75.00	51.21	77.32	69.84	0.4	75.19	66.87	77.14	69.64
0.3	75.14	51.43	77.19	69.79	0.5	74.99	66.62	77.34	69.90
0.4	75.22	51.54	77.09	69.68	0.6	74.79	66.36	77.74	70.47
0.5	75.32	51.69	76.95	69.35	0.7	74.59	66.06	77.93	70.73
0.6	75.45	51.80	76.87	69.36	0.8	74.39	65.79	78.13	70.99
0.7	75.56	51.96	76.72	69.10	0.9	74.20	65.44	78.13	70.99
					1.0	73.99	65.12	78.32	71.24

TABLE 3. Illustrates the approximate solution iterations in column 2, residual error in column 3, and CPU time required by the numerical technique GMADM to determine the approximate solution of the system of generalized fuzzy intuitionistic differential equations used in Example 1 in column 4. Whenever the error of all methods is taken into account, we can conclude that the GMADM has a better convergence behavior and is more stable than the ADM and TSM, respectively.

Table 3: Error comparison			
Paramater	Iterations	Error	CPU-time
GMADM	4	0.1E-15	1.12451
ADM	8	5.8E-08	2.14510
TSM	7	3.1E-05	2.10325

$$\begin{aligned}
 & - \frac{1}{68000}(-17 + \sqrt{17})(92000 + 1429\alpha) \\
 & - 6000\sqrt{17}e^{\frac{1}{40}(-9+\sqrt{17})t} \\
 & \times \sqrt{17} - \frac{1}{68000}(17 + \sqrt{17})(92000 \\
 & + 1429\alpha + 6000\sqrt{17})e^{-\frac{1}{40}(9+\sqrt{17})t} \\
 & \times \sqrt{17} - \frac{3}{68000}(-17 + \sqrt{17})(92000 + 1429\alpha \\
 & - 6000\sqrt{17})e^{\frac{1}{40}(-9+\sqrt{17})t} \\
 & + \frac{3}{68000}(17 + \sqrt{17})(92000 + 1429\alpha \\
 & + 6000\sqrt{17})e^{-\frac{1}{40}(9+\sqrt{17})t}), \\
 (\tilde{x}_2(t, \alpha), \tilde{y}_2(t, \alpha)) \\
 & = \left(\frac{1}{66000}(-33 + \sqrt{33})(-98000 + 1429\alpha \right. \\
 & \left. + 3000\sqrt{33})e^{\frac{1}{40}(-9+\sqrt{33})t} \right. \\
 & \left. - \frac{1}{66000}(33 + \sqrt{33})(-98000 + 1429\alpha \right. \\
 & \left. - 3000\sqrt{33})e^{-\frac{1}{40}(9+\sqrt{33})t}, \right. \\
 & \left. \frac{1}{132000}(-33 + \sqrt{33})(-98000 + 1429\alpha \right. \\
 & \left. + 3000\sqrt{33})e^{\frac{1}{40}(-9+\sqrt{33})t} \sqrt{33} \right.
 \end{aligned}$$

$$\begin{aligned}
 & \left. + \frac{1}{132000}(33 + \sqrt{33})(-98000 + 1429\alpha \right. \\
 & \left. - 3000\sqrt{33})e^{-\frac{1}{40}(9+\sqrt{33})t} \sqrt{33} \right. \\
 & \left. + \frac{1}{44000}(-33 + \sqrt{33})(-98000 + 1429\alpha \right. \\
 & \left. + 3000\sqrt{33})e^{\frac{1}{40}(-9+\sqrt{33})t} \right. \\
 & \left. - \frac{1}{44000}(33 + \sqrt{33})(-98000 + 1429\alpha \right. \\
 & \left. - 3000\sqrt{33})e^{-\frac{1}{40}(9+\sqrt{33})t}), \\
 (\tilde{x}_1(t, \beta), \tilde{y}_1(t, \beta)) \\
 & = \left(\frac{1}{132}(-33 + \sqrt{33})(-192 + 5\beta + 6\sqrt{33})e^{\frac{1}{40}(-9+\sqrt{33})t} \right. \\
 & \left. - \frac{1}{132}(33 + \sqrt{33})(-192 + 5\beta - 6\sqrt{33})e^{-\frac{1}{40}(9+\sqrt{33})t}, \right. \\
 & \left. \frac{1}{264}(-33 + \sqrt{33})(-192 + 5\beta + 6\sqrt{33})e^{\frac{1}{40}(-9+\sqrt{33})t} \sqrt{33} \right. \\
 & \left. + \frac{1}{264}(33 + \sqrt{33})(-192 + 5\beta - 6\sqrt{33})e^{-\frac{1}{40}(9+\sqrt{33})t} \sqrt{33} \right. \\
 & \left. + \frac{1}{88}(-33 + \sqrt{33})(-192 + 5\beta + 6\sqrt{33})e^{\frac{1}{40}(-9+\sqrt{33})t} \sqrt{33} \right. \\
 & \left. - \frac{1}{88}(33 + \sqrt{33})(-192 + 5\beta - 6\sqrt{33})e^{-\frac{1}{40}(9+\sqrt{33})t}), \right. \\
 (\tilde{x}_2(t, \beta), \tilde{y}_2(t, \beta)) \\
 & = \left(-\frac{1}{132}(-33 + \sqrt{33})(193 + 5\beta - 6\sqrt{33})e^{\frac{1}{40}(-9+\sqrt{33})t} \right. \\
 & \left. + \frac{1}{132}(33 + \sqrt{33})(193 + 5\beta + 6\sqrt{33})e^{-\frac{1}{40}(9+\sqrt{33})t}, \right. \\
 & \left. - \frac{1}{264}(-33 + \sqrt{33})(193 + 5\beta - 6\sqrt{33})e^{\frac{1}{40}(-9+\sqrt{33})t} \sqrt{33} \right. \\
 & \left. - \frac{1}{264}(33 + \sqrt{33})(193 + 5\beta + 6\sqrt{33})e^{-\frac{1}{40}(9+\sqrt{33})t} \sqrt{33} \right. \\
 & \left. - \frac{1}{88}(-33 + \sqrt{33})(193 + 5\beta - 6\sqrt{33})e^{\frac{1}{40}(-9+\sqrt{33})t} \right. \\
 & \left. + \frac{1}{88}(33 + \sqrt{33})(193 + 5\beta + 6\sqrt{33})e^{-\frac{1}{40}(9+\sqrt{33})t}), \right.
 \end{aligned}$$

Table 2, $\overset{\star}{x}_1(t, \alpha)$, $\overset{\star}{x}_2(t, \alpha)$, $\overset{\star}{y}_1(t, \alpha)$ and $\overset{\star}{y}_2(t, \alpha)$ represents exact solution of the membership functions of the Example1 for $\alpha \in [0, 0.7]$ and $\overset{\star}{x}_1(t, \beta)$, $\overset{\star}{x}_2(t, \beta)$, $\overset{\star}{y}_1(t, \beta)$ and $\overset{\star}{y}_2(t, \beta)$ represents exact solution of non-membership function of the Example1 for $\beta \in [0.2, 1.0]$.

Example 2 Coupled Oscillators: Consider a mechanical system [44], [45], [46], [47] constituting of two masses $m_1 = 1Kg$ and $m_2 = 1Kg$ that are free to slide over a friction less horizontal surface. The masses are attached to one another, and to two rigid walls, with the help of three springs. The spring constants for this system are $k_1 = 1Nm^{-1}$, $k_2 = 2Nm^{-1}$ and $k_3 = 1Nm^{-1}$. The instantaneous state of the system is conveniently specified by the $\overset{\star}{x}(t)$ and $\overset{\star}{y}(t)$ respectively. Thus, the equations of motions of two masses are as follows:

$$\begin{cases} m \frac{d^2 \overset{\star}{x}(t)}{dt^2} = -k_1 \overset{\star}{x} - k_2 (\overset{\star}{x} - \overset{\star}{y}), \\ m \frac{d^2 \overset{\star}{y}(t)}{dt^2} = -k_3 \overset{\star}{y} - k_2 (\overset{\star}{y} - \overset{\star}{x}). \end{cases} \quad (37)$$

Using the given data, we get;

$$\begin{cases} \frac{d^2 \overset{\star}{x}(t)}{dt^2} = 2\overset{\star}{y} - 3\overset{\star}{x}, \\ \frac{d^2 \overset{\star}{y}(t)}{dt^2} = 2\overset{\star}{x} - 3\overset{\star}{y}, \end{cases} \quad (38)$$

with initial conditions,

$$\begin{cases} \overset{\star}{x}(0) = \langle (2, 4, 8, 15; 0.6); (1, 4, 8, 18; 0.3) \rangle, \\ \overset{\star}{x}'(0) = \langle (2, 5, 8, 10; 0.6); (1, 5, 8, 12; 0.3) \rangle, \\ \overset{\star}{y}(0) = \langle (8, 9, 10, 11; 0.6); (7, 9, 10, 12; 0.3) \rangle, \\ \overset{\star}{y}'(0) = \langle (11, 12, 13, 14; 0.6); (10, 12, 13, 15; 0.3) \rangle. \end{cases} \quad (39)$$

By taking (α, β) -cut of (38), (39), we get the following equations:

$$\begin{cases} \frac{d^2 \overset{\star}{x}(t, \alpha)}{dt^2} = 2\overset{\star}{y} - 3\overset{\star}{x}, \\ \overset{\star}{x}_1(0, \alpha) = 3.33\alpha + 2, \overset{\star}{x}'_1(0, \alpha) = 5\alpha + 2, \\ \frac{d^2 \overset{\star}{y}(t, \alpha)}{dt^2} = 2\overset{\star}{x} - 3\overset{\star}{y}, \\ \overset{\star}{y}_1(0, \alpha) = 1.67\alpha + 8, \overset{\star}{y}'_1(0, \alpha) = 1.67\alpha + 11, \end{cases} \quad (40)$$

$$\begin{cases} \frac{d^2 \overset{\star}{x}(t, \alpha)}{dt^2} = 2\overset{\star}{y} - 3\overset{\star}{x}, \\ \overset{\star}{x}_2(0, \alpha) = 15 - 11.67\alpha, \overset{\star}{x}'_2(0, \alpha) = 10 - 3.33\alpha, \\ \frac{d^2 \overset{\star}{y}(t, \alpha)}{dt^2} = 2\overset{\star}{x} - 3\overset{\star}{y}, \\ \overset{\star}{y}_2(0, \alpha) = -1.67\alpha + 11, \overset{\star}{y}'_2(0, \alpha) = -1.67\alpha + 14, \end{cases} \quad (41)$$




FIGURE 2. Exact solution of $(\overset{\star}{x}_{1k}, \overset{\star}{y}_{1k})$ of generalized trapezoidal intuitionistic fuzzy number.




FIGURE 3. Exact solution of $(\overset{\star}{x}_{2k}, \overset{\star}{y}_{2k})$ of generalized trapezoidal intuitionistic fuzzy number.

$$\begin{cases} \frac{d^2 \overset{\star}{x}(t, \beta)}{dt^2} = 2\overset{\star}{y} - 3\overset{\star}{x}, \\ \overset{\star}{x}_1(0, \beta) = -4.29\beta + 5.29, \overset{\star}{x}'_1(0, \beta) = 6.71 - 0.85\beta, \\ \frac{d^2 \overset{\star}{y}(t, \beta)}{dt^2} = 2\overset{\star}{x} - 3\overset{\star}{y}, \\ \overset{\star}{y}_1(0, \beta) = 9 - 2.86\beta, \overset{\star}{y}'_1(0, \beta) = 12 - 2.86\beta, \end{cases} \quad (42)$$

$$\begin{cases} \frac{d^2 \overset{\star}{x}(t, \beta)}{dt^2} = 2\overset{\star}{y} - 3\overset{\star}{x}, \\ \overset{\star}{x}_2(0, \beta) = 3.71 + 14.29\beta, \overset{\star}{x}'_2(0, \beta) = 6.29 + 5.71\beta, \\ \frac{d^2 \overset{\star}{y}(t, \beta)}{dt^2} = 2\overset{\star}{x} - 3\overset{\star}{y}, \\ \overset{\star}{y}_2(0, \beta) = 9.14 + 2.86\beta, \overset{\star}{y}'_2(0, \beta) = 12.14 + 2.86\beta. \end{cases} \quad (43)$$

Here $\overset{\star}{L} = \frac{d^2}{dt^2}$ and by taking $\overset{\star}{L}^{-1}(\cdot) = \int_0^t \int_0^t (\cdot) dt dt$ on both sides of (40), (41), (42) and (43), and using the initial conditions we obtain;

$$\begin{cases} \overset{\star}{x}_1(t, \alpha) = \int_0^t (3\overset{\star}{x}_1(u, \alpha)(-t + u) + 2\overset{\star}{y}_1(u, \alpha)(t - u)) du \\ \quad + 2 + 5\alpha t + 3.33\alpha + 2t, \\ \overset{\star}{y}_1(t, \alpha) = \int_0^t (2\overset{\star}{x}_1(u, \alpha)(t - u) + 3\overset{\star}{y}_1(u, \alpha)(-t + u)) du \\ \quad + 8 + 1.67\alpha t + 1.67\alpha + 11t, \end{cases} \quad (44)$$

Figure 2-3: Shows exact solution of the system of generalized trapezoidal intuitionistic fuzzy initial value problem used




FIGURE 4. Exact solution of $(\hat{x}_{1_k}^*(t, \alpha), \hat{y}_{1_k}^*(t, \alpha))$ of generalized trapezoidal intuitionistic fuzzy number.




FIGURE 6. Exact solution of $(\hat{x}_{1_k}^*(t, \beta), \hat{y}_{1_k}^*(t, \beta))$ of generalized trapezoidal intuitionistic fuzzy number.




FIGURE 5. Exact solution of $(\hat{x}_{2_k}^*(t, \alpha), \hat{y}_{2_k}^*(t, \alpha))$ of generalized trapezoidal intuitionistic fuzzy number.




FIGURE 7. Exact solution of $(\hat{x}_{2_k}^*(t, \beta), \hat{y}_{2_k}^*(t, \beta))$ of generalized trapezoidal intuitionistic fuzzy number.

in example 1.

$$\begin{cases} \hat{x}_2(t, \alpha) = \int_0^t (3\hat{x}_2(u, \alpha)(-t + u) + 2\hat{y}_2(u, \alpha)(t - u))du \\ \quad + 15 - 3.33\alpha t - 11.67\alpha + 10t, \\ \hat{y}_2(t, \alpha) = \int_0^t (2\hat{x}_2(u, \alpha)(t - u) + 3\hat{y}_2(u, \alpha)(-t + u))du \\ \quad + 11 - 1.67\alpha t - 1.67\alpha + 14t, \end{cases} \quad (45)$$

$$\begin{cases} \hat{x}_1(t, \beta) = \int_0^t (3\hat{x}_1(u, \beta)(-t + u) + 2\hat{y}_1(u, \beta)(t - u))du \\ \quad + 5.29 - 0.85\beta t + 6.71t - 4.29\beta, \\ \hat{y}_1(t, \beta) = \int_0^t (2\hat{x}_1(u, \beta)(t - u) + 3\hat{y}_1(u, \beta)(-t + u))du \\ \quad + 9 - 2.86\beta t - 2.86\beta + 12t, \end{cases} \quad (46)$$

$$\begin{cases} \hat{x}_2(t, \beta) = \int_0^t (3\hat{x}_2(u, \beta)(-t + u) + 2\hat{y}_2(u, \beta)(t - u))du \\ \quad + 3.71 + 5.71\beta t + 6.29t + 14.29\beta, \\ \hat{y}_2(t, \beta) = \int_0^t (2\hat{x}_2(u, \beta)(t - u) + 3\hat{y}_2(u, \beta)(-t + u))du \\ \quad + 9.14 + 2.86\beta t + 12.14t + 2.86\beta. \end{cases} \quad (47)$$

Figures 3-7: Displays the membership and non-membership functions, as well as the exact and approximate

solutions, for the generalized trapezoidal intuitionistic fuzzy initial value problem used in example 2.

Now by using GMADM the solution of (44), (45), (46) and (47), can be expressed as;

$$\begin{cases} \hat{x}_{1_0}(t, \alpha) = 2 + t(5\alpha + 2) + 3.33\alpha, \\ \hat{y}_{1_0}(t, \alpha) = 8 + t(1.67\alpha + 11) + 1.67\alpha, \\ \hat{x}_{1_1}(t, \alpha) = -1.943333334\alpha t^3 + 2.666666667t^3 \\ \quad + 5.00000000t^2 - 3.325000000\alpha t^2, \\ \hat{y}_{1_1}(t, \alpha) = 2.498333334\alpha t^3 - 4.166666668t^3 \\ \quad - 8.00000000t^2 + 4.155000000\alpha t^2, \\ \hat{x}_{1_{k+1}}(t, \alpha) = \int_0^t (3\hat{x}_{1_k}(u, \alpha)(-t + u) \\ \quad + 2\hat{y}_{1_k}(u, \alpha)(t - u))du, k \geq 1, \\ \hat{y}_{1_{k+1}}(t, \alpha) = \int_0^t (2\hat{x}_{1_k}(u, \alpha)(t - u) \\ \quad + 3\hat{y}_{1_k}(u, \alpha)(-t + u))du, k \geq 1. \end{cases} \quad (48)$$

$$\left\{ \begin{aligned}
 \overset{*}{x}_{2_0}(t, \alpha) &= 15 + t(-3.33\alpha + 10) - 11.67\alpha, \\
 \overset{*}{y}_{2_0}(t, \alpha) &= 11 + t(-1.67\alpha + 14) - 1.67\alpha, \\
 \overset{*}{x}_{2_1}(t, \alpha) &= 1.108333334\alpha t^3 - 0.333333334t^3 \\
 &\quad - 11.50000000t^2 + 15.83500000\alpha t^2, \\
 \overset{*}{y}_{2_1}(t, \alpha) &= -1.385000000\alpha t^3 - 0.333333334t^3 \\
 &\quad + 13.50000000t^2 - 20.83500000\alpha t^2, \\
 \overset{*}{x}_{2_{k+1}}(t, \alpha) &= \int_0^t (3\overset{*}{x}_{2_k}(u, \alpha)(-t + u) \\
 &\quad + 2\overset{*}{y}_{2_k}(u, \alpha)(t - u))du, k \geq 1, \\
 \overset{*}{y}_{2_{k+1}}(t, \alpha) &= \int_0^t (2\overset{*}{x}_{2_k}(u, \alpha)(t - u) \\
 &\quad + 3\overset{*}{y}_{2_k}(u, \alpha)(-t + u))du, k \geq 1.
 \end{aligned} \right. \tag{49}$$

$$\left\{ \begin{aligned}
 \overset{*}{x}_{1_0}(t, \beta) &= 5.29 + t(-0.85\beta + 6.71) - 4.29\beta, \\
 \overset{*}{y}_{1_0}(t, \beta) &= 9 + t(-2.86\beta + 12) - 2.86\beta, \\
 \overset{*}{x}_{1_1}(t, \beta) &= -0.528333333\beta t^3 + 0.645000000t^3 \\
 &\quad + 1.065000000t^2 + 3.575000000\beta t^2, \\
 \overset{*}{y}_{1_1}(t, \beta) &= 0.863333334\beta t^3 - 1.526666667t^3 \\
 &\quad - 2.920000000t^2 - 4.290000000\beta t^2, \\
 \overset{*}{x}_{1_{k+1}}(t, \beta) &= \int_0^t (2\overset{*}{x}_{1_k}(u, \beta)(t - u) \\
 &\quad + 3\overset{*}{y}_{1_k}(u, \beta)(-t + u))du, k \geq 1, \\
 \overset{*}{y}_{1_{k+1}}(t, \beta) &= \int_0^t (2\overset{*}{x}_{1_k}(u, \beta)(t - u) \\
 &\quad + 3\overset{*}{y}_{1_k}(u, \beta)(-t + u))du, k \geq 1.
 \end{aligned} \right. \tag{50}$$

$$\left\{ \begin{aligned}
 \overset{*}{x}_{2_0}(t, \beta) &= 3.71 + t(5.71\beta + 6.29) + 14.29\beta, \\
 \overset{*}{y}_{2_0}(t, \beta) &= 9.14 + t(2.86\beta + 12.14) + 2.86\beta, \\
 \overset{*}{x}_{2_1}(t, \beta) &= -1.901666667\beta t^3 + 0.901666667t^3 \\
 &\quad + 3.575000000t^2 - 18.57500000\beta t^2, \\
 \overset{*}{y}_{2_1}(t, \beta) &= 2.376666667\beta t^3 - 1.876666667t^3 \\
 &\quad - 6.290000000t^2 + 24.29000000\beta t^2, \\
 \overset{*}{x}_{2_{k+1}}(t, \beta) &= \int_0^t (3\overset{*}{x}_{2_k}(u, \beta)(-t + u) \\
 &\quad + 2\overset{*}{y}_{2_k}(u, \beta)(t - u))du, k \geq 1, \\
 \overset{*}{y}_{2_{k+1}}(t, \beta) &= \int_0^t (2\overset{*}{x}_{2_k}(u, \beta)(t - u) \\
 &\quad + 3\overset{*}{y}_{2_k}(u, \beta)(-t + u))du, k \geq 1.
 \end{aligned} \right. \tag{51}$$

By solving (48), (49), (50) and (51), we get the approximate solution after four iterations as follows:

$$\begin{aligned}
 &(\overset{*}{x}_1(t, \alpha), \overset{*}{y}_1(t, \alpha)) \\
 &= (2 + 5\alpha t + 2t + 3.33\alpha - 1.943333334\alpha t^3 \\
 &\quad + 2.666666667t^3 + 5.00000000t^2 - 3.325000000\alpha t^2 \\
 &\quad + 0.541333334\alpha t^5 - 0.816666667t^5 - 2.583333330t^4 \\
 &\quad + 1.523749999\alpha t^4 - 0.0750198413\alpha t^7 \\
 &\quad + 0.1134920635t^7 \\
 &\quad + 0.502777777t^6 - 0.295513888\alpha t^6 \\
 &\quad + 0.00607264110\alpha t^9 \\
 &\quad - 0.00918761024t^9 - 0.0523313492t^8 \\
 &\quad + 0.0307552083\alpha t^8, \\
 &8 + 1.67\alpha t + 11t + 1.67\alpha + 2.498333334\alpha t^3 \\
 &\quad - 4.166666668t^3 \\
 &\quad - 8.00000000t^2 + 4.155000000\alpha t^2 - 0.763416667\alpha t^5 \\
 &\quad + 1.158333334t^5 + 3.66666667t^4 - 2.147083332\alpha t^4 \\
 &\quad + 0.1060853175\alpha t^7 - 0.1605158731t^7 - 0.711111110t^6 \\
 &\quad + 0.417874999\alpha t^6 - 0.00858799053\alpha t^9 \\
 &\quad + 0.0129932761t^9 \\
 &\quad + 0.0740079365t^8 - 0.0434942956\alpha t^8), \\
 &(\overset{*}{x}_2(t, \alpha), \overset{*}{y}_2(t, \alpha)) \\
 &= (15 - 3.33\alpha t + 10t - 11.67\alpha + 1.108333334\alpha t^3 \\
 &\quad - 0.333333334t^3 - 11.50000000t^2 + 15.83500000\alpha t^2 \\
 &\quad - 0.304750000\alpha t^5 + 0.01666666662t^5 + 5.12500000t^4 \\
 &\quad - 7.43125000\alpha t^4 + 0.0422162699\alpha t^7 \\
 &\quad - 0.00198412699t^7 \\
 &\quad - 0.993055554t^7 + 1.442263887\alpha t^6 \\
 &\quad - 0.00341724537\alpha t^9 \\
 &\quad + 0.000159832452t^9 + 0.1033482144t^8 \\
 &\quad - 0.150104911\alpha t^8, \\
 &11 - 1.67\alpha t + 14t - 1.67\alpha - 1.385000000\alpha t^3 \\
 &\quad - 0.333333334t^3 + 13.50000000t^2 - 20.83500000\alpha t^2 \\
 &\quad + 0.429416667\alpha t^5 - 0.01666666682t^5 - 7.20833333t^4 \\
 &\quad + 10.48708333\alpha t^4 - 0.0596964286\alpha t^7 \\
 &\quad + 0.00277777778t^7 \\
 &\quad + 1.404166665t^6 - 2.03954167\alpha t^6 + 0.00483269953\alpha t^9 \\
 &\quad - 0.000225970018t^9 - 0.146155754t^8 \\
 &\quad + 0.212280011\alpha t^8), \\
 &(\overset{*}{x}_1(t, \beta), \overset{*}{y}_1(t, \beta)) \\
 &= (5.29 - 85\beta t + 6.71t - 4.29\beta - 0.528333333\beta t^3 \\
 &\quad + 0.645000000t^3 + 1.065000000t^2 + 3.575000000\beta t^2 \\
 &\quad + 0.1655833334\beta t^5 - 0.2494166672t^5 - 0.752916666t^4 \\
 &\quad - 1.608749999\beta t^4 - 0.0230257937\beta t^7 \\
 &\quad + 0.0348630952t^7 \\
 &\quad + 0.1476249998t^6 + 0.311819444\beta t^6 \\
 &\quad + 0.00186405974\beta t^9 \\
 &\quad - 0.00282277888t^9 - 0.0153687996t^8 \\
 &\quad - 0.0324516369\beta t^8, \\
 &9 - 2.86\beta t + 12t - 2.86\beta + 0.863333334\beta t^3 \\
 &\quad - 1.526666667t^3
 \end{aligned}$$

$$\begin{aligned}
 & - 2.92000000t^2 - 4.290000000\beta t^2 - 0.2351666672\beta t^5 \\
 & + 0.358000000t^5 + 1.085000000t^4 + 2.264166666\beta t^4 \\
 & + 0.0325674603\beta t^7 - 0.0493253968t^7 - 0.208888889t^6 \\
 & - 0.440916666\beta t^6 - 0.00263618827\beta t^9 \\
 & + 0.00399206349t^9 \\
 & + 0.0217351191t^8 + 0.0458933532\beta t^8), \\
 (\hat{x}_2(t, \beta), \hat{y}_2(t, \beta)) \\
 & = (3.71 + 5.71\beta t + 6.29t + 14.29\beta - 1.901666667\beta t^3 \\
 & + 0.901666667t^3 + 3.57500000t^2 - 18.57500000\beta t^2 \\
 & + 0.522916667\beta t^5 - 0.322916667t^5 - 1.942083333t^4 \\
 & + 8.69208333\beta t^4 - 0.0724384921\beta t^7 + 0.0450575397t^7 \\
 & + 0.378486111t^6 - 1.686819440\beta t^6 \\
 & + 0.00586361883\beta t^9 \\
 & - 0.00364801036t^9 - 0.0393960814t^8 \\
 & + 0.175556795\beta t^8, \\
 & 9.14 + 2.86\beta t + 12.14t + 2.86\beta + 2.376666667\beta t^3 \\
 & - 1.876666667t^3 - 6.29000000t^2 + 24.29000000\beta t^2 \\
 & - 0.736833334\beta t^5 + 0.461833333t^5 + 2.764166670t^4 \\
 & - 12.26416666\beta t^4 + 0.1024325397\beta t^7 \\
 & - 0.0637420635t^7 \\
 & - 0.535361111t^6 + 2.38536111\beta t^6 \\
 & - 0.00829238317\beta t^9 \\
 & + 0.00515911596t^9 + 0.0557147819t^8 \\
 & - 0.248274306\beta t^8),
 \end{aligned}$$

In Table 3, $\hat{x}_1(t, \alpha), \hat{x}_2(t, \alpha), \hat{y}_1(t, \alpha)$ and $\hat{y}_2(t, \alpha)$ represents analytical solution of the membership functions of the Example1 for $\alpha \in [0, 0.6]$ and $\hat{x}_1(t, \beta), \hat{x}_2(t, \beta), \hat{y}_1(t, \beta)$ and $\hat{y}_2(t, \beta)$ represents analytical solution of non-membership function of the Example2 for $\beta \in [0.3, 1.0]$.

The following is the mathematical and exact solution to Example 2 using the classical method, as shown in Figure 4(a-d) and Table 4:

$$\begin{aligned}
 & (\hat{x}_1(t, \alpha), \hat{y}_1(t, \alpha)) \\
 & = ((\frac{667}{200}\alpha + \frac{13}{2})\sin(t) + (5 + \frac{5}{2}\alpha)\cos(t) \\
 & + \frac{9}{1000}\sqrt{5}(-100 + 37\alpha) \\
 & \times \sin(\sqrt{5}t) + (-3 + \frac{83}{100}\alpha)\cos(\sqrt{5}t), (\frac{667}{200}\alpha + \frac{13}{2})\sin(t)
 \end{aligned}$$

$$\begin{aligned}
 & + (5 + \frac{5}{2}\alpha)\cos(t) - \frac{9}{1000}\sqrt{5}(-100 + 37\alpha)\sin(\sqrt{5}t) \\
 & - (-3 + \frac{83}{100}\alpha)\cos(\sqrt{5}t)), \\
 & (\hat{x}_2(t, \alpha), \hat{y}_2(t, \alpha)) \\
 & = (-\frac{1}{500}\sqrt{5}(200 + 83\alpha)\sin(\sqrt{5}t) + (2 - 5\alpha)\cos(\sqrt{5}t) \\
 & + (-\frac{5}{2}\alpha + 12)\sin(t) + (13 - \frac{667}{100}\alpha)\cos(t), \\
 & \frac{1}{500}\sqrt{5}(200 + 83\alpha) \\
 & \times \sin(\sqrt{5}t) - (2 - 5\alpha)\cos(\sqrt{5}t) + (-\frac{5}{2}\alpha + 12)\sin(t) \\
 & + (13 - \frac{667}{100}\alpha)\cos(t)), \\
 & (\hat{x}_1(t, \beta), \hat{y}_1(t, \beta)) \\
 & = (\frac{1}{1000}\sqrt{5}(-529 + 201\beta)\sin(\sqrt{5}t) \\
 & + (-\frac{371}{200} - \frac{143}{200}\beta)\cos(\sqrt{5}t) \\
 & + (-\frac{371}{200}\beta + \frac{1871}{200})\sin(t) + (\frac{1429}{200} - \frac{143}{40}\beta)\cos(t), \\
 & - \frac{1}{1000}\sqrt{5} \\
 & \times (-529 + 201\beta)\sin(\sqrt{5}t) - (-\frac{371}{200} - \frac{143}{200}\beta)\cos(\sqrt{5}t) \\
 & + (-\frac{371}{200}\beta + \frac{1871}{200})\sin(t) + (\frac{1429}{200} - \frac{143}{40}\beta)\cos(t)), \\
 & (\hat{x}_2(t, \beta), \hat{y}_2(t, \beta)) \\
 & = ((\frac{857}{200}\beta + \frac{1843}{200})\sin(t) + (\frac{257}{40} + \frac{343}{40}\beta)\cos(t) + \frac{3}{200}\sqrt{5} \\
 & \times (-39 + 19\beta)\sin(\sqrt{5}t) + (-\frac{543}{200} + \frac{1143}{200}\beta)\cos(\sqrt{5}t), \\
 & (\frac{857}{200}\beta + \frac{1843}{200})\sin(t) + (\frac{257}{40} + \frac{343}{40}\beta)\cos(t) \\
 & - \frac{3}{200}\sqrt{5}(-39 + 19\beta)\sin(\sqrt{5}t) \\
 & - (-\frac{543}{200} + \frac{1143}{200}\beta)\cos(\sqrt{5}t)).
 \end{aligned}$$

In Table 4, $\hat{x}_1(t, \alpha), \hat{x}_2(t, \alpha), \hat{y}_1(t, \alpha)$ and $\hat{y}_2(t, \alpha)$ represents exact solution of the membership functions of the Example2 for $\alpha \in [0, 0.6]$ and $\hat{x}_1(t, \beta), \hat{x}_2(t, \beta), \hat{y}_1(t, \beta)$ and $\hat{y}_2(t, \beta)$ represents exact solution of non-membership function of the Example2 for $\beta \in [0.3, 1.0]$.

Example 3: Consider the first order non-homogenous system of intuitionistic fuzzy differential equation as follows:

$$\left. \begin{aligned}
 \frac{dx}{dt} &= 2x(t) + 3y(t) - 7, \\
 \frac{dy}{dt} &= -x(t) - 2y(t) + 5.
 \end{aligned} \right\} \quad (52)$$

TABLE 4. Illustrates the approximation to the generalized trapezoidal intuitionistic fuzzy initial value problem from Example 2 for t=1.

α	$\tilde{x}_1(t, \alpha)$	$\tilde{y}_1(t, \alpha)$	$\tilde{x}_2(t, \alpha)$	$\tilde{y}_2(t, \alpha)$	β	$\tilde{x}_1(t, \beta)$	$\tilde{y}_1(t, \beta)$	$\tilde{x}_2(t, \beta)$	$\tilde{y}_2(t, \beta)$
0	4.408	10.87	17.38	32.20	0.3	7.746	15.51	11.57	22.28
0.1	4.891	11.62	16.42	30.54	0.4	7.348	14.77	12.85	24.44
0.2	5.377	12.38	15.45	28.89	0.5	6.950	14.02	14.13	26.60
0.3	5.863	13.14	14.48	27.24	0.6	6.552	13.27	15.41	28.77
0.4	6.350	13.89	13.51	25.59	0.7	6.154	12.52	16.69	30.93
0.5	6.836	14.65	12.54	23.93	0.8	5.757	11.77	17.97	33.09
0.6	7.322	15.40	11.57	22.28	0.9	5.359	11.02	19.25	35.25
					1.0	4.961	10.288	20.53	37.41

TABLE 5. Illustrates the exact solution to the generalized trapezoidal intuitionistic fuzzy initial value problem from Example 2 for t=1.

α	$\tilde{x}_1(t, \alpha)$	$\tilde{y}_1(t, \alpha)$	$\tilde{x}_2(t, \alpha)$	$\tilde{y}_2(t, \alpha)$	β	$\tilde{x}_1(t, \beta)$	$\tilde{y}_1(t, \beta)$	$\tilde{x}_2(t, \beta)$	$\tilde{y}_2(t, \beta)$
0	4.382	10.63	17.23	17.09	0.3	7.699	11.40	11.49	14.37
0.1	4.864	10.90	16.27	16.64	0.4	7.305	10.99	12.75	15.02
0.2	5.347	11.18	15.32	16.19	0.5	6.911	10.57	14.02	15.67
0.3	5.829	11.46	14.36	15.73	0.6	6.519	10.16	15.28	16.32
0.4	6.313	11.73	13.40	15.28	0.7	6.126	9.750	16.56	16.98
0.5	6.795	12.01	12.45	14.83	0.8	5.732	9.336	17.80	17.62
0.6	7.276	12.28	11.49	14.37	0.9	5.337	8.932	19.08	18.27
					1.0	4.945	8.513	20.34	18.92

TABLE 6. Illustrates the approximate solution iterations in column 2, residual error in column 3, and CPU time required by the numerical technique GMADM to determine the approximate solution of the system of generalized fuzzy intuitionistic differential equations used in Example 2 in column 4. Whenever the error of all methods is taken into account, we can conclude that the GMADM has a better convergence behavior and is more stable than the ADM and TSM, respectively.

Parameter	Iterations	Error	CPU-time
GMADM	4	0.2E-19	1.00145
ADM	7	3.1E-07	1.94512
TSM	6	4.2E-09	1.14532

with initial conditions

$$\left. \begin{aligned} \tilde{x}(0) &= \langle (2, 4, 8, 15; 0.6); (1, 4, 8, 18; 0.3) \rangle, \\ \tilde{y}(0) &= \langle (2, 5, 8, 10; 0.6)(1, 5, 8, 12; 0.3) \rangle. \end{aligned} \right\} \quad (53)$$

By taking (α, β) -cuts of (52) and (53), we get,

$$\left. \begin{aligned} \frac{d\tilde{x}_1(t, \alpha)}{dt} &= 2\tilde{x}_1(t, \alpha) + 3\tilde{y}_1(t, \alpha) - 7, \\ \tilde{x}_1(0, \alpha) &= 3.33\alpha + 2, \\ \frac{d\tilde{y}_1(t, \alpha)}{dt} &= -\tilde{x}_1(t, \alpha) - 2\tilde{y}_1(t, \alpha) + 5, \\ \tilde{y}_1(0, \alpha) &= 5\alpha + 2, \end{aligned} \right\} \quad (54)$$

$$\left. \begin{aligned} \frac{d\tilde{x}_2(t, \alpha)}{dt} &= 2\tilde{x}_2(t, \alpha) + 3\tilde{y}_2(t, \alpha) - 7, \\ \tilde{x}_2(0, \alpha) &= -11.67\alpha + 15, \\ \frac{d\tilde{y}_2(t, \alpha)}{dt} &= -\tilde{x}_2(t, \alpha) - 2\tilde{y}_2(t, \alpha) + 5, \\ \tilde{y}_2(0, \alpha) &= -3.33\alpha + 10, \end{aligned} \right\} \quad (55)$$

$$\left. \begin{aligned} \frac{d\tilde{x}_1(t, \beta)}{dt} &= 2\tilde{x}_1(t, \beta) + 3\tilde{y}_1(t, \beta) - 7, \\ \tilde{x}_1(0, \beta) &= -4.29\beta + 5.29, \\ \frac{d\tilde{y}_1(t, \beta)}{dt} &= -\tilde{x}_1(t, \beta) - 2\tilde{y}_1(t, \beta) + 5, \\ \tilde{y}_1(0, \beta) &= -0.85\beta + 6.71, \end{aligned} \right\} \quad (56)$$

$$\left. \begin{aligned} \frac{d\tilde{x}_2(t, \beta)}{dt} &= 2\tilde{x}_2(t, \beta) + 3\tilde{y}_2(t, \beta) - 7, \\ \tilde{x}_2(0, \beta) &= 14.29\beta + 3.71, \\ \frac{d\tilde{y}_2(t, \beta)}{dt} &= -\tilde{x}_2(t, \beta) - 2\tilde{y}_2(t, \beta) + 5, \\ \tilde{y}_2(0, \beta) &= 5.71\beta + 6.29. \end{aligned} \right\} \quad (57)$$

Here $L = \frac{d}{dt}$ and by taking $L^{-1}(\cdot) = \int_0^t (\cdot) dt$ on both sides of (54), (55), (56) and (57), and using the initial conditions we obtain;

$$\left. \begin{aligned} \tilde{x}_1(t, \alpha) &= \int_0^t (2\tilde{x}_1(u, \alpha) + 3\tilde{y}_1(u, \alpha)) du \\ &\quad + 3.33\alpha + 2 - 7t, \end{aligned} \right\} \quad (58)$$

$$\left. \begin{aligned} \tilde{y}_1(t, \alpha) &= \int_0^t (-\tilde{x}_1(u, \alpha) - 2\tilde{y}_1(u, \alpha)) \\ &\quad + 5\alpha + 2 + 5t, \\ \tilde{x}_2(t, \alpha) &= \int_0^t (2\tilde{x}_2(u, \alpha) + 3\tilde{y}_2(u, \alpha)) du \\ &\quad - 11.67\alpha + 15 - 7t, \end{aligned} \right\} \quad (59)$$

$$\left. \begin{aligned} \tilde{y}_2(t, \alpha) &= \int_0^t (-\tilde{x}_2(u, \alpha) - 2\tilde{y}_2(u, \alpha)) \\ &\quad - 3.33\alpha + 10 + 5t, \end{aligned} \right\}$$

$$\left. \begin{aligned} \tilde{x}_1(t, \beta) &= \int_0^t (2\tilde{x}_1(u, \beta) + 3\tilde{y}_1(u, \beta)) du \\ &\quad + 5.29 - 4.29\beta - 7t, \end{aligned} \right\} \quad (60)$$

$$\left. \begin{aligned} \tilde{y}_1(t, \beta) &= \int_0^t (-\tilde{x}_1(u, \beta) - 2\tilde{y}_1(u, \beta)) \\ &\quad + 6.71 - 0.85\beta + 5t, \end{aligned} \right\}$$

$$\left. \begin{aligned} \tilde{x}_2(t, \beta) &= \int_0^t (2\tilde{x}_2(u, \beta) + 3\tilde{y}_2(u, \beta)) du \\ &\quad + 14.29\beta + 3.71 - 7t, \end{aligned} \right\} \quad (61)$$

$$\left. \begin{aligned} \tilde{y}_2(t, \beta) &= \int_0^t (-\tilde{x}_2(u, \beta) - 2\tilde{y}_2(u, \beta)) + 5.71\beta \\ &\quad + 6.29 + 5t. \end{aligned} \right\}$$

Now by using GMADM we get;

$$\begin{cases} \overset{*}{x}_{1_0}(t, \alpha) = 3.33\alpha + 2, \\ \overset{*}{y}_{1_0}(t, \alpha) = 5\alpha + 2, \\ \overset{*}{x}_{1_1}(t, \alpha) = 21.66000000\alpha t + 3t, \\ \overset{*}{y}_{1_1}(t, \alpha) = -13.33000000\alpha t - t, \\ \overset{*}{x}_{1_{k+1}}(t, \alpha) = \int_0^t (2\overset{*}{x}_{1_k}(u, \alpha) + 3\overset{*}{y}_{1_k}(u, \alpha))du, k \geq 1, \\ \overset{*}{y}_{1_{k+1}}(t, \alpha) = \int_0^t (-\overset{*}{x}_{1_k}(u, \alpha) - 2\overset{*}{y}_{1_k}(u, \alpha))du, k \geq 1. \end{cases} \quad (62)$$

$$\begin{cases} \overset{*}{x}_{2_0}(t, \alpha) = -11.67\alpha + 15, \\ \overset{*}{y}_{2_0}(t, \alpha) = -3.33\alpha + 10, \\ \overset{*}{x}_{2_1}(t, \alpha) = -33.33000000\alpha t + 53t, \\ \overset{*}{y}_{2_1}(t, \alpha) = 18.33000000\alpha t - 30t, \\ \overset{*}{x}_{2_{k+1}}(t, \alpha) = \int_0^t (2\overset{*}{x}_{2_k}(u, \alpha) + 3\overset{*}{y}_{2_k}(u, \alpha))du, k \geq 1, \\ \overset{*}{y}_{2_{k+1}}(t, \alpha) = \int_0^t (-\overset{*}{x}_{2_k}(u, \alpha) - 2\overset{*}{y}_{2_k}(u, \alpha))du, k \geq 1, \end{cases} \quad (63)$$

$$\begin{cases} \overset{*}{x}_{1_0}(t, \beta) = 5.29 - 4.29\beta, \\ \overset{*}{y}_{1_0}(t, \beta) = 6.71 - 0.85\beta, \\ \overset{*}{x}_{1_1}(t, \beta) = 23.71000000t - 11.13000000\beta t, \\ \overset{*}{y}_{1_1}(t, \beta) = -13.71000000t + 5.990000000\beta t, \\ \overset{*}{x}_{1_{k+1}}(t, \beta) = \int_0^t (2\overset{*}{x}_{1_k}(u, \alpha) + 3\overset{*}{y}_{1_k}(u, \alpha))du, k \geq 1, \\ \overset{*}{y}_{1_{k+1}}(t, \beta) = \int_0^t (-\overset{*}{x}_{1_k}(u, \alpha) - 2\overset{*}{y}_{1_k}(u, \alpha))du, k \geq 1. \end{cases} \quad (64)$$

$$\begin{cases} \overset{*}{x}_{2_0}(t, \beta) = 3.71 + 14.29\beta, \\ \overset{*}{y}_{2_0}(t, \beta) = 6.29 + 5.71\beta, \\ \overset{*}{x}_{2_1}(t, \beta) = 19.29000000t + 45.71000000\beta t, \\ \overset{*}{y}_{2_1}(t, \beta) = -11.29000000t - 25.71000000\beta t, \\ \overset{*}{x}_{2_{k+1}}(t, \beta) = \int_0^t (2\overset{*}{x}_{2_k}(u, \alpha) + 3\overset{*}{y}_{2_k}(u, \alpha))du, k \geq 1, \\ \overset{*}{y}_{2_{k+1}}(t, \beta) = \int_0^t (-\overset{*}{x}_{2_k}(u, \alpha) - 2\overset{*}{y}_{2_k}(u, \alpha))du, k \geq 1. \end{cases} \quad (65)$$




FIGURE 8. Exact solution of $(\overset{*}{x}_{1_k}(t, \alpha), \overset{*}{y}_{1_k}(t, \alpha))$ of generalized trapezoidal intuitionistic fuzzy number.




FIGURE 9. Exact solution of $(\overset{*}{x}_{2_k}(t, \alpha), \overset{*}{y}_{2_k}(t, \alpha))$ of generalized trapezoidal intuitionistic fuzzy number.




FIGURE 10. Exact solution of $(\overset{*}{x}_{1_k}(t, \beta), \overset{*}{y}_{1_k}(t, \beta))$ of generalized trapezoidal intuitionistic fuzzy number.

Figures 8-11: Displays the membership and non-membership functions, as well as the exact and approximative solutions, for the generalised trapezoidal intuitionistic fuzzy initial value problem used in example 3.




FIGURE 11. Exact solution of $(\hat{x}_{2k}^*(t, \beta), \hat{y}_{2k}^*(t, \beta))$ of generalized trapezoidal intuitionistic fuzzy number.

By solving the (62), (63), (64) and (65), we get the approximate solution after three iteration as follows:

$$\begin{aligned}
 &(\hat{x}_1(t, \alpha), \hat{y}_1(t, \alpha)) \\
 &= (3.33\alpha + 2 + 3t + 21.66000000\alpha t + 1.500000000t^2 \\
 &\quad + 1.665000000\alpha t^2 + 0.500000000t^3 + 3.610000000\alpha t^3 \\
 &\quad + 0.125000000t^4 + 0.1387500007\alpha t^4, 5\alpha + 2 - t \\
 &\quad - 13.33000000\alpha t + 2.500000000\alpha t^2 - 0.500000000t^2 \\
 &\quad - 0.166666666t^3 - 2.221666666\alpha t^3 \\
 &\quad - 0.04166666670t^4 \\
 &\quad + 0.2083333329\alpha t^4), \\
 &(\hat{x}_2(t, \alpha), \hat{y}_2(t, \alpha)) \\
 &= (-11.67\alpha + 15 + 53t - 33.33000000\alpha t + 8.000000000t^2 \\
 &\quad - 5.835000000\alpha t^2 + 8.833333330t^3 - 5.554999998\alpha t^3 \\
 &\quad + 0.6666666650t^4 - 0.486249990\alpha t^4, \\
 &\quad - 3.33\alpha + 10 - 30t \\
 &\quad + 18.33000000\alpha t - 1.665000000\alpha t^2 + 3.50000000t^2 \\
 &\quad - 5.000000000t^3 + 3.055000000\alpha t^3 + 0.2916666675t^4 \\
 &\quad - .1387500006\alpha t^4), \\
 &(\hat{x}_1(t, \beta), \hat{y}_1(t, \beta)) \\
 &= (-4.29\beta + 5.29 + 23.71000000t - 11.13000000\beta t \\
 &\quad + 3.145000000t^2 - 2.145000000\beta t^2 + 3.951666665t^3 \\
 &\quad - 1.854999999\beta t^3 + 0.262083332t^4 \\
 &\quad - 0.1787499999\beta t^4, \\
 &\quad - 0.85\beta + 6.71 - 13.71000000t + 5.990000000\beta t \\
 &\quad - 0.425000000\beta t^2 + 1.855000000t^2 - 2.284999999t^3 \\
 &\quad + 0.9983333329\beta t^3 + 0.1545833333t^4 \\
 &\quad - 0.03541666665\beta t^4), \\
 &(\hat{x}_2(t, \beta), \hat{y}_2(t, \beta)) \\
 &= (14.29\beta + 3.71 + 19.29000000t + 45.71000000\beta t \\
 &\quad + 2.355000000t^2 + 7.145000000\beta t^2 + 3.214999999t^3
 \end{aligned}$$

$$\begin{aligned}
 &+ 7.618333330\beta t^3 + 0.1962499999t^4 \\
 &+ 0.5954166665\beta t^4, \\
 &5.71\beta + 6.29 - 11.29000000t - 25.71000000\beta t \\
 &+ 2.855000000\beta t^2 + 1.645000000t^2 - 1.881666666t^3 \\
 &- 4.284999998\beta t^3 + 0.1370833333t^4 \\
 &+ 0.2379166666\beta t^4).
 \end{aligned}$$

The following is the mathematical and exact solution to Example 3 using the classical method, as shown in Figure 4(a-d) and Table 6:

$$\begin{aligned}
 \hat{x}_1(t, \alpha) &= -\frac{1833}{200}\alpha e^{-t} + (3 + \frac{2499}{200}\alpha)e^t - 1, \\
 \hat{y}_1(t, \alpha) &= \frac{1833}{200}\alpha e^{-t} - \frac{1}{3}(3 + \frac{2499}{200}\alpha)e^t + 3\}, \\
 \hat{x}_2(t, \alpha) &= (\frac{69}{2} - \frac{45}{2}\alpha)e^t + (-\frac{37}{2} + \frac{1083}{100}\alpha)e^{-t} - 1, \\
 \hat{y}_2(t, \alpha) &= -(-\frac{37}{2} + \frac{1083}{100}\alpha)e^{-t} \\
 &\quad - \frac{1}{3}(\frac{69}{2} - \frac{45}{2}\alpha)e^t + 3\}, \\
 \hat{x}_1(t, \beta) &= (-\frac{871}{100} + \frac{171}{50}\beta)e^{-t} + (15 - \frac{771}{100}\beta)e^t - 1, \\
 \hat{y}_1(t, \beta) &= -(-\frac{871}{100} + \frac{171}{50}\beta)e^{-t} \\
 &\quad - \frac{1}{3}(15 - \frac{771}{100}\beta)e^t + 3, \\
 \hat{x}_2(t, \beta) &= (12 + 30\beta)e^t + (-\frac{729}{100} - \frac{1571}{100}\beta)e^{-t} - 1, \\
 \hat{y}_2(t, \beta) &= -(-\frac{729}{100} - \frac{1571}{100}\beta)e^{-t} \\
 &\quad - \frac{1}{3}(12 + 30\beta)e^t + 3\},
 \end{aligned}$$

IV. RESULTS AND DISCUSSION

We discuss how the computing efficiency, stability, and residual error robustness of the proposed modified technique, GMADM, outperforms the ADM and TSM approaches.

- Tables 1 to 9 illustrate how GMADM, a recently created approach, is more reliable and consistent than ADM and TSM. While solving a system of fuzzy intuitionistic differential equations, it is observed that GMADM converges more quickly and accurately than ADM and TSM.
- Tables 3, 6, and 9 clearly demonstrate that GMADM is superior to ADM and TSM in terms of iterations, residual error, and CPU time.
- Figures 2-11 compare the numerical simulation of our recently modified family GMADM to a precise solution of the generalized trapezoidal intuitionistic fuzzy initial value problem used in Example 1-3 respectively.
- Figures 2-3, 4-11 illustrate the precise and approximate solutions for the membership and non-membership functions of the generalized trapezoidal intuitionistic

TABLE 7. Illustrates the approximation to the generalized trapezoidal intuitionistic fuzzy initial value problem from Example 3 for $t=1$.

Table 7: Approximate solution of numerical example at $t=1$.

α	$\hat{x}_1(t, \alpha)$	$\hat{y}_1(t, \alpha)$	$\hat{x}_2(t, \alpha)$	$\hat{y}_2(t, \alpha)$	β	$\hat{x}_1(t, \beta)$	$\hat{y}_1(t, \beta)$	$\hat{x}_2(t, \beta)$	$\hat{y}_2(t, \beta)$
0	7.125	0.292	85.50	-21.21	0.3	30.48	-5.572	51.37	-11.46
0.1	10.17	-0.493	79.81	-19.58	0.4	28.52	-5.004	58.91	-13.58
0.2	13.21	-1.277	74.13	-17.96	0.5	26.56	-4.436	66.45	-15.70
0.3	16.25	-2.061	68.44	-16.33	0.6	24.60	-3.867	73.98	-17.82
0.4	19.29	-2.846	62.75	-14.71	0.7	22.64	-3.301	81.52	-19.93
0.5	22.33	-3.630	57.06	-13.08	0.8	20.68	-2.733	89.05	-22.05
0.6	25.37	-4.414	51.37	-11.46	0.9	18.72	-2.165	96.59	-24.17
					1.0	16.76	-1.597	104.13	-26.29

TABLE 8. Illustrates the exact solution to the generalized trapezoidal intuitionistic fuzzy initial value problem from Example 3 for $t=1$.

Table 8: Exact solution of numerical example at $t=1$.

α	$\hat{x}_1(t, \alpha)$	$\hat{y}_1(t, \alpha)$	$\hat{x}_2(t, \alpha)$	$\hat{y}_2(t, \alpha)$	β	$\hat{x}_1(t, \beta)$	$\hat{y}_1(t, \beta)$	$\hat{x}_2(t, \beta)$	$\hat{y}_2(t, \beta)$
0	7.155	0.282	85.98	-21.45	0.3	30.66	-5.669	51.67	-11.61
0.1	10.21	-0.513	80.26	-19.81	0.4	28.69	-5.096	59.25	-13.75
0.2	13.27	-1.308	74.54	-18.17	0.5	26.72	-4.523	66.82	-15.89
0.3	16.33	-2.103	68.82	-16.53	0.6	24.75	-3.950	74.40	-18.03
0.4	19.39	-2.899	63.10	-14.89	0.7	22.78	-3.378	81.97	-20.17
0.5	22.45	-3.693	57.39	-13.25	0.8	20.81	-2.805	89.55	-22.31
0.6	25.51	-4.488	51.67	-11.61	0.9	18.84	-2.232	97.13	-24.45
					1.0	16.87	-1.659	104.7	-26.59

TABLE 9. Illustrates the approximate solution iterations in column 2, residual error in column 3, and CPU time required by the numerical technique GMADM to determine the approximate solution of the system of generalized fuzzy intuitionistic differential equations used in Example 3 in column 4. Whenever the error of all methods is taken into account, we can conclude that the GMADM has a better convergence behavior and is more stable than the ADM and TSM, respectively.

Table 9: Error comparison

Paramater	Iterations	Error	CPU-time
GMADM	3	0.6E-18	0.0145
ADM	9	4.7E-06	0.1646
TSM	8	0.9E-05	0.1531

fuzzy system of initial value problems used in example 1-3, respectively.

- The numerical results obtained in Tables 1-2,4-6,7-8, and Figures 1-11 clearly demonstrate that the exact and approximate solutions are matched up to 30 decimal places using GMADM, 7 decimal places using ADM, and 9 decimal places using TSM. The numerical simulation of our methods demonstrates unequivocally how much superior our method is to ADM and TSM.

V. CONCLUSION

In this work, Generalized Modified Adomian Decomposition Method have been utilized for computing the approximate solution of the linear system of generalized trapezoidal intuitionistic fuzzy initial value problems. We used the initial conditions as generalized trapezoidal intuitionistic fuzzy numbers. We have applied this procedure to brine tank problems and coupled oscillators. Moreover, by comparing the approximate results with exact solution, we have shown that this method is more reliable. Future studies will therefore focus on the solution of systems of higher order generalized tripezodial intuitionistic fuzzy system of differential equations as well as a system of nonlinear first order differential equations and their application [48], [49], [50], [51], [52], [53] in a more generalized fuzzy environment utilizing GMADM.

COMPETING INTERESTS

The authors confirm that there is no competing interests between them.

REFERENCES

- [1] L. A. Zadeh, "Fuzzy sets," *Inf. Control*, vol. 8, no. 3, pp. 338–353, Jun. 1965.
- [2] A. Kandel and W. J. Byatt, "Fuzzy processes," *Fuzzy Sets Syst.*, vol. 4, no. 2, pp. 117–152, Jan. 1980.
- [3] A. Ashraf, K. Ullah, A. Hussain, and M. Bari, "Interval-valued picture fuzzy Maclaurin symmetric mean operator with application in multiple attribute decision-making," *Rep. Mech. Eng.*, vol. 3, no. 1, pp. 301–317, Apr. 2022.
- [4] S. Kumar, S. R. Maity, and L. Patnaik, "Optimization of wear parameters for Duplex-TiAlN coated MDC-K tool steel using fuzzy MCDM techniques," *Oper. Res. Eng. Sci., Theory Appl.*, vol. 5, no. 3, pp. 40–67, Jul. 2022.
- [5] R. Alikhani and M. Mostafazadeh, "First order linear fuzzy differential equations with fuzzy variable coefficients," *Comput. Methods Differ. Equ.*, vol. 9, no. 1, pp. 1–21, Jan. 2021.
- [6] C. Vasavi, G. S. Kumar, T. S. Rao, and B. V. AppaRao, "Application of fuzzy differential equations for cooling problems," *Int. J. Mech. Eng. Technol.*, vol. 8, no. 12, pp. 712–721, Dec. 2017.
- [7] S. S. Devi and K. Ganesan, "Modelling electric circuit problem with fuzzy differential equations," *J. Phys., Conf. Ser.*, vol. 1377, Jan. 2019, Art. no. 012024.
- [8] J. Ahmad, A. Iqbal, and Q. M. UL Hassan, "Study of nonlinear fuzzy integro differential equations using mathematical methods and applications," *Int. J. Fuzzy Log. Intell. Syst.*, vol. 21, no. 1, pp. 76–85, Mar. 2021.
- [9] A. Sadeghi, A. I. Md Ismail, and A. F. Jameel, "Solving system of fuzzy differential equation," *Int. Math. Forum*, vol. 6, no. 42, pp. 2087–2100, 2011.
- [10] J. J. Buckley, T. Feuring, and Y. Hayashi, "Linear systems of first order ordinary differential equations: Fuzzy initial conditions," *Soft Comput.*, vol. 6, pp. 415–421, Sep. 2002.
- [11] M. S. Hashemi, J. Malekinaged, and H. R. Marasi, "Series solution of the system of fuzzy differential equations," *Adv. Fuzzy Syst.*, vol. 2012, no. 1, pp. 1–16, Jan. 2012.
- [12] K. T. Atanassov, "Intuitionistic fuzzy sets," *Fuzzy Sets Syst.*, vol. 20, pp. 87–96, Aug. 1986.
- [13] M. R. Seikh, P. K. Nayak, and M. Pal, "Generalized triangular fuzzy numbers in intuitionistic fuzzy environment," *Int. J. Eng. Res. Develop.*, vol. 5, no. 1, pp. 8–13, Nov. 2012.
- [14] S. Melliani and L. S. Chadli, "Introduction to intuitionistic fuzzy differential equations," *Notes IFS*, vol. 7, pp. 39–42, Apr. 2001.

- [15] B. B. Amma, S. Melliani, and L. S. Chadli, "Numerical solution of intuitionistic fuzzy differential equations by Euler and Taylor methods," *Notes Intuitionistic Fuzzy Sets*, vol. 22, no. 2, pp. 71–86, Jan. 2016.
- [16] M. Gulzar, M. H. Mateen, D. Alghazzawi, and N. Kausar, "A novel application of complex intuitionistic fuzzy sets in group theory," *IEEE Access*, vol. 8, pp. 196075–196085, 2020.
- [17] M. Gulzar, D. Alghazzawi, M. H. Mateen, and N. Kausar, "A certain class of t-intuitionistic fuzzy subgroups," *IEEE Access*, vol. 8, pp. 163260–163268, 2020, doi: [10.1109/ACCESS.2020.3020366](https://doi.org/10.1109/ACCESS.2020.3020366).
- [18] M. Gulzar, D. Alghazzawi, M. H. Mateen, and M. Premkumar, "On some characterization of Q-complex fuzzy sub-rings," *J. Math. Comput. Sci.*, vol. 22, no. 3, pp. 295–305, Aug. 2020.
- [19] O. Akin and S. Bayeg, "Intuitionistic fuzzy initial value problems—An application," *J. Math. Statist.*, vol. 48, pp. 1682–1694, Jun. 2019.
- [20] O. Akin and S. Bayeg, "System of intuitionistic fuzzy differential equations with intuitionistic fuzzy initial values," *Notes Intuitionistic Fuzzy Sets*, vol. 24, no. 4, pp. 141–171, Nov. 2018.
- [21] S. P. Mondal and T. K. Roy, "First order homogenous ordinary differential equation with initial value as triangular intuitionistic fuzzy number," *J. Uncertain. Math. Sci.*, vol. 2014, pp. 1–17, Jan. 2014.
- [22] S. P. Mondal and T. K. Roy, "System of differential equations with initial value as triangular intuitionistic fuzzy number and its application," *Int. J. Appl. Comput. Math.*, vol. 1, pp. 449–474, Sep. 2015.
- [23] S. P. Mondal and T. K. Roy, "Generalized intuitionistic fuzzy Laplace transform and its application in electric circuit," *J. Appl. Eng. Math.*, vol. 5, no. 1, pp. 30–45, Jan. 2015.
- [24] B. C. Saw and S. B. Hazra, " $\alpha, \beta, \gamma, \delta$ parametric form for solving intuitionistic fuzzy linear system of equations," *Adv. Fuzzy Math.*, vol. 16, no. 1, pp. 1–15, May 2021.
- [25] G. Adomian, "Convergent series solution of nonlinear equations," *J. Comput. Appl. Math.*, vol. 11, pp. 225–230, Oct. 1984.
- [26] G. Adomian, "A review of the decomposition method in applied mathematics," *J. Math. Anal. Appl.*, vol. 135, no. 2, pp. 501–544, Nov. 1988.
- [27] A.-M. Wazwaz, "A reliable modification of Adomian decomposition method," *Appl. Math. Comput.*, vol. 102, no. 1, pp. 77–86, Jul. 1999.
- [28] J. J. Buckley and T. Feuring, "Fuzzy initial value problem for Nth-order linear differential equations," *Fuzzy Sets. Syst.*, vol. 121, no. 2, pp. 247–255, Jul. 2001.
- [29] N. Gasilova, S. E. Amrahov, and C. G. gluFatullayev, "A geometric approach to solve fuzzy linear systems of differential equations," *Appl. Math. Inf. Sci.*, vol. 5, no. 3, pp. 484–499, Oct. 2011.
- [30] M. S. Hashemi, M. K. Mirnia, and S. Shahmorad, "Solving fuzzy linear systems by using the schur complement when coefficient matrix is an M-matrix," *Iranian J. Fuzzy Syst.*, vol. 5, no. 3, pp. 15–29, May 2008.
- [31] S. P. Mondal and T. K. Roy, "First order linear non homogeneous ordinary differential equation in fuzzy environment," *Math. Theory. Model.*, vol. 3, no. 1, pp. 85–95, Dec. 2013.
- [32] S. Melliani, R. M. Ettoussi, and L. S. Chadli, "Solution of intuitionistic fuzzy differential equations by successive approximations method," *Notes Intuitionistic Fuzzy Sets*, vol. 21, no. 2, pp. 51–62, Jun. 2015.
- [33] V. Nirmala and P. S. Chenthur, "Numerical approach for solving intuitionistic fuzzy differential equation," *Appl. Math. Sci.*, vol. 9, no. 367, pp. 3337–3346, Dec. 2015.
- [34] S. Biswas and T. K. Roy, "Adomian decomposition method for fuzzy differential equations with linear differential operator," *J. Inform. Comput. Sci.*, vol. 11, no. 4, pp. 243–250, May 2016.
- [35] G. Uthra, K. Thangavelu, and R. M. Umamageswari, "An optimal solution for generalized trapezoidal intuitionistic fuzzy transportation problem," *Adv. Fuzzy Math.*, vol. 12, no. 3, pp. 763–770, Feb. 2017.
- [36] V. Ulucay, I. Deli, and M. Sahin, "Intuitionistic trapezoidal fuzzy multi-numbers and its application to multi-criteria decision-making problems," *Complex Intell. Syst.*, vol. 5, no. 1, pp. 65–78, Mar. 2019.
- [37] S. Banerjee and T. K. Roy, "Arithmetic operations on generalized trapezoidal fuzzy number and its applications," *Turkish J. Fuzzy Syst.*, vol. 3, no. 1, pp. 16–44, Jan. 2012.
- [38] D. Chakraborty, D. K. Jana, and T. K. Roy, "Arithmetic operations on generalized intuitionistic fuzzy number and its applications to transportation problem," *Opsearch*, vol. 52, no. 3, pp. 431–471, Sep. 2015.
- [39] A. Raza, S. Farid, M. Amir, M. Yasir, and R. M. Danyal, "A study on convergence analysis of Adomian decomposition method applied to different linear and non-linear equations," *Interface J. Sci. Eng. Res.*, vol. 11, no. 5, pp. 1358–1378, May 2020.
- [40] S. S. Ray, "New approach for general convergence of the Adomian decomposition method," *World Appl. Sci. J.*, vol. 32, no. 11, pp. 2264–2268, Jan. 2014.
- [41] E. A. Az-Zo'bi and K. Al-Khaled, "A new convergence proof of the Adomian decomposition method for a mixed hyperbolic elliptic system of conservation laws," *Appl. Math. Comp.*, vol. 217, no. 8, pp. 4248–4256, Dec. 2010.
- [42] A. Hassan and Y. Zakari, "Application of first order differential equation in temperature problems," *Appl. Differ. Equ.*, vol. 16, pp. 1–6, Jul. 2018.
- [43] S. Pak, "Solitary wave solutions for the RLW equation by He's semi inverse method," *Interface J. Non-line. Sci. Numer. Sim.*, vol. 10, no. 4, pp. 505–508, Apr. 2009.
- [44] Y. Praroopa, A. S. Reddy, and P. B. Rao, "Review concept of application of differential equations," *Int. J. Eng. Res. Appl.*, vol. 1, no. 3, pp. 34–37, Apr. 2016.
- [45] M. Yavuz and T. Abdeljawad, "Nonlinear regularized long-wave models with a new integral transformation applied to the fractional derivative with power and Mittag-Leffler kernel," *Adv. Difference Equ.*, vol. 2020, no. 1, pp. 1–18, 2020.
- [46] M. Shams, N. Kausar, S. Kousar, D. Pamucar, E. Ozbilge, and B. Tantai, "Computationally semi-numerical technique for solving system of intuitionistic fuzzy differential equations with engineering applications," *Adv. Mech. Eng.*, vol. 13, no. 12, pp. 1–17, Dec. 2022.
- [47] O. S. Fard, "An iterative scheme for the solution of generalized system of linear fuzzy differential equations," *World Appl. Sci. J.*, vol. 7, pp. 1597–1604, Jan. 2009.
- [48] M. Yavuz, "Novel solution methods for initial boundary value problems of fractional order with conformable differentiation," *Int. J. Optim. Control, Theories Appl.*, vol. 8, no. 1, pp. 1–7, Jan. 2018.
- [49] N. Rafiq, N. Yaqoob, N. Kausar, M. Shams, N. A. Mir, Y. U. Gaba, and N. Khan, "Computer-based fuzzy numerical method for solving engineering and real-world applications," *Math. Prob. Eng.*, vol. 2021, pp. 1–13, Nov. 2021.
- [50] M. Shams, N. Kausar, P. Agarwal, S. Momani, and M. A. Shah, "Highly efficient numerical scheme for solving fuzzy system of linear and non-linear equations with application in differential equations," *Appl. Math. Sci. Eng.*, vol. 30, no. 1, pp. 777–810, Dec. 2022.
- [51] M. Yavuz, "Dynamical behaviors of separated homotopy method defined by conformable operator," *Konuralp J. Math.*, vol. 7, no. 1, pp. 1–6, Jan. 2019.
- [52] A. M. Elmoasry Shams, N. Yaqoob, N. Kausar, Y. U. G. Gaba, and N. Rafiq, "Numerical scheme for finding roots of interval-valued fuzzy non-linear equation with application in optimization," *J. Funct. Spaces*, vol. 2021, no. 1, pp. 1–16, Jan. 2021.
- [53] M. Dirik, "Detection of counterfeit banknotes using genetic fuzzy system," *J. Fuzzy Extension Appl.*, vol. 3, no. 4, pp. 302–312, Oct. 2022.



MUDASSIR SHAMS received the Ph.D. degree in computational mathematics (numerical analysis) from Riphah International University, Islamabad, Pakistan, in 2022. His current research interests include computational mathematics, in particular, numerical analysis, computational fluid dynamics, integral inequalities, iterative methods for nonlinear algebraic and transcendental equations, numerical techniques for solving initial value problems, and analytical and numerical techniques for solving fuzzy initial and boundary value problems. He is a reviewer of various international journals.

NASREEN KAUSAR received the Ph.D. degree in mathematics from Quaid-i-Azam University, Islamabad, Pakistan. She is currently an Associate Professor of mathematics with Yildiz Technical University, Istanbul, Turkey. Her research interests include the numerical analysis and numerical solutions of the ordinary differential equations (ODEs), partial differential equations (PDEs), Volterra integral equations, as well as associative and commutative, nonassociative and noncommutative fuzzy algebraic structures and their applications.

KHULUD ALAYYASH received the B.Sc. degree (Hons.) in mathematics from King Khaled University, Saudi Arabia, the M.Sc. degree in applied computation and bio-mathematics from Heriot-Watt University, U.K., and the Ph.D. degree in mathematics of solid mechanics (multiscale modeling, limit states analysis, optimization, finite-element modeling, and structural analysis) from Cardiff University, U.K. She is currently a Treasurer and a member of the Society for Industrial and Applied Mathematics (SIAM) Chapter, Cardiff University (build cooperation between mathematics and the worlds of science and technology through our publications, research, and community). She is also an Assistant Professor.

MOHAMMED M. AL-SHAMIRI received the Ph.D. degree in geometric topological algebra from Menoufia University, Menoufia, Egypt, in 2008. He is currently an Associate Professor of mathematics with King Khalid University, Saudi Arabia. He is also an Associate Professor with Ibb University, Yemen. His research interests include topological spaces, topological geometry, fuzzy topology, graph and knot and fuzzy graph and fuzzy knot, geometrical transformations (folding, retraction, and deformation retract), fuzzy group, fuzzy ring, fuzzy module, fuzzy field, and fuzzy decision support systems.



NAYYAB ARIF thesis work focused on analytical techniques for solving generalized trapezoidal intuitionistic fuzzy initial value problems. She is about to enroll in the Ph.D. degree. Her research interests include fuzzy set theory and its applications. She researches analytical methods which solve fuzzy initial value problems.



RASHAD ISMAIL received the Ph.D. degree in scientific computing from Assiut University, Egypt, in 2011. He is currently an Assistant Professor with King Khalid University, Saudi Arabia. He is also an Assistant Professor of scientific computing with Ibb University, Yemen. His research interests include graph theory and its applications, graph domination, meta-heuristics, data mining, computer networks, fuzzy, fuzzy graph, and fuzzy decision support systems.

...