

Chapman University

Chapman University Digital Commons

Sociology Faculty Articles and Research

Sociology

3-30-2023

Obesity Heterogeneity by Neighborhood Context in a Largely Latinx Sample

Ashley W. Kranjac

Chapman University, kranjac@chapman.edu

Dinko Kranjac

University of La Verne, dinko@laverne.edu

Zeev N. Kain

University of California, Irvine, zkain@hs.uci.edu

Louis Ehwerhemuepha

Children's Hospital of Orange County

Brooke N. Jenkins

Chapman University, bjenkins@chapman.edu

Follow this and additional works at: https://digitalcommons.chapman.edu/sociology_articles



Part of the [Inequality and Stratification Commons](#), [Latina/o Studies Commons](#), [Medicine and Health Commons](#), [Other Public Health Commons](#), [Pediatrics Commons](#), [Place and Environment Commons](#), and the [Race and Ethnicity Commons](#)

Recommended Citation

Kranjac, A.W., Kranjac, D., Kain, Z.N. et al. Obesity Heterogeneity by Neighborhood Context in a Largely Latinx Sample. *J. Racial and Ethnic Health Disparities* (2023). <https://doi.org/10.1007/s40615-023-01578-6>

This Article is brought to you for free and open access by the Sociology at Chapman University Digital Commons. It has been accepted for inclusion in Sociology Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Obesity Heterogeneity by Neighborhood Context in a Largely Latinx Sample

Comments

This article was originally published in *Journal of Racial and Ethnic Health Disparities* in 2023.
<https://doi.org/10.1007/s40615-023-01578-6>

Creative Commons License




This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Copyright

The authors



Obesity Heterogeneity by Neighborhood Context in a Largely Latinx Sample

Ashley W. Kranjac^{1,2} · Dinko Kranjac³ · Zeev N. Kain^{2,4,5} · Louis Ehwerhemuepha⁶ · Brooke N. Jenkins^{2,4,7} 

Received: 7 December 2022 / Revised: 27 February 2023 / Accepted: 19 March 2023
© The Author(s) 2023

Abstract

Neighborhood socioeconomic context where Latinx children live may influence body weight status. Los Angeles County and Orange County of Southern California both are on the list of the top ten counties with the largest Latinx population in the USA. This heterogeneity allowed us to estimate differential impacts of neighborhood environment on children's body mass index *z*-scores by race/ethnicity using novel methods and a rich data source. We geocoded pediatric electronic medical record data from a predominantly Latinx sample and characterized neighborhoods into unique residential contexts using latent profile modeling techniques. We estimated multilevel linear regression models that adjust for comorbid conditions and found that a child's place of residence independently associates with higher body mass index *z*-scores. Interactions further reveal that Latinx children living in Middle-Class neighborhoods have higher BMI *z*-scores than Asian and Other Race children residing in the most disadvantaged communities. Our findings underscore the complex relationship between community racial/ethnic composition and neighborhood socioeconomic context on body weight status during childhood.

Keywords Children · Electronic medical records · Latent profile analysis · Latinx · Neighborhoods · Obesity

Introduction

Childhood obesity is a major public health challenge in the USA [1]. The prevalence of obesity among children aged 2–19 varies by race/ethnicity, and the number of Latinx children and adolescents who are presenting with obesity is increasing

[2–4]. Latinx youth represent one-quarter of the entire population of children under the age of 18 in the USA and are disproportionately burdened by obesity [5, 6]. A continuing rise in the prevalence of obesity among Latinx children is a serious concern because an excess of body fat is associated with adverse effects on health that may persist into adulthood such as hypertension, prediabetes and diabetes mellitus, sleep apnea, asthma, and depression [1, 6–8]. Although the prevalence varies among Latinx Americans of different ethnic backgrounds regardless of origin group, Latinx adults are more likely to be diagnosed with these medical conditions, as compared to their non-Latinx White counterparts [9–13].

Unhealthy weight gain develops from a chronic, positive energy balance through an interplay of genetic, biological, behavioral, socioeconomic, and environmental factors [14–16]. For example, underlying medical causes such as poisoning, infections, respiratory or heart disease, and immune system, central nervous system, or endocrine system dysfunction may lead to obesity [17–23]. Also, children from low socioeconomic status (SES) families are more likely to be diagnosed with obesity, and, predictably, this is true among Latinx youth in general and in California, specifically [24, 25]. Relatedly, the consumption of high-calorie foods with little-to-no nutritional value and the

✉ Brooke N. Jenkins
bjenkins@chapman.edu

¹ Department of Sociology, Chapman University, Orange, CA, USA

² Center for Stress & Health, University of California School of Medicine, Irvine, CA, USA

³ Department of Psychology, Institute for Mental Health and Psychological Well-Being, University of La Verne, La Verne, CA, USA

⁴ Department of Anesthesiology and Perioperative Care, University of California, Irvine, CA, USA

⁵ Yale Child Study Center, Yale University, New Haven, CT, USA

⁶ Computational Research, Children's Health of Orange County, Orange, CA, USA

⁷ Department of Psychology, Chapman University, One University Drive, Orange, CA 92866, USA

chronic psychosocial stress brought on by having limited economic resources influence weight trajectories among Latinx children [26–28]. Moreover, the neighborhood socioeconomic environment where Latinx children live influences body weight status, and neighborhoods with predominately racial/ethnic minority residents have higher levels of poverty and lower levels of education, which are independently associated with childhood obesity [29–35]. Notably, data indicate that living in either a low-SES household or a low-SES neighborhood is enough to increase the risk of child obesity [36]. Multiple domains of the social neighborhood environment also are relevant to the development of overweight and obesity [37]. For example, social cohesion and collective socialization and trust in Latinx communities are some of the more pertinent factors because they can impact norms regarding diet and physical activity [38]. Regarding the home and family environment, the parents of Latinx youth tend to underestimate their child’s weight status and perceive their child as “normal” weight even when the child is classified as overweight or obese according to body mass index (BMI) categories [39]. Taken together, these multiple influences enhance the risk for obesity among children in general, as well as in Latinx children specifically, a population where prevalence is increasing especially rapidly [2–6].

Obesity is a multifactorial disease, and thus, we need to appreciate fully the relevance of how socioenvironmental factors like residential context combine to associate with increased body weight. As but a single example, findings from the *Moving to Opportunities Study* conducted by the Department of Housing and Urban Development indicate that relocation to a low-poverty census tract led to lower obesity prevalence [40]. One way to address the significance of multifactorial neighborhood-level determinants of obesity is to use novel methods on unique populations to draw out the extent to which distinctive residential contexts give rise to social patterning that produces variation in obesity prevalence [31, 32]. Indeed, our data source allows us to overcome some of the methodological challenges plaguing the current literature, including an objective collection of height/weight measures, access to medical diagnoses, adequate sample size and number of children per residential tract, and racial/ethnic diversity [41]. Specifically, we geocoded pediatric medical record data for residents in Southern California, a geographical area with the largest Latinx population in the USA [42]. Moreover, as a geographic setting, Los Angeles County has the nation’s largest Latinx population, while Orange County ranks tenth among the counties with the largest Latinx population in the USA [42]. We took advantage of this heterogeneity to estimate the differential impacts of neighborhood context on children’s BMI z -scores. First, we explored the relationship between neighborhood context and BMI z -scores. Given past work demonstrating an association between neighborhood socioeconomic status and children’s

body weight, we expected that neighborhood disadvantage would associate with heightened BMI z -scores. In addition, we expected race/ethnicity would moderate the association between neighborhood context and BMI z -scores, and that Latinx children will have higher body weight relative to other race/ethnicities irrespective of contextual factors.

Methods

Data Sources

We use a compilation of data from multiple sources. We extracted electronic medical records (EMR) data from children ($n = 53,735$) between the ages of 0 and 18 admitted between June 2013 and June 2018 as inpatients across a variety of care settings. EMR data were excluded for pediatric cancer patients ($n = 1319$), those without complete body mass index (BMI) measures ($n = 3921$), and outlier patients for the length of stay ($n = 529$) [40]. Patient addresses from the remaining EMR data were geocoded and linked to the corresponding residential Census tract. To capture the appropriate time period for analysis, neighborhood measures were assigned to children temporally by first taking the child’s address from the electronic medical record at the time of their height and weight measurement. Then, we used 5-year ACS estimates for the census tracts which “surround” the timing of the child’s records. In this way, the 5-year ACS estimates characterize the child’s area within that 5-year period. As previously done, we used Census tracts to represent neighborhoods [43]. Social and economic indicators were extracted from the 2014–2018 American Community Survey (ACS) data. We excluded observations unmatched to Federal Information Processing Series (FIPS) Codes ($n = 7610$), those living in Census tracts with fewer than 20 children per tract ($n = 32,234$), and children living outside of California ($n = 90$), resulting in a total sample size of 8092 children nested within 672 census tracts or neighborhoods.

Variables

The key outcome is age- and sex-specific BMI z -score (i.e., the number of standard deviation (SD) units that the child’s BMI deviates from the age- and sex-normed mean reference value, based on the 2000 Center for Disease Control (CDC) Growth Charts: USA) [44, 45]. We included all covariates available to us from the EMR to represent the child and familial characteristics. Sociodemographic characteristics include age at the time of visit, sex (1, girl; ref.; 2, boy), race/ethnicity (1, Latinx, ref.; 2, non-Latinx White; 3, non-Latinx Black; 4, Asian; 5, Other Race), and insurance type (1, Private, ref.; 2, Public (e.g., Medicare or MediCal)) as a proxy for socioeconomic status (SES) [31, 46–48]. Other measures include

admission type (1, elective, ref.; 2, emergency; 3, other (i.e., urgent care or accident)); and diagnosis based on ICD9/10 codes (0, absence of diagnosis; 1, presence of diagnosis), including bacterial and viral infection (A00-A99), diseases of the blood and blood-forming organs, and disorders involving immune mechanisms (D50-D89), endocrine, nutritional and metabolic diseases (E00-E89), ref., diseases of the circulatory system (I00-I99), nervous system diseases (G00-G99), congenital malformations, deformations and chromosomal abnormalities (Q00-Q99), diseases of the digestive system (K00-K95), diseases of the genitourinary system (N00-N99), diseases of the musculoskeletal system (M00-M99), diseases of the respiratory system (J00-J99), injury, poisoning, and certain other external consequences (S00-T88), and other diagnoses (H00-H59, L00-L99, O00-O9, P00-P96, Z00-Z99). We also included length of stay (mean = 3.82; Std. Dev. = 5.95) and patient maximum pain assessed by healthcare providers throughout the child's hospital stay using several developmentally and situationally appropriate measurement tools, including faces, legs, activity, cry, and consolability scale, faces pain scale, numeric rating scale, and neonatal pain, agitation and sedation scale (range from 0, no pain to 10, severe pain). Social and economic indicators of the child's neighborhood of residence come from the ACS and include community-level education (% adults with < 12 years of education; % adults with 12 years of education; % adults with > 12 years and < 16 years of education; % adults with 16 years of education; % adults with 18 years of education; % adults with > 18 years and < 21 years of education), median community-level income, percent of female-headed households, percent receiving public assistance, percent in poverty, percent of homes in the tract that are rented, and racial/ethnic composition measured by percent of major racialized categories (% non-Latino White; % non-Latino Black; % Latinx; % Asian; % Other Race). After testing the cross-level assumption that the random effect of the intercept is correlated with a level 1 measure, using a Hausman test, we uncovered that some unobservable neighborhood characteristics relegated to the error term are correlated with some observable patient characteristics. We tested for correlations between neighborhoods and patient-level characteristics and found that the offending explanatory measures were race/

ethnicity and insurance type. Then, we computed neighborhood aggregates of these patient-level characteristics (i.e., means of race/ethnicity and insurance type) to control for neighbor effects and included these measures at level 2 in our fully specified models [49].

Statistical Analysis

We first used latent profile analysis (LPA) to group neighborhoods into clusters based on the above-listed socioeconomic indicators extracted from the ACS, which are often used to define a child's neighborhood of residence [50, 51]. Using MPlus 8.8 software, we estimated a 1-profile model in a type I covariance structure and fit successive models with an increasing number of profiles to characterize neighborhoods [52, 53]. We used entropy and theoretically driven evidence to select the most appropriate number of profiles and further evaluated model fit with AIC, BIC, a-BIC, and loglikelihood values to identify the most parsimonious model. Specifically, our substantive neighborhood cluster interpretations are based on theoretical neighborhood stratification observed across the USA (i.e., upper class, middle class, working class, and disadvantaged communities). The stability of our 4-profile solution was verified by model fit statistics wherein we stopped adding profiles once the model fit indicators began to increase and the entropy levels began to decrease (Table 1). Given our data, analyses indicated that neighborhoods are most appropriately captured by a 4-profile solution (Table 1). Figure 1 displays the patterning of neighborhood-level characteristics by LPA-generated neighborhood profiles. We standardized all measures for this figure by converting median household income to the ratio of income to poverty using 2016-inflation adjusted dollars for a 3-person household.


We then test the influence of distinct neighborhoods on children's BMI z-scores using multilevel linear regression modeling with Stata 16.1 software [54]. This technique treats level-1 children as nested within level-2 neighborhoods and neutralizes the lack of independence of data within higher groups. Our modeling approach uses adaptive quadrature to adjust for problems that otherwise downwardly bias estimated standard errors, including different sample sizes for level-1 and level-2 units, clustering within

Table 1 Model fit & usefulness information for LPAs with 1–5 latent profiles

Classes	AIC	BIC	a-BIC	LL	LL <i>p</i> -value	Entropy
1	2949364.575	2949654.352	2949539.946	-1474646.29	0.000	1.00
2	2866031.981	2866474.696	2866299.908	-1432960.99	0.000	0.93
3	2816267.306	2816862.958	2816627.789	-1408059.65	0.000	0.92
4	2781335.073	2782083.663	2781788.113	-1390574.54	0.000	0.95
5	2815388.178	2816289.707	2815933.774	-1407582.09	0.333	0.91

Source: Data are from electronic medical records and the 2014–2018 American Community Survey. *Note:* Bolded row indicates the most parsimonious model

Fig. 1 Descriptive neighborhood-level characteristics by LPA-generated neighborhood profiles



neighborhoods, variable numbers of cases within level-2 units, and heteroscedastic error terms [55]. We performed a series of conditional models that first included the covariates of a child- and family-level predictors (age, sex, race/ethnicity, insurance type, admission type, diagnosis, length of stay, patient-reported pain levels) to test the influence of child and family factors on children's BMI z -scores. In the next set of models, we included the LPA-constructed neighborhoods and neighborhood averages of race/ethnicity and insurance status at the neighborhood level (and a neighborhood-level error component) along with the child- and family-level predictors and an individual error term. In our last set of models, we included interactions between the LPA-constructed neighborhoods and race/ethnicity at the child/family level to determine whether race/ethnicity moderates the association between neighborhood disadvantage and the expected heightened BMI z -scores for Latinx children.

Results


We display how the four neighborhood profiles cluster across the area of study in Fig. 2. We assigned the following labels: Advantaged, Middle-Class, Working-Class, and Disadvantaged based on the descriptive characteristics. The advantaged neighborhood profile is largely concentrated in east Orange County, north-west Riverside County, and north-west San Diego County regions. As shown in Table 2, Advantaged neighborhoods have the highest median household income (\$130,461), the highest overall levels of education (31% of residents have at least 16 years of education), and the lowest percentage of people living in poverty (8%). The Disadvantaged communities make up the areas

around north-west Orange County and central Riverside County regions and score the worst on nearly every indicator. These communities have the lowest median household income (\$41,470), the lowest education levels (48% of adult residents lack a high school degree), and the highest proportion of the population in poverty (29%).

We show descriptive information for the child and family characteristics overall, and by neighborhood context in Table 3. The average overall BMI is 0.38 SDs above the national reference. However, the majority of children fall at or below the 67th percentile, or in the "Healthy Weight" category. Children living in Advantaged neighborhoods have 0.02 SDs below the mean while those residing in Disadvantaged neighborhoods have 0.65 SDs above the mean. The mean age for the entire sample was 9.80 years, with children in Advantaged neighborhoods slightly older than those in other neighborhood types. In Fig. 3, we show that, although there is some clustering of children of specific race/ethnicity across neighborhoods, each racial/ethnic group is represented within each neighborhood. As displayed in Table 3, children admitted to the emergency department and presenting with diseases of the digestive system are slightly over-represented in Disadvantaged neighborhoods compared to those residing in other neighborhoods.

Results from our multilevel linear regression models predicting BMI z -scores are shown in Table 4. Model 1 partitions the total variation in BMI z -scores into within- and between-neighborhood variance. Model 2 only included the child/family-level characteristics (age, sex, race/ethnicity, insurance type, admission type, diagnosis, length of stay, patient-reported pain levels). Model 3 includes the LPA-generated neighborhoods. Fully specified Model 4 includes the interaction terms

Fig. 2 Neighborhood profiles by census tracts, Southern California



by race/ethnicity and neighborhood profiles. The random effects estimated across all models indicate significant variation in pediatric BMI z -scores across neighborhoods.

In Model 1, intraclass correlation from an unconditional model reveals that around 20% of the variation in children's BMI z -scores is attributed between neighborhoods. In Model 2, we see that non-Latinx white (-0.32 ; 95% CI: $-0.44 - -0.19$), non-Latinx black (-0.34 ; 95% CI: $-0.58 - -0.09$), and Asian children (-0.56 ; 95% CI: $-0.72 - -0.39$), relative to Latinx, have lower BMI z -scores. Each additional year in age significantly increased BMI z -scores by 0.03 SD units (95% CI: 0.02–0.03). Publicly insured children (0.15; 95% CI: 0.05–0.26) and those with diseases of the genitourinary system (0.19; 95% CI: 0.04–0.35) and higher pain levels (0.04; 95% CI: 0.03–0.05) have higher BMI z -scores, whereas children with longer lengths of stay (-0.01 ; 95% CI: $-0.02 - -0.01$), congenital malformations, deformations and chromosomal abnormalities (-0.33 ; 95% CI: $-0.46 - -0.20$), and diseases of the musculoskeletal system and connective tissues (-0.15 ; 95% CI: $-0.28 - -0.01$) have lower BMI z -scores. In Model 3, accounting for neighborhood profiles slightly attenuates the racial/ethnic and socioeconomic differences in BMI z -scores indicating that some of the lower BMI z -scores for non-Latinx white, non-Latinx black, and Asian children, relative to Latinx, and higher BMI z -scores for publicly insured children, are due to the neighborhood environment. Model 3 largely mirrors Model 2 in other child/family-level factors and further shows that higher

levels of neighborhood disadvantage independently associate with higher BMI z -scores. More specifically, children housed in Disadvantaged neighborhoods have 0.19 SDs higher BMI (95% CI: 0.02–0.36) compared to those living in Middle-Class communities. Finally, we examined whether race/ethnicity moderates the influence of neighborhood context on BMI z -scores. Model 4 indicates an interaction between neighborhoods and race/ethnicity. Specifically, we find that Latinx children living in Middle-Class neighborhoods have higher BMI z -scores than Asian (-0.79 ; 95% CI: $-1.47 - -0.12$) and Other Race (-0.47 ; 95% CI: $-0.89 - -0.05$) children living in the most disadvantaged communities.

Discussion

This is the first study to date that employed LPA to detail how neighborhood context matters differently for children's body weight status in a predominantly Latinx sample. We extracted EMR data from pediatric patients living in Southern California, a geographic setting with the largest Latinx population in the USA [42]. Importantly, we controlled for comorbid conditions like endocrine, nutritional, and metabolic diseases, as well as chromosomal, cardiovascular, pulmonary, neurological, musculoskeletal, and digestive system disorders that otherwise would bias estimates [7, 14–23]. Specifically, we generated neighborhood typologies based on social and economic indicators extracted from the ACS

Table 2 Descriptive neighborhood-level characteristics by LPA-generated neighborhood profiles

	Advantaged		Middle-Class		Working-Class		Disadvantaged		Diff
		95% CI		95% CI		95% CI		95% CI	
Socioeconomic proportions									
Median income (in \$10 K)	130.46	(128.64–133.32)	79.83	(78.86–81.36)	56.78	(56.14–57.78)	41.47	(40.67–42.72)	<.001
% adults < 12 years of education	4.83	(4.70–5.04)	6.96	(6.41–6.97)	24.54	(24.32–25.95)	47.63	(47.28–48.96)	<.001
% adults = 12 years of education	9.23	(9.19–9.67)	20.96	(20.62–21.44)	28.10	(27.89–28.48)	25.00	(24.69–25.49)	<.001
% adults > 12 and < 16 years of education	21.05	(21.02–21.86)	36.73	(36.27–36.94)	28.04	(29.05–31.05)	16.30	(16.18–20.04)	<.001
% adults = 16 years of education	36.25	(36.03–36.60)	22.06	(21.66–22.67)	12.05	(11.78–12.48)	6.30	(6.26–6.72)	<.001
% adults = 18 years of education	18.86	(18.58–19.31)	8.25	(8.08–8.52)	4.07	(3.95–4.24)	2.23	(2.15–2.34)	<.001
% adults > 18 and < 21 years of education	4.74	(4.59–4.97)	2.83	(2.77–2.92)	1.70	(1.65–1.78)	1.37	(1.29–1.49)	<.001
% adults = 21 years of education	5.04	(4.92–5.23)	2.21	(2.16–2.29)	1.50	(1.44–1.60)	1.17	(1.16–1.42)	<.001
% receiving public assistance	3.98	(3.87–4.16)	7.31	(7.11–7.62)	15.29	(14.96–15.82)	25.09	(24.54–26.21)	<.001
% female-headed households	5.02	(4.92–5.19)	7.19	(7.05–7.42)	12.19	(11.94–12.57)	17.43	(17.01–18.10)	<.001
% of residents in poverty	7.84	(7.61–8.20)	10.60	(10.35–10.99)	17.96	(17.59–18.53)	29.03	(28.30–30.18)	<.001
% of homes renter occupied	35.68	(34.81–37.06)	64.66	(35.72–37.67)	51.93	(51.13–53.17)	66.46	(65.43–68.09)	<.001
Racial/ethnic composition									
% Non-Latinx White	67.58	(67.00–68.50)	58.39	(57.81–59.29)	31.41	(30.76–34.43)	20.11	(20.38–23.26)	<.001
% Non-Latinx Black	4.17	(4.54–5.09)	4.48	(4.17–4.95)	5.72	(5.32–5.35)	5.69	(5.26–5.37)	<.01
% Latinx	1.79	(1.52–1.22)	14.14	(13.67–14.89)	40.73	(40.43–46.77)	55.93	(50.67–56.37)	<.001
% Asian	21.89	(21.37–22.72)	14.96	(14.52–15.64)	12.82	(12.35–13.56)	7.73	(6.89–7.93)	<.001
% Other Race	4.47	(4.32–4.70)	8.22	(8.02–8.55)	9.45	(9.72–9.60)	10.58	(10.89–15.65)	<.001
Neighborhoods <i>n</i> =	94		198		218		162		
Children <i>n</i> =	1159		1749		2265		2838		

Source: Data are from the 2013–2018 electronic medical records and the 2014–2018 American Community Survey (ACS)

Significance is evaluated using one-way MANOVA with the neighborhood variables as the dependent variables and LPA neighborhood type as the independent variable

and found that the residential socioeconomic environment is associated with heightened BMI *z*-score. Our finding is consistent with published studies, but other researchers failed to account for underlying diagnoses that could introduce bias. Thus, in line with our expectations, using a diverse sample and accounting for important comorbid conditions, we show that a child's neighborhood of residence independently associates with higher BMI *z*-scores. This could be driven by the overall higher incidence of overweight and obesity in the USA among Latinx youth, combined with compounding neighborhood-level determinants such as the absence of a health-promoting infrastructure that led to the greater prevalence of higher BMI [2–4, 31–37].

Our findings support previous reports that race/ethnicity moderates the association between LPA-constructed neighborhoods and children's body weight. Specifically, here, we found that Latinx children living in more advantaged neighborhoods had higher BMI *z*-scores than Asian and Other children residing in the most disadvantaged communities. These results are troublesome given that a wealth of literature supports the notion that with affluence a child's risk of

presenting with obesity declines. Vis-à-vis individual-level indicators, the variation in children's body weight associates with age, sex, race/ethnicity, SES, physical activity level, screen-time viewing, and sleep duration/bedtime [56–59]. Family-level, obesity-relevant factors include family physical activity, family mealtimes, food insecurity, and the income-to-need ratio [60–62]. The relationship among individual characteristics, in-home practices, and child obesity, however, is complicated by the influence of neighborhood-level descriptors [31–37]. For example, neighborhood median income modifies the association between a child's BMI *z*-score and proximity to fast-food restaurants [63]. More specifically, the proximity to "unhealthy" food establishments has a stronger adverse effect on body weight status in lower-income neighborhoods, compared to the effect on residents of more affluent communities [63]. Thus, we need to optimize and implement a range of long-term, multicomponent intervention programs that focus on the early prevention of obesity. For example, a 3-year-long multilevel approach to reduce obesity among low-income, primarily Latinx children in Northern California, did show some

Table 3 Means and standard deviations (SD) for independent and dependent variables overall and by neighborhood profile

	Overall		Advantaged		Middle-Class		Working-Class		Disadvantaged		Sig
	Mean or %	SD	Mean or %	SD	Mean or %	SD	Mean or %	SD	Mean or %	SD	
Dependent variable											
BMI z-scores	0.38	1.49	-0.02	1.31	0.17	1.65	0.41	1.49	0.65	1.41	<.01
Independent variables											
Sociodemographic											
Age	9.80	4.90	10.34	4.83	9.44	5.01	9.77	4.82	9.86	4.91	<.01
Sex											
Girl	48.00	0.50	40.38	0.49	49.80	0.50	48.96	0.50	50.00	0.50	
Boy	52.00	0.50	59.62	0.49	50.20	0.50	51.04	0.50	50.00	0.50	
Race/ethnicity											
Latinx	56.90	0.50	9.58	0.29	30.65	0.46	69.58	0.46	82.70	0.38	<.001
Non-Latinx White	15.24	0.36	33.91	0.47	28.30	0.45	7.68	0.27	4.97	0.22	<.001
Non-Latinx Black	3.42	0.18	3.80	0.19	4.46	0.21	5.87	0.24	0.70	0.08	
Asian	9.76	0.30	19.50	0.40	19.56	0.40	6.53	0.25	2.40	0.15	<.001
Other Race	14.68	0.35	33.22	0.47	17.04	0.38	10.33	0.30	9.23	0.27	<.05
Health insurance											
Private	28.92	0.45	77.57	0.42	47.28	0.49	15.89	0.37	7.61	0.27	<.001
Public	71.08	0.45	22.43	0.42	52.72	0.49	84.11	0.37	92.39	0.27	<.001
Patient characteristics											
Admission type											
Elective	32.13	0.47	36.84	0.48	43.97	0.50	28.52	0.45	25.33	0.44	<.05
Emergency	35.15	0.48	28.90	0.45	27.90	0.45	37.75	0.49	40.35	0.49	<.05
Other	32.72	0.49	34.25	0.47	28.13	0.45	33.73	0.47	34.32	0.47	
Diagnosis											
Bacterial and viral infection	10.58	0.31	10.09	0.30	8.75	0.28	11.79	0.32	11.03	0.31	<.05
Diseases of the blood	15.90	0.37	17.69	0.38	19.44	0.40	17.84	0.38	11.56	0.32	<.001
Endocrine and metabolic	17.47	0.38	15.53	0.36	21.10	0.41	18.37	0.39	14.69	0.35	<.001
Circulatory system	6.59	0.25	6.13	0.24	6.40	0.24	7.64	0.27	6.10	0.24	
Nervous system	16.82	0.37	17.95	0.38	20.81	0.41	17.00	0.38	13.78	0.34	<.001
Congenital and chromosomal	11.96	0.32	11.22	0.32	13.21	0.34	11.52	0.32	11.95	0.32	
Digestive system	19.43	0.40	15.70	0.36	16.87	0.37	20.22	0.40	21.92	0.41	<.001
Genitourinary system	6.06	0.24	6.04	0.24	4.75	0.21	6.75	0.25	6.38	0.24	
Musculoskeletal system	9.03	0.29	8.46	0.28	8.06	0.27	11.08	0.31	8.28	0.28	<.01
Respiratory system	17.85	0.38	17.00	0.38	18.07	0.39	18.59	0.39	17.62	0.38	
Trauma, injury, poisoning	9.56	0.29	9.92	0.30	6.23	0.24	10.82	0.31	10.54	0.31	<.001
Other diagnoses	35.26	0.48	39.78	0.49	31.79	0.47	34.75	0.48	35.66	0.48	<.01
Length of stay	3.67	5.35	3.44	5.02	3.27	4.99	3.93	5.60	3.84	5.47	<.01
Pain-level	4.48	3.51	4.12	3.49	4.01	3.42	4.74	3.58	4.68	3.47	<.01

Source: Data are from the 2013–2018 electronic medical records and the 2014–2018 American Community Survey (ACS)


Asterisks indicate significance difference evaluated using two-tailed independent means *t*-test

promise for reducing weight gain [64]. Also, it is noteworthy that combined nutrition–physical activity initiatives, like *The California Endowment's Healthy Eating, Active Communities* program, can be effective at preventing and reducing childhood obesity in low-income communities [65].

Published studies do offer insight into community-level factors that enhance susceptibility to obesity among children from socially vulnerable groups. For example, neighborhood

safety influences the amount of physical activity and screen time [66, 67]. Also, poor adherence to physical activity recommendations among low-income Latinx women is a predictor of excessive gestational weight gain, which is independently associated with infant birthweight percentile [68, 69]. With regard to children, racial/ethnic differences exist across different groups in terms of physical activity and sedentary behaviors overall, and Latinx, compared to

Fig. 3 Distribution of children across LPA neighborhood contexts by race/ethnicity



non-Latinx Black and non-Latinx White, children were found to participate in the least physical activity [70–73]. Crucially, the neighborhood context does impact girls and boys in a different way in terms of health benefits [31]. Relatedly, similar findings have been reported with regard to race/ethnicity. In Los Angeles County, for example, WIC-participating children across levels of neighborhood poverty are less likely to develop obesity, yet the prevalence of obesity remains higher among Latinx WIC-enrolled children compared to those who are non-Latinx White, non-Latinx Black, or Asian [74, 75]. Additionally, the neighborhood's social and cultural fabric is yet another obesogenic factor. Indeed, the interplay of social group norms and networks like peers, school teachers, and digital mass media exposure informs acceptable ranges of body size and contributes to weight gain among Latinx youth [76–78]. Overall, data show that there is an immense disparity in the rates of pediatric overweight and obesity among racial/ethnic groups [4, 79]. Hence, intervention programs and public policies should include a multicomponent approach that targets parental behaviors, family patterns, household conditions, and the built and social infrastructure of communities [64, 65].

Our study, though informative, is not without limitations. We are constrained by the limited individual- and family-level variables available in the EMR data that are typically used for administrative purposes. For example, prior work indicates that children of immigrant parents present with higher levels of overweight, but we do not have information on immigration status nor origin group [80, 81]. Still, we included all covariates available to us in the EMR data

known to associate with children's body weight. Moreover, we use a public–private insurance dichotomization as a proxy measure for SES. We acknowledge that it is not ideal to use insurance type as a proxy for SES, but publicly provided health care coverage like Medicaid is only obtainable by children who meet stringent income criteria, with the exception of those with certain medical conditions [82]. Still, insurance coverage does have reasonable validity and reliability when used as a marker for individual-level SES [31, 32, 46–48]. Other scholars should test the reliability of our findings by using a more comprehensive set of individual- and family-level covariates known to associate with children's body weight, including but not limited to nativity status, cultural differences, or parental education and income. Similarly, the cross-sectional nature of our data that are restricted to Los Angeles County and Orange County in Southern California limits the scope of our analyses. Nonetheless, this geographic setting has the largest Latinx population in the USA, which provides a unique opportunity to estimate the differential impacts of comprehensive neighborhood profiles on BMI *z*-scores in a diverse group of children [42]. Also, we used census tracts to represent neighborhoods. Census tracts are by no means a perfect operationalization of residential environments, but they remain a useful spatial entity available to us in the approximation of a neighborhood [30–32, 43]. Certainly, using addresses instead of zip codes provides a more robust spatial unit of analysis [37]. Finally, we examined the influence of distinct neighborhoods on children's BMI *z*-scores. Although BMI is a simple measure, BMI categories are reasonably good

Table 4 Multilevel linear regression models predicting BMI z -scores; $N=8011$ in 672 census tracts

	Model 1		Model 2		Model 3		Model 4	
	Coeff	95% CI	Coeff	95% CI	Coeff	95% CI	Coeff	95% CI
Intercept	0.36***	0.29–0.42	0.09	–0.07–0.26	–0.05	–0.35–0.73	–0.14	–0.43–0.16
Patient-level								
Sociodemographic								
Age			0.03***	0.02–0.03	0.03***	–0.14–0.02	0.03***	0.02–0.03
Sex (girl, ref)								
Boy			–0.07	–0.15–0.01	–0.06	–0.14–0.02	–0.06	–0.14–0.02
Race/ethnicity (Latinx, ref)								
Non-Latinx White			–0.32***	–0.44––0.19	–0.30***	–0.46––0.13	–0.15	–0.45–0.15
Non-Latinx Black			–0.34**	–0.58––0.09	–0.39**	–0.69––0.08	–0.25	–0.83–0.33
Asian			–0.56***	–0.72––0.39	–0.48***	–0.70––0.27	–0.23	–0.65–0.20
Other Race			–0.10	–0.22–0.02	–0.07	–0.23–0.10	0.11	–0.22–0.45
Patient characteristics								
Health insurance (private provider, ref)								
Public provider			0.15**	0.05–0.26	0.10*	–0.00–0.22	0.09	–0.03–0.21
Admission type (elective, ref)								
Emergency			0.04	–0.06–0.15	0.04	–0.06–0.14	0.04	–0.06–0.14
Other			–0.10	–0.15–0.07	–0.05	–0.15–0.06	–0.05	–0.15–0.06
Diagnosis (endocrine, nutritional, and metabolic diseases, ref.)								
Bacterial and viral infection			0.07	–0.05–0.19	0.07	–0.05–0.20	0.08	–0.04–0.20
Diseases of the blood			–0.12	–0.24–0.01	–0.11	–0.23–0.01	–0.11	–0.23–0.01
Circulatory system			0.02	–0.13–0.18	0.03	–0.13–0.19	0.04	–0.12–0.19
Nervous system			0.05	–0.07–0.16	0.08	–0.04–0.19	0.08	–0.04–0.20
Congenital and chromosomal			–0.33***	–0.46––0.20	–0.35***	–0.48––0.21	–0.35***	–0.48–0.22
Digestive system			–0.02	–0.12–0.07	–0.02	–0.12–0.07	–0.04	–0.13–0.06
Genitourinary system			0.19*	0.04–0.35	0.16*	0.01–0.32	0.16*	0.01–0.32
Musculoskeletal system			–0.15*	–0.28––0.01	–0.14*	–0.28––0.01	–0.15*	–0.28––0.01
Respiratory system			–0.09	–0.20–0.02	–0.09	–0.19–0.02	–0.09	–0.19–0.02
Trauma, injury, poisoning			0.05	–0.09–0.19	0.05	–0.09–0.19	0.06	–0.08–0.19
Other diagnoses			0.05	–0.05–0.15	0.05	–0.05–0.14	0.05	–0.05–0.14
Length of stay			–0.01***	–0.02––0.01	–0.01***	–0.02––0.01	–0.01**	–0.02––0.01
Pain levels			0.04***	0.03–0.05	0.04***	0.02–0.05	0.03***	0.02–0.05
Neighborhood-level								
Neighborhood (Middle-Class, ref)								
Advantaged					–0.21	–0.42–0.01	0.01	–0.43–0.45
Working-Class					0.06	–0.12–0.23	0.12	–0.12–0.35
Disadvantaged					0.21*	0.01–0.40	0.35**	0.11–0.60
Neighborhood average race/ ethnicity					0.07	–0.01–0.15	0.05	–0.37–0.13
Neighborhood average health insurance					0.05	–0.23–0.33	0.07	–0.21–0.35
Interaction effects (Latinx*Middle-Class, ref.)								
Non-Latinx White*Advantaged							–0.28	–0.82–0.26
Non-Latinx White*Working- Class							–0.20	–0.62–0.23
Non-Latinx White*Disadvantaged							–0.10	–0.56–0.36
Non-Latinx Black*Advantaged							–0.27	–1.22–0.68
Non-Latinx Black*Working- Class							–0.16	–0.90–0.58

Table 4 (continued)

	Model 1		Model 2		Model 3		Model 4	
	Coeff	95% CI	Coeff	95% CI	Coeff	95% CI	Coeff	95% CI
Non-Latinx							0.10	−0.92–1.11
Black*Disadvantaged								
Asian*Advantaged							−0.31	−0.95–0.34
Asian*Working-Class							−0.17	−0.71–0.37
Asian*Disadvantaged							−0.76*	−1.43–−0.08
Other Race*Advantaged							−0.27	−0.82–0.29
Other Race*Working-Class							0.04	−0.39–0.46
Other Race*Disadvantaged							−0.44*	−0.86–−0.01
Random effects								
Intercept	0.43***	0.36–0.51	0.26***	0.20–0.33	0.42***	0.34–0.51	0.41***	0.34–0.50
BMI z-scores	1.77***	1.71–0.82	1.79***	1.73–1.87	1.65***	1.58–1.72	1.64***	1.58–1.71

Source: Data are from the 2013–2018 electronic medical records and the 2014–2018 ACS

*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

for diagnosing pediatric obesity, especially when height and weight measures are collected objectively [83, 84]. Indeed, for our current study, height and weight were measured by trained healthcare workers.

Here, we sought to understand the pathways that underly the development of increased body weight in children by applying novel methods to a rich data source. Specifically, we used geocoded EMR data from pediatric patients to identify potential strategic targets for obesity prevention among Latinx youth residing in neighborhoods located in Los Angeles County and Orange County of Southern California. Our findings underscore the complex relationship between community racial/ethnic composition and body weight status during childhood. Indeed, here, we confirmed the significance of the place of residence on children's body weight by focusing on the differential influence of race/ethnicity and neighborhood socioeconomic context. The results of our analyses raise critical questions for future research and represent opportunities for policy to help children maintain a healthy weight.

Author Contribution AWK and DK conducted the statistical analyses, geocoded the data, drafted the manuscript, and revised the article for intellectual content. ZNK and BNJ contributed to the conception and design of the larger study, acquisition of the data, and reviewed and revised the article for intellectual content. LE extracted and pre-processed the data and reviewed and revised the article for intellectual content. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Funding Open access funding provided by SCELCC, Statewide California Electronic Library Consortium The authors received financial support for the research from the Kay Family Foundation Data Analytics Grant.

Data Availability Patient data are protected and are not available due to data privacy laws. Neighborhood data are available from the United States Census Bureau at <https://www.census.gov/programs-surveys/acs/>.

Declarations

Ethics Approval This study was performed in line with the principles of the Institutional Review Boards at the Children's Hospital of Orange County (IRB NetID: 1359937–13) and Chapman University (IRB #: IRB-23–47).

Competing Interests ZNK serves as a consultant for Edwards Lifesciences, Medtronic, and Huron consulting and is the President of the American College of Perioperative Medicine. All other authors have no conflicts of interest to report.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Sanyaolu A, Okorie C, Qi X, et al. Childhood and adolescent obesity in the United States: a public health concern. *Glob Pediatr Health*. 2019;6:2333794X19891305.
2. Stierman B, Afful J, Carroll MD, et al. National Health and Nutrition Examination Survey 2017–March 2020 prepandemic data files development of files and prevalence estimates for selected health outcomes. *Natl Health Stat Report*. 2021;158:1–20.
3. Ogden CL, Fryar CD, Martin CB, et al. Trends in obesity prevalence by race and Hispanic origin – 1999–2000 to 2017–2018. *JAMA*. 2020;324(12):1208–10.

4. Hales CM, Carroll MD, Fryar CD, et al. Prevalence of obesity among adults and youth: United States, 2015–2016. In: NCHS data brief, no. 288:1–8. National Center for Health Statistics; 2017.
5. Lopez MH, Krogstad JM, Flores A. Key facts about young Latinos, one of the nation's fastest-growing populations. Washington: National Survey of Latinos. Pew Research Center; 2018. <https://www.pewresearch.org/facttank/2018/09/13/key-facts-about-young-latinos/>
6. Skinner AC, Ravanbakht SN, Skelton JA, et al. Prevalence of obesity and severe obesity in US children, 1999–2016. *Pediatrics*. 2018;141:e20173459.
7. Sahoo K, Sahoo B, Choudhury AK, et al. Childhood obesity: causes and consequences. *J Family Med Prim Care*. 2015;4(2):187–92.
8. Dietz WH, Robinson TN. Overweight children and adolescents. *N Engl J Med*. 2005;352(20):2100–9.
9. Thomas IC, Allison MA. Hypertension in Hispanics/Latinos: epidemiology and considerations for management. *Curr Hypertens Rep*. 2019;21(6):43.
10. Cartwright K. Social determinants of the Latinx diabetes health disparity. *Soc Sci Med*. 2021;15:100869.
11. Roncoroni J, Okun M, Hudson A. Systematic review: sleep health in the US Latinx population. *Sleep*. 2022;45(7):zsac092.
12. Gold DR, Wright R. Population disparities in asthma. *Annu Rev Public Health*. 2005;26:89–113.
13. Wassertheil-Smoller S, Arredondo EM, Cai JW, et al. Depression, anxiety, antidepressant use, and cardiovascular disease among Hispanic men and women of different national backgrounds: results from the Hispanic Community Health Study/Study of Latinos. *Ann Epidemiol*. 2014;24(11):822–30.
14. Warrington NM, Beaumont RN, Horikoshi M, et al. Maternal and fetal genetic effects on birth weight and relevance to cardiometabolic risk factors. *Nat Genet*. 2019;51(5):804–14.
15. Bell JF, Zimmerman FJ. Shortened nighttime sleep duration in early life and subsequent childhood obesity. *Arch Pediatr Adolesc Med*. 2010;164:840–5.
16. Jabeile H, Kelly AS, O'Malley GO, et al. Obesity in children and adolescents: epidemiology, causes, assessment, and management. *Lancet Diabetes Endocrinol*. 2022;10(5):351–65.
17. Roberts JR, Karr CJ. Pesticide exposure in children. *Pediatrics*. 2012;130(6):e1765–88.
18. Li D-K, Chen H, Ferber J, et al. Infection and antibiotic use in infancy and risk of childhood obesity: a longitudinal birth cohort study. *Lancet Diabetes Endocrinol*. 2017;5(1):18–25.
19. di Palma E, Filice E, Cavallo A, et al. Childhood obesity and respiratory diseases: which link? *Children (Basel)*. 2021;8(3):177.
20. Pinto NM, Marino BS, Wernovsky G, et al. Obesity is a common comorbidity in children with congenital and acquired heart disease. *Pediatrics*. 2007;120(5):e1157–64.
21. Dai W, Liu X, Su H, et al. Influence of adipose tissue immune dysfunction on childhood obesity. *Cytokine Growth Factor Rev*. 2022;65:27–38.
22. Haliloglu B, Bereket A. Hypothalamic obesity in children: pathophysiology to clinical management. *J Pediatr Endocrinol Metab*. 2015;28(5–6):503–13.
23. Arslan N, Erdur B, Aydin A. Hormones and cytokines in childhood obesity. *Indian Pediatr*. 2010;47(10):829–39.
24. Ayala-Marin AM, Iguacel I, Miguel-Etayo PD, et al. Consideration of social disadvantages for understanding and preventing obesity in children. *Front Public Health*. 2020;8:423.
25. O'Lawrence, Martinez L, Castelo G, et al. The latest review of childhood obesity among Hispanic and Latinx populations in California. *Diab Res Open Access*. 2020;2(2):22–30.
26. Foster BA, Linville D, Miller-Bedell ER, et al. Food security and feeding behaviors in low-income, Latinx families with preschool-aged children. *Public Health Nutr*. 2022;25(12):1–6. <https://doi.org/10.1017/S1368980022001884>.
27. McCurdy K, Gans KM, Risica PM, et al. Food insecurity, food parenting practices, and child eating behaviors among low-income Hispanic families of young children. *Appetite*. 2022;169:105857.
28. Carter MA, Dubois L. Neighbourhoods and child adiposity: a critical appraisal of the literature. *Health Place*. 2010;16:616–28.
29. Trasande L, Cronk C, Durkin M, et al. Environment and obesity in the National Children's Study. *Environ Health Perspect*. 2009;117:159–66.
30. Rogers R, Eagle TF, Sheetz A, et al. The relationship between childhood obesity, low socioeconomic status, and race/ethnicity: lessons from Massachusetts. *Child Obes*. 2015;11(6):691–5.
31. Kranjac AW, Boyd C, Kimbro RT, et al. Neighborhoods matter; but for whom? Heterogeneity of neighborhood disadvantage on child obesity by sex. *Health Place*. 2021;68:102534.
32. Kranjac AW, Denney JT, Kimbro RT, et al. Child obesity and the interaction of family and neighborhood socioeconomic context. *Popul Res Policy Rev*. 2019;38:347–69.
33. Malacarne D, Handakas E, Robinson O, et al. The built environment as determinant of childhood obesity: a systematic literature review. *Obes Rev*. 2022;23(1):e13385.
34. Johnson KA, Jones-Smith J, Curriero FC, et al. Low-income black and Hispanic children's neighborhood food environment and weight trajectories in early childhood. *Acad Pediatr*. 2020;20(6):784–92.
35. Shrewsbury V, Wardle J. Socioeconomic status and adiposity in childhood: a systematic review of cross-sectional studies 1990–2005. *Obesity (Silver Spring)*. 2008;16:275–84.
36. Miller M, Saldarriaga EM, Jones-Smith JC. Household socioeconomic status modifies the association between neighborhood SES and obesity in a nationally representative sample of first grade children in the United States. *Prev Med Rep*. 2020;20:101207.
37. Carroll-Scott A, Gilstad-Hayden K, Rosenthal L, et al. Disentangling neighborhood contextual associations with child body mass index, diet, and physical activity: the role of built, socioeconomic, and social environments. *Soc Sci Med*. 2013;95:106–14.
38. Cradock AL, Kawachi I, Colditz GA, et al. Neighborhood social cohesion and youth participation in physical activity in Chicago. *Soc Sci Med*. 2009;68(3):427–35.
39. Sadeghi B, Schaefer S, Tserogounis IE, et al. Prevalence and perception of childhood obesity in California's farmworker communities. *J Community Health*. 2017;42:377–84.
40. Ludwig J, Sanbonmatsu L, Gennetian L, et al. Neighborhoods, obesity, and diabetes — a randomized social experiment. *N Engl J Med*. 2011;365(16):1509–19.
41. Donaldson CD, Kain ZN, Ehwerhemuepha L, et al. Anxiety and mood disorders impacting physician opioid prescribing in the pediatric hospital setting. *J Clin Psychol Med S*. 2021;28:757–70.
42. "Hispanic population growth and dispersion across U.S. counties, 1980–2020". Pew Research Center tabulation of decennial census data: P.L.94–171 data for. 2000, 2010 and 2020; STF-1a data for 1980. Washington, D.C.: Pew Research Center; 1990. p. 2022.
43. Massey DS, Gross AB, Shibuya K. Migration, segregation, and the geographic concentration of poverty. *Am Sociol Rev*. 1994;59(3):425–45.
44. Must A, Anderson SE. Body mass index in children and adolescents: considerations for population-based approaches. *Int J Obes*. 2006;30:590–4.
45. Wang Y, Chen HJ. Use of percentiles and Z-scores in anthropometry. In: Preedy VR, editor. *Handbook of anthropometry: Physical measures of human form in health and disease*. New York: Springer Science Business Media; 2012. https://doi.org/10.1007/978-1-4419-1788-1_2.
46. Kranjac AW, Kranjac D, Fortier MA, et al. Surgical patients' hospital experience scores: neighborhood context conceptual framework. *Ann Surg*. 2021;2(1):e037–040.
47. Kranjac AW, Denney JT, Kimbro RT, et al. Neighborhood and social environmental influences on child chronic disease prevalence. *Popul Environ*. 2018;40(2):93–114.

48. Kranjac AW, Kimbro RT, Denney JT, et al. Comprehensive neighborhood portraits of child asthma disparities. *Matern Child Health J.* 2017;21(7):1552–62.
49. Hanchane S, Mostafa T. Endogeneity problems in multilevel estimation of education production functions: An analysis using PISA data. Centre for Learning and Life Chances in Knowledge Economies and Societies; 2010. p. 2021. <http://www.llakes.org>.
50. Lazarsfeld PF, Henry NW. Latent structure analysis. Boston: Houghton Mill; 1968.
51. Jencks C, Mayer SE. The social consequences of growing up in a poor neighborhood. pg. 111–186 in *Inner-city poverty in the United States*. Vol. 111. Washington: National Academy Press; 1990.
52. Muthén LK, Muthén BO. Mplus user's guide. Eighth ed. Los Angeles: Muthén & Muthén; 1998–2017.
53. Johnson SK. Latent profile transition analyses and growth mixture models: A very non-technical guide for researchers in child and adolescent development. *New Dir Child Adoles.* 2021:111–39. <https://doi.org/10.1002/cad.20398>.
54. Rabe Hesketh S, Skrondal A. Multilevel and longitudinal modeling using stata. *Stata Press.* 2008; <https://doi.org/10.1111/j.1541-0420.2008.01138.15.x>.
55. StataCorp. Stata statistical software: release 16. College Station, TX: StataCorp LLC; 2019.
56. Hales CM, Fryar CD, Carroll MD, et al. Trends in obesity and severe obesity prevalence in U.S. youth and adults by sex and age, 2007–2008 to 2015–2016. *JAMA.* 2018;319(16):1723–5.
57. Vazquez CE, Cubbin C. Socioeconomic status and childhood obesity: a review of literature from the past decade to inform intervention research. *Curr Obes Rep.* 2020;9:562–70.
58. Laurson KR, Lee JA, Gentile DA, et al. Concurrent associations between physical activity, screen time, and sleep duration with childhood obesity. *ISRN Obes.* 2014;204540 <https://doi.org/10.1155/2014/204540>. eCollection 2014
59. Golley RK, Maher CA, Matricciani L, et al. Sleep duration or bedtime? Exploring the association between sleep timing behaviour, diet and BMI in children and adolescents. *Int J Obes.* 2013;37(4):546–51.
60. Xu H, Wen LM, Rissel C. Associations of parental influences with physical activity and screen time among young children: a systematic review. *J Obes.* 2015:546925. <https://doi.org/10.1155/2015/546925>.
61. Eisenmann JC, Gundersen C, Lohman BJ, et al. Is food insecurity related to overweight and obesity in children and adolescents? A summary of studies, 1995–2009. *Obes Rev.* 2011;12(5):e73–83.
62. Fiese BH, Hammons A, Grigsby-Toussaint D. Family mealtimes: a contextual approach to understanding childhood obesity. *Econ Hum Biol.* 2012;10(4):365–74.
63. Fiechtner L, Sharifi M, Sequist T, et al. Food environments and childhood weight status: effects of neighborhood median income. *Child Obes.* 2015;11(3):260–8.
64. Robinson TN, Matheson D, Wilson DM, et al. A community-based, multi-level, multi-setting, multi-component intervention to reduce weight gain among low socioeconomic status latinx children with overweight or obesity: the Stanford GOALS randomised controlled trial. *Lancet Diabetes Endocrinol.* 2021;9(6):336–49.
65. Samuels S, Craypo L, Boyle M, et al. The California Endowment's Health Eating, Active Communities program: a midpoint review. *Am J Public Health.* 2010;100(11):2114–23.
66. Franzini L, Elliott MN, Cuccaro P, et al. Influences of physical and social neighborhood environments on children's physical activity and obesity. *Am J Public Health.* 2009;99(2):271–8.
67. Burdette HL, Whitaker RC. A national study of neighborhood safety, outdoor play, television viewing, and obesity in preschool children. *Pediatrics.* 2005;116(3):657–62.
68. Dolin CD, Gross RS, Deierlein AL, et al. Predictors of gestational weight gain in a low-income Hispanic population: sociodemographic characteristics, health behaviors, and psychosocial stressors. *Int J Environ Res Public Health.* 2020;17(1):352.
69. Elwan D, Olveda R, Medrano R, et al. Excess pregnancy weight gain in Latinas: impact on infant's adiposity and growth hormones at birth. *Prev Med Rep.* 2021;22:101341.
70. Barr-Anderson DJ, Flynn JJ, Dowda M, et al. The modifying effects of race-ethnicity and socioeconomic status on the change in physical activity from elementary to middle school. *J Adolesc Health.* 2017;61(5):562–70.
71. Fakhouri TH, Hughes JP, Brody DJ, et al. Physical activity and screen-time viewing among elementary school-aged children in the United States from 2009–2010. *JAMA Pediatr.* 2013;167(3):223–9.
72. Basterfield L, Jones AR, Parkinson KN, et al. Physical activity, diet and BMI in children aged 6–8 years: a cross-sectional analysis. *BMJ Open.* 2014;4(6):e005001.
73. Janssen I, Leblanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act.* 2013;7:40.
74. Chaparro MP, Whaley SE, Anderson CE, et al. The role of income and neighborhood poverty in the association between the 2009 Special Supplemental Nutrition Program for Women, Infants and Children (WIC) food package change and child obesity among WIC-participating children in Los Angeles County, 2003–2016. *Public Health Nutr.* 2021;24(13):4212–9.
75. Pan L, Freedman DS, Sharma AJ, et al. Trends in obesity among participants aged 2–4 years in the special supplemental nutrition program for women, infants, and children – United States, 2000–2014. *MMWR Morb Mortal Wkly Rep.* 2016;65:1256–60.
76. Kilanowski JF. Influences on healthy-eating decisions making in Latino adolescent children of migrant and seasonal agricultural workers. *J Pediatr Health Care.* 2016;30:224–30.
77. Bishop W, Chavarin C, Gonzales HA, et al. Healthy eating and physical activity among low-income Hispanic adolescents and a school-based intervention. *Child Youth Serv Rev.* 2018;86:151–6.
78. Romo LF, Mirales-Rios R, Hurtado A. Chultural, media, and peer influences on body beauty perceptions of Mexican American adolescent girls. *J Adolesc Res.* 2016;31:474–501.
79. Ogden C, Carroll M, Fryar C, et al. Prevalence of obesity among adults and youth: United States, 2011–2014. *NCHS Data Brief.* 2015;219:1–8.
80. Balistreri KS, Van Hook J. Trajectories of overweight among US school children: a focus on social and economic characteristics. *Matern Child Health.* 2011;15:610–9.
81. Balistreri KS, Van Hook J. Socioeconomic status and body mass index among Hispanic children of immigrants. *Am J Public Health.* 2009;99:2238–46.
82. Rosenbaum S. Medicaid. *N Engl J Med.* 2002;346:635–40.
83. Simmonds M, Burch J, Llewellyn A, et al. The use of measures of obesity in childhood for predicting obesity and the development of obesity-related diseases in adulthood: a systematic review and meta-analysis. *Health Technol Assess.* 2015;19(43):1–336.
84. Flegal KM, Ogden CL, Fryar C, et al. Comparisons of self-reported and measured height and weight, BMI, and obesity prevalence from National Surveys:1999–2016. *Obesity (Silver Spring).* 2019;27(10):1711–9.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.