



**Manchester
Metropolitan
University**

Wilson, Caitlin, Janes, Gillian ORCID logoORCID: <https://orcid.org/0000-0002-1609-5898>, Lawton, Rebecca and Benn, Jonathan (2023) The types and effects of feedback for emergency ambulance staff: a systematic mixed studies review and meta-analysis. *BMJ Quality and Safety*. ISSN 2044-5415

Downloaded from: <https://e-space.mmu.ac.uk/631606/>

Version: Published Version

Publisher: BMJ Publishing Group

DOI: <https://doi.org/10.1136/bmjqs-2022-015634>

Usage rights: Creative Commons: Attribution 4.0

Please cite the published version

<https://e-space.mmu.ac.uk>



OPEN ACCESS

Types and effects of feedback for emergency ambulance staff: a systematic mixed studies review and meta-analysis

Caitlin Wilson ^{1,2,3} Gillian Janes ⁴ Rebecca Lawton ^{1,3}
Jonathan Benn ^{1,3}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjqs-2022-015634>).

¹School of Psychology, University of Leeds, Leeds, UK

²Research and Development Department, Yorkshire Ambulance Service NHS Trust, Wakefield, UK

³Yorkshire Quality and Safety Research Group, Bradford Institute for Health Research, Bradford, UK

⁴Department of Nursing, Manchester Metropolitan University, Manchester, UK

Correspondence to

Mrs Caitlin Wilson, University of Leeds School of Psychology, Leeds LS2 9JT, UK; c.wilson60@nhs.net

Royal College of Emergency Medicine Annual Virtual Scientific Conference (13th & 14th October 2020, online); College of Paramedics National Annual Research Conference (26th November 2020, online); HSR UK (7th July 2021, online).

Received 12 October 2022
Accepted 13 March 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

To cite: Wilson C, Janes G, Lawton R, *et al.* *BMJ Qual Saf* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bmjqs-2022-015634

ABSTRACT

Background Extensive research has been conducted into the effects of feedback interventions within many areas of healthcare, but prehospital emergency care has been relatively neglected. Exploratory work suggests that enhancing feedback and follow-up to emergency medical service (EMS) staff might provide staff with closure and improve clinical performance. Our aim was to summarise the literature on the types of feedback received by EMS professionals and its effects on the quality and safety of patient care, staff well-being and professional development.

Methods A systematic review and meta-analysis, including primary research studies of any method published in peer-reviewed journals. Studies were included if they contained information on systematic feedback to emergency ambulance staff regarding their performance. Databases searched from inception were MEDLINE, Embase, AMED, PsycINFO, HMIC, CINAHL and Web of Science, with searches last updated on 2 August 2022. Study quality was appraised using the Mixed Methods Appraisal Tool. Data analysis followed a convergent integrated design involving simultaneous narrative synthesis and random effects multilevel meta-analyses.

Results The search strategy yielded 3183 articles, with 48 studies meeting inclusion criteria after title/abstract screening and full-text review. Interventions were categorised as audit and feedback (n=31), peer-to-peer feedback (n=3), postevent debriefing (n=2), incident-prompted feedback (n=1), patient outcome feedback (n=1) or a combination thereof (n=4). Feedback was found to have a moderate positive effect on quality of care and professional development with a pooled effect of $d=0.50$ (95% CI 0.34, 0.67). Feedback to EMS professionals had large effects in improving documentation ($d=0.73$ (0.00, 1.45)) and protocol adherence ($d=0.68$ (0.12, 1.24)), as well as small effects in enhancing cardiac arrest performance ($d=0.46$ (0.06, 0.86)), clinical decision-making ($d=0.47$ (0.23, 0.72)), ambulance times ($d=0.43$ (0.12, 0.74)) and survival rates ($d=0.22$ (0.11, 0.33)). The between-study heterogeneity variance was estimated at $\sigma^2=0.32$ (95% CI 0.22, 0.50), with an I^2 value of 99% (95% CI 98%, 99%), indicating substantial statistical heterogeneity.

Conclusion This review demonstrated that the evidence base currently does not support a clear single point estimate of the pooled effect of feedback to EMS staff as a single intervention type due to study heterogeneity. Further research is needed to provide guidance and frameworks supporting better design and evaluation of feedback interventions within EMS.

PROSPERO registration number CRD42020162600.

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Within the wider healthcare context, interventions based on audit and feedback are well researched, with systematic reviews suggesting that feedback results in small to moderate improvements to patient care through enhancing clinical performance. Existing reviews have not included studies within prehospital emergency care, despite a growing body of research suggesting that feedback to emergency medical service (EMS) personnel could improve staff well-being and patient safety.

WHAT THIS STUDY ADDS

⇒ This review summarises the literature on the types of feedback received by EMS personnel and its effects on quality of care and professional development.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ The review findings suggest that feedback to EMS professionals may improve performance and patient care; and highlight the need for more sophisticated feedback designs and robust evaluations within EMS.

INTRODUCTION

The phenomenon of feedback is well researched within healthcare, including, for example, debriefing, patient experience feedback and feedback from incident reporting.^{1–3} The feedback type investigated most frequently in clinical settings is audit and feedback, also known as clinical performance feedback, which is defined as: ‘any summary of clinical performance

over a specified time period'.⁴ A Cochrane review published in 2012 suggested that audit and feedback results in small to moderate improvements in patient care by enhancing healthcare professionals' compliance with desired clinical practice.⁵

Patient outcome feedback is also increasingly used to support reflection and self-evaluation of clinical performance.⁶ This information enables clinicians to gain insight and knowledge, and may improve diagnostic ability and prompt other behaviour changes.⁷ It may also improve calibration, that is, the relationship between a clinician's confidence in their diagnostic accuracy and the true accuracy of the diagnosis, thereby improving patient care and patient safety.⁶⁻⁸

Feedback and follow-up on patients and performance is particularly difficult for emergency medical service (EMS) personnel. Although their work involves engaging and communicating with other parts of the health service during handover and referral of patients, EMS operate largely in silos with a geographically dispersed workforce delivering discrete episodes of care without direct supervision.⁹ This disconnect from the subsequent care team and episodic nature of autonomous EMS work means that EMS staff struggle to obtain feedback.

Extensive research has been conducted into the effects of feedback interventions within many healthcare settings; however, in 2012 relatively little work had been undertaken within EMS, as illustrated by the lack of EMS studies eligible for inclusion in the Cochrane review of audit and feedback.⁵ Since then, a growing body of literature has been published on feedback within EMS suggesting that enhancing feedback and follow-up for EMS staff might improve clinical performance and staff well-being,¹⁰⁻¹² but this has not yet been synthesised comprehensively.

Our aim was to summarise the literature on the types of feedback received by EMS professionals and their effects on quality and safety of patient care. The following questions were addressed:

- ▶ What are the intervention types, design elements, potential mechanisms and reported effects of feedback interventions to EMS professionals?
- ▶ How do EMS professionals perceive current feedback provision?
- ▶ What are the reported contextual factors (eg, barriers, facilitators, opportunities) for effective feedback to EMS staff?

METHODS

The study was defined as a 'systematic mixed studies review', as preliminary searches identified important quantitative, qualitative and mixed-methods studies of relevance to the review aims.¹³ The review was registered on PROSPERO (CRD42020162600) and the protocol published a priori.¹⁴

Search strategy

The search strategy was developed in collaboration with an ambulance service librarian and included three facets: ambulance staff, feedback and feedback content (online supplemental appendix 1). Ovid (MEDLINE, Embase, AMED, PsycINFO, HMIC), EBSCO (CINAHL) and Web of Science were searched from their respective inception dates and searches last updated on 2 August 2022.

Study inclusion and exclusion criteria

English language primary studies of qualitative, quantitative and mixed-methods methodology were included if they described, evaluated or discussed feedback to EMS professionals. Reasons for excluding non-English studies were pragmatic (ie, lack of funding for translation, reducing the number of retrieved articles and therefore screening time) and theoretical (ie, the review targeted the Anglo-American EMS system involving paramedics rather than physicians). Morrison *et al*¹⁵ and Moher *et al*¹⁶ provided reassurance that this exclusion would not introduce bias in systematic reviews within medicine and healthcare.

Feedback was defined as the systematic provision of information to EMS professionals regarding their performance using process metrics, patient outcome metrics or both types. Grey literature was excluded along with articles focused on automated feedback by devices (eg, automated external defibrillators) and feedback based on hypothetical data as these were considered distinct cases of feedback with their own existing literature syntheses.¹⁷⁻²⁵ Educational settings were excluded as the review focused on real-world performance.

Study selection

CW independently screened the full set of titles and abstracts retrieved in two passes: (1) setting (prehospital emergency care) and (2) intervention (feedback to EMS professionals). Emily Parker independently screened a random 10% subset and disagreements were resolved through discussion. Although independent review of all search results by two reviewers is recommended to reduce the probability of missing relevant studies, dual screening a small percentage of records is an acceptable alternative that pragmatically balances thoroughness and resource use.^{26 27} Cohen's kappa was calculated to determine the level of agreement between reviewers for screening (strong agreement for setting $k_s=0.85$, moderate agreement for intervention $k_i=0.78$). CW assessed eligibility of articles selected for full-text review with JB verifying articles excluded at this stage.

Study quality assessment

CW appraised the methodological quality of individual studies using the Mixed Methods Appraisal Tool.²⁸ This tool allowed for the appraisal of qualitative research,

randomised controlled trials, non-randomised studies, quantitative descriptive studies and mixed-methods studies within a single framework. JB confirmed the ratings assigned to each study and disagreements were resolved through discussion.

Data extraction

CW performed data extraction using a comprehensive, standardised extraction template that was tested and iterated by the review team (online supplemental appendix 2). A second researcher (GJ, RL or JB) checked the data extraction forms for accuracy and detail for a total of 20% (n=10) of studies. CW contacted four authors for missing data to enable inclusion in the meta-analysis, with one providing additional data, two unable to provide data due to study methods or governance issues and one not responding. We included the study where additional data were provided in the meta-analysis, while effect sizes of the remaining three studies were reported descriptively.

Data synthesis

Studies were divided into three a priori defined categories:

1. Empirical studies reporting feedback interventions within EMS, subcategorised into evaluative studies and descriptive case studies.
2. Empirical non-interventional studies of feedback within EMS.
3. Empirical studies in which feedback to EMS was one, but not the sole or primary focus of the study—these studies were not included in the analysis but are introduced in the discussion to highlight opportunities for enhanced feedback interventions.

The resulting groupings contained quantitative, qualitative and mixed-methods studies. As the review questions could be addressed by both qualitative and quantitative data, a convergent integrated approach was chosen, which involved synthesis occurring simultaneously (*convergent*) and data being transformed to combine quantitative and qualitative data (*integrated*).^{13 29 30} Specifically, we transformed quantitative data into qualitative data by converting it into textual descriptions to be included in our narrative synthesis.²⁹ An example of this would be 41.7% of respondents in the Cash *et al*³¹ survey reporting to have received patient outcome feedback, compared with 54.7% receiving medical care feedback, which we converted to ‘patient outcome feedback was less frequent than clinical performance feedback’.

We performed a narrative synthesis for data relating to feedback types and design elements, EMS professionals’ perceptions of feedback provision and key contextual factors. In line with guidance for narrative synthesis³² and best practices within the broader field of implementation science, we used theory to further understand the mechanisms of effective feedback.^{33 34} To describe feedback design elements we used clinical

performance feedback intervention theory (CP-FIT), which is a prominent and widely adopted theory within the implementation science literature based on a meta-synthesis of 73 feedback interventions.³⁵ Our own qualitative work on feedback to emergency ambulance staff found CP-FIT to have good face validity when exploring causal mechanisms at an abstract level, for example, ‘processing and reflection’.¹²

To identify potential mechanisms at a more detailed level for this review, we used behaviour change theory, which aims to identify active ingredients of interventions seeking to change behaviour³⁶ and has previously been used to synthesise evidence from audit and feedback interventions.^{37 38} CW identified potential causal mechanisms deductively using an established list of mechanisms of action from behaviour change theory,³⁶ with a researcher experienced in behaviour change theory coding (RL) verifying the assigned codes of a 20% (n=10) random sample. Disagreements were resolved through discussion. Once agreement was reached, CW reviewed the remaining studies.

Lastly, we analysed feedback effects by conducting post hoc random-effects multilevel meta-analyses of quantitative evaluative studies, including tabulation and aggregation of standardised mean differences (Cohen’s *d*) and corresponding 95% CIs using the package ‘metafor’ in R V.4.1.3.³⁹ Cohen’s *d* is an effect size that evaluates the difference between two means and is interpreted as small (*d*=0.2), medium (*d*=0.5) or large (*d*=0.8).⁴⁰ Cohen’s *d* is particularly useful in meta-analyses of studies that assess the same (continuous) outcome but measure it using different scales, as it allows each study result to be standardised to a uniform scale and therefore allows studies to be combined.²⁷ Effect sizes were extracted directly, approximated from median and IQRs or calculated from raw data.^{27 41–44} Individual meta-analyses were conducted for each outcome category, as well as a combined meta-analysis to quantify the effects of feedback overall on EMS care quality and safety.⁴⁵ A multilevel approach was chosen for the overall meta-analysis to account for including multiple effect sizes from individual studies.⁴⁶ Where studies reported multiple effect sizes within a single outcome category, the largest effect size was chosen, in line with a ‘proof of concept’ approach.⁴⁷ Relationships in the data were meta-analysed using subgroup analyses informed by CP-FIT’s desirable feedback intervention design elements, feedback type and study quality.³⁵

Between-study heterogeneity was assessed using χ^2 test, I^2 , τ^2 and prediction intervals (PIs), following guidance to use multiple measures to characterise statistical heterogeneity in meta-analyses due to the relative strengths and weaknesses of the different measures available.^{27 48} The χ^2 test (Cochran’s *Q*) is the traditional test to detect whether heterogeneity is present in meta-analyses and examines the null hypothesis that all studies are evaluating the same effect.⁴⁹

Cochran's Q has low power to detect differences with small numbers of studies and high power when there are many studies, so alpha was set to $p < 0.10$ for the individual meta-analyses of each outcome category and $p < 0.05$ for the overall meta-analysis.²⁷ To describe the extent of between-study variability due to heterogeneity, the I^2 statistic was calculated and interpreted as: not important (0%–40%), moderate (30%–60%), substantial (50%–90%) and considerable (75%–100%).²⁷ I^2 can be biased in small and large meta-analyses,^{50 51} so τ^2 was used to quantify the variance of the true effect sizes underlying our data in a way that was insensitive to the number of studies.⁴⁸ Due to our overall meta-analysis being a multilevel analysis, between-study heterogeneity variance is denoted as σ^2 instead of τ^2 .⁴⁸ As τ^2 and σ^2 are difficult to interpret, we used their results to calculate 95% PIs, which provide an estimate of the expected range of true effects in future studies using the same scale as the effect size metric (ie, Cohen's d).^{27 52}

Quality of evidence

As this review's convergent synthesis design required data transformation and integration, the certainty associated with the whole body of literature was not assessed and instead individual study quality reported. This aligns with recent guidance, warning against using Grading of Recommendations Assessment, Development and Evaluation (GRADE)/Confidence in Evidence from Reviews of Qualitative research (CERQual) in mixed-studies systematic reviews.²⁹

RESULTS

Our cumulative search identified 4891 results (figure 1). Once duplicates were removed, 3183 titles and abstracts were screened for inclusion. Of these, 2195 were excluded based on the setting and 611 based on the intervention criteria. The remaining 377 articles underwent full-text review, resulting in 329 exclusions and a total of 48 articles included in the analysis.

Study characteristics

An overview of included studies is presented in table 1 with a detailed breakdown in online supplemental appendix 3. There were 42 interventional studies and six non-interventional studies. Studies were published between 1989 and 2022 with 73% (n=35) published in the last 10 years, and originated most frequently in the USA (n=19, 40%). The most common study design was cross-sectional (n=30, 63%). Most studies were situated within paramedic emergency services (n=39, 81%), but emergency operations centres (n=6, 13%) were also represented.

Quality assessment

All studies were assessed as having clear research questions and collected data addressing these (online

supplemental appendices 3 and 4). The qualitative and mixed-methods studies were of high quality. Several quantitative descriptive studies (n=5, 10%) omitted reporting on non-response bias, and were therefore of high-to-moderate quality, with the remaining being of high quality. Many of the quantitative non-randomised studies were of moderate quality due to confounders not being accounted for in the study design and analysis (n=23, 48%), as well as not reporting on representativeness of participants (n=6, 13%) and intervention fidelity (n=30, 63%).

Feedback intervention types and designs within EMS (interventional studies)

The interventional studies (n=42) described 37 unique EMS feedback interventions. Sources of feedback included researchers (n=17, 40%), ambulance service managers (n=14, 33%), peers (n=6, 14%) and external databases (n=1, 2%). Feedback was provided to organisations (n=8, 19%), ambulance crews/teams (n=16, 38%), individual clinicians (n=10, 24%) or a combination (n=7, 17%). An example of an organisational-level intervention provided ambulance services with feedback on their care bundle performance for myocardial infarction and stroke via monthly teleconferences and weekly control charts of performance data.⁵³ Weston *et al*, on the other hand, provided individual paramedics with a single page of written feedback on their performance in cardiac arrests, including chest compression depth and rate.⁵⁴

Feedback was delivered as a stand-alone initiative (n=17, 40%), or as part of wider organisational (n=12, 29%) or educational (n=13, 31%) interventions. An example of a stand-alone initiative was the provision of hospital-directed feedback to an EMS organisation on compliance with state protocols and documentation.⁵⁵ Other studies described feedback being implemented alongside training⁵⁶ or new equipment and guidelines.⁵⁷

The clinical topic was varied with the most frequent being cardiac arrest (n=13, 31%), myocardial infarction (n=10, 24%), stroke (n=3, 7%) and trauma (n=2, 5%). Other studies included multiple conditions or all patients (n=12, 29%), with the remaining focusing on paediatrics (n=1, 2%) and non-conveyed patients (n=1, 2%). Overall, interventional studies described six different feedback types according to the typology developed in our previously published exploratory interview study¹²: audit and feedback (n=31, 74%), postevent debriefing (n=2, 5%), peer-to-peer feedback (n=3, 7%), incident-prompted feedback (n=1, 2%), audit and patient outcome feedback (n=4, 10%) and patient outcome feedback (n=1, 2%) (table 2).

Feedback was provided on individual (n=18, 43%), aggregate (n=10, 24%) or individual and aggregate patient cases (n=6, 14%) and in all but one study consisted of 'push' model feedback, that is, not in response to recipients actively seeking feedback.⁵⁸ Only




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram.

six studies (14%) reported that recipients were offered a comparison or benchmark alongside the feedback. The format of feedback interventions included written (n=14, 33%), verbal (n=10, 24%) or a combination of strategies (n=5, 12%). Visual elements were seldom reported but included line charts, viewing recordings of resuscitation performance and green/red highlighting on written reports. Frequency varied from after each call to only once or every 6 months, with lag time similarly varying from 48 hours to 3 months.

Many of the design characteristics discussed in the audit and feedback literature, for example, timeliness, user-friendly designs and inclusion of patient-level data, were not explicitly considered and no studies compared different feedback types. No studies reported using existing feedback theory (eg, CP-FIT); although one paper referenced audit and feedback literature,⁵⁹

five used quality improvement methods,^{53 60–62} and a further paper acknowledged the challenge of sustainability within implementation science.⁶³

Outcomes and effectiveness of feedback interventions (interventional studies)

The feedback interventions reported in the evaluative interventional studies (n=36) measured a number of quality and safety outcomes, including process metrics—such as ambulance times (n=13, 36%, eg, arrival times, on-scene times, call-to-needle times), protocol adherence (n=12, 12%), cardiac arrest performance (n=10, 28%), clinical decision-making including non-conveyance decisions (n=6, 17%) and documentation (n=5, 14%)—as well as patient outcomes such as survival rates (n=8, 22%) and patient satisfaction (n=1, 3%). The remaining descriptive case

Table 1 Characteristics of included studies

Characteristic	Interventional feedback studies (n=42)		Non-interventional feedback studies (n=6)		Total (N=48)	
	n	%*	n	%*	n	%*
Year						
1989–1999	6	14	1	17	7	15
2000–2009	3	7	–	–	3	6
2010–2019	26	62	2	33	28	58
2020–2022	7	17	3	50	10	21
Country						
USA	16	38	3	60	19	40
Germany	6	14	–	–	6	13
UK	4	10	2	33	6	13
Sweden	3	7	–	–	3	6
Netherlands	2	5	–	–	2	4
Australia	2	5	–	–	2	4
Other or multiple countries	9	21	1	17	10	21
Study design						
Cohort study	4	10	–	–	4	8
Case–control study	1	2	–	–	1	2
Cross-sectional study	30	71	–	–	30	63
Quantitative descriptive study	4	10	3	50	7	15
Qualitative study	1	2	3	50	4	8
Mixed-methods study	2	5	–	–	2	4
Context						
Paramedic emergency services	33	79	6	100	39	81
Emergency operations centre	6	14	–	–	6	13
Ambulance service organisation	2	5	–	–	2	4
Other or multiple contexts	1	2	–	–	1	2

*Sections of individual columns may not add up to exactly 100% due to rounding.

studies (n=6) simply demonstrated that feedback was possible in the prehospital emergency care setting, for example, Stella *et al*,⁵⁸ who reported how incident reporting data could be used to provide feedback to EMS professionals.⁵⁸

Of the 36 evaluative interventional studies, 30 (83%) provided sufficient data to calculate standardised effect sizes and were tabulated according to study outcomes (figure 2). All 30 studies were quantitative and non-randomised with the majority providing before-and-after comparison. Feedback to EMS professionals was found to have statistically significant positive effects in 73% (n=40) of the 55 extracted outcome measures, with the remaining outcome measures indicating non-significant effects in a positive direction (n=7, 13%), negative direction (n=4, 7%) or no effect (n=4, 7%). Outcome measures are listed in online supplemental appendix 3 and included, for example: stroke care bundle delivery (*protocol adherence*),⁵³ satisfaction level of patients (*patient satisfaction*),⁶⁴ survival to hospital discharge (*survival rates*),⁶⁵ adequate documentation pertaining to physical examination (*documentation*),⁶⁶ patients transported directly to the catheterisation laboratory (*clinical decision-making*),⁶⁷ chest compressions per minute (*cardiac arrest performance*)⁶² and EMS run time (*ambulance times*).⁶⁸

Overall, feedback had a moderate positive effect with a standardised mean difference for all outcome measures of $d=0.50$ (95% CI 0.34, 0.67). There was evidence of substantial between-study heterogeneity ($I^2=99%$, 95% CI 98%, 99%) at a statistically significant level ($Q=2650.26$, $p<0.001$). Variance of between-study heterogeneity was estimated at $\sigma^2=0.32$ (95% CI 0.22, 0.50), resulting in a wide PI (–0.64, 1.64), indicated in dotted lines on either side of the pooled effect size in figure 2. Our calculated PI illustrates the uncertainty in the pooled estimate for EMS feedback effects and crosses the line of no effect, so the true effect in future studies may be negative, null or higher than our pooled estimate.⁵²

Separate meta-analyses for each outcome category revealed that feedback had a large effect in improving documentation ($d=0.73$ (0.00, 1.45)) and protocol adherence ($d=0.68$ (0.12, 1.24)), as well as small effects in enhancing cardiac arrest performance ($d=0.46$ (0.06, 0.86)), clinical decision-making ($d=0.47$ (0.23, 0.72)), ambulance times ($d=0.43$ (0.12, 0.74)) and survival rates ($d=0.22$ (0.11, 0.33)). Heterogeneity was substantial for all outcome categories except for survival rates ($I^2=35.2%$) and clinical decision-making ($I^2=48.2%$), which fell into the moderate range of the I^2 index. Our results indicated moderate between-study

Table 2 Characteristics of feedback to EMS professionals identified in this systematic review

Feedback type	Feedback on professional practice			Patient outcome feedback		
	Audit and feedback	Postevent debriefing	Peer-to-peer feedback	Incident-prompted feedback	Audit and patient outcome feedback	Patient outcome feedback
Illustrative example	Feedback to EMS staff and ambulance stations by the research team on adherence to corticosteroid administration ⁶⁰	Feedback to EMS staff in a group setting following a resuscitation attempt using data collected from audit forms and defibrillator downloads with input from other EMS staff and the clinical training team ¹²⁷	Feedback to an EMS professional in a group setting on an everyday professional situation described by the feedback recipient with other EMS professionals offering their reflections and ideas on how to best achieve efficient professional care ⁷³	Feedback to EMS staff and ambulance stations on the observed trends of reported patient safety incidents, including patient outcomes and clinical management ⁵⁸	Feedback to EMS staff on their performance of EMS stroke quality measures, ambulance times and the patients' final diagnosis ⁶³ at hospital discharge	Feedback to EMS dispatchers by EMS staff on scene regarding patients' primary condition when the ambulance arrived on scene, non-transport by the ambulance and level of priority ⁵⁹
Feedback interventions reviewed, n (%)	31 (74)	2 (5)	3 (7)	1 (2)	4 (10)	1 (2)
Data	(Most commonly) Aggregated	Individual or aggregate	Individual	Individual	Individual or aggregate	Individual
Content	Clinical performance metrics and adherence to these	EMS treatment and management plan	The views of peers on the EMS treatment and management plan	Analysis of EMS patient safety incidents	Clinical performance metrics and outcomes of patients	Outcomes of patients (eg, diagnosis, hospital treatment, management plan)
Set-up	Formal	Formal	Formal or informal	Formal	Formal	Formal or informal
Comparison	Explicit benchmarks	May include explicit benchmarks	Indirect comparison with other clinicians	No explicit comparison	May include explicit benchmarks	No explicit comparison
(Anticipated) Effects	Change clinician behaviour: improve clinical performance	Change clinician behaviour: improve clinical performance, provide support	Change clinician behaviour: improve patient management skills, provide peer support	Change clinician behaviour: reduce patient safety incidents	Change clinician behaviour: improve clinical performance, improve clinicians' diagnostic ability	Change clinician behaviour: improve clinicians' diagnostic ability and patient management skills
EMS, emergency medical service.						




Figure 2 Forest plot of standardised mean differences and corresponding 95% CIs for feedback interventions in emergency medical service (EMS). RE, random effects.

heterogeneity at a statistically significant level for clinical decision-making ($Q=9.84, p=0.080$), but not for survival rates ($Q=9.66, p=0.209$). This was supported by the PI for clinical decision-making not including the null point, suggesting future studies will demonstrate positive effects, whereas the PI for survival rates included no effect. One study⁶⁴ suggested feedback to EMS personnel may improve patient satisfaction, but was low quality and effect sizes did not reach statistical significance. The number of studies in the subgroup analyses (online supplemental appendix 5) was small and results not statistically significant.

The remaining six evaluative interventional studies were not included in the meta-analysis due to three not providing sufficient data to calculate effect sizes,^{60 69 70} two not including a comparison^{71 72} and one conducting qualitative analysis.⁷³ The unstandardised effect sizes of the three studies with insufficient data indicated positive effects in the areas of protocol adherence, ambulance times and clinical decision-making (online supplemental appendix 6).^{60 69 70} The other three studies provided descriptive data on subjective self-reported measures, such as job satisfaction (*'making clinical shifts more enjoyable'*,⁷¹ *'increased motivation'*⁷²), team climate (*'improved relationships with colleagues'*,^{71 73} *'increased organisational commitment'*⁷²) and clinical decision-making (*'avoid repeating experienced colleagues' mistakes'*,⁷³ *'improved confidence'*^{71 73}).

Mechanisms of feedback effects (interventional and non-interventional studies)

Mechanisms of EMS feedback effects were poorly reported in interventional and non-interventional studies, with none exploring how mechanisms linked to study outcomes. Figure 3 presents deductive analysis of causal mechanisms for reported feedback effects

based on coding to an established list of 26 mechanisms from behaviour change theory.³⁶

Some mechanisms, such as knowledge, skills and feedback processes, were identified across different feedback types, while others, such as professional role and identity, environmental context and decision processes, were unique to specific feedback types. Audit and feedback interventions appeared to achieve their effects by influencing intentions and goals; while interventions, also including patient outcome information, appeared to include beliefs about consequences and decision processes. Meanwhile, post-event debriefing, peer-to-peer feedback and patient experience feedback interventions included causal mechanisms used by audit and patient outcome feedback, such as social influences and emotions. Without inferring causal mechanisms from limited information, no potential mechanisms could be identified for nine studies, including the incident-prompted feedback intervention.

EMS professionals' evaluation of feedback provision (non-interventional studies)

The non-interventional studies all reported that EMS professionals were dissatisfied with current feedback provision and desired more feedback, particularly patient outcome feedback.^{10-12 31 74 75} Concerns were that feedback was skewed towards the negative^{10 75} and only triggered by highly traumatic incidents,¹¹ while EMS professionals desired routine, frequent and high-quality feedback.¹⁰⁻¹² One UK study suggested that EMS professionals desired 'pull'-type feedback, that is, initiated by individual clinicians, delivered electronically or involving staff mediators.¹¹ In addition to the feedback types identified in interventional studies (table 2), non-interventional studies also discussed patient experience feedback, which involved EMS


Mechanisms of action	Illustrative example	Audit and feedback (n=31)	Postevent debriefing (n=2)	Peer-to-peer feedback (n=4)	Patient experience feedback (n=2)	Audit and patient outcome feedback (n=9)	Patient outcome feedback (n=1)
Knowledge	a 30-minute, web-based training module directed at improving stroke-related knowledge ⁶³	53 56 60 62 69 128 133	127	71		12 63 68	59
Skills	Every resuscitation attempt was recorded and reviewed in a team debriefing session with a special focus on the quality of chest compression ¹²⁷	57 60 130 131 134	127	71		10 135	
Social/professional role and identity	Participants learned to meet the demands of the profession and visualise the manner in which the individual should improve his/her care work ⁷³			73			
Beliefs about capabilities	Allows paramedics to be more confident in their ability to treat patients in an isolated care episode by understanding the wider patient experience ³¹			71		11 12	
Beliefs about consequences	Feedback stops you from "wondering around curious all the time whatever happened to that patient, once again, did I do the right thing?" ¹⁰					10 12 63	59
Reinforcement	One trust also implemented a financial incentive scheme linked to their feedback system ⁵³	53 70				135	
Intentions	Showing interruptions to chest compressions visually on-screen was particularly important to induce a change in performance ¹³⁶	136					
Goals	Dispatcher training included all the goals and learning objectives to be achieved ¹³⁷	54 56 67 70 130 131 133 137 144					
Memory, attention and decision processes	Improvement of diagnostic skills by 'enabling patterns of clinical presentation to be learned' ³¹					11 68 135	
Environmental context and resources	Educational resource cards were created and dispersed for EMS providers outlining corticosteroid dosing for paediatric patients ⁶⁰	53 60 69					
Social influences	"A partner must be able to point out your weaknesses and assist in strengthening (them)" ⁷¹		127	71 73		12	
Emotion	Participants thought that receiving feedback may positively impact on their mental wellbeing by affirming that they had done the best they could do ³²			71 73 74	12 74	10 12	
Motivation	Constructive feedback is provided to individuals with emphasis on why certain behaviours may be detrimental to patient care ¹²⁹	72 129				12 75	
Self-image	"Out there, you have to make a lot of decisions on your own and it is not always easy. So you need a forum, where you can get confirmation of whether or not you acted correctly in the situation" ⁷³			73 74	12 74		
Feedback processes	When feedback is given with explained suggestions for corrections, the health care staff can integrate the information and improve performance ¹⁴⁵	60	61 127	73 145		12 63	
Optimism, behavioural regulation, norms, subjective norms, attitude towards the behaviour, needs, values, social learning/imitation, behavioural cueing, general attitudes/beliefs and perceived susceptibility or vulnerability were not represented and are therefore not listed in this table.		Number of studies: 					

Figure 3 Potential mechanisms of feedback effects by feedback type.

professionals receiving feedback from patients or relatives on the care they provided.

Key contextual factors for feedback interventions (non-interventional studies)

The main barriers reported were practical, such as the lack of linkage between EMS and in-hospital information systems⁶⁸ and the time and resources required for generating feedback.⁷⁵ Potential social barriers were also highlighted, including how staff interact with and respond to feedback, especially within EMS culture where feedback may be perceived as linked to disciplinary action or being of questionable value.^{10 12} A reported ethical barrier was sharing of confidential patient details, especially for non-hospital-affiliated EMS agencies in the USA.^{10 75}

Potential facilitators for effective feedback interventions were use of electronic health records to reduce demands of information capture and feedback delivery, thereby promoting sustainability.^{11 76 77} It was further reported that having clearly defined performance measures may support the provision of high-quality feedback.⁶⁸ One study described a shift in EMS culture towards being more accepting of discussing mistakes, which may encourage clinicians to engage with feedback.¹⁰

DISCUSSION

The present review aimed to address a gap in existing evidence synthesis for feedback within EMS by summarising the literature on the types and effects of feedback received by EMS professionals. Previous reviews in the prehospital emergency care setting have focused on automated feedback from defibrillators^{17–20} and debriefing after simulation.^{21–25} Further to this, a non-systematic literature review from 2018 descriptively summarised a selection of the published literature relating to EMS clinical feedback.⁷⁸ The present review sought to overcome previous reviews' limitations by not restricting inclusion to randomised controlled trials, taking a broader view on feedback, drawing on explanatory theory and employing systematic search and evidence synthesis methods.

The reviewed interventional studies indicate that the source, content and mode of feedback interventions within EMS vary greatly, while design elements are poorly reported. Feedback interventions within EMS targeted and measured a number of quality and safety outcomes, including protocol adherence, patient satisfaction, survival rates, documentation, clinical decision-making, cardiac arrest performance and ambulance times. The meta-analyses of 30 evaluative interventional studies revealed substantial heterogeneity, so the moderate positive effect overall ($d=0.50$ (0.34, 0.67)) must be treated with caution, although it is comparable with recent systematic reviews within audit and feedback reporting ($d=0.40$).^{5 47} Our analysis indicated high heterogeneity for the majority of outcome categories, but found only moderate

heterogeneity for clinical decision-making, which suggests existing evidence may more consistently support the positive point estimate in this outcome category. The high proportion of positive effects across individual studies and pooled effects for clinical decision-making suggests feedback to EMS personnel has a positive impact; however, the current evidence does not support a single point estimate of the aggregated effect of EMS feedback across multiple intervention types and outcomes. The high levels of clinical and statistical heterogeneity indicate a clear need to further examine different feedback types and identify which mechanisms are most effective within EMS.

Reviewed studies indicate that the current design of feedback interventions within EMS is for the large part atheoretical, which is a significant barrier to understanding feedback design and effectiveness. Analysis using behaviour change theory suggested there may be unique mechanisms for different feedback types, that is, audit and feedback (*intention/goals*), patient outcome feedback (*beliefs about consequences*) and peer-to-peer feedback (*professional role/self-image*), though we were unable to demonstrate a statistical difference between interventions due to small numbers of studies in each subgroup.

Implications and recommendations

Interventions within EMS that explore patient outcome and patient experience feedback are needed as these were desired by staff in non-interventional studies yet under-represented in interventional studies. Considerable opportunity exists to enhance feedback provision to EMS personnel through data linkage and integrated data sets that span service boundaries.^{76 77} Similarly, enhancing existing electronic record systems might facilitate disaggregation of data to an individual level and enable targeted queries that support 'pull model' feedback, thereby making feedback more relevant to the recipient.^{11 76} As part of designing and implementing feedback interventions, information governance and ethical concerns linked to patient confidentiality should be addressed by bringing together relevant stakeholders.^{11 12}

Beyond the literature included in this review, the proliferation of unpublished local feedback initiatives within EMS suggests that a review of current practice is needed to understand whether the published evidence is being implemented in current feedback provision. Poor reporting of intended mechanisms of action and a lack of theoretical underpinning indicate limited understanding of feedback mechanisms, consistent with the wider literature.^{79 80}

The broader literature suggests specific opportunities for enhanced feedback to EMS personnel, such as combined audit and patient outcome feedback to improve decision-making^{81–84} for specific patient presentations (eg, cardiac arrest,^{85–93} myocardial infarction,^{94–96} stroke,^{97 98} abdominal pain,⁹⁹ paediatrics,¹⁰⁰ trauma^{101 102}). Furthermore, patient outcome feedback may improve staff mental health and learning^{103–109} for

patients not conveyed to hospital,¹¹⁰ with non-specific complaints,¹¹¹ with significant differences between EMS and in-hospital diagnoses¹¹² and patients referred to the coroner,¹¹³ while increased audit and feedback of particular skills (eg, intubation,¹¹⁴ ultrasound¹¹⁵) or situations (eg, handover,¹¹⁶ triage¹¹⁷) may improve performance. Lastly, increasing peer-to-peer and patient outcome feedback for certain staff groups may provide peer support and improve patient management skills for newly qualified¹¹⁹ and specialist paramedics.^{121–126}

Strengths and limitations

This is the first systematic review to investigate feedback within EMS, by adopting a broad definition of feedback and a rigorous methodology, which was published and registered a priori. The search strategy was developed from Cochrane reviews in audit and feedback, with the addition of subject matter expertise to increase sensitivity to the prehospital emergency care setting. The inter-rater reliability check indicated good reliability, with the lower kappa value for the intervention criterion potentially being explained by its more complex nature. Despite adopting a rigorous, systematic approach, the search strategy may not have identified every relevant article. Excluding articles without full-text English language translations may have removed potentially relevant studies and as with all systematic reviews, findings are subject to publication bias.

Although we were able to conduct a meta-analysis of a subset of quantitative evaluative studies (n=30, 83%), heterogeneity and variations in reporting quality limit the ability to recommend prioritisation of any specific feedback intervention or design feature. Our meta-analysis synthesised the available evidence, which did not include any randomised controlled trials and indicated high levels of heterogeneity; therefore, evidence for effects must be treated with caution despite using robust synthesis methods. Due to feedback interventions within EMS frequently being reported as part of multifaceted interventions with interacting components, it was not always possible to identify clearly whether a potential mechanism related to the feedback aspect of the intervention.

CONCLUSIONS

In summary, this review demonstrated that the evidence base currently does not support a clear single point estimate of the pooled effect of feedback within EMS. Our meta-analysis revealed moderate positive effects but substantial heterogeneity across a range of quality and safety outcomes, including protocol adherence, patient satisfaction, survival rates, documentation, clinical decision-making, cardiac arrest performance and ambulance times. Viewed in the context of the existing audit and feedback literature, feedback within EMS is still in its infancy. Further research is needed to provide guidance and frameworks supporting better design and evaluation of feedback interventions within EMS in order to strengthen the evidence base. There is a clear need to

develop a consensus about the key active components and mechanisms in feedback to EMS professionals.

Twitter Caitlin Wilson @999_Caitlin, Gillian Janes @DrGillianJanes, Rebecca Lawton @LawtonRebecca and Jonathan Benn @JonathanBenn8

Acknowledgements The authors acknowledge Matt Holland, Library and Knowledge Service for NHS Ambulance Services in England, for assistance with developing the search string, and Emily Parker, School of Psychology, University of Leeds, for being the second reviewer for the screening process.

Contributors CW conceived the study and designed its protocol under supervision and guidance from GJ, RL and JB. CW led on data collection and analysis, with contributions and supervision from GJ, RL and JB. JB was the main supervisor for this project. CW drafted the article and all authors contributed substantially to its revision and approved the final version. CW acts as guarantor.

Funding This research was funded by the National Institute for Health Research (NIHR) Yorkshire and Humber Patient Safety Translational Research Centre (NIHR Yorkshire and Humber PSTRC).

Disclaimer The views expressed are those of the authors and not necessarily those of the NIHR, or the Department of Health and Social Care.

Competing interests None declared.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Caitlin Wilson <http://orcid.org/0000-0002-9854-4289>
Gillian Janes <http://orcid.org/0000-0002-1609-5898>
Rebecca Lawton <http://orcid.org/0000-0002-5832-402X>
Jonathan Benn <http://orcid.org/0000-0001-5919-9905>

REFERENCES

- 1 Tannenbaum SI, Greulich PE. The debrief imperative: building teaming competencies and team effectiveness. *BMJ Qual Saf* 2023;32:125–8.
- 2 Marsh C, Peacock R, Sheard L, *et al*. Patient experience feedback in UK hospitals: what types are available and what are their potential roles in quality improvement (Qi)? *Health Expect* 2019;22:317–26.

- 3 Benn J, Koutantji M, Wallace L, *et al.* Feedback from incident reporting: information and action to improve patient safety. *Qual Saf Health Care* 2009;18:11–21.
- 4 Jamtvedt G, Young JM, Kristoffersen DT, *et al.* Does telling people what they have been doing change what they do? A systematic review of the effects of audit and feedback. *Qual Saf Health Care* 2006;15:433–6.
- 5 Ivers N, Jamtvedt G, Flottorp S, *et al.* Audit and feedback: effects on professional practice and healthcare outcomes. *Cochrane Database Syst Rev* 2012;2012:CD000259.
- 6 Cifra CL, Sittig DF, Singh H. Bridging the feedback gap: a sociotechnical approach to informing clinicians of patients' subsequent clinical course and outcomes. *BMJ Qual Saf* 2021;30:591–7.
- 7 Fernandez Branson C, Williams M, Chan TM, *et al.* Improving diagnostic performance through feedback: the diagnosis learning cycle. *BMJ Qual Saf* 2021;30:1002–9.
- 8 Lavoie C, Croskerry P, *et al.* Outcome feedback and patient safety. In: Croskerry P, Cosby KS, Schenkel SM, eds. *Patient Safety in Emergency Medicine*. Philadelphia, PA, USA: Lippincott Williams & Wilkins, 2009: 251–8.
- 9 McCann L. *The paramedic at work: a sociology of a new profession*. Oxford: Oxford University Press, 2022.
- 10 Morrison L, Cassidy L, Welsford M, *et al.* Clinical performance feedback to paramedics: what they receive and what they need. *AEM Educ Train* 2017;1:87–97.
- 11 Eaton-Williams P, Mold F, Magnusson C. Exploring paramedic perceptions of feedback using a phenomenological approach. *Br Paramed J* 2020;5:7–14.
- 12 Wilson C, Howell A-M, Janes G, *et al.* The role of feedback in emergency ambulance services: a qualitative interview study. *BMC Health Serv Res* 2022;22:296.
- 13 Pluye P, Hong QN, Vedel I. Toolkit for mixed studies reviews (V3). 2016. Available: <http://toolkit4mixedstudiesreviews.pbworks.com> [Accessed 16 Dec 2019].
- 14 Wilson C, Janes G, Lawton R, *et al.* The types and effects of feedback received by emergency ambulance staff: protocol for a systematic mixed studies review with narrative synthesis. *IJES* 2021;10:247–65.
- 15 Morrison A, Polisen J, Husereau D, *et al.* The effect of english-language restriction on systematic review-based meta-analyses: a systematic review of empirical studies. *Int J Technol Assess Health Care* 2012;28:138–44.
- 16 Moher D, Pham B, Klassen TP, *et al.* What contributions do languages other than english make on the results of meta-analyses? *J Clin Epidemiol* 2000;53:964–72.
- 17 Yeung J, Meeks R, Edelson D, *et al.* The use of CPR feedback/prompt devices during training and CPR performance: a systematic review. *Resuscitation* 2009;80:743–51.
- 18 An M, Kim Y, Cho WK. Effect of smart devices on the quality of CPR training: a systematic review. *Resuscitation* 2019;144:145–56.
- 19 Kirkbright S, Finn J, Tohira H, *et al.* Audiovisual feedback device use by health care professionals during CPR: a systematic review and meta-analysis of randomised and non-randomised trials. *Resuscitation* 2014;85:460–71.
- 20 Wang S-A, Su C-P, Fan H-Y, *et al.* Effects of real-time feedback on cardiopulmonary resuscitation quality on outcomes in adult patients with cardiac arrest: a systematic review and meta-analysis. *Resuscitation* 2020;155:82–90.
- 21 Abellsson A, Rystedt I, Suserud B-O, *et al.* Mapping the use of simulation in prehospital care—a literature review. *Scand J Trauma Resusc Emerg Med* 2014;22:22.
- 22 Mundell WC, Kennedy CC, Szostek JH, *et al.* Simulation technology for resuscitation training: a systematic review and meta-analysis. *Resuscitation* 2013;84:1174–83.
- 23 Garden AL, Le Fevre DM, Waddington HL, *et al.* Debriefing after simulation-based non-technical skill training in healthcare: a systematic review of effective practice. *Anaesth Intensive Care* 2015;43:300–8.
- 24 Levett-Jones T, Lapkin S. A systematic review of the effectiveness of simulation debriefing in health professional education. *Nurse Educ Today* 2014;34:e58–63.
- 25 Sahu S, Lata I. Simulation in resuscitation teaching and training, an evidence based practice review. *J Emerg Trauma Shock* 2010;3:378–84.
- 26 Stoll CRT, Izadi S, Fowler S, *et al.* The value of a second reviewer for study selection in systematic reviews. *Res Synth Methods* 2019;10:539–45.
- 27 Higgins J, Thomas J. *Cochrane handbook for systematic reviews of interventions*. Cochrane, 2022.
- 28 Hong QN, Fàbregues S, Bartlett G, *et al.* The mixed methods appraisal tool (MMAT) version 2018 for information professionals and researchers. *EFI* 2018;34:285–91.
- 29 Stern C, Lizarondo L, Carrier J, *et al.* Methodological guidance for the conduct of mixed methods systematic reviews. *JBIM Evid Synth* 2020;18:2108–18.
- 30 Lizarondo L, Stern C, Carrier J, *et al.* Chapter 8: mixed methods systematic reviews. In: Aromataris E, Munn Z, eds. *Joanna Briggs Institute Reviewer's Manual*. The Joanna Briggs Institute, 2017.
- 31 Cash RE, Crowe RP, Rodriguez SA, *et al.* Disparities in feedback provision to emergency medical services professionals. *Prehosp Emerg Care* 2017;21:773–81.
- 32 Popay J, Roberts H, Sowden A, *et al.* Guidance on the conduct of narrative synthesis in systematic reviews: a product from the ESRC methods programme. Lancaster, UK University of Lancaster; 2006. Available: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.178.3.100&rep=rep1&type=pdf> [Accessed 13 Dec 2019].
- 33 Ivers NM, Sales A, Colquhoun H, *et al.* No more “business as usual” with audit and feedback interventions: towards an agenda for a reinvigorated intervention. *Implement Sci* 2014;9:14.
- 34 Ivers NM, Grimshaw JM, Jamtvedt G, *et al.* Growing literature, stagnant science? Systematic review, meta-regression and cumulative analysis of audit and feedback interventions in health care. *J Gen Intern Med* 2014;29:1534–41.
- 35 Brown B, Gude WT, Blakeman T, *et al.* Clinical performance feedback intervention theory (CP-FIT): a new theory for designing, implementing, and evaluating feedback in health care based on a systematic review and meta-synthesis of qualitative research. *Implement Sci* 2019;14:40.
- 36 Carey RN, Connell LE, Johnston M, *et al.* Behavior change techniques and their mechanisms of action: a synthesis of links described in published intervention literature. *Ann Behav Med* 2019;53:693–707.
- 37 Gardner B, Whittington C, McAteer J, *et al.* Using theory to synthesise evidence from behaviour change interventions: the example of audit and feedback. *Soc Sci Med* 2010;70:1618–25.
- 38 Tuti T, Nzinga J, Njoroge M, *et al.* A systematic review of electronic audit and feedback: intervention effectiveness and use of behaviour change theory. *Implement Sci* 2017;12:61.

- 39 Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw* 2010;36:1–48.
- 40 Cohen J. *Statistical power analysis for the behavioural sciences*. New York: Academic Press, 1969.
- 41 Lenhard W, Lenhard A. Computation of effect sizes: psychometrica. 2016. Available: https://www.psychometrica.de/effect_size.html [Accessed 24 Feb 2022].
- 42 Wilson DB. Practical meta-analysis effect size calculator. 2022. Available: <https://www.campbellcollaboration.org/escalc/html/EffectSizeCalculator-Home.php> [Accessed 24 Feb 2022].
- 43 Lin H. Effect size converter. 2022. Available: <https://www.escal.site/> [Accessed 24 Feb 2022].
- 44 Wan X, Wang W, Liu J, *et al*. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014;14:135.
- 45 Borenstein M, Hedges LV, Higgins J, *et al*. *When does it make sense to perform a meta-analysis? Introduction to meta-analysis*. Chichester: John Wiley & Sons, 2009: 357–64.
- 46 Assink M, Wibbelink CJM. Fitting three-level meta-analytic models in R: a step-by-step tutorial. *TQMP* 2016;12:154–74.
- 47 Hysong SJ. Meta-analysis: audit and feedback features impact effectiveness on care quality. *Med Care* 2009;47:356–63.
- 48 Harrer M, Cuijpers P, Furukawa TA, *et al*. *Doing meta-analysis with R: a hands-on guide*. 1st ed. Boca Raton, FL and London: Chapman & Hall/CRC Press, 2021.
- 49 West SL, Gartlehner G, Mansfield AJ, *et al*. AHRQ methods for effective health care. In: *Comparative Effectiveness Review Methods: clinical Heterogeneity*. Rockville (MD): Agency for Healthcare Research and Quality (US), 2010.
- 50 von Hippel PT. The heterogeneity statistic I (2) can be biased in small meta-analyses. *BMC Med Res Methodol* 2015;15:35.
- 51 Rücker G, Schwarzer G, Carpenter JR, *et al*. Undue reliance on I (2) in assessing heterogeneity may mislead. *BMC Med Res Methodol* 2008;8:79.
- 52 Int'Hout J, Ioannidis JPA, Rovers MM, *et al*. Plea for routinely presenting prediction intervals in meta-analysis. *BMJ Open* 2016;6:e010247.
- 53 Siriwardena AN, Shaw D, Essam N, *et al*. The effect of a national quality improvement collaborative on prehospital care for acute myocardial infarction and stroke in England. *Implement Sci* 2014;9:17.
- 54 Weston BW, Jasti J, Mena M, *et al*. Self-assessment feedback form improves quality of out-of-hospital CPR. *Prehosp Emerg Care* 2019;23:66–73.
- 55 Choi B, Tsai D, McGillivray CG, *et al*. Hospital-directed feedback to emergency medical services improves prehospital performance. *Stroke* 2014;45:2137–40.
- 56 Hardeland C, Skåre C, Kramer-Johansen J, *et al*. Targeted simulation and education to improve cardiac arrest recognition and telephone assisted CPR in an emergency medical communication centre. *Resuscitation* 2017;114:21–6.
- 57 Hopkins CL, Burk C, Moser S, *et al*. Implementation of pit crew approach and cardiopulmonary resuscitation metrics for out-of-hospital cardiac arrest improves patient survival and neurological outcome. *J Am Heart Assoc* 2016;5:e002892.
- 58 Stella J, Bartley B, Jennings P. Introduction of a prehospital critical incident monitoring system -- final results. *Prehosp Disaster Med* 2010;25:515–20.
- 59 Lindström V, Karlsten R, Falk A-C, *et al*. Feasibility of a computer-assisted feedback system between dispatch centre and ambulances. *Eur J Emerg Med* 2011;18:143–7.
- 60 Riney LC, Schwartz H, Murtagh Kurowski E, *et al*. Improving administration of prehospital corticosteroids for pediatric asthma. *Pediatr Qual Saf* 2021;6:e410.
- 61 Lukas R-P, Gräsner JT, Seewald S, *et al*. Chest compression quality management and return of spontaneous circulation: a matched-pair registry study. *Resuscitation* 2012;83:1212–8.
- 62 Olasveengen TM, Tomlinson A-E, Wik L, *et al*. A failed attempt to improve quality of out-of-hospital CPR through performance evaluation. *Prehosp Emerg Care* 2007;11:427–33.
- 63 Oostema JA, Chassee T, Baer W, *et al*. Brief educational intervention improves emergency medical services stroke recognition. *Stroke* 2019;50:1193–200.
- 64 Persse DE, Key CB, Baldwin JB. The effect of a quality improvement feedback loop on paramedic-initiated nontransport of elderly patients. *Prehosp Emerg Care* 2002;6:31–5.
- 65 Bobrow BJ, Spaite DW, Vadeboncoeur TF, *et al*. Implementation of a regional telephone cardiopulmonary resuscitation program and outcomes after out-of-hospital cardiac arrest. *JAMA Cardiol* 2016;1:294–302.
- 66 Joyce SM, Dutkowski KL, Hynes T. Efficacy of an EMS quality improvement program in improving documentation and performance. *Prehosp Emerg Care* 1997;1:140–4.
- 67 Scholz KH, Hilgers R, Ahlersmann D, *et al*. Contact-to-balloon time and door-to-balloon time after initiation of a formalized data feedback in patients with acute ST-elevation myocardial infarction. *Am J Cardiol* 2008;101:46–52.
- 68 Daudelin DH, Sayah AJ, Kwong M, *et al*. Improving use of prehospital 12-lead ECG for early identification and treatment of acute coronary syndrome and ST-elevation myocardial infarction. *Circ Cardiovasc Qual Outcomes* 2010;3:316–23.
- 69 Gropen TI, Gazi M, Minor M, *et al*. Centrally guided identification of patients with large vessel occlusion: lessons from trauma systems. *J Stroke Cerebrovasc Dis* 2019;28:2388–97.
- 70 DelliFraine J, Langabeer J, Segrest W, *et al*. Developing an ST-elevation myocardial infarction system of care in dallas county. *Am Heart J* 2013;165:926–31.
- 71 Gevers L, Makkink A. Peer-assisted learning in the prehospital educational setting in South Africa. *AJHP* 2020;12:6.
- 72 Noble HE, Scott JW, Nyinawankusi JD, *et al*. The impact of data feedback on continuous quality improvement projects in rwanda: a mixed methods analysis. *Afr J Emerg Med* 2020;10:578–84.
- 73 Brink P, Bäck-Petersson S, Sernert N. Group supervision as a means of developing professional competence within pre-hospital care. *Int Emerg Nurs* 2012;20:76–82.
- 74 Mock EF, Wrenn KD, Wright SW, *et al*. Feedback to emergency medical services providers: the good, the bad, and the ignored. *Prehosp Disaster Med* 1997;12:145–8.
- 75 McGuire SS, Luke A, Klassen AB, *et al*. It's time to talk to prehospital providers: feedback disparities among ground-based emergency medical services providers and its impact on job satisfaction. *Prehosp Disaster Med* 2021;36:486–94.
- 76 Porter A, Badshah A, Black S, *et al*. Electronic health records in ambulances: the era multiple-methods study. *Health Serv Deliv Res* 2020;8:1–140.
- 77 Cuk S, Wimmer H, Powell LM. Problems associated with patient care reports and transferring data

- between ambulance and hospitals from the perspective of emergency medical technicians. *Issues Inf Syst* 2017;18:16–26.
- 78 Eaton-Williams P, Mold F, Magnusson C. Effective clinical feedback provision to ambulance clinicians: a literature review. *J Paramed Pract* 2020;12:109–17.
- 79 Colquhoun HL, Carroll K, Eva KW, *et al.* Advancing the literature on designing audit and feedback interventions: identifying theory-informed hypotheses. *Implement Sci* 2017;12:117.
- 80 Gude WT, Brown B, van der Veer SN, *et al.* Clinical performance comparators in audit and feedback: a review of theory and evidence. *Implement Sci* 2019;14:39.
- 81 Ulrich Hansen M, Vejzovic V, Zdravkovic S, *et al.* Ambulance nurses' experiences of using prehospital guidelines for patients with acute chest pain—a qualitative study. *Int Emerg Nurs* 2022;63:101195.
- 82 O'Hara R, Johnson M, Hirst E, *et al.* A qualitative study of decision-making and safety in ambulance service transitions. *Health Serv Deliv Res* 2014;2:1–138.
- 83 Hörberg A, Lindström V, Kalén S, *et al.* Striving for balance—a qualitative study to explore the experiences of nurses new to the ambulance service in sweden. *Nurse Educ Pract* 2017;27:63–70.
- 84 Wihlborg J, Edgren G, Johansson A, *et al.* Reflective and collaborative skills enhances ambulance nurses' competence—a study based on qualitative analysis of professional experiences. *Int Emerg Nurs* 2017;32:20–7.
- 85 Bång A, Ortgren P-O, Herlitz J, *et al.* Dispatcher-assisted telephone CPR: a qualitative study exploring how dispatchers perceive their experiences. *Resuscitation* 2002;53:135–51.
- 86 Bobrow BJ, Vadeboncoeur TF, Clark L, *et al.* Establishing Arizona's statewide cardiac arrest reporting and educational network. *Prehosp Emerg Care* 2008;12:381–7.
- 87 Davis DR, Fisher R, Buono C, *et al.* Predictors of intubation success and therapeutic value of paramedic airway management in a large, urban EMS system. *Prehosp Emerg Care* 2006;10:356–62.
- 88 Dyson K, Bray JE, Smith K, *et al.* Paramedic resuscitation competency: A survey of australian and new zealand emergency medical services. *Emerg Med Australas* 2017;29:217–22.
- 89 Henry K, Murphy A, Willis D, *et al.* Out-of-hospital cardiac arrest in cork, ireland. *Emerg Med J* 2013;30:496–500.
- 90 Larsson R, Engström Å. Swedish ambulance nurses' experiences of nursing patients suffering cardiac arrest. *Int J Nurs Pract* 2013;19:197–205.
- 91 van Diepen S, Abella BS, Bobrow BJ, *et al.* Multistate implementation of guideline-based cardiac resuscitation systems of care: description of the heartrescue project. *Am Heart J* 2013;166:647–53.
- 92 Yost D, Phillips RH, Gonzales L, *et al.* Assessment of CPR interruptions from transthoracic impedance during use of the LUCAS (TM) mechanical chest compression system. *Resuscitation* 2012;83:961–5.
- 93 Lee SCL, Mao DR, Ng YY, *et al.* Emergency medical dispatch services across pan-asian countries: a web-based survey. *BMC Emerg Med* 2020;20:1.
- 94 Bradley EH, Curry LA, Spatz ES, *et al.* Hospital strategies for reducing risk-standardized mortality rates in acute myocardial infarction. *Ann Intern Med* 2012;156:618–26.
- 95 Landman AB, Spatz ES, Cherlin E, *et al.* Abstract 246: emergency medical services and hospital collaboration in the care of patients with acute myocardial infarction. *Circ Cardiovasc Qual Outcomes* 2012;5.
- 96 Martin LK, Lewis VJ, Clark D, *et al.* Frontline barriers to effective paramedic and emergency nursing STEMI management: clinician perspectives. *Australas Emerg Care* 2020;23:126–36.
- 97 Brunton L, Boaden R, Knowles S, *et al.* Pre-hospital stroke recognition in a UK centralised stroke system: a qualitative evaluation of current practice. *Br Paramed J* 2019;4:31–9.
- 98 Magdon-Ismail Z, Benesch C, Cushman JT, *et al.* Establishing recommendations for stroke systems in the thrombectomy era: the upstate new york stakeholder proceedings. *Stroke* 2017;48:2003–6.
- 99 Tegelerberg A, Muntlin Å, Juhlin C, *et al.* Engagement under difficult conditions: caring for patients with acute abdominal pain across the acute-care chain: a qualitative study. *Int Emerg Nurs* 2020;52:100910.
- 100 Brown SA, Hayden TC, Randell KA, *et al.* Improving pediatric education for emergency medical services providers: a qualitative study. *Prehosp Disaster Med* 2017;32:20–6.
- 101 Hoogervorst EM, van Beeck EF, Goslings JC, *et al.* Developing process guidelines for trauma care in the netherlands for severely injured patients: results from a delphi study. *BMC Health Serv Res* 2013;13:79.
- 102 Repas SJ, McCarthy M, Parikh P, *et al.* Trauma-specific performance improvement activities for emergency medical services providers: a statewide perspective. *J Surg Res* 2022;279:474–9.
- 103 Whitley TW, Benson NH, Allison EJ, *et al.* Occupational stress and job satisfaction among flight nurses. A survey of national flight nurses association members. *Hosp Aviat* 1989;8:6–8.
- 104 Forslund K, Kihlgren A, Kihlgren M. Operators' experiences of emergency calls. *J Telemed Telecare* 2004;10:290–7.
- 105 Lateef F. Job-related evaluation of post-traumatic stress (JETS). *Singap Nurs J* 2005;32:37–41.
- 106 Minnie L, Goodman S, Wallis L. Exposure to daily trauma: the experiences and coping mechanism of emergency medical personnel. A cross-sectional study. *Afr J Emerg Med* 2015;5:12–8.
- 107 Crowe RP, Fernandez AR, Pepe PE, *et al.* The association of job demands and resources with burnout among emergency medical services professionals. *J Am Coll Emerg Physicians Open* 2020;1:6–16.
- 108 Holmberg M, Nørgaard J, Eriksson M, *et al.* Dyadic teams and nursing care: a critical incident study of nurses in the emergency medical service. *J Clin Nurs* 2020;29:3743–53.
- 109 Smith MP, Kaji A, Young KD, *et al.* Presentation and survival of prehospital apparent sudden infant death syndrome. *Prehosp Emerg Care* 2005;9:181–5.
- 110 Lederman J, Löfvenmark C, Djärv T, *et al.* Assessing non-conveyed patients in the ambulance service: a phenomenological interview study with swedish ambulance clinicians. *BMJ Open* 2019;9:e030203.
- 111 Ivic R, Vicente V, Kurland L, *et al.* Pre-hospital emergency nurse specialist's experiences in caring for patients with non-specific chief complaints in the ambulance - A qualitative interview study. *Int Emerg Nurs* 2022;63:101178.
- 112 Brichko L, Jennings P, Bain C, *et al.* Selecting cases for feedback to pre-hospital clinicians—a pilot study. *Aust Health Rev* 2016;40:306–10.
- 113 Ter Avest E, McWhirter E, Dunn S, *et al.* Prehospital death after traumatic cardiac arrest: time for better feedback? *Air Med J* 2019;38:78–81.

- 114 Kunkes T, Makled B, Norfleet J, *et al.* Understanding the cognitive demands, skills, and assessment approaches for endotracheal intubation: cognitive task analysis. *JMIR Perioper Med* 2022;5:e34522.
- 115 Donovan JK, Burton SO, Jones SL, *et al.* Use of point-of-care ultrasound by intensive care paramedics to assess respiratory distress in the out-of-hospital environment: a pilot study. *Prehosp Emerg Care* 2022;2022:1–7.
- 116 Jaynes CL, Cook P, Farmer R, *et al.* Assessing satisfaction and quality in the EMS/HEMS working relationship. *Air Med J* 2013;32:338–42.
- 117 Meisel ZF, Shea JA, Peacock NJ, *et al.* Optimizing the patient handoff between emergency medical services and the emergency department. *Ann Emerg Med* 2015;65:310–7.
- 118 Jönsson K, Fridlund B. A comparison of adherence to correctly documented triage level of critically ill patients between emergency department and the ambulance service nurses. *Int Emerg Nurs* 2013;21:204–9.
- 119 Hörberg A, Jirwe M, Kalén S, *et al.* We need support! A delphi study about desirable support during the first year in the emergency medical service. *Scand J Trauma Resusc Emerg Med* 2017;25:89.
- 120 Hörberg A, Kalén S, Jirwe M, *et al.* Treat me nice! -a cross-sectional study examining support during the first year in the emergency medical services. *Scand J Trauma Resusc Emerg Med* 2018;26:92.
- 121 Eaton G, Happs I, Tanner R. Designing and implementing an educational framework for advanced paramedic practitioners rotating into primary care in north wales. *Educ Prim Care* 2021;32:289–95.
- 122 Fjällman C, Hilli Y, Jonasson L-L. Home health nurses' views on participating in pre-ambulance emergency treatment: a qualitative descriptive study. *Nord J Nurs Res* 2021;41:92–100.
- 123 Brayman C, Hobbs B, Hill W, *et al.* ICU without walls-interprofessional high acuity response teams (harts) improve access to higher level of care in rural and remote communities. *Can J Respir Ther* 2012;48:14–9.
- 124 Donald M, Paterson B. Introduction of an electronic debrief and governance tool in prehospital care. *Emerg Med J* 2007;24:363–6.
- 125 Akula VB, Hedli LC, Van Meurs K, *et al.* Neonatal transport in california: findings from a qualitative investigation. *J Perinatol* 2020;40:394–403.
- 126 Heffernan E, Keegan D, Clarke B, *et al.* Quality improvement in A crisis: a qualitative study of experiences and lessons learned from the irish national ambulance service response to the COVID-19 pandemic. *BMJ Open* 2022;12:e057162.
- 127 Clarke S, Lyon RM, Short S, *et al.* A specialist, second-tier response to out-of-hospital cardiac arrest: setting up TOPCAT2. *Emerg Med J* 2014;31:405–7.
- 128 Ebbs P, Middleton PM, Bonner A, *et al.* Do clinical safety charts improve paramedic key performance indicator results? (a clinical improvement programme evaluation). *Emerg Med J* 2012;29:596–7.
- 129 Eckstein M, Alo K. The effect of a quality improvement program on paramedic on-scene times for patients with penetrating trauma. *Acad Emerg Med* 1999;6:191–5.
- 130 Hubner P, Lobmeyr E, Wallmüller C, *et al.* Improvements in the quality of advanced life support and patient outcome after implementation of a standardized real-life post-resuscitation feedback system. *Resuscitation* 2017;120:38–44.
- 131 Park JH, Shin SD, Ro YS, *et al.* Implementation of a bundle of utstein cardiopulmonary resuscitation programs to improve survival outcomes after out-of-hospital cardiac arrest in a metropolis: a before and after study. *Resuscitation* 2018;130:124–32.
- 132 Tanaka Y, Taniguchi J, Wato Y, *et al.* The continuous quality improvement project for telephone-assisted instruction of cardiopulmonary resuscitation increased the incidence of bystander CPR and improved the outcomes of out-of-hospital cardiac arrests. *Resuscitation* 2012;83:1235–41.
- 133 Tödt T, Thylén I, Alfredsson J, *et al.* Strategies to reduce time delays in patients with acute coronary heart disease treated with primary PCI -- the stop watch study: a multistage action research project. *BMJ Open* 2013;3:e003493.
- 134 Lyon RM, Clarke S, Milligan D, *et al.* Resuscitation feedback and targeted education improves quality of pre-hospital resuscitation in Scotland. *Resuscitation* 2012;83:70–5.
- 135 Walters G, D'Auria D, Glucksman E. Automated external defibrillators: implications for training qualified ambulance staff. *Ann Emerg Med* 1992;21:692–7.
- 136 Bleijenberg E, Koster RW, de Vries H, *et al.* The impact of post-resuscitation feedback for paramedics on the quality of cardiopulmonary resuscitation. *Resuscitation* 2017;110:1–5.
- 137 Clawson JJ, Cady GA, Martin RL, *et al.* Effect of a comprehensive quality management process on compliance with protocol in an emergency medical dispatch center. *Ann Emerg Med* 1998;32:578–84.
- 138 O'Connor RE, Megargel RE. The effect of a quality improvement feedback loop on paramedic skills, charting, and behavior. *Prehosp Disaster Med* 1994;9:35–8.
- 139 Scholz KH, Friede T, Meyer T, *et al.* Prognostic significance of emergency department bypass in stable and unstable patients with ST-segment elevation myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2020;9:34–44.
- 140 Scholz KH, Lengenfelder B, Jacobshagen C, *et al.* Long-term effects of a standardized feedback-driven quality improvement program for timely reperfusion therapy in regional STEMI care networks. *Eur Heart J Acute Cardiovasc Care* 2020;2048872620907323.
- 141 Scholz KH, Maier SKG, Jung J, *et al.* Reduction in treatment times through formalized data feedback: results from a prospective multicenter study of ST-segment elevation myocardial infarction. *JACC Cardiovasc Interv* 2012;5:848–57.
- 142 Scholz KH, Meyer T, Lengenfelder B, *et al.* Patient delay and benefit of timely reperfusion in ST-segment elevation myocardial infarction. *Open Heart* 2021;8:e001650.
- 143 Scott JW, Nyinawankusi JD, Enumah S, *et al.* Improving prehospital trauma care in rwanda through continuous quality improvement: an interrupted time series analysis. *Injury* 2017;48:1376–81.
- 144 Swor RA, Hoelzer M. A computer-assisted quality assurance audit in a multiprovider EMS system. *Ann Emerg Med* 1990;19:286–90.
- 145 Bahouth H, Abramov R, Bodas M, *et al.* The feedback form and its role in improving the quality of trauma care. *Int J Environ Res Public Health* 2022;19:1866.