

University of Groningen

A circadian clock in *Saccharomyces cerevisiae*

Eelderink-Chen, Zheng; Mazzotta, Gabriella; Sturre, Marcel; Bosman, Jasper; Roenneberg, Till; Mellow, Martha

Published in:

Proceedings of the National Academy of Sciences of the United States of America

DOI:

[10.1073/pnas.0907902107](https://doi.org/10.1073/pnas.0907902107)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Eelderink-Chen, Z., Mazzotta, G., Sturre, M., Bosman, J., Roenneberg, T., & Mellow, M. (2010). A circadian clock in *Saccharomyces cerevisiae*. *Proceedings of the National Academy of Sciences of the United States of America*, 107(5), 2043-2047. <https://doi.org/10.1073/pnas.0907902107>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supporting Information

Eelderink-Chen et al. 10.1073/pnas.0907902107

SI Materials

Entrainment of Oscillators. The first “modern” record of entrainment of oscillators is attributed to the Dutch scientist, Christian Huygens (1). He was attempting to solve the problem of keeping accurate time at sea and thus had multiple pendula swinging simultaneously. As he lay ill in bed one day (it was the custom to have one’s bed off to the side of the living area), he noted that the pendula on the adjacent wall were synchronized. When he perturbed them they would resynchronize after some minutes. This held only for pendula on the same wall. They could thus communicate with each other and entrain each other.

This phenomenon is comparable to circadian rhythms synchronizing to the external zeitgeber cycle. The physical cycles need to communicate in some way with the biological ones (those of the circadian clock). Most entrainment mechanisms have been defined using light as a zeitgeber; it is an important zeitgeber for most circadian systems and it is experimentally facile to use. In the case of the work here, the external cycle uses temperature as a zeitgeber. There is virtually nothing known concerning the mechanisms by which temperature entrains circadian rhythms. It may be changes in biochemical reactions in the cell or it may be via specialized temperature sensors, akin to photoreceptors. Regardless of the mechanisms, temperature is apparently also a universal zeitgeber for circadian systems. In the case of homeotherms, low-amplitude temperature cycles are effective synchronizers (2). In the case of poikilotherms, the experienced temperature cycles are typically much higher.

Temperature acts to synchronize the metabolism of yeast in a highly systematic way like what was first demonstrated by Hoffmann, when he put lizards into temperature cycles of different length, showing that they would entrain later as cycles became shorter (3, 4). This phenomenon turns out to be one of the circadian rules that even contributes to the explanation of chronotype, the distinct entrained phase of an individual. In the general population, there is a distribution of chronotypes (5) and this is thought to be due to differences in genetic background resulting in differences in free-running period (and probably other circadian properties, such as zeitgeber input pathways, as well). Indeed, in the 1970s, the concept of zeitgeber strength as it regulates chronotype was demonstrated in hamsters (6).

Obviously, the environmental cycle is the cycle to which the biological one must adjust. If the biological clock would simply respond with a switch-like mechanism (becoming active when the sun comes up, for instance), then the organism would need no biological clock. A circadian clock, however, allows differential entrainment according to season, for instance, thus allowing gating of seasonal behaviors such as reproduction (7, 8). The experiments described in this paper use these basic ideas, namely that the absence of a clock should yield driven responses and, conversely, the presence of a clock should yield entrainment in a similar manner as has been demonstrated for other circadian systems. We have applied short temperature cycles, which should show a biological clock entraining to a later phase (3, 4, 9, 10). We have altered the zeitgeber strength, which would be predicted to change the entrained phase of a circadian system (5, 10).

1. Huygens C (1673) *Horologium Oscillatorium sive de motu pendulorum* (Apud F. Muguet, Parisiis).
2. Brown SA, Zimbrunn G, Fleury-Olela F, Preitner N, Schibler U (2002) Rhythms of mammalian body temperature can sustain peripheral circadian clocks. *Curr Biol* 12: 1574–1583.
3. Hoffmann K (1963) Zur Beziehung zwischen Phasenlage und Spontanfrequenz bei der endogenen Tagesperiodik. *Z Naturforsch* 18b:154–157.
4. Hoffmann K, ed (1968) *Temperaturzyklen als Zeitgeber der circadianen Periodik* (Verh. Deut. Zool. Ges. Innsbruck, Austria), pp 265–275.
5. Roenneberg T, et al. (2004) A marker for the end of adolescence. *Curr Biol* 14: R1038–R1039.
6. Pittendrigh CS, Daan S (1976) A functional analysis of circadian pacemakers in nocturnal rodents. IV. Entrainment: Pacemaker as clock. *J Comp Physiol [A]* 106:291–331.
7. Elliott JA (1976) Circadian rhythms and photoperiodic time measurement in mammals. *Fed Proc* 35:2339–2346.
8. Bünning E (1936) Die endogene Tagesperiodik als Grundlage der photoperiodischen Reaktion. *Ber Dtsch Bot Ges* 54:590–608.
9. Mellow M, Brunner M, Roenneberg T (1999) Assignment of circadian function for the *Neurospora* clock gene frequency. *Nature* 399:584–586.
10. Roenneberg T, Dragovic Z, Mellow M (2005) Demasking biological oscillators: Properties and principles of entrainment exemplified by the *Neurospora* circadian clock. *Proc Natl Acad Sci USA* 102:7742–7747.




Fig. 53. Oscillations following release to constant conditions. In two experiments, the pH did not change substantially on release from a 24-h temperature cycle to constant conditions. In these cases, the period of the nonentrained, free-running oscillation was closer to 24 h and did not appear to be unstable (although it did damp rapidly). These two examples contrast what is more typically observed in the fermentor cultures (see Fig. 3 and note the reproducibility therein between experiments). (A) 18/25 °C temperature cycle with a release to 25 °C. (B) 21/28 °C cycle with a release to 28 °C.




Fig. 54. RNA expression during entrainment with temperature. MEP2 (A) and GAP1 (B) mRNA levels were determined in a 24-h 21/28 °C temperature cycle. In each graph, two separate experiments are shown, with MEP2/GAP1 RNA as dashed and dotted lines (representing the two experiments). The oscillations in proton concentrations are shown in the same graphs (solid lines). The lines with squares as markers correspond to the same experiment; the lines with triangles and diamonds are samples from the same experiment. The gray and open bars at the Top of the graphs indicate cold and warm phases of the entraining cycle, respectively. Tubulin was used for normalization; for each time series, the high value is set to 1.