

University of Groningen

## Formal normal form of $A_k$ slow-fast systems

Jardón-kojakhmetov, Hildeberto

*Published in:*  
Comptes Rendus Mathématique

*DOI:*  
[10.1016/j.crma.2015.06.009](https://doi.org/10.1016/j.crma.2015.06.009)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2015

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*  
Jardón-kojakhmetov, H. (2015). Formal normal form of  $A_k$  slow-fast systems. *Comptes Rendus Mathématique*, 353(9), 795-800. <https://doi.org/10.1016/j.crma.2015.06.009>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Ordinary differential equations/Dynamical systems

Formal normal form of  $A_k$  slow–fast systems*Forme normale formelle des systèmes lents–rapides de type  $A_k$* 

Hildeberto Jardón-Kojakhmetov

Johann Bernoulli Institute for Mathematics and Computer Science, University of Groningen, P.O. Box 407, 9700 AK, Groningen, The Netherlands

## ARTICLE INFO

## Article history:

Received 18 March 2015

Accepted after revision 25 June 2015

Available online 16 July 2015

Presented by the Editorial Board

## ABSTRACT

An  $A_k$  slow–fast system is a particular type of singularly perturbed ODE. The corresponding slow manifold is defined by the critical points of a universal unfolding of an  $A_k$  singularity. In this note we propose a formal normal form of  $A_k$  slow–fast systems.

© 2015 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

## R É S U M É

Un système lent–rapide de type  $A_k$  est une équation différentielle ordinaire singulièrement perturbée avec une structure particulière. La variété lente correspondante est définie par les points critiques d'un déploiement universel d'une singularité de type  $A_k$ . Dans cette note, nous proposons une forme normale formelle des systèmes lents–rapides de type  $A_k$ .

© 2015 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

## 1. Introduction

A slow–fast system (SFS) is a singularly perturbed ODE usually written as

$$\begin{aligned}\dot{x} &= f(x, z, \varepsilon) \\ \varepsilon \dot{z} &= g(x, z, \varepsilon)\end{aligned}\tag{1}$$

where  $x \in \mathbb{R}^m$ ,  $z \in \mathbb{R}^n$  and  $0 < \varepsilon \ll 1$  is a small parameter, and where the over-dot denotes the derivative with respect to a time parameter  $t$ . Slow–fast systems are often used as mathematical models of phenomena that occur in two time scales. A couple of classical examples of real life phenomena that were modeled by an SFS are Zeeman's heartbeat and nerve-impulse models [17]. For  $\varepsilon \neq 0$ , we can define a new time parameter  $\tau$  by  $t = \varepsilon\tau$ . With this new time  $\tau$ , we can write (1) as

$$\begin{aligned}x' &= \varepsilon f(x, z, \varepsilon) \\ z' &= g(x, z, \varepsilon),\end{aligned}\tag{2}$$

where the prime denotes the derivative with respect to  $\tau$ . An important geometric object in the study of SFSs is the *slow manifold*, which is defined by

E-mail address: h.jardon.kojakhmetov@gmail.com.

$$S = \{(x, z) \in \mathbb{R}^m \times \mathbb{R}^n \mid g(x, z, 0) = 0\}. \quad (3)$$

When  $\varepsilon = 0$ , the manifold  $S$  serves as the phase space of (1) and as the set of equilibrium points of (2). In the rest of the document, we prefer to work with an SFS written as (2). Furthermore, to avoid working with an  $\varepsilon$ -parameter family of vector fields as in (2), we extend (2) by adding the trivial equation  $\varepsilon' = 0$ . To be more precise, we treat a  $C^\infty$ -smooth vector field defined as follows.

**Definition 1.1** ( *$A_k$  slow-fast system*). Let  $k \in \mathbb{N}$  with  $k \geq 2$ . An  $A_k$  slow-fast system (for short  $A_k$ -SFS) is a vector field  $X$  of the form

$$X = \varepsilon(1 + \varepsilon f_1) \frac{\partial}{\partial x_1} + \sum_{i=2}^{k-1} \varepsilon^2 f_i \frac{\partial}{\partial x_i} - (G_k - \varepsilon f_k) \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}, \quad (4)$$

where  $G_k = z^k + \sum_{i=1}^{k-1} x_i z^{i-1}$  and where each  $f_i = f_i(x_1, \dots, x_{k-1}, z, \varepsilon)$  is a  $C^\infty$ -smooth function vanishing at the origin.

Multi-scale models described by an  $A_k$ -SFS are of interest as they exhibit local, fast transitions between stable states, e.g., [8,13,14,17].

**Remark 1.1.** The slow manifold associated with an  $A_k$ -SFS is defined by

$$S = \left\{ (x, z) \in \mathbb{R}^k \mid z^k + \sum_{i=1}^{k-1} x_i z^{i-1} = 0 \right\}. \quad (5)$$

The manifold  $S$  can be regarded as the critical set of the universal unfolding of a smooth function with an  $A_k$  singularity at the origin [1,3]. Hence the name  $A_k$ -SFS.

Observe that the origin is a non-hyperbolic equilibrium point of  $X$  and thus, it is not possible to study its local dynamics with the classical Geometric Singular Perturbation Theory [5]. In this case, the blow-up technique [4,9] can be applied to desingularize the SFS. This methodology has been successfully used in many cases, e.g., [2,7,10,11,15,16], where many of these deal with an  $A_k$ -SFS with fixed  $k = 2$  or  $k = 3$ . Briefly speaking, the blow-up technique consists in an appropriate change of coordinates under which the induced vector field is regular or has simpler singularities (hyperbolic or partially hyperbolic).

In this paper we propose a normal form of  $A_k$ -SFSs given by Definition 1.1. In this normalization, the unknown functions  $f_i$  of (4) are eliminated. As it is shown below, the structure of the  $A_k$ -SFS plays an important role in the normalization process. Moreover, this normalization greatly simplifies the local analysis of systems given by (4), as shown in [6,7].

## 2. Formal normal form of an $A_k$ slow-fast system

We regard the vector field  $X$  of Definition 1.1 as  $X = F + P$ , where  $F$  and  $P$  are smooth vector fields called “the principal part” and “the perturbation” respectively. That is:

$$F = \varepsilon \frac{\partial}{\partial x_1} + \sum_{i=2}^{k-1} 0 \frac{\partial}{\partial x_i} - G_k \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}, \quad P = \sum_{i=1}^{k-1} \varepsilon^2 f_i \frac{\partial}{\partial x_i} + \varepsilon f_k \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}. \quad (6)$$

The idea of the rest of the document is motivated by [12]. In short, we want to formally simplify the expression of  $X$  by eliminating the perturbation  $P$ . The terminology used below is that of [12].

The vector field  $F$  is quasihomogeneous of type  $r = (k, k-1, \dots, 1, 2k-1)$  and quasidegree  $k-1$  [1,12]. From now on, we fix the type of quasihomogeneity  $r$ . A quasihomogeneous object of type  $r$  will be called  $r$ -quasihomogeneous.

**Definition 2.1** (*Good perturbation*). Let  $F$  be an  $r$ -quasihomogeneous vector field of quasidegree  $k-1$ . A good perturbation  $X$  of  $F$  is a smooth vector field  $X = F + P$ , where  $P = P(x_1, \dots, x_{k-1}, z, \varepsilon)$  satisfies the following conditions:

- $P$  is a smooth vector field of quasiorder greater than  $k-1$ ,
- $P = \sum_{i=1}^{k-1} P_i \frac{\partial}{\partial x_i} + P_k \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}$ , with  $P|_{\varepsilon=0} = 0$ .

*Notation* By  $\mathcal{P}_\delta$  we denote the space of  $r$ -quasihomogeneous polynomials (in  $k+1$  variables) of quasidegree  $\delta$ . By  $\mathcal{H}_\gamma$  we denote the space of  $r$ -quasihomogeneous vector fields (in  $\mathbb{R}^{k+1}$ ) of quasidegree  $\gamma$  and such that for all  $U \in \mathcal{H}_\gamma$  we have  $U = \sum_{i=1}^k U_k \frac{\partial}{\partial x_i} + 0 \frac{\partial}{\partial x_{k+1}}$ . The formal series expansion of a function  $f$  is denoted by  $\hat{f}$ .

**Definition 2.2** (The inner product  $\langle \cdot, \cdot \rangle_{r,\delta}$  [12]). Let  $x = (x_1, \dots, x_n)$ , and  $s, q \in \mathbb{N}^n$ . Let  $f, g \in \mathcal{P}_\delta$ , that is  $f = \sum_{(r,s)=\delta} f_s x^s$ , where  $f_s \in \mathbb{R}$ ,  $x^s = x_1^{s_1} \cdots x_n^{s_n}$ ; and similarly for  $g$ . Then the inner product  $\langle \cdot, \cdot \rangle_{r,\delta}$  is defined as

$$\langle f, g \rangle_{r,\delta} = \sum_{(r,s)=\delta} f_s g_s \frac{(s!)^r}{\delta!}, \tag{7}$$

where  $(s!)^r = (s_1!)^{r_1} \cdots (s_n!)^{r_n}$ , and where  $(r, s)$  denotes the dot product  $r \cdot s$ . So for monomials one has

$$\langle x^s, x^q \rangle_{r,\delta} = \begin{cases} \frac{(s_1!)^{r_1} \cdots (s_n!)^{r_n}}{\delta!} & \text{if } s = q \text{ with } (s, r) = \delta, \\ 0 & \text{otherwise.} \end{cases} \tag{8}$$

Accordingly, for vector fields: let  $X = \sum_{i=1}^n X_i \frac{\partial}{\partial x_i} \in \mathcal{H}_\delta$ , and  $Y = \sum_{i=1}^n Y_i \frac{\partial}{\partial x_i} \in \mathcal{H}_\delta$ . Then

$$\langle X, Y \rangle_{r,\delta} = \sum_{i=1}^n \langle X_i, Y_i \rangle_{r,\delta+r_i}. \tag{9}$$

**Definition 2.3** (The operators  $d, d^*$  and  $\square$  [12]). The operator  $d : \mathcal{H}_\gamma \rightarrow \mathcal{H}_{\gamma+k-1}$  (associated with  $F$ ) is defined by  $d(U) = [F, U]$  for any  $U \in \mathcal{H}_\gamma$ , where  $[\cdot, \cdot]$  denotes the Lie bracket. The operator  $d^*$  is the adjoint operator of  $d$  with respect to the inner product of Definition 2.2. This is, given  $U \in \mathcal{H}_\gamma, V \in \mathcal{H}_{\gamma+k-1}$  we have

$$\langle d(U), V \rangle_{r,\gamma+k-1} = \langle U, d^*(V) \rangle_{r,\gamma}. \tag{10}$$

For any quasidegree  $\beta > k - 1$ , the self adjoint operator  $\square_\beta : \mathcal{H}_\beta \rightarrow \mathcal{H}_\beta$  is defined by  $\square_\beta(U) = dd^*(U)$  for all  $U \in \mathcal{H}_\beta$ .

**Definition 2.4** (Resonant vector field [12]).

- We say that a vector field  $U \in \mathcal{H}_\beta$  is resonant if  $U \in \ker \square_\beta$ .
- A formal vector field is called resonant if all its quasihomogeneous components are resonant.

**Definition 2.5** (Normal form [12]). A good perturbation  $X = F + R$  of  $F$  is a normal form with respect to  $F$  if  $R$  is resonant.

It is important to note the following.

**Lemma 2.1.**  $\ker \square_\beta = \ker d^*|_{\mathcal{H}_\beta}$ .

**Proof.** Let  $\alpha = k - 1$ , then  $d : \mathcal{H}_\gamma \rightarrow \mathcal{H}_{\gamma+\alpha}$  and  $d^* : \mathcal{H}_{\gamma+\alpha} \rightarrow \mathcal{H}_\gamma$ . Due to the fact that  $d^*$  is the adjoint of  $d$ , we have the decomposition  $\mathcal{H}_\gamma = \text{Im } d^*|_{\mathcal{H}_{\gamma+\alpha}} \oplus \ker d|_{\mathcal{H}_\gamma}$ . Now let  $U \in \mathcal{H}_{\gamma+\alpha} = \mathcal{H}_\beta$ , then  $\square_\beta(U) = dd^*(U) = 0$  if and only if  $d^*U \in \ker d$ . Furthermore,  $d^*U \in \text{Im } d^*$ . That is  $d^*U \in \text{Im } d^* \cap \ker d$ . However  $\text{Im } d^*$  and  $\ker d$  are orthogonal. Then  $\square_\beta(U) = 0$  if and only if  $d^*U = 0$ .  $\square$

We now recall a result of [12] (Proposition 4.4); we only adapt it for the present context.

**Theorem 2.1** (Formal normal form [12]). Let  $X = F + P$  be a good perturbation of  $F$  as in Definition 2.1. Then there exists a formal diffeomorphism  $\hat{\Phi}$  such that  $\hat{\Phi}$  conjugates  $\hat{X}$  to a vector field  $F + R$ , where  $R$  is a resonant formal vector field in the sense of Definition 2.4.

Finally, we present our result: we prove that the resonant vector field  $R$  in Theorem 2.1 associated with  $F$  given by (6) is  $R = 0$ .

**Theorem 2.2.** Let  $X = F + P$  be a good perturbation of the vector field

$$F = \varepsilon \frac{\partial}{\partial x_1} + \sum_{i=2}^{k-1} 0 \frac{\partial}{\partial x_i} - \left( z^k + \sum_{j=1}^{k-1} x_j z^{j-1} \right) \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}. \tag{11}$$

Then, there exists a formal diffeomorphism  $\hat{\Phi}$  that conjugates  $\hat{X}$  with  $F$ , this is  $\hat{\Phi}_* \hat{X} = F$ .

**Proof.** From Theorem 2.1 and Lemma 2.1 we will show that if  $P \in \ker d^*|_{\mathcal{H}_{\geq k}}$  then  $P = 0$ . To simplify the notation, let  $\alpha \geq k$ ,  $P \in \mathcal{H}_\alpha$ ,  $\beta = \alpha - k + 1$ , and  $Q \in \mathcal{H}_\beta$ ; and let  $x = (x_1, \dots, x_{k-1}, z, \varepsilon) = (x_1, \dots, x_{k-1}, x_k, x_{k+1})$ . If  $D$  is an operator, its adjoint with respect to the inner product Definition 2.2 is always denoted as  $D^*$ . We start with the inner product (Definition 2.2)

$$\langle d(Q), P \rangle_{r,\alpha} = \langle Q, d^*(P) \rangle_{r,\beta}. \tag{12}$$

We can write  $d(Q) = \sum_{i=1}^{k+1} F(Q_i) - Q(F_i)$ , where  $F(Q_i) = \sum_{j=1}^{k+1} F_j \frac{\partial Q_i}{\partial x_j}$  and similarly for  $Q(F_i)$ , then

$$\begin{aligned} \langle d(Q), P \rangle_{r,\alpha} &= \sum_{i=1}^{k+1} \langle F(Q_i) - Q(F_i), P_i \rangle_{r,\beta} = \sum_{i=1}^{k+1} \langle F(Q_i), P_i \rangle_{r,\alpha+r_i} - \langle Q(F_i), P_i \rangle_{r,\alpha+r_i} \\ &= \sum_{i=1}^{k+1} \langle Q_i, F^*(P_i) \rangle_{r,\beta+r_i} - \langle Q(F_i), P_i \rangle_{\alpha+r_i} = \sum_{i=1}^{k+1} \langle Q_i, F^*(P_i) \rangle_{r,\beta+r_i} - \sum_{j=1}^{k+1} \langle Q_j, \left( \frac{\partial F_i}{\partial x_j} \right)^* (P_i) \rangle_{\beta+r_j} \\ &= \sum_{i=1}^{k+1} \langle Q_i, F^*(P_i) - \sum_{j=1}^{k+1} \left( \frac{\partial F_j}{\partial x_i} \right)^* (P_j) \rangle_{\beta+r_i} \end{aligned} \tag{13}$$

Comparing (13) to  $\langle Q, d^*(P) \rangle_{r,\beta}$  we can write

$$d^*(P) = \begin{bmatrix} F^* - \left( \frac{\partial F_1}{\partial x_1} \right)^* & - \left( \frac{\partial F_2}{\partial x_1} \right)^* & \dots & - \left( \frac{\partial F_{k+1}}{\partial x_1} \right)^* \\ - \left( \frac{\partial F_1}{\partial x_2} \right)^* & F^* - \left( \frac{\partial F_2}{\partial x_2} \right)^* & \dots & - \left( \frac{\partial F_{k+1}}{\partial x_2} \right)^* \\ \vdots & \vdots & \ddots & \vdots \\ - \left( \frac{\partial F_1}{\partial x_{k+1}} \right)^* & - \left( \frac{\partial F_2}{\partial x_{k+1}} \right)^* & \dots & F^* - \left( \frac{\partial F_{k+1}}{\partial x_{k+1}} \right)^* \end{bmatrix} \begin{bmatrix} P_1 \\ P_2 \\ \vdots \\ P_{k+1} \end{bmatrix}. \tag{14}$$

Plugging in the expressions of  $F$  and  $P$  into (14) we get

$$d^*(P) = \begin{bmatrix} F^* & 0 & \dots & 0 & 1 & 0 \\ 0 & F^* & \dots & 0 & z^* & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & F^* & (z^{k-1})^* & 0 \\ 0 & 0 & \dots & 0 & F^* + Z^* & 0 \\ -1 & 0 & \dots & 0 & 0 & F^* \end{bmatrix} \begin{bmatrix} P_1 \\ P_2 \\ \vdots \\ P_{k-1} \\ P_k \\ 0 \end{bmatrix} = 0, \tag{15}$$

where  $Z^* = \left( kz^{k-1} + \sum_{i=2}^{k-1} (i-1)x_i z^{i-2} \right)^*$ . Note that (15) implies  $P_1 = P_k = 0$  and  $F^*(P_j) = 0$  for all  $j = 2, \dots, k-1$ .

**Remark 2.1.** For  $k = 2$  the result is trivial: we have  $F = \varepsilon \frac{\partial}{\partial x_1} - (z^2 + x_1) \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}$ . Therefore  $d^*(P) = 0$  is written as

$$d^*(P) = \begin{bmatrix} F^* & 1 & 0 \\ 0 & F^* + 2z^* & 0 \\ -1 & 0 & F^* \end{bmatrix} \begin{bmatrix} P_1 \\ P_2 \\ 0 \end{bmatrix} = 0, \tag{16}$$

which immediately implies  $P_1 = P_2 = 0$ .

Now, we study  $F^*(P_j) = 0$ . Recall that  $P = P(x_1, \dots, x_{k-1}, z, \varepsilon)$  is not any vector field, but it has the property that  $P(x_1, \dots, x_{k-1}, z, 0) = 0$ . That is, we can write

$$P = \sum_{i=1}^{k-1} \varepsilon \bar{P}_i \frac{\partial}{\partial x_i} + \varepsilon \bar{P}_k \frac{\partial}{\partial z} + 0 \frac{\partial}{\partial \varepsilon}, \tag{17}$$

where  $\bar{P}_j \in \mathcal{P}_{\alpha+r_j-2k+1}$ . This is because the (quasihomogeneous) weight of  $\varepsilon$  is  $2k-1$ . Now, since it is complicated to work with the adjoint, we first rewrite the problem  $F^*(\varepsilon \bar{P}_j) = 0$ . We then prove that  $F^*(\varepsilon \bar{P}_j) = 0$  implies that  $\bar{P}_j = 0$ .

Note that  $F^*(\varepsilon \bar{P}_j) = 0$  is equivalent to  $\langle Q, F^*(\varepsilon \bar{P}_j) \rangle_{\beta+r_j} = 0$  for all  $Q \in \mathcal{P}_{\beta+r_j}$ . Next, we use the definition of  $F^*$  that is

$$\langle Q, F^*(\varepsilon \bar{P}_j) \rangle_{r,\beta+r_j} = \langle F(Q), \varepsilon \bar{P}_j \rangle_{r,\alpha+r_j} = 0. \tag{18}$$

We will now show that if  $\langle F(Q), \varepsilon \bar{P}_j \rangle_{r,\alpha+r_j} = 0$  for all  $Q \in \mathcal{P}_{\beta+r_j}$ , then  $\bar{P}_j = 0$ . Note that by (18), this is the same as proving that  $F^*(\varepsilon \bar{P}_j) = 0$  implies  $\bar{P}_j = 0$ . First, we choose an element  $x^q$  of the basis of  $\mathcal{P}_{\beta+r_j}$ ; this is

$$x^q = x_1^{q_1} \dots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon^{q_{k+1}}, \quad (r, q) = \beta + r_j, \tag{19}$$

then, it follows that

$$F(x^q) = q_1 x_1^{q_1-1} \dots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon^{q_{k+1}+1} - \left( z^k + \sum_{i=1}^{k-1} x_i z^{i-1} \right) q_k x_1^{q_1} \dots x_{k-1}^{q_{k-1}} z^{q_k-1} \varepsilon^{q_{k+1}}. \tag{20}$$

Let us write  $\varepsilon \bar{P}_j \in \mathcal{P}_{\alpha+r_j}$  as

$$\varepsilon \bar{P}_j = \varepsilon \sum_{(r,p)=\alpha+r_j-2k+1} a_p x_1^{p_1} \dots x_{k-1}^{p_{k-1}} z^{p_k} \varepsilon^{p_{k+1}}, \tag{21}$$

where  $a_p \in \mathbb{R}$ . We now proceed by recursion on the exponent of  $\varepsilon$ . Let  $q_{k+1} = 0$ , then the inner product  $\langle F(Q), \varepsilon \bar{P}_j \rangle_{\alpha+r_j}$  has only one term, since  $F(Q)$  has only one monomial containing  $\varepsilon$ . That is

$$\langle F(Q), \varepsilon \bar{P}_j \rangle_{\alpha+r_j} |_{q_{k+1}=0} = \langle q_1 x_1^{q_1-1} \dots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon, \varepsilon a_p x_1^{p_1} \dots x_{k-1}^{p_{k-1}} z^{p_k} \rangle_{r,\alpha+r_j} = 0. \tag{22}$$

We naturally consider  $q_1 > 0$ . If  $q_1 = 0$ , then the equality is automatically satisfied. Recalling Definition 2.2 of the inner product, the equality (22) means that

$$\langle q_1 x_1^{q_1-1} \dots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon, \varepsilon a_p x_1^{p_1} \dots x_{k-1}^{p_{k-1}} z^{p_k} \rangle_{r,\alpha+r_j} = q_1 a_p \frac{(q!)^r}{(\alpha+r_j)!} = 0, \tag{23}$$

and therefore from (22) we have

$$a_p = a_{q_1-1, p_2, \dots, p_k, 1} = 0, \tag{24}$$

for all  $q_1 > 0, p_2, \dots, p_k \geq 0$  (naturally, also satisfying the degree condition  $(r, p) = \alpha + r_j$ ). Next, let  $q_{k+1} = 1$ . Then

$$F(x^q) = q_1 x_1^{q_1-1} \dots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon^2 - \left( z^k + \sum_{i=1}^{k-1} x_i z^{i-1} \right) q_k x_1^{q_1} \dots x_{k-1}^{q_{k-1}} z^{q_k-1} \varepsilon. \tag{25}$$

Once again, the inner product  $\langle F(Q), \varepsilon \bar{P}_j \rangle_{r,\alpha+r_j}$  has only one term, now this is due to the fact that all coefficients  $a_p$  of monomials containing  $\varepsilon$  are zero due to (24). Then

$$\langle F(Q), \varepsilon \bar{P}_j \rangle_{\alpha+r_j} |_{q_{k+1}=1} = \langle q_1 x_1^{q_1-1} \dots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon^2, \varepsilon a_p x_1^{p_1} \dots x_{k-1}^{p_{k-1}} z^{p_k} \varepsilon \rangle_{r,\alpha+r_j} = 0. \tag{26}$$

Therefore, similarly as above, we have the condition

$$a_p = a_{q_1-1, p_2, \dots, p_k, 2} = 0, \tag{27}$$

for all  $q_1 > 0, p_2, \dots, p_k \geq 0$  (naturally, also satisfying the degree condition  $(r, p) = \alpha + r_j$ ). By recursion arguments, assume  $q_{k+1} = n$  and that all the coefficients

$$a_p = a_{p_1, p_2, \dots, p_k, m} = 0, \quad \forall m \leq n. \tag{28}$$

Then, again, the inner product  $\langle F(Q), \varepsilon \bar{P}_j \rangle_{r,\alpha+r_j}$  has only one term, namely

$$\langle F(Q), \varepsilon \bar{P}_j \rangle_{\alpha+r_j} |_{q_{k+1}=n} = \langle q_1 x_1^{q_1-1} \dots x_{k-1}^{q_{k-1}} z^{q_k} \varepsilon^{n+1}, \varepsilon a_p x_1^{p_1} \dots x_{k-1}^{p_{k-1}} z^{p_k} \varepsilon^n \rangle_{r,\alpha+r_j} = 0. \tag{29}$$

The latter implies:

$$a_p = a_{q_1-1, p_2, \dots, p_k, n+1} = 0. \tag{30}$$

This finishes the proof of  $\langle F(Q), \varepsilon \bar{P}_j \rangle_{r,\alpha+r_j} = 0$ , which implies  $\bar{P}_j = 0$  for all  $j = 2, \dots, k-1$ .  $\square$

**Remark 2.2.** Theorem 2.2, together with Borel's lemma [3], implies that an  $A_k$ -SFS  $X = F + P$  is smoothly conjugate to a smooth vector field  $Y = F + H$  where  $H$  is flat at the origin. The benefits of this normal form are exploited in [6,7].

**Acknowledgements**

The author gratefully acknowledges Henk W. Broer, Robert Roussarie, and Laurent Stolovitch for fruitful discussions and valuable comments and suggestions. This work is partially supported by the CONACYT postgraduate grant 214759-310040.

## References

- [1] V.I. Arnold, S.M. Gusein-Zade, A.N. Varchenko, *Singularities of Differentiable Maps*, vol. I, Birkhäuser, Basel, Switzerland, 1985.
- [2] H.W. Broer, T.J. Kaper, M. Krupa, Geometric desingularization of a cusp singularity in slow-fast systems with applications to Zeeman's examples, *J. Dyn. Differ. Equ.* 25 (4) (2013) 925–958.
- [3] T. Bröcker, *Differentiable Germs and Catastrophes*, Lecture Note Series, vol. 17, Cambridge University Press, Cambridge, UK, 1975.
- [4] F. Dumortier, R. Roussarie, *Canard Cycles and Center Manifolds*, vol. 121, American Mathematical Society, 1996.
- [5] N. Fenichel, Geometric singular perturbation theory, *J. Differ. Equ.* 31 (1) (1979) 53–98.
- [6] H. Jardón Kojakhmetov, *Classification of constrained differential equations embedded in the theory of slow fast systems:  $A_k$  singularities and geometric desingularization*, PhD thesis, Rijksuniversiteit Groningen, 2015, 164 p.
- [7] H. Jardón-Kojakhmetov, Henk W. Broer, R. Roussarie, *Analysis of a slow-fast system near a cusp singularity*, submitted to *J. Differ. Equ.* (2015).
- [8] Ilona Kosiuk, Peter Szmolyan, Scaling in singular perturbation problems: blowing up a relaxation oscillator, *SIAM J. Appl. Dyn. Syst.* 10 (4) (2011) 1307–1343.
- [9] M. Krupa, P. Szmolyan, Extending geometric singular perturbation theory to non-hyperbolic points: fold and canard points in two dimensions, *SIAM J. Math. Anal.* 33 (2001) 286–314.
- [10] M. Krupa, P. Szmolyan, Relaxation oscillation and canard explosion, *J. Differ. Equ.* 174 (2001) 312–368.
- [11] M. Krupa, M. Wechselberger, Local analysis near a folded saddle-node singularity, *J. Differ. Equ.* 248 (12) (2010) 2841–2888.
- [12] É. Lombardi, L. Stolovitch, Normal forms of analytic perturbations of quasihomogeneous vector fields: rigidity, invariant analytic sets and exponentially small approximation, *Ann. Sci. Éc. Norm. Supér.* (4) 43 (2010) 659–718.
- [13] S. Rinaldi, M. Scheffer, Geometric analysis of ecological models with slow and fast processes, *Ecosystems* 3 (6) (2000) 507–521.
- [14] M.R. Rose, R. Harmsen, Ecological outbreak dynamics and the cusp catastrophe, *Acta Biotheor.* 30 (4) (1981) 229–253.
- [15] P. Szmolyan, M. Wechselberger, Canards in  $\mathbb{R}^3$ , *J. Differ. Equ.* 177 (2) (2001) 419–453.
- [16] S. van Gils, M. Krupa, P. Szmolyan, Asymptotic expansions using blow-up, *Z. Angew. Math. Phys.* 56 (8) (2005) 369–397.
- [17] E.C. Zeeman, *Differential equations for the heart beat and nerve impulse*, in: *Towards a Theoretical Biology*, vol. 4, Edinburgh University Press, Edinburgh, UK, 1972, pp. 8–67.