

University of Groningen

The relationship between gait dynamics and future cognitive decline

Kikkert, Lisette H J; Vuillerme, Nicolas; van Campen, Jos P; Appels, Bregje A; Hortobágyi, Tibor; Lamoth, Claudine J C

Published in:
International Psychogeriatrics

DOI:
[10.1017/S1041610217002770](https://doi.org/10.1017/S1041610217002770)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kikkert, L. H. J., Vuillerme, N., van Campen, J. P., Appels, B. A., Hortobágyi, T., & Lamoth, C. J. C. (2018). The relationship between gait dynamics and future cognitive decline: A prospective pilot study in geriatric patients. *International Psychogeriatrics*, 30(9), 1301-1309. <https://doi.org/10.1017/S1041610217002770>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The relationship between gait dynamics and future cognitive decline: a prospective pilot study in geriatric patients

Lisette H. J. Kikkert,^{1,2,3} Nicolas Vuillerme,^{2,4} Jos P. van Campen,³ Bregje A. Appels,⁵ Tibor Hortobágyi¹ and Claudine J. C. Lamoth¹

¹University of Groningen, University Medical Centre Groningen, Center for Human Movement Sciences, Groningen, The Netherlands

²Université Grenoble Alpes, EA AGEIS, Grenoble, France

³Department of Geriatric Medicine, MC Slotervaart Hospital, Amsterdam, The Netherlands

⁴Institut Universitaire de France, Paris, France

⁵Department of Medical Psychology and Hospital Psychiatry, MC Slotervaart Hospital, Amsterdam, The Netherlands

ABSTRACT

Background: Walking ability recently emerged as a sub-clinical marker of cognitive decline. Hence, the relationship between baseline gait and future cognitive decline was examined in geriatric patients. Because a “loss of complexity” (LOC) is a key phenomenon of the aging process that exhibits in multiple systems, we propose the idea that age- and cognition-related LOC may also become manifested in gait function. The LOC theory suggests that even healthy aging is associated with a (neuro)physiological breakdown of system elements that causes a decline in variability and an overall LOC. We used coordination dynamics as a conceptual framework and hypothesized that a LOC is reflected in dynamic gait outcomes (e.g. gait regularity, complexity, stability) and that such outcomes could increase the specificity of the gait-cognition link.

Methods: 19 geriatric patients (age 80.0 ± 5.8) were followed for 14.4 ± 6.6 months. An iPod collected three-dimensional (3D) trunk accelerations while patients walked for 3 minutes. Cognition was evaluated with the Mini-Mental State Examination (MMSE) and the Seven-Minute screen (7MS) test. The Reliable Change Index (RCI) quantified the magnitude of cognitive change. Spearman’s Rho coefficients (ρ) indexed correlations between baseline gait and future cognitive change.

Results: Seven patients showed reliable cognitive decline (“Cognitive Decline” group), and 12 patients remained cognitively stable (“Cognitive Stable” group) over time. Future cognitive decline was correlated with a more regular ($\rho = 0.579^*$) and predictable ($\rho = 0.486^*$) gait pattern, but not with gait speed.

Conclusions: The increase in gait regularity and predictability possibly reflects a LOC due to age- and cognition-related (neuro)physiological decline. Because dynamic versus traditional gait outcomes (i.e. gait speed and (variability of) stride time) were more strongly correlated with future cognitive decline, the use of wearable sensors in predicting and monitoring cognitive and physical health in vulnerable geriatric patients can be considered promising. However, our results are preliminary and do require replication in larger cohorts.

Key words: geriatric patients, frailty, cognitive impairment, gait analysis, non-linear dynamics, prediction, loss of complexity

Introduction

Medical developments have substantially extended human lifespan. An increase in age, however, comes hand-in-hand with co-morbidities, such as cognitive decline, muscle weakness, frailty, polypharmacy, and falling (Inouye *et al.*, 2007).

Hence, we can anticipate that an increase in the number of “older old adults” will need specialized geriatric care (i.e. geriatric patients) to slow functional decline. Cognitive impairment is a frequent geriatric condition substantially affecting independence, mobility, and quality of life. If applied in time, tailored interventions could delay disease onset and perhaps extend the asymptomatic phase. Predicting future cognitive loss is therefore important in this vulnerable population, in which walking ability has recently emerged as a non-invasive sub-clinical marker that predicts cognitive decline (Kikkert *et al.*, 2016).

Correspondence should be addressed to: Lisette H. J. Kikkert, University of Groningen, University Medical Centre Groningen, Center for Human Movement Sciences, Ant Deusinglaan 1, 9713 AV Groningen, The Netherlands. Phone: +31(0)50 363 2710. Email: l.h.j.kikkert@umcg.nl. Received 29 Jun 2017; revision requested 12 Sep 2017; revised version received 17 Oct 2017; accepted 7 Nov 2017.

The view that walking is no longer considered an automatic task supports its potential to serve as an early marker of cognitive decline. Because brain areas affected by cognitive impairment partly overlap with brain areas activated during walking (Holtzer *et al.*, 2014), subtle, pre-clinical changes in gait could be precursors of evolving cognitive impairment. Such an overlap gave rise to the concepts of the “Motoric Cognitive Risk” syndrome (Verghese *et al.*, 2013), a “Gait Phenotype” of cognitive decline (Allali *et al.*, 2016), and the “Motor Signature” of cognitive decline (Montero-Odasso *et al.*, 2014). Longitudinal studies confirmed the close relation between gait and cognition (Beauchet *et al.*, 2016; Kikkert *et al.*, 2016), with the majority of prediction studies focussing on gait speed as the main predictor of cognitive decline (Kikkert *et al.*, 2016). Individuals who developed Mild Cognitive Impairment (MCI) or dementia presented with a slower baseline gait speed (0.91 and 0.80 m/s, respectively) as compared to individuals who remained cognitively intact (1.11 m/s) (Kikkert *et al.*, 2016). While a slow walking speed provides an important marker of multiple adverse outcomes (e.g. falling, hospitalization, and even survival) (Studenski *et al.*, 2011), more delicate gait measures could unravel why geriatric patients walk slow, with final gait speed being the cumulative result of interactions between multiple, subtle gait functions. Hence, we hypothesize that those gait details could increase the specificity of the gait–cognition link. Key principles of the aging neuro-musculo-skeletal system (NMSS) provide a theoretical framework for the latter hypothesis.

Although age- and pathology-related declines in neural, sensory motor, cognitive, and muscular function, i.e. declines in NMSS, are generally examined separately, such a view limits our understanding of the aging NMSS as a whole (Sleimen-Malkoun *et al.*, 2014). Because vulnerable geriatric patients show degradation in multiple interacting systems, we propose the idea to place the gait–cognition link into a more encompassing perspective to better understand the coupling and coordination between elements of the NMSS (i.e. gait and cognition). To this idea, we consider a key phenomenon of the aging NMSS, namely the “loss of complexity” (LOC). The LOC theory is derived from the field of non-linear dynamics, and suggests that even healthy aging is associated with a (neuro)physiological breakdown of system elements that causes a loss of overall complexity (Lipsitz and Goldberger, 1992). This LOC, in turn, leads to a reduced adaptive capacity and most likely to poor functional outcomes, such as frailty and an increased fall risk (Lipsitz, 2002). Indeed, frail versus healthy old adults and

fallers versus non-fallers are characterized by a global LOC (Lipsitz, 2002), and a LOC indicates transitions from healthy to frail aging (Lang *et al.*, 2009). (Neuro)physiological decline and cognitive impairment may add to this LOC, and possibly becomes manifested in gait function. A loss of gait complexity is characterized by an increased gait regularity/predictability, in which perfect regular signals would resemble a sine function. Coordination dynamics constitute a conceptual framework to quantify complexity caused by physiological breakdown because it can capture changes of functional status over time by means of, for example, indexing coupling and self-organization properties (Sleimen-Malkoun *et al.*, 2014).

Trunk accelerations have demonstrated their ability to quantify non-linear coordination dynamics of gait function, including self-affinity, regularity, complexity, variability, and stability (Lamoth *et al.*, 2002; Riva *et al.*, 2013; Kobsar *et al.*, 2014; Rispens *et al.*, 2014). Such gait dynamics describe overall gait coordination and the ability to overcome or adequately respond to perturbations. Cross-sectionally, gait dynamics distinguished old adults with and without cognitive impairment, with cognitively impaired old adults presenting with a less variable and less-stable gait pattern (Lamoth *et al.*, 2011). Longitudinally, adding gait dynamics to clinical tests increased the accuracy of a fall prediction model by 14% (van Schooten *et al.*, 2015), and the specificity of a fall classification model from 60% to 80% (Kikkert *et al.*, 2017). However, less is known about the longitudinal link between gait dynamics and cognitive decline. In addition to more traditional outcomes, such as gait speed, it is quite possible that an age- and pathology-related LOC is expressed in detailed dynamic gait outcomes derived from non-linear analyses. Gait dynamics could therefore assist to index and predict long-term cognitive change and provide insights into (unconscious) strategies old adults use to compensate for this anticipated cognitive loss. Hence, the aim of the present prospective pilot study was to examine the relationship between baseline gait function and future cognitive decline, and to identify indicators of future cognitive decline in terms of a geriatric patient’s gait. We hypothesize that a LOC of the aging NMSS translates to gait function, and is reflected in measures that quantify dynamic aspects of gait.

Methods

Patients

In total, 70 geriatric patients were recruited from the geriatric diagnostic day clinic of the MC

Slotervaart hospital in Amsterdam between January 2015 and July 2016 (mean age 80 ± 6.6 ; 53% women). Patients were referred by a general practitioner based on general or specific decline, and underwent extensive screening for physical and cognitive functioning during their seven-hour visit. Inclusion criteria were as follows: age 65 years or older. Exclusion criteria were as follows: (1) inability to walk for three minutes without a walking aid, (2) the presence of neurological disorders other than dementia related (e.g. Parkinson's disease), (3) having neurological or orthopedic disabilities that limit mobility function (e.g. recent surgery), and (4) inability to speak and understand Dutch. Only patients at risk for (further) cognitive deterioration were invited for a re-evaluation in approximately one year. However, this referral depended on subjective and objective evaluations of the clinical geriatrician who treated the patient. Hence, 19 patients were included in the present pilot study. While all patients were at risk for cognitive impairment, 10 out of 19 patients actually had a diagnosis of MCI at baseline. The protocol was approved by the Medical Ethical Committee of the MC Slotervaart hospital. Because some of the test results could directly be used by the involved clinical geriatrician (e.g. gait speed, hand grip strength, and frailty), the tests were part of a standard clinical evaluation when a researcher was present who could administer the measurements. It was therefore not necessary to obtain informed consent from patients.

Demographic information, including age, height, weight, and Body Mass Index (BMI), were extracted from medical records. The Charlson Comorbidity Index (CCI) quantified the number and severity of comorbidities, and polypharmacy was denoted when patients used >4 medications. Cognitive performance was evaluated with the Mini-Mental State Examination (MMSE) and with the seven-minute screen (7MS) test. The 7MS assessed memory function using the Benton's Temporal Orientation (BTO) (range 0–113) and the Enhanced Cued Recall (ECR) test (range 0–16), and executive and visuospatial function using the animal verbal fluency (range 0–45) and clock-drawing test (range 0–14). A logistic regression formula based on the four sub-tests resulted in a total 7MS-score, with a score of 0 corresponding to a 50% chance that a patient has dementia, and negative and positive scores corresponding to a lower and higher than 50% chance to have dementia, respectively. We refer to our previous study for references of the cognitive screening batteries (Kikkert *et al.*, 2017). The Reliable Change Index (RCI) quantified the magnitude of cognitive change and to determine whether the changes in cognition were clinically meaningful and




Figure 1. (Colour online) Syntax of the study protocol. Abbreviations used in Figure 1: BMI = Body Mass Index; CCI = Charlson Comorbidity Index; MMSE = Mini-Mental State Examination; 7MS = Seven-Minute Screen.

statistically reliable, with a RCI of >1.96 indicating a significant change between baseline and follow-up outcomes ($p < 0.05$) (Jacobson and Truax, 1991). Patients who showed a reliable decline in both, MMSE and 7MS-score (average RCI score >1.96), were coded as the “Cognitive Decline” group. Patients who remained cognitively stable or showed non-significant changes in cognition formed the “Cognitive Stable” group. Figure 1 illustrates the study protocol.

Procedures and pre-processing

At baseline, patients walked for three minutes at habitual gait speed on a 10-m long course marked with cones. Accelerations of the lower trunk (near the level of the third lumbar vertebra) were registered using a built-in accelerometer of an iPod touch G4 (iOS 6, Apple Inc.; sample frequency ± 100 Hz). The validity of gait and standing posture parameters from trunk accelerations as indicated by intra-class correlation (ICC) was high (ICC = 0.85–0.99), and test–retest reliability was good (ICC = 0.81–0.97) in old adults, under varying conditions (Kosse *et al.*, 2015). A custom-made application “iMoveDetection” stored the acceleration signals in anterior-posterior (AP), medio-lateral (ML), and vertical (V) directions (Kosse *et al.*, 2015). The signals were then

Table 1. Descriptions and formula of the quantified gait outcomes

	GAIT OUTCOME	FORMULA	DESCRIPTION
(1)	Gait speed (m/s)	$\text{Speed} = \frac{\text{Distance (m)}}{\text{Time walked (s)}}$	Average walking speed.
(2)	Stride time (s)	$\text{Stride time} = \frac{\text{Time walked (s)}}{\text{Number of strides}}$	Stride time reflects the average time from heel strike-to-heel strike of the ipsilateral foot.
(3)	CoV stride time (%)	$\text{CoV} = \frac{\text{Standard deviation}}{\text{Mean}} \times 100\%$	The coefficient of variation of stride time is the standardized measure of dispersion of the average stride time.
(4)	Root mean square	$\text{RMS} = \sqrt{\frac{x^1 + x^2 + \dots + x^n}{n}}$	The RMS quantifies the magnitude of amplitude variability.
(5)	Step and stride regularity	$\text{AC}(t) = \frac{1}{N- t } \sum_{i=1}^{N- t } x_i x_{i+t}$	Step and stride regularity indicate the regularity of steps and strides, respectively. Autocorrelation refers to the correlation of a time series with its own past and future values. Perfect regular steps or strides will adopt a value of 1 (Kobsar <i>et al.</i> , 2014).
(6)	Multi-scale sample entropy	$\text{MSE}(N, m, \tau, \gamma) = -\ln\left[\frac{A^{m+1}(\gamma)}{A^m(\gamma)}\right]$	Mscale-En reflects the degree of predictability of a gait pattern. A complete predictable signal has an Mscale-En value of 0 (Costa <i>et al.</i> , 2003).
(7)	Maximal Lyapunov exponent	$\lambda_i = \lim_{n \rightarrow \infty} \frac{1}{t} \log_2 \frac{p_i(t)}{p_i(0)}$	Maximal Lyapunov exponent indicates local dynamic stability, i.e. the ability to resist perturbations. A larger λ_{\max} reflects a less local dynamic stability (Cignetti <i>et al.</i> , 2012).

transferred via blue tooth to an online platform and analyzed with custom MATLAB software (version 2014b, The MathWorks Inc.). All times series were de-trended, corrected for horizontal tilt, and low-pass filtered using a fourth-order Butterworth filter (cut-off frequency of 10 Hz).

Gait outcomes

We previously identified gait outcomes that characterized the same population of geriatric patients with and without cognitive impairment in a cross-sectional study (submitted), and therefore included those outcomes in the present longitudinal analysis. In addition, average stride time and the Coefficient of Variation (CoV) of stride time were computed. Time indices of foot contacts were manually identified from peaks in the AP acceleration signals to calculate stride-related gait outcomes. The magnitude of the resultant vector of the AP, ML, and V signals was calculated according to the following equation: $R = \sqrt{\text{AP}^2 + \text{ML}^2 + \text{V}^2}$. The dynamic gait outcomes were quantified over

this “summary” signal. Table 1 describes stride-related (outcomes 1–3) and dynamic (outcomes 4–7) gait outcomes.

Statistical analysis

Statistical analyses were performed using SPSS version 24. Group differences (“Cognitive Decline” vs. “Cognitive Stable”) were examined with the Mann–Whitney’s U test, and within-group differences (baseline vs. follow-up) with a repeated measure analysis of variance (ANOVA). Correlations between baseline gait and longitudinal cognitive change in MMSE and 7MS scores were computed using Spearman’s Rho correlation coefficients (ρ) for non-linear data. Perfect monotonic relations would reveal a ρ of 1 or -1 . Significance was set at 0.05.

Results

In total, 19 patients were followed for 14.4 ± 6.6 months (age 80.0 ± 5.8 ; 63% women). Based on

Table 2. Patient demographics (mean±SD) by final cognitive state. Differences were evaluated with the Mann–Whitney’s U test, with significant differences between groups indicated in bold

	BASELINE		p	FOLLOW-UP		p
	COGNITIVE DECLINE (n = 7)	COGNITIVE STABLE (n = 12)		COGNITIVE DECLINE (n = 7)	COGNITIVE STABLE (n = 12)	
Age (years)	80.9 ± 6.4	79.5 ± 5.6	0.37	NA	NA	
Height (cm)	166.0 ± 7.3	164.6 ± 9.2	0.53	NA	NA	
Weight (kg)	65.3 ± 10.7	69.7 ± 14.4	0.55	NA	NA	
BMI	23.6 ± 2.3	25.9 ± 5.6	0.40	23.8 ± 2.3	25.9 ± 5.9	0.35
CCI	3.3 ± 1.5	2.2 ± 1.3	0.13	3.3 ± 1.5	2.5 ± 1.7	0.24
Medications	6.1 ± 3.4	6.2 ± 3.3	0.93	5.0 ± 4.2	6.1 ± 4.1	0.40
MMSE	25.7 ± 3.7	26.3 ± 2.8	0.83	22.3 ± 6.2	26.4 ± 3.4	0.05*
7MS	8.3 ± 16.7	0.0 ± 4.6	1.00	16.1 ± 23.9	0.3 ± 4.3	0.01*

*p < 0.05.

BMI = Body Mass Index; CCI = Charlson Comorbidity Index; MMSE = Mini-Mental State Examination; 7MS = Seven-Minute Screen; NA = Not Applicable.

average RCI scores of MMSE and 7MS, 7 of 19 patients (37%) showed significant cognitive decline over time and were retrospectively classified as the “Cognitive Decline” group (average follow-up time was 11.3±2.0 months). The other 12 patients remained cognitively stable or slightly increased in cognitive function and formed the “Cognitive Stable” group (average follow-up time was 16.2±1.7 months). At baseline, the groups were comparable in terms of age, height, weight, BMI, CCI, the number of medication used, MMSE-score, and 7MS-score (all p > 0.05). At follow-up, the groups only differed in MMSE (p = 0.05) and 7MS (p = 0.01). The MMSE (p = 0.05) and the 7MSE scores (p = 0.01) decreased in the “Cognitive Decline” group but not in the “Cognitive Stable” group (MMSE: p = 1.0; 7MS: p = 0.83). Other outcomes did not significantly change (p > 0.05) in either group (Table 2).

Relationship between baseline gait and future cognitive change

Table 3 presents correlations between baseline gait outcomes and changes in cognitive function 14 months later. Stride-related outcomes (gait speed, stride time, CoV of stride time) did not correlate with changes in cognition ($\rho < 0.3$). Trunk outcomes showed modest correlations with changes in cognition, with a decline in cognition corresponding to more regular steps ($\rho = 0.579^*$) and strides ($\rho = 0.347$), and a more predictable gait ($\rho = -0.484^*$). Figure 2 shows the correlations between gait outcomes (step regularity, stride regularity, and multi-scale sample entropy) and changes in cognition in each group. Correlations between baseline gait and cognition in MMSE and

Table 3. Spearman’s Rho (ρ) correlations between baseline gait and future change in cognition in 19 geriatric patients. The degree of decline in cognition is expressed using the Reliable Change Index (RCI), combining MMSE and 7MS-scores

	CHANGE IN COGNITION (AVERAGE OF MMSE AND 7MS)
Stride-related outcomes	
Gait speed	0.073
Stride time	0.051
Coefficient of variation of stride time	-0.260
Trunk outcomes	
Root mean square	-0.065
Step regularity	0.579*
Stride regularity	0.347
Multi-scale sample entropy	-0.484*
Maximal Lyapunov exponent	0.196

*p < 0.05.

7MS separately were in the same direction, in which some gait outcomes related more to change in MMSE, while others were more closely linked to change in 7MS. For example, stride regularity correlated more strongly with Δ MMSE ($\rho = 0.339$) than with Δ 7MS ($\rho = 0.230$), while multi-scale sample entropy correlated more strongly with 7MS ($\rho = -0.509^*$) as compared to MMSE ($\rho = -0.321$).

Discussion

The present prospective pilot study examined whether baseline gait characteristics predicted




Figure 2. Correlations between baseline gait outcomes and future decline in cognition (MMSE+7MS). The degree of cognitive change was quantified with the Reliable Change Index (RCI), with an RCI > 1.96 indicating significant cognitive change over time ($p < 0.05$). Each symbol denotes an individual patient. Patients whose MMSE and 7MS decreased are colored in orange ($n = 7$) and those patients who remained cognitively stable are colored in blue ($n = 12$).

* $p < 0.05$.

changes in cognition 14.4 ± 6.6 months later in 19 geriatric patients (mean age 80.0 ± 5.8). The results revealed that a more regular and predictable gait pattern correlated with future cognitive decline in geriatric patients admitted to an outpatient diagnostic clinic. We discuss these results from a theoretical perspective in terms of the “LOC hypothesis,” and from a clinical perspective in terms of how smart devices that extract gait details could possibly facilitate the prediction of cognitive decline and the development of early tailored interventions.

While population-based studies reported that gait speed predicts cognitive decline (Kikkert

et al., 2016; Savica *et al.*, 2017), the present analyses did not confirm these results. Patient characteristics may account for the discrepant data. While previous studies focused on relatively young and healthy old adults with a mean age around 65, our geriatric patients had a mean age of 80. Geriatric syndromes that are also linked to gait slowing (e.g. muscle weakness, polypharmacy, falling, osteoporosis-related factors) (de Groot *et al.*, 2014; 2016) may have caused a slower gait, also in the group that remained cognitively stable. Gait speed alone might thus not be specific enough to predict future cognitive decline in geriatric patients that suffer from multi-system degeneration

(de Groot *et al.*, 2016). These results are in agreement with a recent population study that concluded that a slow baseline gait speed was only modestly related to future cognitive decline, and provided no early marker of clinical progression to dementia (Hooghiemstra *et al.*, 2017). The short follow-up period in the present study could also account for the discrepancies relative to previous studies, as our average follow-up time versus the average follow-up time of a systematic review were 1.2 and 4.5 years, respectively.

Cognitive decline over time correlated with a more regular (higher autocorrelation coefficients) and more predictable (lower multi-scale sample entropy) baseline gait pattern. Because increases in stride regularity and predictability during gait reflect a decline in gait complexity (Costa *et al.*, 2003), we confirm our hypothesis that the “LOC hypothesis” (Lipsitz and Goldberger, 1992) could provide a theoretical framework to relate those findings to health status. The LOC theory suggests that a deterioration in age-related and pathological physiological functioning leads to a breakdown of system elements, causing a decline in variability and overall complexity (Lipsitz and Goldberger, 1992). This LOC, in turn, relates to a reduced adaptive capacity and most likely relate to poor functional outcomes, such as an increased fall risk (Lipsitz, 2002). Our results are in agreement with postural control studies that reported a reduced complexity of postural fluctuations (as quantified by the multi-scale sample entropy) in older adults with sensory impairments as compared to age-matched controls (Manor *et al.*, 2010). A sample of frail and pre-frail older adults (who closely compare to the present sample of geriatric patients) exhibited less postural complexity as compared to non-frail controls (Kang *et al.*, 2009). Physiological decline related to cognitive loss possibly adds to this LOC and the association between cognitive decline and higher gait regularity and predictability might thus reflect a LOC of the aging system. Although increased regularity and predictability may seem beneficial, such a strategy reduces the ability to resist and recover from perturbations and may actually promote gait instability and increase the risk of falling (Sleimen-Malkoun *et al.*, 2014). The latter is in agreement with studies that reported a decline in gait stability in old adults with dementia as compared to age-matched controls (Lamoth *et al.*, 2011). However, given the limited sample size and explorative nature of the present study, the results should be taken with caution and need replication in larger cohorts.

The finding that trunk outcomes versus more traditional gait outcomes (i.e. gait speed) were more closely correlated to future cognitive decline

provides support for the use of wearable sensors in clinics. Indeed, the compact size, relatively low cost, and ease of operation facilitate the incorporation of sensor technologies in gait analysis. Based on the present preliminary results, it seems that the use of dynamical systems in gait analyses could make prediction of cognitive decline more accurate as compared to the use of traditional gait outcomes, such as gait speed. Except for its use in the prediction of cognitive decline, monitoring gait dynamics is encouraged because such details also relate to one’s ability to act independently and autonomously. Intervention strategies could, in turn, specifically tailor gait functions in an aim to remain activities of daily living and to reduce fall risk. Although studies that examined the effects of exercise on gait and cognition showed contradictory results, a recent population-based study highlighted that transitions in gait as well as in cognitive function were mutable and reversible over a nine-year period, even in the oldest old (Qualls *et al.*, 2017). However, despite technological and clinical advantages of incorporating gait analysis derived from wearable sensors, future studies should confirm the clinical utility and the predictive ability of such technologies, and applications should be built to translate gait details to clinical outcomes.

Procedures at the MC Slotervaart hospital are highly patient oriented. While this provided us with an extensive characterization of this vulnerable population in terms of demographic, physical, and social information, standard procedures at the hospital also placed some difficulties. For example, even though the protocol for a follow-up appointment is set to be at one year, the actual follow-up period always depends on organizational and patient factors. This resulted in a large variation in follow-up time between patients (follow-up period is 14.4 ± 6.6 months), which can be considered a limitation of the present study. For the same reasons, we were unable to measure gait function at follow-up. Because there is a clear theoretical and experimental basis for the relationship between gait and cognitive impairment (Beauchet *et al.*, 2016; Kikkert *et al.*, 2016), we expected that gait would have been changed in the patient group who presented with significant cognitive decline over time, and that correlations between gait and cognition would have become stronger. A recent study highlighted the sensitivity of changes in gait speed over a follow-up period of one year in healthy old adults aged 75. This study showed that 25% of healthy old adults showed a gait speed decline of more than 0.1 m/s per year (Brach *et al.*, 2011). We therefore expected that those changes in mobility may also be reflected in elements of mobility that underlie gait speed, i.e.

gait dynamics. Finally, the small sample size places a limitation of the present study.

In conclusion, the present pilot study revealed that a more regular and predictable gait pattern was correlated with future cognitive decline in geriatric patients admitted to an outpatient diagnostic clinic. Those results could reflect a LOC of the aging NMSS. In addition to traditional outcomes, such as gait speed, trunk outcomes derived from wearable sensors are promising indicators of cognitive as well as physical decline. Hence, we recommend the incorporation of a non-invasive detailed gait analysis in predicting, diagnosing, and monitoring health status in vulnerable geriatric patients. However, our results and interpretations are preliminary and need replication in larger cohorts, as for now, our conclusions are based on a small sample size and a relatively short follow-up period.

Conflict of interest

None.

Description of authors' roles

CJCL, TH, NV, and LHJK created the concept and design of the study. LHJK collected the data. BAA and JPVC were responsible for clinical aspects, such as patient inclusion, diagnoses, and ethical issues. LHJK analyzed the data, and the results were interpreted by CJCL, TH, and LHJK. LHJK wrote a draft of the full manuscript. CJCL, TH, NV, BAA, and JPVC critically revised the manuscript and approved the final version.

Acknowledgments

This work was supported by the French National Research Agency in the framework of the "Investissements d'avenir" program (ANR-10-AIRT-05 and ANR-15-IDEX-02).

References

Allali, G. *et al.* (2016). Gait phenotype from mild cognitive impairment to moderate dementia: results from the GOOD initiative. *European Journal of Neurology*, 23, 527–541. doi: [10.1111/ene.12882](https://doi.org/10.1111/ene.12882).

Beauchet, O. *et al.* (2016). Poor gait performance and prediction of dementia: results from a meta-analysis. *Journal of the American Medical Directors Association*, 17, 482–490. doi: [10.1016/j.jamda.2015.12.092](https://doi.org/10.1016/j.jamda.2015.12.092).

Brach, J. S., Perera, S., Van Swearingen, J. M., Hile, E. S., Wert, D. M. and Studenski, S. A. (2011). Challenging gait conditions predict 1-year decline in gait speed in older adults with apparently normal gait. *Physical Therapy*, 91, 1857–1864. doi: [10.2522/ptj.20100387](https://doi.org/10.2522/ptj.20100387).

Cignetti, F., Decker, L. M. and Stergiou, N. (2012). Sensitivity of the Wolf's and Rosenstein's algorithms to evaluate local dynamic stability from small gait data sets. *Annals of Biomedical Engineering*, 40, 1122–1130. doi: [10.1007/s10439-011-0474-3](https://doi.org/10.1007/s10439-011-0474-3).

Costa, M., Peng, C. -L., Goldberger, A. and Hausdorff, J. M. (2003). Multiscale entropy analysis of human gait dynamics. *Physica A: Statistical Mechanics and its Applications*, 330, 53–60. doi: <https://doi.org/10.1016/j.physa.2003.08.022>.

de Groot, M. H., van Campen, J. P., Kosse, N. M., de Vries, O. J., Beijnen, J. H. and Lamoth, C. J. (2016). The association of medication-use and frailty-related factors with gait performance in older patients. *PLoS One*, 11, e0149888. doi: [10.1371/journal.pone.0149888](https://doi.org/10.1371/journal.pone.0149888).

de Groot, M. H., van der Jagt-Willems, H. C., van Campen, J. P., Lems, W. F., Beijnen, J. H. and Lamoth, C. J. (2014). A flexed posture in elderly patients is associated with impairments in postural control during walking. *Gait & Posture*, 39, 767–772. doi: [10.1016/j.gaitpost.2013.10.015](https://doi.org/10.1016/j.gaitpost.2013.10.015).

Holtzer, R., Epstein, N., Mahoney, J. R., Izzetoglu, M. and Blumen, H. M. (2014). Neuroimaging of mobility in aging: a targeted review. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 69, 1375–1388. doi: [10.1093/gerona/glu052](https://doi.org/10.1093/gerona/glu052).

Hooghiemstra, A. M. *et al.* (2017). Gait speed and grip strength reflect cognitive impairment and are modestly related to incident cognitive decline in memory clinic patients with subjective cognitive decline and mild cognitive impairment: findings from the 4C study. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 72, 846–854. doi: [10.1093/gerona/glx003](https://doi.org/10.1093/gerona/glx003).

Inouye, S. K., Studenski, S., Tinetti, M. E. and Kuchel, G. A. (2007). Geriatric syndromes: clinical, research, and policy implications of a core geriatric concept. *Journal of the American Geriatrics Society*, 55, 780–791. doi: [JGS1156 \[pii\]](https://doi.org/10.1111/j.1532-5415.2007.01111.x).

Jacobson, N. S. and Truax, P. (1991). Clinical significance: a statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59, 12–19.

Kang, H. G. *et al.* (2009). Frailty and the degradation of complex balance dynamics during a dual-task protocol. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 64, 1304–1311. doi: [10.1093/gerona/64.11.1304](https://doi.org/10.1093/gerona/64.11.1304).

Kikkert, L. H. J. *et al.* (2017). Gait dynamics to optimize fall risk assessment in geriatric patients admitted to an outpatient diagnostic clinic. *PLoS One*, 12, e0178615. doi: [10.1371/journal.pone.0178615](https://doi.org/10.1371/journal.pone.0178615).

Kikkert, L. H., Vuillerme, N., van Campen, J. P., Hortobagyi, T. and Lamoth, C. J. (2016). Walking ability to predict future cognitive decline in old adults: a scoping review. *Ageing Research Reviews*, 27, 1–14. doi: [S1568-1637\(16\)30009-5 \[pii\]](https://doi.org/10.1016/j.arr.2016.05.005).

Kobsar, D., Olson, C., Paranjape, R., Hadjistavropoulos, T. and Barden, J. M. (2014). Evaluation of age-related differences in the stride-to-stride fluctuations, regularity and symmetry of gait using a waist-mounted tri-axial accelerometer. *Gait & Posture*, 39, 553–557. doi: [10.1016/j.gaitpost.2013.09.008](https://doi.org/10.1016/j.gaitpost.2013.09.008).

- Kosse, N. M., Caljouw, S., Vervoort, D., Vuillerme, N. and Lamoth, C. J.** (2015). Validity and reliability of gait and postural control analysis using the tri-axial accelerometer of the ipod touch. *Annals of Biomedical Engineering*, 43, 1935–1946. doi: [10.1007/s10439-014-1232-0](https://doi.org/10.1007/s10439-014-1232-0).
- Lamoth, C. J., Beek, P. J. and Meijer, O. G.** (2002). Pelvis-thorax coordination in the transverse plane during gait. *Gait & Posture*, 16, 101–114. doi: [S0966636201001461](https://doi.org/S0966636201001461) [pii].
- Lamoth, C. J., van Deudekom, F. J., van Campen, J. P., Appels, B. A., de Vries, O. J. and Pijnappels, M.** (2011). Gait stability and variability measures show effects of impaired cognition and dual tasking in frail people. *Journal of Neuroengineering and Rehabilitation*, 8, 2-0003-8-2. doi: [10.1186/1743-0003-8-2](https://doi.org/10.1186/1743-0003-8-2).
- Lang, P. O., Michel, J. P. and Zekry, D.** (2009). Frailty syndrome: a transitional state in a dynamic process. *Gerontology*, 55, 539–549. doi: [10.1159/000211949](https://doi.org/10.1159/000211949).
- Lipsitz, L. A.** (2002). Dynamics of stability: the physiologic basis of functional health and frailty. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 57, B115–125.
- Lipsitz, L. A. and Goldberger, A. L.** (1992). Loss of ‘complexity’ and aging: potential applications of fractals and chaos theory to senescence. *JAMA*, 267, 1806–1809. doi: [10.1001/jama.1992.03480130122036](https://doi.org/10.1001/jama.1992.03480130122036) [doi].
- Manor, B. et al.** (2010). Physiological complexity and system adaptability: evidence from postural control dynamics of older adults. *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 109, 1786–1791. doi: [10.1152/jappphysiol.00390.2010](https://doi.org/10.1152/jappphysiol.00390.2010).
- Montero-Odasso, M. et al.** (2014). The motor signature of mild cognitive impairment: results from the gait and brain study. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 69, 1415–1421. doi: [10.1093/gerona/glu155](https://doi.org/10.1093/gerona/glu155).
- Qualls, C. et al.** (2017). Reversible states of physical and/or cognitive dysfunction: a 9-Year longitudinal study. *The Journal of Nutrition, Health & Aging*, 21, 271–275. doi: [10.1007/s12603-017-0878-3](https://doi.org/10.1007/s12603-017-0878-3) [doi].
- Rispens, S. M., Pijnappels, M., van Schooten, K. S., Beek, P. J., Daffertshofer, A. and van Dieen, J. H.** (2014). Consistency of gait characteristics as determined from acceleration data collected at different trunk locations. *Gait & Posture*, 40, 187–192. doi: [10.1016/j.gaitpost.2014.03.182](https://doi.org/10.1016/j.gaitpost.2014.03.182).
- Riva, F., Toebes, M. J., Pijnappels, M., Stagni, R. and van Dieen, J. H.** (2013). Estimating fall risk with inertial sensors using gait stability measures that do not require step detection. *Gait & Posture*, 38, 170–174. doi: [10.1016/j.gaitpost.2013.05.002](https://doi.org/10.1016/j.gaitpost.2013.05.002).
- Savica, R. et al.** (2017). Comparison of gait parameters for predicting cognitive decline: the mayo clinic study of aging. *Journal of Alzheimer’s Disease: JAD*, 55, 559–567. doi: [JAD160697](https://doi.org/JAD160697) [pii].
- Sleimen-Malkoun, R., Temprado, J. J. and Hong, S. L.** (2014). Aging induced loss of complexity and dedifferentiation: consequences for coordination dynamics within and between brain, muscular and behavioral levels. *Frontiers in Aging Neuroscience*, 6, 140. doi: [10.3389/fnagi.2014.00140](https://doi.org/10.3389/fnagi.2014.00140).
- Studenski, S. et al.** (2011). Gait speed and survival in older adults. *JAMA*, 305, 50–58. doi: [10.1001/jama.2010.1923](https://doi.org/10.1001/jama.2010.1923).
- van Schooten, K. S., Pijnappels, M., Rispens, S. M., Elders, P. J., Lips, P. and van Dieen, J. H.** (2015). Ambulatory fall-risk assessment: amount and quality of daily-life gait predict falls in older adults. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 70, 608–615. doi: [10.1093/gerona/glu225](https://doi.org/10.1093/gerona/glu225).
- Verghese, J., Wang, C., Lipton, R. B. and Holtzer, R.** (2013). Motoric cognitive risk syndrome and the risk of dementia. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 68, 412–418.