

University of Groningen

## The impact of coronary revascularisation on endothelial and platelet function

Amoroso, Giovanni

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2001

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Amoroso, G. (2001). *The impact of coronary revascularisation on endothelial and platelet function*. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## Chapter 9

# Continued cigarette smoking after coronary artery bypass surgery reduces endothelium-dependent vasodilatation in internal thoracic artery grafts

Giovanni Amoroso<sup>1</sup>  
Massimo A. Mariani<sup>2</sup>  
Rene' A. Tio<sup>1</sup>  
Jan G. Grandjean<sup>2</sup>

<sup>1</sup>Department of Cardiology, Thoraxcenter, University Hospital of Groningen, The Netherlands

<sup>2</sup>Cardiothoracic Department, University of Pisa, Italy

## ABSTRACT

Cigarette smoking is known to promote endothelial dysfunction, thus it can be responsible for an impaired endothelium-dependent vasomotility in arterial grafts late after coronary surgery.

### Methods and results

20 consecutive patients (mean age 64.5 yrs.), previously submitted to coronary bypass surgery with the internal thoracic artery, underwent quantitative angiography of the implanted graft at long-term follow-up (mean time 2.5 yrs.). To assess both endothelium-dependent and -independent vasomotility, angiograms were acquired before and after selective infusions of acetylcholine ( $10^{-6}$  mmol/ml) and nitroglycerine (500 mg). The predictive value of risk factors, including previous and continued smoking, for an impairment in endothelium-dependent vasomotility was assessed.

Continued smoking ( $p=0.038$ ), but not a previous history of smoking ( $p=0.55$ ) was the only predictor of a reduced endothelium-dependent vasodilation. While previous smokers and non-smokers showed a similar response to acetylcholine, current smokers showed a reduced endothelium-dependent vasodilation vs. non-smokers ( $94.8\pm 2.6\%$  vs.  $99.6\pm 2.3\%$  of the maximal vasodilative capacity,  $p=0.001$ ).

### Conclusions

Although maintained, the vasodilative response to acetylcholine appear reduced in internal thoracic artery grafts of patients who continued smoking long-term after coronary bypass surgery. Whether this could affect the long term outcome of these patients has to be further investigated.

## INTRODUCTION

Cigarette smoking has a deleterious effect on coronary arteries since it is responsible for atherosclerosis, thrombogenicity, and vasoconstriction<sup>1</sup>. Cigarette smoking also worsens the outcome of patients after coronary surgery (CABG) by jeopardising the integrity of venous grafts<sup>2</sup>. Currently, there is a tendency to use the internal thoracic artery (ITA) as a first-choice conduit for CABG. The ITA shows an excellent angiographic patency, providing a good long-term event-free survival<sup>3</sup>. To investigate the effect of previous and/or continued cigarette smoking on this graft, we studied ITA's vasomotor response to acetylcholine, an endothelium-dependent vasodilator, by means of quantitative angiography.

### List of abbreviations:

CABG: coronary artery bypass surgery  
ITA: internal thoracic artery  
LVEF: left ventricle ejection fraction  
ECG: electrocardiogram  
ACE: angiotensin converting enzyme  
MI: myocardial infarction

## METHODS

Twenty consecutive patients (mean age:  $64.5 \pm 7.5$  yrs), with no evidences of angina and good left ventricular function (LVEF > 40%) were submitted to an angiographic follow-up (mean time:  $2.5 \pm 0.9$  yrs) after CABG with pedicled ITA. Patients gave informed consent and the protocol was approved by the Institutional Review Board. Among the study group, 10/20 patients (previous smokers) had a history of smoking before CABG: six of them (current smokers) were continuing to smoke at the time of follow-up. No patient started smoking after CABG.

Baseline angiograms of the ITA graft were acquired in a fixed radiological view, then acetylcholine was injected in the ITA in three increasing dosages ( $10^{-8}$ ,  $10^{-7}$  and  $10^{-6}$  mmol/ml x 3 minutes each) under ECG and blood pressure monitoring. Angiograms were repeated within 1 minute after each infusion and after a subsequent administration of nitroglycerine (500  $\mu$ g). End-diastolic frames were chosen and the mean diameter of the proximal part of the ITA was calculated by quantitative analysis (Quantocor QCA 3.0, Pie Medical Imaging, Maastricht, The Netherlands) for baseline, post-acetylcholine (highest dose achieved) and post-nitroglycerine sequences.

As nitrates are considered an endothelium-independent vasodilative agent, post-nitroglycerine mean diameter was taken as the maximal vasodilative capacity of the ITA: the endothelial contribution to vasodilation was then calculated for each patient as the ratio (in percentage) between post-acetylcholine (highest dose achieved) and post-nitroglycerine mean diameters, normalised for baseline diameters.

Statistical analysis was performed by means of the Statistical Package for the Social Science version 8.0 for Windows (SPSS Inc., Carey, NC). Descriptive statistics are expressed as mean values  $\pm$  SD or as percentages. Univariate regression analysis was performed to evaluate the influence of clinical features, medications, or risk factors on endothelium-dependent vasodilation, included as a continuous variable. Variables with a probability value of less than 0.20 were selected for multivariable regression analysis. Analysis of variance was performed for assessing statistical differences between groups (previous and current, smokers and non-smokers).

## RESULTS

All patients tolerated the highest acetylcholine dose ( $10^{-6}$  mmol/mL) without significant side effects.


At univariate analysis, continued smoking ( $p=0.038$ ), but not a previous history of smoking before surgery ( $p=0.55$ ) was the only predictor of an endothelium-dependent vasodilation impairment, thus a multivariable analysis was not performed. Smokers and non-smokers showed similar characteristics regarding clinical features, medications or risk factors (Table I).

Previous smokers vs. non-smokers showed comparable baseline values and maximal vasodilative capacity of the ITA and the endothelium-dependent vasodilation was not significantly different between the two groups ( $97.0 \pm 3.9\%$  vs.  $98.9 \pm 2.2\%$  of the maximal vasodilative capacity;  $p:0.20$ ). Current smokers vs. non-smokers

**Table 1.** Clinical features, medications and other risk factors for previous and current, smokers and non-smokers.

	Previous		Current	
	Smokers (n=10)	Non-smokers (n=10)	Smokers (n=6)	Non-smokers (n=14)
Age (years)	64.4±5.8	64.5±9	64.5±7.3	64.5±7.8
Follow-up (months)	25.1±18.3	35.4±19.8	21.8±17.7	34.5±19.3
Sequential anastomosis	3 (30%)	4 (40%)	3 (50%)	4 (29%)
LVEF (%)	51.5±9.2	51.9±9.5	51.4±8.9	52.1±9.7
ACE inhibitors use	3 (30%)	4 (40%)	3 (50%)	4 (29%)
Statins use	3 (30%)	5 (50%)	2 (33%)	6 (43%)
History of:				
Hypercholesterolemia	3 (30%)	6 (60%)	2 (33%)	7 (50%)
Systemic hypertension	3 (30%)	3 (30%)	2 (33%)	4 (29%)
Diabetes mellitus	2 (20%)	0	1 (16%)	1 (7%)
MI in the grafted area	2 (20%)	2 (20%)	1 (16%)	3 (21%)

Legend: LVEF: left ventricular ejection fraction; MI: myocardial infarction.

**Figure 1.** Endothelium-dependent vasodilation (as percentage of the maximal vasodilative capacity) of ITA grafts in smokers and non-smokers.

showed also comparable baseline values and maximal vasodilative capacity of the ITA, but the endothelium-dependent vasodilation was significantly lower in current smokers than in non-smokers ( $94.8 \pm 2.6$  % vs.  $99.6 \pm 2.3$  %;  $p:0.001$  "95% CI: -7.1 to -2.1") (Figure 1).

## DISCUSSION

This study demonstrates that, though ITAs maintain a vasodilative response to acetylcholine infusion in the presence of coronary atherosclerosis and after CABG, continued smoking, but not a previous history of smoking before surgery, predicts a reduction in endothelium-dependent vasodilatation of the arterial graft.

Cigarette smoking is known to reduce the production of nitric oxide, alter the rheology at the endothelial surface and upregulate the expression of leukocyte adhesion molecules. Smoking can enhance the oxidation of circulating low-density lipoproteins, thus leading to the development of atherosclerosis. It is also able to promote thrombosis onto the dysfunctional endothelium. In the clinical setting this translates as an increased incidence of cardiac adverse events <sup>4</sup>.

Previous studies have reported that smokers have a worse outcome after CABG, due to an increased hazard for perioperative complications, graft occlusion and re-intervention <sup>2</sup>. Referring to the unfavourable results and hypothesising a waste of health care resources, some authors have even suggested not offering CABG to smokers <sup>5</sup>.

Although smoking can be held responsible for myocardial and coronary damages occurring before surgery, a previous history of smoking before CABG does not seem a mayor determinant for a future impairment of the arterial graft integrity. In a previous study we found that both endothelium-dependent and independent vasomotility of arterial conduits are correlated with the age of the patient but do not deteriorate over time after surgery <sup>6</sup>. Even in the presence of diffuse atherosclerosis, and differently from coronary arteries, which show an aberrant vasoconstrictive response to acetylcholine, ITAs vasodilate regardless of any risk factor, including smoking habit before surgery.

However, in patients who continued smoking after surgery the endothelium-dependent vasodilatation is mildly but significantly impaired: whether this finding might negatively influence the long-term outcome in those patients receiving arterial grafts is questionable, also due to the small dimensions of our study group. Nevertheless, an effort should be made in convincing patients who undergo CABG of the beneficial effect of quitting smoking with the aim of preserving the integrity of the implanted grafts.

## References

1. Zhu BQ, Parmley WW. Hemodynamic and vascular effects of active and passive smoking. *Am Heart J* 1995;130:1270-5.
2. Voors AA, van Brussel BL, Plokker HW, et al. Smoking and cardiac events after venous coronary bypass surgery. A 15-year follow-up study. *Circulation* 1996;93:42-7.
3. Cameron A, Davis KB, Green G, Schaff HV. Coronary bypass surgery with internal-thoracic-artery grafts - Effects on survival over a 15-year period. *N Engl J Med* 1996;334:216-9
4. Pepine CJ, Schlaifer JD, Mancini GB, Pitt B, O'Neill BJ, Haber HE. Influence of smoking status on progression of endothelial dysfunction. TREND Investigators. Trial on Reversing Endothelial Dysfunction. *Clin Cardiol* 1998;21:331-4.
5. Underwood MJ, Bailey JS. Coronary bypass surgery should not be offered to smokers. *BMJ* 1993;306:1047-9.
6. Amoroso G, Tio RA, Mariani MA, et al. Functional integrity and aging of the left internal thoracic artery after coronary artery bypass surgery. *J Thorac Cardiovasc Surg* 2000; 120:313-8.