

## University of Groningen

### Protocol Letter

Humbert-Vidan, Laia; Hansen, Christian R; Fuller, Clifton D; Petit, Steven; van der Schaaf, Arjen; van Dijk, Lianne V; Verduijn, Gerda M; Langendijk, Hans; Muñoz-Montplet, Carles; Heemsbergen, Wilma

*Published in:*  
Radiotherapy and Oncology

*DOI:*  
[10.1016/j.radonc.2022.09.014](https://doi.org/10.1016/j.radonc.2022.09.014)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2022

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Humbert-Vidan, L., Hansen, C. R., Fuller, C. D., Petit, S., van der Schaaf, A., van Dijk, L. V., Verduijn, G. M., Langendijk, H., Muñoz-Montplet, C., Heemsbergen, W., Witjes, M., Mohamed, A. S. R., Khan, A. A., Marruecos Querol, J., Oliveras Cancio, I., Patel, V., King, A. P., Johansen, J., & Guerrero Urbano, T. (2022). Protocol Letter: A multi-institutional retrospective case-control cohort investigating PREDiction models for mandibular OsteoRadioNecrosis in head and neck cancer (PREDMORN). *Radiotherapy and Oncology*, 176, 99-100. <https://doi.org/10.1016/j.radonc.2022.09.014>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Contents lists available at ScienceDirect

# Radiotherapy and Oncology

journal homepage: [www.thegreenjournal.com](http://www.thegreenjournal.com)

## Guideline Letter

### Protocol Letter: A multi-institutional retrospective case-control cohort investigating PREDiction models for mandibular OsteoRadioNecrosis in head and neck cancer (PREDMORN)



Laia Humbert-Vidan<sup>a,b,\*</sup>, Christian R. Hansen<sup>c</sup>, Clifton D. Fuller<sup>d</sup>, Steven Petit<sup>e</sup>, Arjen van der Schaaf<sup>f</sup>, Lisanne V. van Dijk<sup>f</sup>, Gerda M. Verduijn<sup>e</sup>, Hans Langendijk<sup>f</sup>, Carles Muñoz-Montplet<sup>g,h</sup>, Wilma Heemsbergen<sup>e</sup>, Max Witjes<sup>f</sup>, Abdallah S.R. Mohamed<sup>d</sup>, Abdul A. Khan<sup>i</sup>, Jordi Marruecos Querol<sup>h,j</sup>, Irene Oliveras Cancio<sup>j</sup>, Vinod Patel<sup>k</sup>, Andrew P. King<sup>l</sup>, Jørgen Johansen<sup>m</sup>, Teresa Guerrero Urbano<sup>n</sup>

<sup>a</sup> Department of Medical Physics, Guy's and St Thomas' NHS Foundation Trust; <sup>b</sup> School of Cancer and Pharmaceutical Sciences, Comprehensive Cancer Centre, King's College London, London, UK; <sup>c</sup> Department of Clinical Research, University of Southern Denmark, Odense, Denmark; <sup>d</sup> Department of Radiation Oncology, The University of Texas MD Anderson Cancer Centre, Houston, TX, United States; <sup>e</sup> Department of Radiotherapy, Erasmus MC Cancer Institute, University Medical Center Rotterdam, Rotterdam; <sup>f</sup> Department of Radiation Oncology, University Medical Centre Groningen, University of Groningen, Groningen, Netherlands; <sup>g</sup> Department of Medical Physics and Radiation Protection, Catalan Institute of Oncology, Girona, Spain; <sup>h</sup> Department of Medical Sciences, University of Girona, Girona, Spain; <sup>i</sup> Department of Oral and Maxillofacial Surgery, Odense University Hospital, Odense, Denmark; <sup>j</sup> Department of Radiation Oncology, Catalan Institute of Oncology, Girona, Spain; <sup>k</sup> Department of Oral Surgery, Guy's and St Thomas' NHS Foundation Trust; <sup>l</sup> School of Biomedical Engineering and Imaging Sciences, King's College London, London, UK; <sup>m</sup> Department of Oncology, Odense University Hospital, Odense, Denmark; <sup>n</sup> Department of Clinical Oncology, Guy's and St Thomas' NHS Foundation Trust, London, UK

## Background

Osteoradionecrosis (ORN) of the mandible is a rare but severe radiation-induced toxicity observed in 4–8% [1–3] of head and neck cancer (HNC) cases treated with radiotherapy (RT) alone or combined with surgery and/or chemotherapy (CRT). Ionising radiation damages the vascularisation of the mandible. Consequently, necrosis of the bone can develop either spontaneously or triggered by trauma to the mandible bone (e.g. dental extractions, surgery, implants). Bone necrosis occurs because it is not able to heal due to reduced blood supply, hypoxia or hypo-cellularity caused by exposure to radiation [4]. The severity of ORN has been graded by Notani [5] into three categories, with the most severe cases experiencing significant pain levels and even pathological fracture of the mandible. Quality of life can be greatly affected in patients who develop mandibular ORN. Moreover, the management of ORN is often complex [6] and requires costly clinical interventions [7].

Normal tissue complication probability (NTCP) prediction models are a valuable treatment decision support tool that can be used to estimate the risk of individual patients developing a particular normal tissue toxicity outcome. Patient-specific prediction of potential ORN in the treatment of HNC may lead to more individualised treatment with risk-reduction measures such as reduced mandibular radiation dose near extraction sites when possible and more dedicated follow-up regimes for early detection and intervention of ORN. A number of case-control studies have inves-

tigated the correlation between dosimetric, clinical and demographic factors and ORN [1–3,8–15]. Efforts on patient-specific prediction of ORN, however, are more limited. An NTCP model for ORN was recently developed [16] based on mandible dose-volume parameters and clinical variables using multivariable stepwise forward selection regression analysis, resulting in a final model that was based on the D30% of the mandible bone and pre-RT dental extraction. Previous work [17] has shown that it is possible to use deep-learning (DL) methods to predict ORN incidence from 3D radiation dose distribution maps of the mandible.

Due to the low prevalence rate of mandibular ORN, low patient numbers represent a statistical limitation. External validation of models using data from multiple centres is an essential feature of effective model evaluation with a view to clinical translation but is often lacking in research studies [18]. Combining datasets from different institutions may not only improve the generalizability of the models but also highlight correlations between clinical practice and toxicity outcome. The present study is a multi-institutional effort involving six teaching hospitals (details provided in Appendix B in the protocol) and including the largest datasets worldwide to develop, train and validate robust and generalisable NTCP models for mandibular ORN.

## Protocol summary

This is a retrospective case-control study where we hypothesise that mandibular ORN can be accurately predicted on an individual level. To test our hypothesis, we will develop and validate multiple NTCP models based on dosimetric parameters extracted from the mandible dose-volume histogram (DVH) and clinical and demographic variables using different mathematical approaches. Both

\* Corresponding author at: Radiotherapy Physics Department, Guy's Cancer Centre, Great Maze Pond, London SE1 9RT, UK.

E-mail address: [laia.humbert-vidan@kcl.ac.uk](mailto:laia.humbert-vidan@kcl.ac.uk) (L. Humbert-Vidan).

the mandible DVH data and the resulting principal components (PCs) from a preliminary principal component analysis (PCA) will be considered as dosimetric variables in separate models. Data will be retrospectively collected by each centre based on well-described parameters (Appendix A in the protocol) and inclusion/exclusion criteria (Table 1 in the protocol). A case-control approach will be followed with a 2:1 control to ORN case ratio (the selection process of the control cohort is described in Fig. 1 of the protocol).

Multiple NTCP models will be trained, validated and tested applying different modelling approaches (LASSO logistic regression, regularised greedy forest and Bayesian neural network) to clinical and dosimetric variables. The different models will be compared and ranked according to their predictive performance, interpretability, parsimony and clinical applicability. Identification of the best performing NTCP model and the most highly predictive variables will provide valuable information for clinical decision-making and treatment optimisation. Clinically applicable mandible radiation dose constraints for radiation treatment plan optimisation will contribute to reduced complication probability.

### Planned future work

NTCP models have traditionally used DVH parameters and this study will use such DVH metrics, along with clinical factors, as variables. However, in a DVH the spatial localisation, dose gradient and direction information is lost. Spatial dosimetric information is, however, of great clinical interest in the investigation of radiation damage to the organs at risk in the HN region. Some studies [17,27,28] have already successfully implemented DL methods such as convolutional neural networks (CNNs) in NTCP models. The CNNs are trained to automatically predict the toxicity probability by adaptively learning the most predictive deep features directly from the 3D radiation distribution maps. Future work will aim to validate and update the existing DL-based ORN NTCP model [17] through the inclusion of the multi-institutional image dataset containing 3D radiation dose distribution maps.

### Study limitations

The primary limitation of this study results from challenges in the standardisation of retrospective data across different centres and potential biases in the data.

The case control design may result in a dataset where the underrepresented groups, i.e. the primary tumour site groups with low event rates, may be oversampled compared to clinical practice. This might introduce complications in the modelling and interpretation of the models. However, sampling the entire HNC population is not feasible for all institutions. In order to compensate for this sampling bias, the actual total number of events and non-events will be recorded per primary tumour site group in order to accurately predict the proportions.

Follow-up time period differences between institutions make it difficult to perform a time-to-event analysis. However, we will perform time-independence prediction models using fixed rates of ORN at a given interval and will look at continuous right censoring approaches as well as a post hoc analysis.

### Funding

This work is supported by the Radiation Research Unit at the Cancer Research UK City of London Centre Award [C7893/A28990] and by the Guy's Cancer Charity via a donation from the Wilson-Olegario foundation and other donations.

## Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.radonc.2022.09.014>.

## References

- [1] Frankart AJ, Frankart MJ, Cervenka B, Tang AL, Krishnan DG, Takiar V. Osteoradionecrosis: Exposing the evidence not the bone. *Int J Radiation Oncol Biol Phys* 2021;109:1206–18.
- [2] Moon DH, Moon SH, Wang K, Weissler MC, Hackman TG, Zanation AM, et al. Incidence of, and risk factors for, mandibular osteoradionecrosis in patients with oral cavity and oropharynx cancers. *Oral Oncol* 2017;72:98–103.
- [3] De Felice F, Tombolini V, Musio D, Polimeni A. Radiation Therapy and Mandibular Osteoradionecrosis: State of the Art. *Curr Oncol Rep* 2020;22:89. <https://doi.org/10.1007/s11912-020-00954-3>. PMID: 32642937.
- [4] Chen J-A, Wang C-C, Wong Y-K, Wang C-P, Jiang R-S, Lin J-C, et al. Osteoradionecrosis of mandible bone in patients with oral cancer-associated factors and treatment outcomes. *Head Neck* 2016;38:762–8. <https://doi.org/10.1002/hed.23949>.
- [5] Notani K et al. Osteoradionecrosis of the mandible – factors influencing severity. *Asian J Oral Maxillofac Surgery* 2002;14:5–9.
- [6] Patel V, Fenlon M, Di Silvo L, McGurk M. Osteoradionecrosis in the Current Era of Radiation Treatment. *Dental Update* 2022;49:64–7.
- [7] Patel V, Ormondroyd L, Lyons A, McGurk M. The financial burden for the surgical management of osteoradionecrosis. *Br Dent J* 2017;222:177–80.
- [8] Aarup-Kirstensen S, Hansen CR, Forner L, Brink C, Eriksen JG, Johansen J. Osteoradionecrosis of the mandible after radiotherapy for head and neck cancer: risk factors and dose-volume correlations. *Acta Oncol* 2019;58:1373–7.
- [9] MD Anderson Head and Neck Cancer Symptom Working Group. Dose-volume correlates of mandibular osteoradionecrosis in oropharynx cancer patients receiving intensity-modulated radiotherapy: Results from a case-matched comparison. *Radiother Oncol* 2017;124:232–9. <https://doi.org/10.1016/j.radonc.2017.06.026>.
- [10] Kubota H, Miyawaki D, Mukumoto N, Ishihara T, Matsumura M, Hasegawa T, et al. Risk factors for osteoradionecrosis of the jaw in patients with head and neck squamous cell carcinoma. *Radiat Oncol* 2021;16:1. <https://doi.org/10.1186/s13014-020-01701-5>.
- [11] Owosho AA, Tsai CJ, Lee RS, Freymiller H, Kadempour A, Varthi S, et al. The prevalence and risk factors associated with osteoradionecrosis of the jaw in oral and oropharyngeal cancer patients treated with intensity-modulated radiation therapy (IMRT): The Memorial Sloan Kettering Cancer Center experience. *Oral Oncol* 2017;64:44–51.
- [12] Raguse J-D, Hossamo J, Tinhofer I, Hoffmeister B, Budach V, Jamil B, et al. Patient and treatment-related risk factors for osteoradionecrosis of the jaw in patients with head and neck cancer. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2016;121:215–21. <https://doi.org/10.1016/j.oooo.2015.10.006>.
- [13] Nabil S, Samman N. Incidence and prevention of osteoradionecrosis after dental extraction in irradiated patients: a systematic review. *Int J Oral Maxillofac Surg* 2011;40:229–43. <https://doi.org/10.1016/j.ijom.2010.10.005>.
- [14] Gomez DR, Estilo CL, Wolden SL, Zelefsky MJ, Kraus DH, Wong RJ, et al. Correlation of osteoradionecrosis and dental events with dosimetric parameters in intensity-modulated radiation therapy for head-and-neck cancer. *Int J Radiat Oncol Biol Phys* 2011;81:e207–13. <https://doi.org/10.1016/j.ijrobp.2011.02.003>.
- [15] Lee JJ, Koom WS, Lee CG, Kim YB, Yoo SW, Keum KC, et al. Risk factors and dose-effect relationship for mandibular osteoradionecrosis in oral and oropharyngeal cancer patients. *Int J Radiat Oncol Biol Phys* 2009;75:1084–91. <https://doi.org/10.1016/j.ijrobp.2008.12.052>.
- [16] Van Dijk LV, Abusaif AA, Rigert J, Naser MA, Hutcheson KA, Lai SY, et al. Normal Tissue Complication Probability (NTCP) Prediction Model for Osteoradionecrosis of the Mandible in Patients With Head and Neck Cancer After Radiation Therapy: Large-Scale Observational Cohort. *Int J Radiation Oncol Biol Phys* 2021;111:549–58.
- [17] Humbert-Vidan L, Patel V, Andlauer R, King AP, Guerrero Urbano T. Preprint: Prediction of mandibular ORN incidence from 3D radiation dose distribution maps using deep learning. arXiv:2112.11503 [physics.med-ph]; 23 Jun 2022.
- [18] Collins GS, Reitsma JB, Altman DG, Moons KG. Transparent Reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): the TRIPOD statement. *Br J Surg* 2015;102:148–58.
- [27] Men K, Geng H, Zhong H, Fan Y, Lin A, Xiao Y. A deep learning model for predicting xerostomia due to radiotherapy for head-and-neck squamous cell carcinoma in the RTOG 0522 clinical trial. *Int J Radiat Oncol Biol Phys* 2019;105:440–7.
- [28] Ibragimov B, Toesca D, Chang D, Yuan Y, Koong A, Xink L. Development of deep neural network for individualized hepatobiliary toxicity prediction after liver sbrt. *Med Phys* 2018;45:10.