

University of Groningen

## Thiols as markers of redox status in type 1 diabetes mellitus

van Dijk, Peter R; Pasch, Andreas; van Ockenburg-Brunet, Sonja L; Waanders, Femke; Eman Abdulle, A; Muis, Marian J; Hillebrands, J L; Bilo, Henk J G; van Goor, Harry

*Published in:*  
Therapeutic advances in endocrinology and metabolism

*DOI:*  
[10.1177/2042018820903641](https://doi.org/10.1177/2042018820903641)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2020

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

van Dijk, P. R., Pasch, A., van Ockenburg-Brunet, S. L., Waanders, F., Eman Abdulle, A., Muis, M. J., Hillebrands, J. L., Bilo, H. J. G., & van Goor, H. (2020). Thiols as markers of redox status in type 1 diabetes mellitus. *Therapeutic advances in endocrinology and metabolism*, 11. Advance online publication. <https://doi.org/10.1177/2042018820903641>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Thiols as markers of redox status in type 1 diabetes mellitus

Peter R. van Dijk<sup>1</sup>, Andreas Pasch, Sonja L. van Ockenburg-Brunet, Femke Waanders, A. Eman Abdulle, Marian J. Muis, J. L. Hillebrands, Henk J. G. Biló and Harry van Goor

## Abstract

**Introduction:** Type 1 diabetes mellitus (T1DM) is associated with inflammation and the production of reactive oxygen species (ROS). Systemically, free thiols (R-SH) can be oxidized by ROS and circulating R-SH concentrations may directly reflect the systemic redox status. In this study the association between R-SH and clinical parameters of T1DM, including glycated haemoglobin A1c (HbA1c), was investigated. This is of particular interest since thiols are amendable to therapeutic intervention.

**Methods:** As part of a prospective cohort study, data from 216 patients with a mean age of 45 (12) years, 57% male, diabetes duration 22 (16, 30) years and HbA1c of 60 (11) mmol/mol were examined. Baseline data were collected in 2002 and follow-up data in 2018. Cox proportional hazards regression analysis, with age, sex, HbA1c and R-SH, was used to assess prognostic factors for the development of complications.

**Results:** At baseline, the plasma concentration of R-SH was  $281.8 \pm 34.0 \mu\text{M}$ . In addition to a lower concentration of NT-proBNP in the highest R-SH quartile (305–379  $\mu\text{M}$ ) there were no differences in baseline characteristics between the quartiles of R-SH. The Pearson correlation coefficient for R-SH and NT-proBNP was  $-0.290$  ( $p < 0.001$ ). No significant correlation between R-SH and baseline HbA1c ( $r = -0.024$ ,  $p = 0.726$ ) was present. During follow-up, 42 macrovascular and 92 microvascular complications occurred. In Cox regression, R-SH was not a prognostic factor for the development of microvascular [hazard ratio (HR) 0.999 (95% confidence interval (CI) 0.993, 1.005)] and macrovascular [HR 0.993 (95% CI 0.984, 1.002)] complications.

**Conclusions:** In addition to a negative association with NT-proBNP, no relevant relationships between R-SH and parameters of T1DM, including HbA1c, were present in this study.

**Keywords:** free sulfhydryl, glycaemia, oxidative stress, thiols, type 1 diabetes

Received: 16 October 2019; revised manuscript accepted: 17 December 2019.

## Introduction

Systemic oxidative stress is caused by an imbalance between oxidative and antioxidative systems, resulting in a net overload of oxidative free radicals and thereby reactive oxygen species (ROS).<sup>1</sup> As a pathogenic result, these excessive amounts of ROS modify the function and structure of cellular components such as proteins and lipids. This leads to cellular dysfunction, as evidenced by an impairment of energy metabolism, changes in cell signalling, variations in cell cycle

control and an impairment of cell transport mechanisms.<sup>2</sup> Subsequently, a dysfunction in processes such as immune activation and inflammation occur. Hyperglycaemia by itself is able to promote the production of ROS as revealed by an enhancement in lipid peroxidation.<sup>2,3</sup> These processes are considered central in the pathogenesis of type 1 diabetes mellitus (T1DM) related microvascular and macrovascular complications.<sup>4</sup> Despite improvements in the management of glycaemia and cardiovascular risk factors,

*Ther Adv Endocrinol Metab*

2020, Vol. 11: 1–8

DOI: 10.1177/  
2042018820903641

© The Author(s), 2020.  
Article reuse guidelines:  
sagepub.com/journals-  
permissions

Correspondence to:  
**Peter R. van Dijk**  
Department of Internal  
Medicine, University of  
Groningen, University  
Medical Center,  
Hanzeplein 1, 9700 RB,  
HP AA41, Groningen, The  
Netherlands

Isala, Diabetes Centre,  
Zwolle, the Netherlands  
[p.r.van.dijk@umcg.nl](mailto:p.r.van.dijk@umcg.nl)

**Andreas Pasch**  
Calcison AG, Nidau,  
Switzerland

**Sonja L. van Ockenburg-  
Brunet**  
**Henk J. G. Biló**  
Department of  
Internal Medicine,  
University of Groningen,  
University Medical  
Center, Groningen, the  
Netherlands

**Femke Waanders**  
Isala, Department of  
Internal Medicine, Zwolle,  
the Netherlands

**A. Eman Abdulle**  
Division of Vascular  
Medicine, University  
of Groningen,  
University Medical  
Center, Groningen, the  
Netherlands

**Marian J. Muis**  
Isala, Diabetes Centre,  
Zwolle, the Netherlands  
Isala, Department of  
Internal Medicine, Zwolle,  
the Netherlands

**J. L. Hillebrands**  
**Harry van Goor**  
Department of Pathology  
and Medical Biology,  
University of Groningen,  
University Medical  
Center, Groningen, the  
Netherlands

T1DM is still accompanied by a high rate of complications, all of which are associated with mortality as well as reduced health-related quality of life (HRQOL).<sup>5</sup>

Thiols are organic compounds that are characterized by the presence of a sulfhydryl (R-SH) moiety.<sup>6</sup> Thiols exist as proteins with one (or more) free cysteine groups or as low-molecular-weight compounds like glutathione in cells and also in extracellular fluids. Systemically, the concentration of thiols is lower than intracellularly. In serum, albumin is the most abundant thiol.<sup>7</sup> R-SH groups can be oxidized by ROS and by other reactive species such as sulphur and nitrogen species. The plasma concentrations of total R-SH are proposed to reflect the systemic body redox status indicating that a decline in circulating R-SH reveals enhancement of the oxidative tone.<sup>8,9</sup> Free thiols are receptive to therapeutic modulation, for example by cysteine derivatives such as N-acetylcysteine. Evaluation of systemic free thiols in disease conditions may therefore have diagnostic or therapeutic merit.<sup>7,10,11</sup>

Serum free thiol depletion has been reported in patients with cardiovascular disease, including acute myocardial infarction (MI), when compared with controls.<sup>12,13</sup> In stable heart failure patients, higher levels of serum free thiols were associated with favourable diseases outcomes.<sup>9</sup> In addition, high R-SH serum concentrations have been shown to be associated with a beneficial cardiovascular risk profile and a better patient and graft survival in renal transplant recipients.<sup>14</sup> Interestingly, in inflammatory bowel disease (IBD) free thiols are a better reflection of the disease activity as determined by endoscopy, then faecal calprotectin, the classical biomarker for IBD.<sup>15</sup> Recently, increased oxidation of thiols towards the disulphide form was recently found in T1DM patients as compared with healthy subjects.<sup>16</sup> In that analysis, the total thiol concentration was correlated negatively with C-reactive protein and glycated haemoglobin A1c (HbA1c).

Given these results, and the potential of R-SH as a modifiable marker of ROS-mediated damage in the progression of T1DM and associated complications, we hypothesized that high R-SH concentrations are positively associated with clinical parameters of T1DM. Therefore, we studied R-SH in a cohort of T1DM patients.

## Patients and methods

The FANTA study was designed as a prospective, cohort study to investigate several disease factors, including oxidative stress and HRQOL in persons with T1DM. Full study design and the results of HRQOL analysis in a subset of patients have been published in detail previously.<sup>17,18</sup> In brief, from January 1995 to January 1996, consecutive visiting T1DM patients treated at the diabetes outpatient clinic of the Weezenlanden Hospital (nowadays Isala; Zwolle, The Netherlands) were invited to participate in the study. T1DM was defined as starting insulin therapy within 6 months after the first signs of diabetes and before the age of 30 years, or the absence of C-peptide secretion. In total, 293 patients agreed to participate. In the period from 1996 to 2002, a total of 32 patients dropped out of the study. Reasons for dropping out were: moving out of the area or referral to another physician ( $n=12$ ), unknown ( $n=10$ ), lack of interest ( $n=6$ ), death ( $n=2$ ) and incorrect diagnosis of T1DM ( $n=2$ ). For the present analyses, we analysed the 261 patients who were participating in 2002. For 45 patients, no serum samples were available. Therefore, a total of 216 patients were included in the present analysis.

The primary objective of the present study was to investigate the transversal association of serum R-SH concentrations and HbA1c in T1DM. As a secondary outcome, association of R-SH with other clinical characteristics of T1DM was assessed. Furthermore, the longitudinal association of serum R-SH concentrations with the development of microvascular and macrovascular complications was investigated.

At baseline, aliquots of ethylenediaminetetraacetic acid (EDTA) samples were collected and stored at  $-80^{\circ}\text{C}$  (without thawing) until measurement. R-SH were measured as described previously, with minor modifications.<sup>19,20</sup> Briefly, 75  $\mu\text{l}$  serum was diluted 1:4 in 0.1 M Tris buffer (pH 8.2) and transferred to a 96-well plate. Using a Sunrise microplate reader (Tecan AG, Männedorf, Switzerland), background absorption was measured at 412 nm with a reference filter at 630 nm. Subsequently, 10  $\mu\text{l}$  3.8 mM 5,5'-dithiobis(2-nitrobenzoic acid) (Sigma-Aldrich, Zwijndrecht, Netherlands) in 0.1 M phosphate buffer (pH 7) was added to the samples. Following 20 min of incubation at room temperature, absorption was

read again. The concentration of R-SH in the samples was determined by comparing their absorbance readings with a standard curve of L-cysteine (15–1000  $\mu\text{M}$ ; Fluka Biochemika, Buchs, Switzerland) in 0.1 M Tris and 10 mM EDTA (pH 8.2).

Data concerning demographics, mode of therapy, height, weight, presence of complications, blood pressure and laboratory measurements were collected annually during follow-up according to a standardized protocol and standardized forms. For the analysis with respect to complications and vital status follow-up data from the year 2018 was gathered. In addition to the standardized forms an additional search concerning complications and vital status was performed using electronic hospital records. HbA1c is expressed in SI, International Federation of Clinical Chemistry and Laboratory Medicine recommended (as mmol/mol) units.

Macrovascular complications were defined as angina pectoris (AP), peripheral artery disease (PAD), MI, percutaneous transluminal coronary angioplasty (PTCA), coronary artery bypass grafting (CABG), cerebrovascular accident (CVA) or transient ischemic attack (TIA). Microvascular complications were defined as diabetic retinopathy, albuminuria (both microalbuminuria and macroalbuminuria) and diabetic peripheral neuropathy. Microalbuminuria was defined as 20–200 mg/l albumin or an albumin–creatinine ratio between 2.5 and 25 mg/mmol (22–221 mg/g) in men and 3.5 and 35 mg/mmol (31–310 mg/g) in women. Macroalbuminuria was defined as >200 mg/l albumin or a albumin–creatinine ratio greater than 25 mg/mmol and 35 mg/mmol for men and women, respectively.<sup>21</sup> An ophthalmologist determined the presence of diabetic retinopathy biannually. Foot sensibility was tested with 5.07 Semmes–Weinstein monofilaments. Neuropathy was defined as two or more errors in a test of three, affecting at least one foot.

Descriptive summaries included the mean with standard deviation (SD) for normally distributed variables and the median with the interquartile range (25th–75th percentile) for other variables. Baseline data were compared with the chi-squared test in case of categorical data. In case of continuous data, Student's *t* test or Mann–Whitney *U* test were used if the data was distributed normally or skewed, respectively. *Q–Q* plots and histograms were used to determine whether the tested

variable had a normal distribution or not. The Pearson correlation coefficient was used to investigate the relation between baseline R-SH and HbA1c concentrations. To visualize the relation of the tertiles of R-SH with the development of complications and death during follow-up, a Kaplan–Meier curve was constructed. Differences in time until occurrence of any complications were assessed for statistical significance using the log-rank test. Cox proportional hazards regression analysis, with age, sex, HbA1c and R-SH, was used to assess prognostic factors for the development of microvascular or macrovascular complication. A (two-sided) *p* value of less than 0.05 was considered statistically significant.

All analyses were performed using SPSS version 24.0 (SPSS Inc. Chicago, IL, USA). A (two-sided) *p* value of less than 0.05 was considered statistically significant. The study was performed in accordance with the Declaration of Helsinki. Written and oral informed consent was obtained from all patients, and the local medical ethics committee of Isala (METC Isala, Zwolle, the Netherlands) approved the protocol.

## Results

Baseline characteristics of the 216 patients with T1DM are presented in Table 1. The cohort had a mean age of 45.1 (11.6) years, 57% were male, diabetes duration was 22.1 (15.8, 30.4) years and HbA1c concentration was 60 (11) mmol/mol. At baseline, there were 110 (51%) patients with a microvascular complication and 22 (10%) patients with macrovascular complications. Among all patients, R-SH concentrations were normally distributed, with a mean concentration of 281.8 (34.0)  $\mu\text{M}$ .

Concentrations of R-SH did not differ between men and woman [281.5  $\mu\text{M}$  (255.0, 301.8), 287.5  $\mu\text{M}$  (260.0, 309.8), *p* = 0.44]. In addition to a lower concentration of NT-proBNP in the highest quartile of R-SH (305–379  $\mu\text{M}$ ) there were no differences in baseline characteristics between the quartiles of R-SH (Table 1). The Pearson correlation coefficient for R-SH and NT-proBNP was  $-0.290$  (*p* < 0.001). Of note, there was no significant correlation between R-SH and HbA1c at baseline (*r* =  $-0.024$ , *p* = 0.726).

During the follow-up duration of 15.4 (7.4, 15.8) years, one patient died (noncardiovascular cause)

**Table 1.** Baseline characteristics of all patients and per quartile of R-SH.

	All	Quartiles of R-SH			
	n = 216	172–257 $\mu\text{M}$ n = 54	258–284 $\mu\text{M}$ n = 51	285–304 $\mu\text{M}$ n = 58	305–379 $\mu\text{M}$ n = 53
<b>Demographics</b>					
Age (years)	45.1 (11.6)	46.3 (7.3)	44.6 (9.5)	44.4 (12.5)	45.2 (15.4)
Diabetes duration (years)	22.1 (15.8, 30.4)	23.4 (15.2, 31.3)	21.4 (16.4, 32.9)	20.7 (14.6, 30.1)	24.3 (16.5, 29.5)
Male gender (n)	123 (57.4)	33 (61.1)	32 (62.7)	34 (58.6)	25 (47.2)
BMI (kg/m <sup>2</sup> )	25.4 (23.3, 28.4)	25.5 (23.1, 28.6)	25.2 (22.8, 28.6)	25.7 (23.3, 29.0)	25.5 (23.6, 28.3)
Systolic blood pressure (mmHg)	130.4 (17.6)	132.6 (19.2)	129.1 (15.2)	131.4 (17.2)	128.3 (18.6)
Diastolic blood pressure (mmHg)	77.3 (10.2)	77.9 (10.1)	76.1 (10.4)	80.0 (10.3)	74.8 (9.3)
Mode of insulin administration: MDI	124 (57.4)	35 (64.8)	24 (47.1)	38 (65.5)	27 (50.9)
Mode of insulin administration: CSII	92 (42.6)	19 (35.2)	27 (52.9)	20 (34.5)	26 (49.1)
Total insulin dose <sup>s</sup>	58.8 (19.6)	60.3 (20.2)	49.5 (16.0)	60.9 (19.5)	39.3 (17.9)
<b>Complications</b>					
Microvascular complications					
Retinopathy (n)	92 (42.6)	26 (48.1)	22 (43.1)	20 (34.5)	24 (45.3)
Neuropathy (n)	24 (11.1)	3 (5.6)	6 (11.8)	7 (12.1)	8 (15.1)
Micro albuminuria (n)	28 (13.0)	10 (18.5)	5 (9.8)	8 (13.8)	5 (9.4)
Macro albuminuria (n)	8 (3.7)	5 (9.3)	0 (0.0)	1 (1.7)	2 (3.8)
Macrovascular events					
Angina pectoris (n)	3 (1.4)	2 (3.7)	0 (0.0)	1 (1.7)	0 (0.0)
MI (n)	3 (1.4)	0 (0.0)	0 (0.0)	1 (1.7)	2 (3.8)
PTCA (n)	3 (1.4)	1 (1.9)	0 (0.0)	2 (3.4)	0 (0.0)
PAD (n)	5 (2.3)	1 (1.9)	2 (3.9)	0 (0.0)	2 (3.8)
CABG (n)	3 (1.4)	0 (0.0)	1 (2.0)	2 (3.4)	0 (0.0)
TIA (n)	2 (0.9)	0 (0.0)	1 (2.0)	1 (1.7)	0 (0.0)
CVA (n)	4 (1.9)	1 (1.9)	1 (2.0)	1 (1.7)	1 (1.9)
<b>Laboratory measurements</b>					
R-SH ( $\mu\text{M}$ )	281.8 (34.0)	237.0 (17.1)	271.4 (7.0)	294.4 (6.3)	323 (15.4)
HbA1c (%)	7.6 (1.0)	7.7 (1.1)	7.5 (1.1)	7.6 (1.1)	7.6 (1.0)
Creatinine ( $\mu\text{mol/l}$ )	85.0 (77.0, 94.0)	84.0 (76.0, 93.5)	86.5 (77.8, 93.0)	84.0 (78.0, 94.0)	85.0 (78.0, 95.0)

(Continued)

**Table 1.** (Continued)

	All	Quartiles of R-SH			
	n = 216	172–257 µM n = 54	258–284 µM n = 51	285–304 µM n = 58	305–379 µM n = 53
eGFR (MDRD, ml/min/1.73m <sup>2</sup> )	126.1 (109.4, 141.7)	105.5 (91.7, 113.4)	105.9 (77.9, 127.9)	111.3 (86.8, 122.4)	103.3 (83.5, 131.0)
Total cholesterol (mmol/l)	4.6 (1.0)	4.6 (1.0)	4.7 (0.9)	4.4 (0.9)	4.8 (1.0)
HDL cholesterol (mmol/l)	1.5 (0.4)	1.6 (0.4)	1.4 (0.4)	1.5 (0.4)	1.6 (0.5)
Total cholesterol:HDL ratio	3.0 (1.0, 3.0)	3.1 (2.3, 3.9)	3.2 (2.6, 4.0)	2.8 (2.3, 3.4)	3.1 (2.6, 4.1)
LDL cholesterol (mmol/l)	2.6 (0.9)	2.5 (1.0)	2.7 (0.7)	2.4 (0.8)	2.7 (0.9)
Triglycerides (mmol/l)	0.9 (0.7, 1.4)	1.0 (0.7, 1.4)	1.0 (0.7, 1.4)	0.8 (0.7, 1.2)	1.0 (0.7, 1.7)
C-reactive protein (mg/l)	2.0 (1.0, 3.0)	2.0 (1.0, 4.3)	1.0 (1.0, 3.0)	2.0 (1.0, 4.0)	2.0 (1.0, 3.0)
NT-proBNP (pmol/l)	40.5 (16.2, 86.2)	73.0 (40.9, 41.7)	45.2 (17.0, 85.7)	29.0 (14.2, 66.9)	25.6 (8.7, 49.4)*

In the second-row, data are presented as number (%), mean (SD) or median (IQR).  
<sup>\*</sup>*p* < 0.05, <sup>§</sup>*n* = 29.  
 BMI, body mass index; CABG, coronary artery bypass grafting; CSII, continuous subcutaneous insulin infusion; CVA, cerebral vascular event; eGFR, estimated glomerular filtration rate; HbA1c, glycated haemoglobin A1c; HDL, high-density lipoprotein; LDL, low-density lipoprotein; MDI, multiple daily injections; MDRD, modification of diet in renal disease; MI, myocardial infarction; NA, not applicable; PAD, peripheral artery disease; PTCA, percutaneous transluminal coronary angioplasty; R-SH, total free thiol groups; TIA, transient ischemic attack.

and 42 patients experienced a total of 76 new macrovascular complications: 17 MI, 17 CVA, 17 PAD, 11 PTCA, 7 CABG, 4 AP and 3 TIA. A total of 92 (43%) patients developed a microvascular complication: neuropathy (*n* = 36), retinopathy (*n* = 55), microalbuminuria (*n* = 19) or macroalbuminuria (*n* = 6). There was no difference in baseline R-SH concentration at baseline between patients with or without the development of a microvascular or macrovascular complication during follow-up: 287.0 (258.0, 306.0) µM for persons without *versus* 281.0 (254.3, 303.3) µM for persons with a microvascular complication. For macrovascular complications these numbers were 270.0 (249.8, 300.2) µM for persons without *versus* 286.0 (261.5, 306.0) µM for persons with macrovascular complications.

Overall, the time until the first event was 11.1 [95% confidence interval (CI) 9.7–12.5] years. As presented in Figure 1 there were no differences between the quartiles of R-SH in the development of (both microvascular and macrovascular) complications during follow-up (*p* = 0.62).


Cox regression analysis demonstrated that R-SH was not a prognostic factor for the development

of both microvascular and macrovascular complications (see Table 2).

## Discussion

There were no relevant relationships at baseline between circulating R-SH concentrations and indices of T1DM management, including HbA1c, and cardiovascular risk factors in this cohort of 216 patients with T1DM. Although a negative correlation of R-SH with NT-proBNP concentrations was present at baseline, we did not observe this for HbA1c.

These findings are opposite to the significant relation between fasting blood glucose, HbA1c, C-reactive protein with disulphide/native thiol and disulphide/total thiol levels found by Ates and colleagues.<sup>16</sup> In their study a shift of dynamic thiol/disulphide homeostasis toward the disulphide form was present in T1DM (*n* = 38) patients as compared with healthy controls (*n* = 38). This shift may be secondary to increased ROS formation owing to (hyperglycaemia induced) inflammation in patients with T1DM. It could be hypothesized that the discrepancy between these and our findings is related to the degree of dysglycaemia; in the



**Figure 1.** Time between start of the study and occurrence of first complication for different quartiles of R-SH. The blue line represents the lowest [172–257  $\mu\text{M}$ ; median time to event 9.4 (95% confidence interval (CI) 7.2–11.6) years], the red [258–284  $\mu\text{M}$ ; 10.2 (95% CI 7.5–13.0) years] and green [285–304  $\mu\text{M}$ ; 12.3 (95% CI 9.4–15.1) years] lines the middle quartile and the orange [305–379  $\mu\text{M}$ ; 12.3 (95% CI 8.5–16.1) years] the highest quartile of free thiols (R-SH).

**Table 2.** Multivariable Cox regression analysis.

	HR (95% CI) for microvascular complications	HR (95% CI) for macrovascular complications
Age	1.025 (1.005, 1.046)	1.071 (1.040, 1.103)
Gender (male)	1.178 (0.762, 1.822)	2.945 (1.292, 6.710)
HbA1c	1.357 (1.100, 1.674)	1.066 (0.787, 1.444)
R-SH	0.999 (0.993, 1.005)	0.993 (0.984, 1.002)

Data are presented as hazard ratio (HR) [95% confidence interval (CI)]. Number of microvascular events: 92. Number of macrovascular events: 42.  
HbA1c, glycated haemoglobin A1c; R-SH, total free thiol groups.

study by Ates and colleagues mean HbA1c was 88mmol/mol as compared with 60mmol/mol in the present study. Another explanation is that thiol metabolism is mostly influenced by short-term oxidative stress caused by acute glycaemic derangements. This may indicate that R-SH merely provide a snapshot of someone's momentary ROS/antioxidant balance, which is then unlikely to correlate with a long-term indicator of glycaemic control such as HbA1c. Therefore, it would be of interest to investigate the within-person relationship between R-SH and within-day glycaemic variability (using continuous glucose measurements) and insulin administration as both instantaneously influences oxidative stress. Unfortunately, the present study lacks this information.

In the present study a negative correlation between R-SH and NT-proBNP, a marker of volume expansion and heart failure, was found. This observation is in agreement with previous studies among persons with heart failure and a renal transplant.<sup>9,14</sup> Of note, this correlation remained present after adjustment for potential confounders (see Table 3). Increased ROS are an important element of the pathophysiology of heart failure, and contributes to the development of myocardial and vascular dysfunction; therefore, it is conceivable that thiols are a reflection of these processes.<sup>22</sup> On the other hand, effects on volume expansion cannot be ruled out. Since T1DM is accompanied by an increased prevalence of heart failure,<sup>23</sup> the nature of the relation

**Table 3.** Multivariate linear regression analysis for R-SH.

	St. beta	95% CI	p
Gender (1 = male)	0.114	-2.2, 17.5	0.129
Creatinine ( $\mu\text{mol/l}$ )	0.051	-0.2, 0.5	0.489
Log BMI ( $\text{kg/m}^2$ )	0.017	-27.9, 35.2	0.807
Log NTproBNP	-0.410	-15.2, -7.6	<0.001
Presence of macrovascular complication (1 = yes)	0.038	-12.2, 21.6	0.588

BMI, body mass index; CI, confidence interval; R-SH, total free thiol groups; St. beta, Standardized beta (regression) coefficient

between R-SH and NT-proBNP needs to be further explored.

For the interpretation of the results of this study, several limitations should be taken into account including the limited sample size. Although this study currently has the largest population with the longest follow-up concerning R-SH in T1DM in the literature, we cannot rule out that the longitudinal analyses were underpowered. In addition, as only one baseline measurement of R-SH was used for these analyses and there was a lack of data regarding other parameters over time (including HbA1c), these analyses should be interpreted with caution. Furthermore, the lack of a nondiabetic reference population, amount of drop-out of the original cohort and the lack of other plasma antioxidant species such as ascorbate, uric acid and small-molecular-weight thiols and markers of inflammation should be mentioned.

In conclusion, no relevant relationships between R-SH and parameters of T1DM care were present in this study. Although the results of this small study needs confirmation, it could be hypothesized that R-SH is more appropriate as an instantaneous, rather than a long-term, read-out of oxidative stress in T1DM.

### Authors' note

Conceptualization: PvD, FW, HvG and HB. Methodology: PvD, SO, AP, AEA,FW, HvG and HB; Software, PvD. Validation: PvD, AP and HvG. Formal Analysis: PvD. Investigation: PvD and FW. Resources: HB, FW, PvD and HvG; Data Curation: PvD; Writing – Original Draft Preparation: PvD. Writing – Review & Editing: PvD, FW, SO, AEA, AP, MM, HB, HvG. Visualization: PvD. Supervision: PvD, HvG and

HB. Project Administration: PvD. Funding Acquisition: HB, FW, PvD and HvG.

### Acknowledgements

The authors want to thank Marian Bulthuis (University Medical Centre, Department of Pathology and Medical Biology, Groningen, the Netherlands) and the Department of Clinical Chemistry of Isala (Zwolle, The Netherlands).

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by grants from the Isala Innovatie and Wetenschapsfonds (grant number INNO1717) and the Zwols Wetenschapsfonds Isala Klinieken. Both the Isala Innovatie and Wetenschapsfonds and Zwols Wetenschapsfonds Isala Klinieken had no role in the study design, data collection, analysis, interpretation, or in the writing of the report.

### Conflict of interest statement

The authors declare that they have no conflict of interest.

### ORCID iD

Peter R. van Dijk  <https://orcid.org/0000-0002-9702-6551>

### References

1. Sies H. Oxidative stress: a concept in redox biology and medicine. *Redox Biol* 2015; 4: 180–183.
2. Rochette L, Zeller M, Cottin Y, *et al.* Diabetes, oxidative stress and therapeutic strategies. *Biochim Biophys Acta* 2014; 1840: 2709–2729.



3. Asmat U, Abad K and Ismail K. Diabetes mellitus and oxidative stress-A concise review. *Saudi Pharm J* 2016; 24: 547–553.
4. Brownlee M. The pathobiology of diabetic complications: a unifying mechanism. *Diabetes* 2005; 54: 1615–1625.
5. Ferranti SD de, Boer IH de, Fonseca V, *et al.* Type 1 diabetes mellitus and cardiovascular disease: a scientific statement from the American heart association and American diabetes association. *Diabetes Care* 2014; 37: 2843–2863.
6. Sen CK and Packer L. Thiol homeostasis and supplements in physical exercise. *Am J Clin Nutr* 2000; 72: 653S–669S.
7. Turell L, Radi R and Alvarez B. The thiol pool in human plasma: the central contribution of albumin to redox processes. *Free Radic Biol Med* 2013; 65: 244–253.
8. Gupta S, Gambhir JK, Kalra O, *et al.* Association of biomarkers of inflammation and oxidative stress with the risk of chronic kidney disease in type 2 diabetes mellitus in North Indian population. *J Diabetes Complications* 2013; 27: 548–552.
9. Koning AM, Meijers WC, Pasch A, *et al.* Serum free thiols in chronic heart failure. *Pharmacol Res* 2016; 111: 452–458.
10. Dröge W. Aging-related changes in the thiol/disulfide redox state: implications for the use of thiol antioxidants. *Exp Gerontol* 2002; 37: 1333–1345.
11. Atkuri KR, Mantovani JJ, Herzenberg LA, *et al.* N-Acetylcysteine—a safe antidote for cysteine/glutathione deficiency. *Curr Opin Pharmacol* 2007; 7: 355–359.
12. Banne AF, Amiri A and Pero RW. Reduced level of serum thiols in patients with a diagnosis of active disease. *J Anti-Aging Med* 2003; 6: 327–334.
13. Kundi H, Ates I, Kiziltunc E, *et al.* A novel oxidative stress marker in acute myocardial infarction; thiol/disulphide homeostasis. *Am J Emerg Med* 2015; 33: 1567–1571.
14. Frenay AS, de Borst MH, Bachtler M, *et al.* Serum free sulfhydryl status is associated with patient and graft survival in renal transplant recipients. *Free Radic Biol Med* 2016; 99: 345–351.
15. Bourgonje AR, Gabriëls RY, de Borst MH, *et al.* Serum free thiols are superior to fecal calprotectin in reflecting endoscopic disease activity in inflammatory bowel disease. *Antioxid Basel Switz* 2019; 8: 351.
16. Ates I, Kaplan M, Yuksel M, *et al.* Determination of thiol/disulphide homeostasis in type 1 diabetes mellitus and the factors associated with thiol oxidation. *Endocrine* 2016; 51: 47–51.
17. Hart HE, Bilo HJ, Redekop WK, *et al.* Quality of life of patients with type I diabetes mellitus. *Qual Life Res* 2003; 12: 1089–1097.
18. van Dijk PR, Logtenberg SJ, Groenier KH, *et al.* Fifteen-year follow-up of quality of life in type 1 diabetes mellitus. *World J Diabetes* 2014; 5: 569–576.
19. Ellman GL. Tissue sulfhydryl groups. *Arch Biochem Biophys* 1959; 82: 70–77.
20. Hu ML, Louie S, Cross CE, *et al.* Antioxidant protection against hypochlorous acid in human plasma. *J Lab Clin Med* 1993; 121: 257–262.
21. International Diabetes Federation Guideline Development Group. Global guideline for type 2 diabetes. *Diabetes Res Clin Pract* 2014; 104: 1–52.
22. Münzel T, Gori T, Keaney JF, *et al.* Pathophysiological role of oxidative stress in systolic and diastolic heart failure and its therapeutic implications. *Eur Heart J* 2015; 36: 2555–2564.
23. Rosengren A, Vestberg D, Svensson AM, *et al.* Long-term excess risk of heart failure in people with type 1 diabetes: a prospective case-control study. *Lancet Diabetes Endocrinol* 2015; 3: 876–885.