



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XX International Grassland Congress

---

## QTL Analysis and Trait Dissection in Ryegrass (*Lolium* Spp.)

T. Yamada

*National Agricultural Research Center for Hokkaido Region, Japan*

J. W. Forster

*La Trobe University, Australia*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium5/35>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## QTL analysis and trait dissection in ryegrasses (*Lolium* spp.)

T. Yamada<sup>1</sup> and J.W. Forster<sup>2</sup>

<sup>1</sup>National Agricultural Research Center for Hokkaido Region, Sapporo 062-8555, Japan,

Email: Toshihiko.Yamada@affrc.go.jp

<sup>2</sup>Primary Industries Research Victoria, Plant Biotechnology Centre, La Trobe University, Bundoora, Victoria 3086, Australia and Molecular Plant Breeding Cooperative Research Centre, Australia

### Key points

1. Molecular marker-based genetic analysis permits the dissection of complex phenotypes through resolution of the locations of pleiotropic and interacting genetic factors.
2. Several QTLs for agronomically important characters such as flowering time, winter hardiness and forage quality have been identified in perennial ryegrass by molecular marker-based map analysis.
3. Some QTLs were putatively orthologous to those for equivalent traits in cereals.
4. The identification of co-location between QTLs and functionally-associated genetic markers is critical for the future implementation of marker-assisted selection programs.

**Keywords:** comparative genomics, flowering time, forage quality, QTL analysis, winter hardiness

### Introduction

The majority of traits of interest to pasture grass breeders, such as dry matter yield, forage quality and environmental stress tolerance, show continuous phenotypic variation and are controlled by a variable number of quantitative trait loci (QTL). Substantial advances have been made in the genetic improvement of plant populations through artificial selection of quantitative traits. Most of this selection has been on the basis of observable phenotype, without knowledge of the genetic architecture of the selected characteristics. In major crop species, the development of high-resolution genetic maps has made it possible to identify the chromosomal regions, or in some instances, the individual sequence variants that are responsible for trait variation. There have been relatively few reports to date of QTL analysis for agronomic traits in forage grasses, due to the absence of a sufficiently well developed genetic map. An enhanced molecular marker-based genetic linkage map of perennial ryegrass (*Lolium perenne* L.) has recently been constructed through the activities of the International *Lolium* Genome Initiative (ILGI), using the p150/112 one-way pseudo-testcross reference population (Jones *et al.*, 2002a,b). Two genetic mapping populations of perennial ryegrass have been independently developed as successors to the p150/112 population, and have been aligned to the reference map using common markers (Armstead *et al.*, 2002, 2004; Faville *et al.*, 2005). These genetic maps contain functionally-associated molecular marker information through the inclusion of gene-associated cleaved amplified polymorphic sequences (CAPS) markers, and both restriction fragment length polymorphism (RFLP) and simple sequence repeat (SSR) markers from expressed sequence tags (ESTs), respectively. In addition, high-density molecular marker-based genetic maps have also recently been constructed for other species of *Lolium* and *Festuca* (Alm *et al.*, 2003; Inoue *et al.*, 2004a; Warnke *et al.*, 2004; Saha *et al.*, 2005). Genetic markers and maps are consequently available for detailed dissection of complex phenotypes to resolve the locations of pleiotropic and interacting genetic factors. In this chapter, recent results from QTL analysis of the p150/112 reference population and another populations of perennial ryegrass are reviewed, and are related to comparative genomics studies with other Poaceae species.

## Flowering time

The timing of flowering during the year is an important adaptive character that impacts on yield and quality in crop and pasture species. Analysis of the molecular basis of several QTLs in *Arabidopsis*, rice, maize and tomato has revealed that in diverse species, orthologous genetic networks can control related complex phenotypes (Paran & Zamir, 2003). For example, QTL cloning studies has revealed that the major flowering-time QTL *ED1* (early day-length insensitive) of the long-day flowering plant *Arabidopsis thaliana* is a novel allele of the blue-light photoreceptor cryptochrome-2 (*CRY2*) (El-Din El-Assal *et al.*, 2001). This allele increased protein stability as a result of a single amino acid substitution that led to early flowering in short days. The genetic control of flowering time control in *Arabidopsis* has been well characterised through the interaction of photoperiod, vernalisation, autonomous and gibberellic acid-dependent pathways (Mouradov *et al.*, 2002; Hayama & Coupland, 2003; Henderson & Dean, 2004; Sung & Amasino, 2004). In rice, a short-day flowering plant, QTL analysis for heading date has been performed using several types of progeny derived from a single cross combination between *Oryza sativa* ssp. *japonica* and *O. sativa* ssp. *indica* cultivars and identified at least 14 QTLs controlling flowering time (Yano *et al.*, 2001). Three heading QTLs (*Hd1*, *Hd3* and *Hd6*) were mapped at high resolution using near isogenic lines (NILs) and were isolated by a map-based cloning approach (Yano *et al.*, 2000; Kojima *et al.*, 2002; Takahashi *et al.*, 2001). *Hd1* was shown encode a homologue of the *Arabidopsis* photoperiod pathway *CONSTANS (CO)* gene (Yano *et al.*, 2000). *Hd3a* was shown to be related to the *Arabidopsis* *FLOWERING LOCUS T (FT)* gene, while *Hd6* was shown to encode the  $\alpha$ -subunit of casein protein kinase 2 (*CK2 $\alpha$* ). Gene interactions in the photoperiod induction pathways of *Arabidopsis* and rice were compared (Izawa *et al.*, 2003).

In cereals, genes that regulate the timing of flowering can be divided three categories based on their interaction with environmental signals: vernalisation response genes (*Vrn*) that regulate flowering using low temperature; photoperiod response genes (*Ppd*) that regulate flowering using day length; and 'earliness' factors that appear to be largely independent of these cues (Laurie, 1997). Conserved genomic locations for genes involved in processes such as vernalisation and photoperiodic induction have been identified between species by comparative genetic studies (Dubcovsky *et al.*, 1998; Laurie *et al.*, 2004). A detailed physical and genetic map of the *Vrn-A<sup>m</sup>1 (Vrn1)* region was constructed for the diploid wheat, *Triticum monococcum* L. and found to be colinear with the corresponding region of rice chromosome 3 (Yan *et al.*, 2003). A candidate gene for *Vrn1* was isolated by positional means, and was identified as a relative of the *Arabidopsis* *apetala-1* gene class (*API*). Allelic variation between spring and winter-type growth habit wheats at the *API* gene was observed only in the promoter region, suggesting that variation of gene expression was the causal factor for differences between these two varietal groups. A second vernalisation gene, designated *Vrn-A<sup>m</sup>2*, was assigned to the distal region of chromosome 5A<sup>m</sup>L within a segment translocated from homoeologous group 4 (Dubcovsky *et al.*, 1998). Yan *et al.* (2004) reported the positional cloning of the *Vrn2* gene, which encodes a dominant repressor of flowering down-regulated by vernalisation. Loss of function at *Vrn2*, whether by natural point mutation or deletion, resulted in lines with spring-type growth habit, which do not require vernalisation to flower.

A single QTL for heading date was observed on linkage group (LG) 4 in the p115/120 reference family of perennial ryegrass (Yamada *et al.*, 2004). However, a number of QTL positions for heading date have been reported from the analysis of single mapping populations in other Poaceae species (Hayes *et al.*, 1993; Laurie *et al.*, 1995; Bezant *et al.*, 1996; Börner *et al.*, 2002). The number of QTLs and their relative importance may vary according to the

origin of the genotypes used to construct mapping families. Studies on geographical populations of *Lolium* species covering the climatic range from the Mediterranean region to northern and central Europe revealed a regular cline in flowering responses to temperature and photoperiod (Cooper, 1960). The heterozygous parent of the p150/112 mapping population was derived from a cross between eastern European (Romanian), southern European (north Italian ecotypes) and northern European ('Melle' or 'S23') genotypes, and might be expected to represent a variety of response genes. In rice, the major heading date QTL *Hd6*, which is associated with inhibition of flowering under long day conditions, is located on chromosome 3 (Yamamoto *et al.*, 2000) and encodes an  $\alpha$ -subunit of CK2 (Takahashi *et al.*, 2001). Comparative genetic mapping studies between rice and wheat based on the colinearity of four common RFLP markers (Kato *et al.*, 1999) have revealed that the rice *Hd6* locus region on chromosome 3 is syntenic with the *VrnA1* region on chromosome 5AL. RFLP-based mapping of the wheat CK2 $\alpha$  gene (*tck2a*) probe detected a genetic locus closely linked (by 1.1 cM) to *VrnA1* (Kato *et al.*, 2002). Two putative CK2 $\alpha$  genes (*Lpck2a-1* and *Lpck2a-2*) genes have been isolated from a cDNA library constructed with mRNA isolated from cold-acclimated crown tissues of *Lolium perenne* using sequence information derived from the *tck2a* gene. The *Lpck2a-1* CAPS marker was assigned to LG 4 of the p115/120 reference family near to the location of the QTL for heading date, while the *Lpck2a-2* CAPS marker was assigned to LG 2 (Shinozuka *et al.*, submitted). The location of the *Lpck2a-1* locus supports the inference of conserved synteny between perennial ryegrass LG 4, the Triticeae homoeologous group 5L chromosomes and the corresponding segment of rice chromosome 3 (Yamada *et al.*, 2004).

A mapping population consisting of 184 F<sub>2</sub> genotypes from a cross between a genotype from the perennial ryegrass synthetic variety 'Veyo' and a genotype from the perennial ryegrass ecotype 'Falster' was measured for vernalisation response as days to heading under artificially controlled condition (Jensen *et al.*, 2005). In total, five QTLs were identified on LGs 2, 4, 6 and 7. A CAPS marker derived from the putative orthologue of the *Triticum monococcum* *VRN1* gene co-located with a major QTL on LG 4 for vernalisation response (Jensen *et al.*, 2005). This data further confirms the presence of flowering time gene orthologues and corresponding QTLs on LG 4.

Heading date QTLs were also identified in one of the second generation reference populations, derived from self-pollination of an F<sub>1</sub> hybrid obtained by crossing individuals from partially inbred lines developed from the two agronomically contrasting cultivars 'Aurora' and 'Perma' (Turner *et al.*, 2001). Genetic mapping of the F<sub>2</sub> (Aurora x Perma) population identified seven linkage groups with a total map length of 628 cM (Armstead *et al.*, 2004), extending the studies of Armstead *et al.* (2002) and consistent with the ILGI reference map (Jones *et al.*, 2002). A major QTL accounting for up to 70% of the variance was identified on LG 7, along with additional small QTLs on LGs 2 and 4 (Armstead *et al.*, 2004). The genomic region associated with the major QTL on LG 7 shows a high degree of conserved synteny with the *Hd3* region of rice chromosome 6.

Two annual ryegrass plants from the cultivar 'Floregon' were crossed with two perennial ryegrass plants from the cultivar 'Manhattan'. From the resultant F<sub>1</sub> populations, two random plants were chosen and crossed to develop a pseudo-F<sub>2</sub> mapping family ( $\psi$ F<sub>2</sub>[MFA-4 x MFB-2]). A total of 235 amplified fragment length polymorphism (AFLP) markers, 81 random amplified polymorphic DNA (RAPD) markers, 16 grass comparative anchor probe RFLPs, 106 SSR markers, 2 isoenzyme loci and 2 morphological characteristics, 8-h flowering and seedling root fluorescence were used to construct a male map 537cM in length and a female

map 712 cM in length, each with 7 LGs (Warnke *et al.*, 2004). Two major QTLs influencing photoperiodic control of flowering were identified in locations syntenic with flowering control regions of the wheat and barley genomes (Warnke *et al.*, 2003a,b).

### Winter hardiness

Winter hardiness is the outcome of a number of interacting factors that may include vernalisation requirement, photoperiod response, low-temperature tolerance and resistance to snow moulds. An understanding of the genetic basis of these component traits permits more efficient selection based on closely linked molecular marker loci. In the Triticeae cereals, QTL analysis has identified a limited number of conserved genome regions as responsible for the winter hardiness character. The most consistently identified region, on homoeologous group 5 chromosomes, contains QTLs for vernalisation response, low temperature tolerance and photoperiod sensitivity (Pan *et al.*, 1994; Cattivelli *et al.*, 2002). These QTL effects have been described as the effects of single loci. Low-temperature tolerance loci on chromosomes 5A, 5B, and 5D of wheat have been given the locus designation *Fr-A1*, *Fr-B1* and *Fr-D1*, respectively (Sutka & Snape, 1989; Snape *et al.*, 1997; Toth *et al.*, 2003). The vernalisation loci have been assigned a *Vrn* prefix, and orthologues of the *T. monococcum Vrn1* locus map to the homoeologous 5 chromosomes (Cattivelli *et al.*, 2002). The *Vrn* gene nomenclature was standardised and map locations were further refined by Dubcovsky *et al.* (1998). RFLP analysis demonstrated that vernalisation requirement and frost resistance are controlled by two different, but tightly linked loci (*Vrn-A1* and *Fr-A1*) on chromosome 5A of wheat (Galiba *et al.*, 1995; Sutka *et al.*, 1999). Because of the observed large effect on frost resistance, molecular marker-assisted selection for the *Vrn-A1-Fr-A1* chromosomal region has been proposed as a method for improvement of cold hardiness in wheat cultivars (Storlie *et al.*, 1998). Wheat NILs with different vernalisation alleles have been evaluated for cold hardiness in order to assess the viability of this strategy (Koemel *et al.*, 2004). Close genetic linkage between the major genes influencing winter hardiness and genes regulating cold-induced sugar production was also observed in wheat (Galiba *et al.*, 1997). In addition, QTLs controlling traits associated with winter hardiness, such as field survival and crown fructan content, were mapped to the long arm of chromosome 5H in a cross between 'winter' x 'spring' barley varieties (Hayes *et al.*, 1993; Pan *et al.*, 1994). Recently, comprehensive measurements of low temperature tolerance and vernalisation requirement were used for analysis of a new 'winter' x 'spring' barley population, and a QTL for accumulation of proteins encoded by COR (COld Regulated) genes on chromosome 5H (*Cor14b*, *tmc-ap3*) was coincident with a QTL for low temperature tolerance (Francia *et al.*, 2004). C-repeat binding factor (CBF) family genes were also mapped in this region (Francia *et al.*, 2004). In *A. thaliana* the transcription factors encoded by CBF family genes have been shown to be key determinants of low temperature tolerance (Thomashow, 1999; Thomashow *et al.*, 2001).

No significant QTLs for winter survival in the field were identified in the reference map of p150/112 (Yamada *et al.*, 2004). However, a QTL for electrical conductivity corresponding to frost tolerance (Dexter *et al.*, 1930, 1932) was located close to a heading date QTL in a region is likely to show conserved synteny with chromosomal regions associated with both winter hardiness and flowering time variation in cereals, as described above.

The F<sub>2</sub> (Aurora x Perma) genetic map population (RASP) was also used to identify QTLs for traits relating winter hardiness, as well as sugar content. Snow mould-resistant varieties accumulate higher levels of fructan and metabolise them at slower rates compared to susceptible varieties (Yoshida *et al.*, 1998). Many of the snow mould fungi, such as *Typhula*

spp., *Microdochium nivale* and *Sclerotinia borealis*, can co-infect on single plants, and their interactions may obscure the respective effects on plant survival (Matsumoto & Araki, 1982; Matsumoto *et al.*, 1982). The use of fungicides with a limited spectrum of activity may clarify these specific effects. *Typhula* snow moulds such as *T. ishikariensis* and *T. incarnata* generally occur in the deep snow environment of the western region of Hokkaido, Japan, including Sapporo. In this environment, control of *S. borealis* and *M. nivale* infections with the fungicide iminoctadine-triacetate is an effective method for evaluation of resistance to *Typhula* snow moulds (Takai *et al.*, 2004). Scores of winter survival were measured in the F<sub>2</sub> (Aurora x Perma) population using this control regime, and QTLs for this trait were identified on LGs 2, 4, 6 and 7. Fructan content was also measured by high performance liquid chromatography (HPLC) using crown tissues from plants grown outdoors in December. QTLs for content of high molecular fructan with more than eight degrees of polymerization (DP) were observed on LGs 1, 2 and 4. QTLs for winter survival in LGs 2 and 4 are close to coincident with the QTLs for high molecular weight fructan content.

Two major QTLs for freezing tolerance (*Fr*) and 4 QTL for winter survival have been identified in the closely related pasture grass species meadow fescue (*Festuca pratensis* Huds.) (Rognli *et al.*, 2002). Comparative mapping with heterologous wheat anchor probes indicated that *Fr4\_1* on LG4 of *F. pratensis* was orthologous to the frost-tolerance loci *Fr1* and *Fr2* in wheat. The QTLs for winter survival, by contrast, were located on LGs 1, 2, 5, and 6 (Rognli *et al.*, 2002).

Fructosyltransferase genes involved in fructan biosynthesis such as 1-SST, 1-FFT and 6G-FFT were isolated from perennial ryegrass and characterized by heterologous expression in the *Pichia pastoris* system (Hisano *et al.*, in preparation). *Lp1-SST* and *Lp1-FFT* mapped to the upper region of LG 7 in the F<sub>2</sub> (Aurora x Perma) genetic map, but failed to show coincidence with any fructan content QTLs. The *Lp1-SST* gene (Chalmers *et al.*, 2003) was also assigned to the equivalent region of LG 7 as a single nucleotide polymorphism (SNP) locus in the F<sub>1</sub> (NA<sub>6</sub> x AU<sub>6</sub>) second generation reference family (Faville *et al.*, 2005). However, *Lp6G-FFT* mapped to LG 3 close to a QTL for low-molecular weight fructan content (Hisano *et al.*, in preparation). As previously described, evidence from comparative genome studies suggests that the upper part of LG4 in perennial ryegrass may contain a region of conserved synteny with the long arms of the Triticeae homoeologous group 5 chromosome. It is possible that allelic variation in regulatory genes such as those for the CBF transcription factor family may contribute to the QTLs for winter survival and fructan content observed on LG 4.

## Herbage quality

Quality is the most important of all agronomic traits for pastures due to the nutritive requirements of grass-fed livestock. The genetic control of nutritive value parameters in pasture species has been reviewed (e.g. Casler, 2001), and genetic variation for specific traits has been established. Digestibility is generally considered to be the most important temperate grass nutritive value trait for either live-weight gain (Wheeler & Corbett, 1989) or dairy production (Smith *et al.*, 1997). Deliberate attempts to improve dry matter digestibility (DMD) in forage crop species have led to rates of genetic gain in the range of 1 - 4.7 % per annum as a proportion of the initial population means (Casler, 2001). Progress in simultaneous improvement of yield and DMD in forage grasses has, however, been variable (Wilkins & Humphreys, 2003).

Forage quality may be directly evaluated by feeding trials using animals, but this approach is costly, laborious and limited for small quantities of herbage from breeding experiments. Indirect methods of assessment include *in vitro* digestibility with rumen liquor (Menke *et al.*, 1979; Tilly and Terry, 1963), enzymatic digestion (De Boever *et al.*, 1986) and chemical analysis of cellular components (van Soest, 1963). The development of near infra-red reflectance spectroscopy (NIRS) analysis for prediction of forage quality has facilitated rapid and non-destructive evaluation of samples from plant breeding programs. NIRS has been used to develop calibrations to predict a wide range of forage quality traits (Marten *et al.*, 1984; Smith and Flinn, 1991) including crude protein (CP) content, estimated *in vivo* dry matter digestibility (IVVDMD), neutral detergent fibre (NDF) content (Smith & Flinn, 1991) and water-soluble carbohydrate (WSC) content (Smith & Kearney, 2000) in perennial ryegrass. NIRS estimates of DMD and related nutritive value traits have been reported in a range of forage systems (e.g. Carpenter & Casler, 1990; Hopkins *et al.*, 1995; Smith *et al.*, 2004).

Lübbstedt *et al.* (1997, 1998) published the first QTL analysis devoted to forage quality in maize. QTLs for cell-wall digestibility and lignification traits in maize were also investigated in two recombinant inbred lines (RIL) progeny by Méchin *et al.* (2001). Cardinal *et al.* (2003) detected 65 QTLs related to fiber and lignin content in maize. The best options for breeding of grasses for improved digestibility was assessed based on a search for genome locations involved in forage quality traits through QTL analysis (Ralph *et al.*, 2004).

Ground herbage samples from genotypes of the p150/112 population were measured for quality traits such as CP, IVVDMD, NDF, estimated metabolisable energy (EstME) and WSC by NIRS analysis (Cogan *et al.*, 2005). A total of 42 QTLs was observed in six different sampling experiments varying by developmental stage (anthesis or vegetative growth), location or year. Coincident QTLs were detected on LGs 3, 5 and 7. The region on LG 3 was associated with variation for all measured traits across various experimental datasets. The region on LG 7 was associated with variation for all traits except CP, and is located in the vicinity of the lignin biosynthesis gene loci *xlpomt1* (caffeic acid-*O*-methyltransferase), *xlpccr1* (cinnamoyl CoA-reductase) and *xlpssrcad2.1* (cinnamyl alcohol dehydrogenase).

WSC provide the most available source of energy for grazing ruminants. In the F<sub>2</sub> (Aurora x Perma) population (RASP), high molecular fructan constituted the major part of the WSC pool was analyzed samples of spring and autumn in tiller bases and leaves with replication of data over years (*ie* collect one replicate each year for several years) (Turner *et al.*, unpublished). Correlation between traits did not always lead to corresponding cluster of QTL and some traits have no reproducible QTL. Tiller base QTL were identified on linkage groups 1 and 5 and leaf QTL on linkage groups 2 and 6, in regions that had previously been identified as important in analyses of single replicates (Humphreys *et al.*, 2003).

Improvements of herbage quality may also be obtained by alteration of the content and ratios of minerals in grasses, to prevent metabolic disorders. Grass tetany (hypomagnesaemia) is caused by low levels of magnesium in the blood of cattle or sheep. Varieties of Italian ryegrass and tall fescue with markedly levels of magnesium have proved to be very effective in maintaining levels of blood magnesium in grazing sheep (Moseley & Baker, 1991) and cattle (Crawford *et al.*, 1998). Milk fever, caused by low blood calcium, produces animal welfare and production problems that could be addressed by reducing potassium content of forage without reducing calcium and magnesium concentrations (Sanchez *et al.*, 1994). Variation in mineral content in grasses may be strongly influenced by genetic factors and is amenable to QTL analysis. Herbage samples of the p150/112 population from four sampling

experiment were analyzed for mineral content (aluminum, calcium, cobalt, copper, iron, magnesium, manganese, molybdenum, nickel, phosphorus, potassium, sodium, sulfur and zinc) by inductively-coupled plasma mass spectroscopy (ICP-MS) and a total of 45 QTLs were identified (Cogan *et al.*, in preparation). QTL clusters were observed on LGs 1, 2, 4 and 5. QTLs for the important trait for control of grass tetany, magnesium content were detected on LGs 2 and 5. Field herbage samples from the F<sub>1</sub> (NA<sub>6</sub> x AU<sub>6</sub>) population were also analysed for mineral content by ICP-MS. A total of 14 QTLs were identified on the NA<sub>6</sub> map, and 9QTLs were identified on the AU<sub>6</sub> genetic map. A number of clustered QTL locations showed coincidence between the two different populations.

### **Morphological traits and other agronomical traits**

QTLs were detected for morphological traits such as plant height, tiller size, leaf length, leaf width, fresh weight at harvest, plant type, spikelet number per spike and spike length using the p150/112 genetic map (Yamada *et al.*, 2004). A number of traits were significantly correlated, and coincident QTL locations were identified. For example, coincident QTLs for plant height, tiller size and leaf length were identified on LG 3. The rice *SD1* semi-dwarfing gene, that launched the 'green revolution', encodes a gibberellin biosynthetic enzyme (*GA20ox*), and was assigned to the long arm of rice chromosome 1 (Sasaki *et al.*, 2002). A CAPS marker developed for the perennial ryegrass orthologous of the *GA20ox* gene was mapped to LG 3 close to the plant height QTL, in a region of conserved synteny with rice chromosome 1 (Kobayashi *et al.*, unpublished). This finding provides further evidence for the utility of the candidate gene-based marker approach.

In Italian ryegrass (*Lolium multiflorum* Lam.), a total of 17 QTLs for six traits related to lodging resistance and heading date were detected by single interval mapping (SIM), while 33 independent QTLs from the male and female parents were detected by composite interval mapping (CIM) (Inoue *et al.*, 2004b). QTLs for plant height were located on LG 1, and for heading date on LGs 4 and 7, potentially in conserved regions with those identified in perennial ryegrass.

### **Conclusions**

This review has demonstrated that although the existing QTL information from in forage grasses is relatively underdeveloped compared to other major crops, the recent establishment of detailed molecular genetic maps is rapidly stimulating QTL analysis and trait dissection. The current genetic maps are largely populated by anonymous genetic markers, with limited diagnostic value. The next generation of molecular genetic markers for forage grasses will be derived from expressed sequences, with an emphasis on functionally-defined genes associated with biochemical and physiological processes that are likely to be correlated with target phenotypic traits (Forster *et al.*, 2004; Faville *et al.*, 2005). Comparative genomics with other Poaceae species such as rice, wheat, barley and maize will support the development of such high value molecular markers through orthologous QTL detection and co-location of candidate genes. Accuracy in phenotypic assessment will be essential for precise QTL detection, and may in future be amenable to automated high-throughput analysis, as for genotyping.



## Acknowledgements

The unpublished research from NARCH described in this paper was carried out by Dr. S. Kobayashi, Mrs. H. Hisano and H. Shinozuka. QTL research was carried out in collaboration with Dr. K. F. Smith, Dr. N.O.I. Cogan and Prof. G. S. Spangenberg of Primary Industries Research Victoria, Australia and Prof. M.O. Humphreys, Drs. I.P. Armstead and L. Turner of Institute of Grassland and Environmental Research, UK. This work was supported in part by Grants-in-Aid for Scientific Research (No. 14360160) from the Ministry of Education, Science, Sports and Culture, Japan.

## References

- Alm, V., C. Fang, C.S. Busso, K.M. Devos, K. Vollan, Z. Grieg & O.A. Rognli (2003). A linkage map of meadow fescue (*Festuca pratensis* Huds.) and comparative mapping with other Poaceae species. *Theoretical and Applied Genetics*, 108, 25-40.
- Armstead, I.P., L.B. Turner, I.P. King, A.J. Cairns & M.O. Humphreys (2002). Comparison and integration of genetic maps generated from F<sub>2</sub> and BC<sub>1</sub>-type mapping populations in perennial ryegrass. *Plant Breeding*, 121, 501-507.
- Armstead, I.P., L.B. Turner, M. Farrell, L. Sköt, P. Gomez, T. Montoya, I.S. Donnison, I.P. King & M.O. Humphreys (2004). Synteny between a major heading-date QTL in perennial ryegrass (*Lolium perenne* L.) and the *Hd3* heading-date locus in rice. *Theoretical and Applied Genetics*, 108, 822-828.
- Bezant, J., D. Laurie, N. Pratchett, J. Chojecki & M. Kearsey (1996). Marker regression mapping of QTL controlling flowering time and plant height in a spring barley (*Hordeum vulgare* L.) cross. *Heredity*, 77, 64-73.
- Börner, A., G.H. Buck-Sorlin, P.M. Hayes, S. Malyshev & V. Korzun (2002). Molecular mapping of major genes and quantitative trait loci determining flowering time in response to photoperiod in barley. *Plant Breeding*, 121, 129-132.
- Cardinal, A.J., M. Lee & K.J. Moore (2003). Genetic mapping and analysis of quantitative trait loci affecting fiber and lignin content in maize. *Theoretical and Applied Genetics*, 106, 866-874.
- Carpenter, J.A. & M.D. Casler (1990). Divergent phenotypic selection response in smooth bromegrass for forage yield and nutritive value. *Crop Science*, 30, 17-22.
- Casler, M.D. (2001). Breeding forage crops for increased nutritive value. *Advances in Agronomy*, 71, 51-107.
- Cattivelli, L., P. Baldi, C. Crosatti, N. Di Fonzo, P. Faccioli, M. Grossi, A.M. Mastrangelo, N. Pecchioni & A.M. Stanca (2002). Chromosome regions and stress-related sequences involved in resistance to abiotic stress in *Triticeae*. *Plant Molecular Biology*, 48, 649-665.
- Chalmers, J., X. Johnson, A. Lidgett & G.C. Spangenberg (2003). Isolation and characterisation of a sucrose:sucrose 1-fructosyltransferase gene from perennial ryegrass (*Lolium perenne* L.). *Journal of Plant Physiology*, 160, 1385-1391.
- Cogan, N.O.I., K.F. Smith, T. Yamada, M.G. Francki, A.C. Vecchies, E.S. Jones, G.C. Spangenberg & J.W. Forster (2005). QTL analysis and comparative genomics of herbage quality traits in perennial ryegrass (*Lolium perenne* L.). *Theoretical and Applied Genetics*, available on-line.
- Cooper, J.P. (1960). Short-day and low-temperature induction in *Lolium*. *Annals Botany*, 24, 232-246.
- Crawford, R.J., M.D. Massie, D.A. Sleper & H.F. Mayland (1998). Use of an experimental high-magnesium tall fescue to reduce grass tetany in cattle. *Journal of Production Agriculture*, 11, 491-496.
- De Boever, J.L., F.X. Cottyn, F.W. Wainman & J.M. Vanacker (1986). The use of an enzymatic technique to predict digestibility, metabolisable and net energy of compound feedstuffs for ruminants. *Animal Feed Science Technology*, 14, 203-214.
- Dexter, S.T., W.E. Tottingham & L.F. Graber (1930). Preliminary results in measuring the hardiness of plants. *Plant Physiology*, 5, 215-223.
- Dexter, S.T., W.E. Tottingham & L.F. Graber (1932). Investigations of the hardiness of plants by measurement of electrical conductivity. *Plant Physiology*, 7, 63-78.
- Dubcovsky, J., D. Lijavetzky, L. Appendino & G. Tranquilli (1998). Comparative RFLP mapping of *Triticum monococcum* genes controlling vernalisation requirement. *Theoretical and Applied Genetics*, 97, 968-975.
- El-Din El-Assal, S., C. Alonso-Blanco, A.J. Peeters, V. Raz & M. Koornneef (2001). A QTL for flowering time in *Arabidopsis* reveals a novel allele of *CRY2*. *Nature Genetics*, 29, 435-440.

- Faville, M.J., A.C. Vecchies, M. Schreiber, M.C. Drayton, L.J. Hughes, E.S. Jones, K.M. Guthridge, K.F. Smith, T. Sawbridge, G.C. Spangenberg, G.T. Bryan & J.W. Forster (2005). Functionally associated molecular genetics marker map construction in perennial ryegrass (*Lolium perenne* L.). *Theoretical and Applied Genetics*, available on-line.
- Forster, J.W., E.S. Jones, J. Batley & K.F. Smith (2004). Molecular marker-based genetic analysis of pasture and turf grasses. A. Hopkin *et al.* (eds.) *Molecular Breeding of Forage and Turf*, 197-238, Kluwer Academic Publishers, the Netherlands.
- Francia, E., F. Rizza, L. Cattivelli, A.M. Stanca, G. Galiba, B. Toth, P.M. Hayes, J.S. Skinner & N. Pecchioni (2004). Two loci on chromosome 5H determine low-temperature tolerance in a 'Nure' (winter) x 'Tremois' (spring) barley map. *Theoretical and Applied Genetics*, 108, 670-680.
- Galiba, G., S.A. Quarrie, J. Sutka, A. Morgounov & J.W. Snape (1995). RFLP mapping of the vernalization (*Vrn-A1*) and frost resistance (*Fr1*) genes on chromosome 5A of wheat. *Theoretical and Applied Genetics*, 90, 1174-1179.
- Galiba, G., I. Kerepesi, J.W. Snape & J. Sutka (1997). Location of gene regulating cold-induced carbohydrate production on chromosome 5A of wheat. *Theoretical and Applied Genetics*, 95, 265-270.
- Hayama, R. & G. Coupland (2003). Shedding light on the circadian clock and photoperiodic control of flowering. *Current Opinion in Plant Biology*, 6, 13-19.
- Hayes, P.M., T. Blake, T.H.H. Chen, S. Tragoonrun, F. Chen, A. Pan & B. Liu (1993). Quantitative trait loci on barley (*Hordeum vulgare* L.) chromosome 7 associated with components of winter hardiness. *Genome*, 36, 66-71.
- Henderson, I. R. & C. Dean (2004). Control of *Arabidopsis* flowering: the chill before the bloom. *Development*, 131: 3829-3838.
- Hopkins, A.A., K.P. Vogel, K.J. Moore, K.D. Johnson & I.T. Carlson (1995). Genotype effects and genotype by environment interactions for traits of elite switchgrass populations. *Crop Science*, 35, 125-132.
- Humphreys, M., L. Turner, L. Skot, L. M. Humphreys, I. King, I. Armstead, I & P. Wilkins (2003). The use of genetic markers in grass breeding. *Czech Journal of Genetics and Plant Breeding*, 39, 112-119. Biodiversity and Genetic Resources as the Bases for Future Breeding. 25th Meeting of the Fodder Crops and Amenity Grasses Section of Eucarpia: Brno, Czech Republic.
- Inoue, M., Z. Gao, M. Hirata, M. Fujimori & H. Cai (2004a). Construction of a high-density linkage map of Italian ryegrass (*Lolium multiflorum* Lam.) using restriction fragment length polymorphism, amplified fragment length polymorphism, and telomeric repeat associated sequence markers. *Genome*, 47, 57-65.
- Inoue, M., Z. Gao & H. Cai (2004b). QTL analysis of lodging resistance and related traits in Italian ryegrass (*Lolium multiflorum* Lam.). *Theoretical and Applied Genetics*, 109, 1579-1585.
- Izawa, T., Y. Takahashi & M. Yano (2003). Comparative biology comes into bloom: genomic and genetic comparison of flowering pathways in rice and *Arabidopsis*. *Current Opinion in Plant Biology*, 6, 113-120.
- Jensen, L.B., J.R. Andersen, U. Frei, Y. Xing, C. Taylor, P.B. Holm & T. Lübstedt (2005). QTL mapping of vernalization response in perennial ryegrass (*Lolium perenne* L.) reveals co-location with an orthologue of wheat *VRN1*. *Theoretical and Applied Genetics*, available on-line.
- Jones, E.S., N.L. Mahoney, M.D. Hayward, I.P. Armstead, J.G. Jones, M.O. Humphreys, I.P. King, T. Kishida, T. Yamada, F. Balfourier, C. Charmet & J.W. Forster (2002a). An enhanced molecular marker-based map of perennial ryegrass (*Lolium perenne* L.) reveals comparative relationships with other Poaceae species. *Genome*, 45, 282-295.
- Jones, E.S., M.D. Dupal, J.L. Dumsday, L.J. Hughes & J.W. Forster (2002b). An SSR-based genetic linkage map for perennial ryegrass (*Lolium perenne* L.). *Theoretical and Applied Genetics*, 105, 577-584.
- Kato, K., H. Miura & S. Sawada (1999). Comparative mapping of the wheat *Vrn-A1* region with the rice *Hd-6* region. *Genome*, 42, 204-209.
- Kato, K., S. Kidou, H. Miura & S. Sawada (2002). Molecular cloning of the wheat *CK2α* gene and detection of its linkage with *Vrn-A1* on chromosome 5A. *Theoretical and Applied Genetics*, 104, 1071-1077.
- Koemel, J.E. Jr., A.C. Guenzi, J.A. Anderson, E.L. Smith (2004) Cold hardiness of wheta near-isogenic lines differing in vernalization alleles. *Theoretical and Applied Genetics*, 109, 839-846.
- Kojima, S., Y. Takahashi, Y. Kobayashi, L. Monna, T. Sasaki, T. Araki & M. Yano (2002). *Hd3a*, a rice ortholog of the *Arabidopsis FT* gene, promotes transition to flowering downstream of Hd1 under short-day condition. *Plant and Cell Physiology*, 43, 1096-1105.
- Laurie, D.A., N. Pratchett, J.H. Bezzant & J.W. Snape (1995). RFLP mapping of five major genes and eight quantitative trait loci controlling flowering time in a winter x spring barley (*Hordeum vulgare* L.) cross. *Genome*, 38, 575-585.
- Laurie, D. A. (1997). Comparative genetics of flowering time. *Plant Molecular Biology*, 35, 167-177.
- Laurie, D.A., S. Griffiths, R.P. Dunford, V. Christodoulou, S.A. Taylor, J. Cockram, J. Beales & A. Turner (2004). Comparative genetics approaches to the identification of flowering time genes in temperate cereals. *Field Crops Research* 90, 87-99.

- Lübberstedt, T., A.E. Melchinger, D. Klein, H. Degenhardt & C. Paul (1997). QTL mapping in testcrosses of European flint lines of maize: II. Comparison of different testers for forage quality traits. *Crop Science*, 37, 1913-1922.
- Lübberstedt, T., A.E. Melchinger, S. Fähr, D. Klein, A. Dally & P. Westhoff (1998). QTL mapping in testcrosses of European flint lines of maize: III. Comparison across populations for forage traits. *Crop Science*, 38, 1278-1289.
- Marten, G.C., G.E. Brink, D.R. Buxton, J.L. Halgerson & J.S. Hornstein (1984). Near infra-red reflectance spectroscopy analysis of forage quality in four legume species. *Crop Science*, 24, 1179-1182.
- Matsumoto, N. & T. Araki (1982). Field observation of snow mold pathogens of grasses under snow cover in Sapporo. *Research Bulletin of Hokkaido National Agricultural Experiment Station*, 135, 1-10.
- Matsumoto, N., T. Sato & T. Araki (1982). Biotype differentiation in the Typhula ishikariensis complex and their allopatry in Hokkaido. *Ann. Phytopath Soc. Japan*, 48, 275-280.
- Méchin, V., O. Argillier, Y. Hebert, E. Guingo, L. Moreau, A. Charcosset & Y. Barrière (2001). Genetic analysis and QTL mapping of cell-wall digestibility and lignification in silage maize. *Crop Science*, 41, 690-697.
- Menke, K.H., L. Raab, A. Salewski, H. Steingass, D. Fritz & W. Schneider (1979) The estimation of the digestibility and metabolisable energy content of ruminant feeding stuffs from the gas production when they are incubated with rumen liquor *in vitro*. *Journal of Agricultural Science (Cambridge)*, 93: 217-222.
- Moseley, G. & D.H. Baker (1991). The efficacy of a high magnesium grass cultivar in controlling hypomagnesaemia in grazing animals. *Grass and Forage Science*, 46, 375-380.
- Mouradov, A., C. F. Cremer & G. Coupland (2002). Control of Flowering Time: Interacting Pathways as a Basis for Diversity. *Plant Cell*, S111-S130 (Suppl.).
- Pan, A., P.M. Hayes, F. Chen, T.H.H. Chen, T. Blake, S. Wright, I. Karsai & Z. Bedö (1994). Genetic analysis of the components of winterhardiness in barley (*Hordeum vulgare* L.). *Theoretical and Applied Genetics*, 89, 900-910.
- Paran, I & D. Zamir (2003). Quantitative traits in plants: beyond the QTL. *TRENDS in Genetics*, 19, 303-306.
- Ralph, J., S. Guillaumie, J.H. Grabber, C. Lapierre & Y. Barrière (2004). Genetic and molecular basis of grass cell-wall biosynthesis and degradability. III. Towards a forage grass ideotype. *Comptes Rendus Biologies*, 327, 467-479.
- Rognli, O.A., V. Alm, C. Busso, C. Fang, A. Larsen, K. Devos, M. Humphreys, K. Vollan & Z. Grieg (2002). Comparative mapping of quantitative trait loci controlling frost and drought tolerance. In: Abstracts of Plant & Animal Genome X Conference, W167. January 12-16, San Diego, CA, USA.
- Saha, M.C., R. Mian, J.C.Zonitzer, K. Chekhovskiy & A. A. Hopkins (2005). An SSR- and AFLP-based genetic linkage map of tall fescue (*Festuca arundinacea* Schreb.). *Theoretical and Applied Genetics*, available. on-line.
- Sanchez, W.K., D.K. Beede & J.A. Cornell (1994). Interactions of sodium potassium, and chloride on lactation, acid-base status, and mineral concentrations. *Journal of Dairy Science*, 77, 1661-1675.
- Sasaki, A., M. Ashikari, M. Ueguchi-Tanaka, H. Itoh, A. Nishimura, D. Swapan, K. Ishiyama, T. Sato, M. Kobayashi, G.S. Khush, H. Kitano & M. Matsuoka (2002). A mutant of gibberellin-synthesis gene in rice. *Nature*, 416: 701-702.
- Smith, K.F. & P.C. Flinn (1991). Monitoring the performance of a broad-based calibration for measuring the nutritive value of two independent populations of pasture using near infra-red reflectance spectroscopy. *Australian Journal of Experimental Agriculture*, 31, 205-210.
- Smith, K.F., K.F.M. Reed & J.Z. Foot (1997). An assessment of the relative importance of specific traits for the genetic improvement of nutritive value in dairy pasture. *Grass and Forage Science*, 52, 167-75.
- Smith, K.F. & G.A. Kearney (2000). The distribution of errors associated with genotype and environment during the prediction of the water-soluble carbohydrate concentration of perennial ryegrass cultivars using near infrared reflectance spectroscopy. *Australian Journal of Agricultural Research*, 51, 481-486.
- Smith, K.F., R.J. Simpson & R.N. Oram (2004). The effects of site and season on the yield and nutritive value of cultivars and half-sib families of perennial ryegrass (*Lolium perenne* L.) *Australian Journal of Experimental Agriculture*, 44, 763-769.
- Snape, J.W., A. Semikhodskii, L. Fish, R.N. Sarma, S.A. Quarrie, G. Galiba & J. Sutka (1997). Mapping frost resistance loci in wheat and comparative mapping with other cereals. *Acta Agron Hungary*, 45, 265-270.
- Storlie, E.W., R.E. Allan & M.K. Walker-Simmons (1998). Effect of the *Vrn1-Fr1* interval on cold hardiness levels in near-isogenic wheta lines. *Crop Science*, 38, 483-488.
- Sung, S. & R. M. Amasino (2004). Vernalization and epigenetics: how plants remember winter. *Current Opinion in Plant Biology*, 7, 4-10.
- Sutka, J. & J.W. Snape (1989). Location of a gene for frost resistance on chromosome 5A of wheat. *Euphytica*, 42, 41-44.
- Sutka, J., G. Galiba, A. Vaguifalvi, B.S. Gill & J.W. Snape (1999). Physical mapping of the *Vrn-A1* and *Fr1* genes on chromosome 5A of wheat using deletion lines. *Theoretical and Applied Genetics*, 99, 199-202.

- Takai, T., Y. Sanada & T. Yamada (2004). Varietal differences of meadow fescue (*Festuca pratensis* Huds.) in resistance to *Typhula* snow mold. *Grassland Science*, 49, 571-576.
- Takahashi, Y., A. Shomura, T. Sasaki & M. Yano (2001). *Hd6*, a rice quantitative trait locus involved in photoperiod sensitivity, encodes the  $\alpha$  subunit of protein kinase CK2. *Proceeding of the National Academy of Sciences of the United States of America*, 98, 7922-7927.
- Thomashow, M.F. (1999). Plant cold acclimation: Freezing tolerance genes and regulatory mechanisms. *Annual Review of Plant Physiology and Plant Molecular Biology*, 50, 571-599.
- Thomashow, M.F., S.J. Gilmour, E.J. Stockinger, K.R. Jaglo-Ottosen & D.G. Zarka (2001). Role of Arabidopsis CBF transcriptional activators in cold acclimation. *Physiologia Plantarum*, 112, 171-175.
- Tilly, J.M.A. & R.A. Terry (1963). A two-stage technique for the *in vitro* digestion of forage crops. *Journal of the British Grassland Society*, 18, 104-111.
- Toth, B., G. Galiba, E. Fehér, J. Sutka & J. W. Snape (2003). Mapping genes affecting flowering time and frost resistance on chromosome 5B of wheat. *Theoretical and Applied Genetics* 107, 509-514.
- Turner, L.B., M.O. Humphreys, A.J. Cairns & C.J. Pollock (2001). Comparison of growth and carbohydrate acculation in seedling of two varieties of *Lolium perenne*. *Journal of Plant Physiology*, 158, 891-897.
- van Soest, P.J. (1963). Use of detergents in the analysis of fibrous feeds. *Journal of the Association of Official Agricultural Chemists*, 46, 825-835.
- Warnke, S., R.E. Barker, S.-C. Sim, G. Jung & J.W. Forster (2003a). Genetic map development and syntenic relationships an annual x perennial ryegrass mapping population. In: Abstract of Plant & Animal Genome XI Conference, W 206. January 11-15, San Diego, CA, USA.
- Warnke, S., R.E. Barker, S.-C. Sim, G. Jung & M.A. R. Mian (2003b). Identification of flowering time QTL's in annual x perennial ryegrass mapping population. In: Abstracts of Molecular Breeding of Forage and Turf, Third International Symposium, p. 122. May 18-22, Dallas, Texas and Ardmore, Oklahoma, USA.
- Warnke, S.E., R.E. Barker, G. Jung, S.-C. Sim, M.A. R. Mian, M.C. Saha, L.A. Brilman, M.P. Dupal & J.W. Forster (2004). Genetic linkage mapping of an annual x perennial ryegrass population. *Theoretical and Applied Genetics*, 109, 294-304.
- Wheeler, J.L. & J.L. Corbett (1989). Criteria for breeding forages of improved nutritive value: results of a Delphi survey. *Grass and Forage Science*, 44, 77-83.
- Wilkins, P.W. & M.O. Humphreys (2003). Progress in breeding perennial forage grasses for temperate agriculture. *Journal of Agricultural Science (Cambridge)*, 140, 129-150.
- Yano, M., Y. Katayose, M. Ashikari, U. Yamanouchi, L. Monna, T. Fuse, T. Baba, K. Yamamoto, Y. Umehara, Y. Nagamura & T. Sasaki (2000). *Hd1*, a major photoperiod sensitivity quantitative trait locus in rice, is closely related to the *Arabidopsis* flowering time gene *CONSTANS*. *Plant Cell*, 12, 2473-2484.
- Yano, M., S. Kojima, Y. Takahashi, H.X. Lin & T. Sasaki (2001). Genetic control of flowering time in rice, a short-day plant. *Plant Physiology*, 127, 1425-1429.
- Yamada, T., E.S. Jones, T. Nomura, H. Hisano, Y. Shimamoto, K.F. Smith, M.D. Hayward & J.W. Forster (2004). QTL analysis of morphological, developmental and winter hardiness-associated traits in perennial ryegrass (*Lolium perenne* L.). *Crop Science*, 44, 925-935.
- Yamamoto, T., H. Lin, T. Sasaki & M. Yano (2000). Identification of heading date quantitative locus *Hd6* characterization of its epistatic interaction with *Hd2* in rice using advanced back-cross progeny. *Genetics*, 154, 885-891.
- Yan, L, A. Loukoianov, G. Tranquilli, M. Helguera, T. Fahima, J. Dubcovsky (2003). Positional cloning of the wheat vernalisation gene *VRN1*. *Proceeding of the National Academy of Sciences of the United States of America*, 100, 6263-6268.
- Yan, L, A. Loukoianov, A. Blechl, G. Tranquilli, W. Ramakrishna, P. SanMiguel, J.L. Bennetzen, V. Echenique, J. Dubcovsky (2004). The wheat *VRN2* gene is a flowering repressor down-regulated by vernalisation. *Science*, 303, 1640-1644.
- Yoshida, M., J. Abe, M. Moriyama, T. Kuwahara (1998). Carbohydrate levels among wheat cultivars varying in freezing tolerance and snow mold resistance during autumn and winter. *Physiologia Plantarum*, 103, 437-444.