




## Role of the BANYULS(BAN) Gene from Arabidopsis Thaliana in Transgenic Alfalfa Expression of Anthocyanins and Proanthocyanidins

S. M. Hesamzadeh Hejazi  
*Research Institute of Forests and Rangelands, IRAN*

Sergio Arcioni  
*Consiglio Nazionale delle Ricerche, Italy*

F. Paolucci  
*Consiglio Nazionale delle Ricerche, Italy*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium5/55>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

## Role of the BANYULS(*BAN*) gene from *Arabidopsis thaliana* in transgenic Alfalfa expression of anthocyanins and proanthocyanidins

S.M. Hesamzadeh Hejazi<sup>1</sup>, S. Arcioni<sup>2</sup> and F. Paolocci<sup>2</sup>

<sup>1</sup>Research institute of forests and rangelands(gene bank),Tehran-IRAN

<sup>2</sup>Istituto di Recherche sul Miglicramento, Genetico della piante Foraggiere, Consiglio Nazionale della Ricerche,Perugia, Italy, Email: smhessamzadeh@rjfr-ac.ir

**Keywords:** *Agrobacterium tumefaciens*, *BAN* gene, condensed tannins, genetic transformation, *Medicago sativa*

**Introduction** Condensed tannins (CTs) are flavonoid oligomers, many of which have beneficial effects on animal (bloat safe) and human health. The *BAN* gene encodes anthocyanidin reductase (ANR), an enzyme proposed to convert anthocyanidins to their corresponding 2,3-cis-flavan-3-ols (Xie *et al.*, 2003). Ectopic expression of *BAN* in Alfalfa transgenic foliage results in accumulation of CTs. Thus, it has been assumed that the *BAN* gene also acts in starter units for the condensation of tannins in Alfalfa.

**Material and methods** Transformation was performed in two genotypes (Rgsy27,P1) of *M. sativa*. The binary vector pBI121.BAN was constructed by removing the *gus* gene and inserting the *BAN* gene into the BamHI and SacI sites of pBI121.1 to put *BAN* expression under control of the CaMV35S promoter and NOS 3' terminator. pBI121.BAN was transformed into *Agrobacterium tumefaciens* EHA105 and LBA4404 by the triparental mating method. To confirm chromosomal insertion of the *BAN* transgene, genomic DNA was extracted from leaves of putative transgenic Rgsy27 and P1(cv.Adriana) plants (Sambrook *et al.*, 1989). The NPTII selectable marker gene in the binary vector was used as a probe after labeling with <sup>32</sup>P-dCTP using the Ready -TO-GO DNA labeling beads (dCTP)Kit. For RT-PCR analysis of *BAN* transcript levels, total RNA isolated from leaf tissue using Nucleo Spin RNA Kit. CTs in the leaves of transgenic plants were visualized by staining tissues in a solution of methanol:6M HCl (1:1) containing 2%(w/v) DMACA for 3 min (Li *et al.*, 1996), then washing 3 times with MilliQ water. For quantitative analysis of CTs in transgenic plants, we used DMACA-HCl and Butanol-HCl assays at A643, A550 and A725.

**Results** We found a simple and efficient method for regeneration-transformation of *Medicago* including the perennial *M. sativa* var.Rgsy27 (2n=4x=32) and *M. sativa* var.Adriana (P1)(2n=4x=32). Development of this method (using 4 new media M1,M2,M3,M4 with low levels of 2,4-D and BAP) produced transgenic plants on all explants within 1.5-2 months which remained at the same ploidy level and maintained fertility.

The presence and structure of the transgene in the plants were analysed by genomic southern blots in the Rgsy27 and P1 genotypes. In each case both an internal fragment of the construct as well as one of the T-DNA borders were probed. The expected size of the internal fragments characteristic for each construct was detected in the plants analysed except for a few transgenic plants where the *BAN* fragment was not present. The number of T-DNA borders detected in these plants, representing the number of T-DNA copies integrated in the plant genome, varied from one to probably three or more. In several of the transformants we found aberrant sized *BAN* bands. This was probably due to the loss of one of the T-DNA BamHI sites, indicating that truncation of some of the T-DNA copies had occurred in these plants. The qualitative expression of the *BAN* gene was studied by RT-PCR and was compared with the expression of LAR and EF-1 genes. The strain LBA4404 was much better than EHA105 for transformation of both genotypes.

The *BAN* gene encodes anthocyanidin reductase (ANR), an enzyme proposed to convert anthocyanidins to their corresponding 2,3-cis-flavan-3ols. Ectopic expression of *BAN* in Alfalfa transgenic foliage results in accumulation of CTs. The percent of tannin in the dry matter of control and transgenic plants were: for P1(control) = 0.418, Rgsy27(control) = 0.259, P1+*BAN* = 0.600 and for Rgsy27+*BAN* = 0.474.

**Conclusions** Preliminary results of our work indicated that in forage legumes BANYULS(*BAN*) could not autonomously induce tannin. However, when tissues are committed for tannin accumulation, the *BAN* gene can strongly increase CT levels.

### References

- Xie, D.Y., S.B., Sharma,N.L.,Paiva, D.Ferreria, R. A., Dixon (2003). Role of anthocyanidin reductase encoded by *BANYULS* in plant flaonoid biosynthesis. SCIENCE, 299: 396-399.
- Sambrook, J., Ef., Fritsch, T., Maniatis,(1989). Molecular cloning : alaboratory manual- cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, ed. 2<sup>nd</sup>.
- Li, Y.-G. Tanner, G.J., Larkin, P.J., (1996). The DMACA-HCL protocol and the threshold proantho cyanidin content for bloat safety in forage legumes. J. Sci. Food Agric. 70: 89-101.