



University of Kentucky
UKnowledge

International Grassland Congress Proceedings


XX International Grassland Congress

Belowground Meristem Populations as Regulators of Grassland Dynamics

Harmony J. Dalgleish
Kansas State University

D. C. Hartnett
Kansas State University

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium3/86>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Belowground meristem populations as regulators of grassland dynamics

H.J. Dalgleish and D.C. Hartnett

Division of Biology, Kansas State University, Manhattan, KS 66506, USA. Email: hjdal@ksu.edu

Keywords: bud, meristem limitation

Introduction Studies of plant populations are critical for linking organism to ecosystem-level phenomena and for understanding mechanisms driving responses to global change. In perennial grasslands, the below-ground population of meristems (the bud bank) plays a fundamental role in local plant population recruitment, persistence and dynamics. We explore two aspects of the bud bank in North American grasslands. It has been hypothesized that low variability in arid biomes is explained by meristem limitation, which constrains responses to pulses of high resource availability. Our research tests this hypothesis by comparing bud-bank populations across six sites in the United States that vary 3-fold in precipitation and 4.5-fold in productivity. In addition, we are examining the effects of management practices, such as fire and grazing, on bud-bank populations using replicated long-term treatments at Konza Prairie LTER site located in north-central Kansas.

Materials and methods At each of six study sites, stem densities, above-ground biomass and bud densities of grasses and forbs were measured within a 25 cm x 25 cm x 15 cm depth sample. We also sampled replicate watersheds in a factorial design at Konza Prairie that have been subjected to annual or four-year burning cycles and are either grazed by Bison (*Bos bison*) or are ungrazed.

Results Data from November 2003 and March 2004 support the hypothesis that bud densities increase from east to west along a precipitation/productivity gradient (Figure 1). Bud densities increased significantly at all sites between sampling dates ($P=0.001$). Letters indicate differences among sites in March at $P<0.05$ (LSD). Grass bud density increased from November 2003 to March 2004, but there were no significant treatment effects (grazing, $P=0.215$ and fire, $P=0.339$). There was a trend of higher grass bud density in the annually burned, ungrazed treatment compared to the four-year burning cycle. However, this trend disappears with the introduction of grazers. Forb bud density did not change from November 2003 to March 2004 and there was a significant treatment interaction ($P=0.001$, Figure 2). Annually burned and grazed tallgrass prairie had the highest bud density, but this difference disappeared in the absence of grazers.




Figure 1 Bud density along a precipitation gradient in central North America




Figure 2 Effects of fire and grazing on forb bud populations

Conclusions Below-ground meristem (bud) density increased with increasing precipitation and productivity. Overall lower bud density in arid grasslands supported the hypothesis that meristem limitation constrains production variability in these grasslands. Grass and forb bud populations had different responses to fire and grazing in tallgrass prairie, with forb bud density increasing in grazed areas while grass bud density remained unchanged. Therefore, demographic mechanisms may be important for forbs in these grasslands.