



Microarray-Based Transcriptome Analysis of the Interaction Between Perennial Ryegrass (*Lolium Perenne*) and the Fungal Endophyte *Neotyphodium Lolii*

S. Felitti

La Trobe University, Australia

P. Tian

Molecular Plant Breeding Cooperative Research Centre, Australia

T. Webster

Victorian Microarray Technology Consortium, Australia

D. Edwards

La Trobe University, Australia

G. C. Spangenberg

Victorian Bioinformatics Consortium, Australia

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium5/63>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

Microarray-based transcriptome analysis of the interaction between perennial ryegrass (*Lolium perenne*) and the fungal endophyte *Neotyphodium lolii*

S.A. Felitti^{1,2}, P. Tian^{1,2}, T. Webster^{1,3}, D. Edwards^{1,2,4} and G.C. Spangenberg^{1,2,3,4}

¹Primary Industries Research Victoria, Plant Biotechnology Centre, La Trobe University, Bundoora, Victoria 3086, Australia ²Molecular Plant Breeding Cooperative Research Centre, Australia ³Victorian Microarray Technology Consortium, Australia ⁴Victorian Bioinformatics Consortium, Australia
Email: silvina.felitti@dpi.vic.gov.au

Keywords: fungal endophytes, perennial ryegrass, expressed sequence tags (ESTs), microarrays, transcriptome

Introduction *Neotyphodium lolii*, *Neotyphodium coenophialum* and *Epichloë festucae* are common symbiotic fungal endophytes of the temperate pasture grasses perennial ryegrass (*Lolium perenne*), tall fescue (*Festuca arundinacea*) and red fescue (*Festuca rubra*), respectively. A genomic resource of 13,964 expressed sequence tags (ESTs), representing 7,585 unique endophyte genes, has been established for *Neotyphodium* and *Epichloë* fungal endophytes.

Materials and methods The endophyte genomic resource established has enabled the design and fabrication of two endophyte-specific cDNA microarrays (NchipTM microarray and EndoChipTM microarray). The NchipTM and EndoChipTM microarrays have been applied to comparative transcriptome analyses of different asexual (*N. coenophialum* and *N. lolii*) and sexual (*E. festucae*) endophyte taxa under various saprophytic growth conditions, leading to the identification of differentially expressed genes.

Results and conclusions The NchipTM and EndoChipTM microarrays permit the interrogation of 3,806 *Neotyphodium* genes (NchipTM microarray), and 4,195 *Neotyphodium* and 920 *Epichloë* genes (EndoChipTM microarray), respectively. They represent tools for high-throughput transcriptome analysis, including genome-specific gene expression studies, profiling of novel endophyte genes, and investigation of the host grass-fungal symbiont interaction. Microarray-based transcriptome analysis was also undertaken in transgenic *Neotyphodium* endophyte expressing a chimeric *gusA* reporter gene [strain FM13 (Lp1/pNOM-101 and pAN7-1) (Murray *et al.*, 1992) kindly provided by B. Scott, Institute of Molecular BioSciences, Massey University, Palmerston North, New Zealand] compared to untransformed control, confirming the power of these tools for applications in characterising genetically modified endophytes. A recent extension of this proof-of-concept research has included the transcriptome analysis of transgene-expressing endophytes to assess gene expression responses to specific genetic modifications (e.g. gain-of-function, knock-outs and knock-downs). Within the context of a comprehensive spatial and temporal systems biology approach, a transcriptomics study of the mutualistic interaction between perennial ryegrass (*L. perenne* L.) and its fungal endophyte (*N. lolii*) was undertaken. In combination with a 15K ryegrass unigenic microarray, the EndoChipTM microarray was applied to the detailed analysis of in planta gene expression in different ryegrass organs using endophyte-infected and endophyte-free ryegrass plants in an isogenic host genetic background. Data derived from endophyte microarray analysis has been validated using both northern hybridisation and RT-PCR analyses.




Figure 1 *Neotyphodium in planta* gene expression microarray-based transcriptome analysis. A) Hierarchical clustering of mean signal values (Euclidean distance) from infected and uninfected ryegrass organs on the Endophyte unigenic microarray. B) Endophyte-infected stem prevalent gene expression

Reference

Murray, F.R., Latch, G.C.M. and Scott, D.B. (1992). Surrogate transformation of perennial ryegrass, *Lolium perenne*, using genetically modified *Acremonium* endophyte. *Mol. Gen. Genet.*, 233, 1-9.